



RESEARCH

Open Access

Rational/antiemotional behaviors in interpersonal relationships and the functional prognosis of patients with rheumatoid arthritis: a Japanese multicenter, longitudinal study

Jun Nagano^{1*}, Takako Morita^{2,3}, Koji Taneichi⁴, Shohei Nagaoka⁵, Sadanobu Katsume⁶, Tomiaki Asai⁷, Masao Yukioka⁸, Kiyoshi Takasugi⁹, Masakazu Kondo¹⁰ and Yasuro Nishibayashi¹¹

Abstract

Background: The repression of negative emotions is a personality factor that received considerable attention in the 1950–60s as being relevant to the onset and course of rheumatoid arthritis (RA). Despite subsequent, repeated criticisms of the cross-sectional nature of the earlier studies, even to date few prospective studies have been reported on this issue. This multicenter study prospectively examined if “rational and antiemotional” behavior (antiemotionality), characterized by an extreme tendency to suppress emotional behaviors and to rationalize negative experiences in conflicting interpersonal situations, is associated with the functional prognosis of patients with RA.

Methods: 532 patients with RA who regularly visited one of eight hospitals/clinics in Japan in 2000 were recruited for study. All completed a self-administered baseline questionnaire about lifestyle and psychosocial factors including antiemotionality. Two years after, 460 (mean age, 56.1 years; 54 men and 406 women) of 471 patients who continued to visit the clinics agreed to take the follow-up questionnaire. The functional status of the patients was evaluated by rheumatologists based on the ACR classification system.

Results: A multiple logistic regression model that included baseline demographic, disease activity/severity-related, therapeutic, and socioeconomic factors as covariates found a tendency toward higher antiemotionality to be related to poorer functional status at follow-up. This relationship was not explained by lifestyle factors.

Conclusions: Antiemotionality may be a prognostic factor for the functional status of patients with RA. This finding sheds light on a seemingly forgotten issue in the care of patients with RA.

Keywords: Rheumatoid arthritis, Psychological stress, Personality, Emotions, Function, Prospective studies

Background

Rheumatoid arthritis (RA) is an autoimmune disease characterized by chronic systemic inflammation that mainly affects joints, which causes a loss of physical functioning. It can lead to severe systemic dysfunction and a premature death [1]. Physicians and patients agree that psychosocial stress (stress) in daily life can affect the inflammatory and functional status of patients with RA, and

some studies have supported this notion [2–4]. Recent advancement in psychoneuroimmunology has helped clarify the mechanisms by which stress affects the disease through interaction between the endocrine, nervous, and immune systems [5–7].

Stress, individual internal experience, can be affected not only by external stressors, but also by interaction between stressors and individual emotional, cognitive, and behavioral responses to them [4]. When a person responds to stressors with a pattern that is repeated on a frequent basis, such a response pattern can be understood within the context of personality. It is difficult for

* Correspondence: jun@artsci.kyushu-u.ac.jp

¹Faculty of Arts and Science, Kyushu University, 6-1 Kasuga Park, Kasuga, Fukuoka 816-8580, Japan

Full list of author information is available at the end of the article

an individual to control external stressors, but it is possible to alter the properties of personality and response style to stressors, and in turn to modify the effects of stressors to the RA disease status. For centuries, clinicians have been impressed by the role of psychosocial factors on RA, and in the 1950s and 1960s, RA came to be viewed as one of the classical psychosomatic diseases [8,9]. Particular personality factors that possibly relate to the onset and course of RA have received considerable attention [10], and a personality characterized by an inability to express aggressive feelings, e.g., “contained hostility” [11] was the one on which many researchers agreed [12]. The nature of these earlier studies included a cross-sectional design, and they were repeatedly criticized, with reviewers concurring on the necessity for prospective studies that allow better causal interpretations [5,9,13,14]. A “neurotic” personality (anxious and depressed), another hypothesis drawn from some earlier cross-sectional studies that used the Minnesota Multiphasic Personality Inventory, was later denied by longitudinal studies [15,16] and re-interpreted as a result rather than a cause of this chronic disease [17]. To the authors’ knowledge, however, even to date few prospective studies have reported on the classical issue of the repression of negative, especially aggressive, emotions as a personality factor associated with the onset/course of RA.

Grossarth-Maticek and colleagues hypothesized that the “rationality and antiemotionality” or “type 5” behavior is a risk factor for the onset of chronic diseases including cancer and RA [18-20]. The rational-antiemotional behavior (antiemotionality) is characterized by an extreme tendency to suppress emotional behaviors and to rationalize negative experiences in conflicting interpersonal situations. Thus, this behavioral pattern largely fits the concept of the repression of negative emotions, whereas it clearly differs from the concept of neurotic personality. It was found, in a cohort study begun in the 1970s, that a high proportion of persons with a highly typical antiemotionality later developed RA [20]. Moreover, it was shown that “Autonomy Training”, a psychotherapeutic intervention method, could alter such a tendency in the direction of healthier behaviors, leading to better physical outcomes [21-23]. A recent study showed that antiemotionality may also be a prognostic factor for lung cancer patients [24]. The present multicenter cohort study examined if antiemotionality is associated with a poorer prognosis for the functional status of patients with RA.

Methods

Subjects

The baseline data for the present study was taken from a multi-center cohort survey of Japanese patients with RA carried out in 2000. It was part of a study supported by the Ministry of Health and Welfare, Japan entitled

“Assessment and improvement of the system for interdisciplinary medical services for RA (AISIMS)” [25]. Rheumatologists, one working at each of the 12 hospitals/clinics that participated in the program, invited patients with RA who regularly visited their hospitals/clinics to cooperate with the AISIMS cohort study. Patient eligibility included age between 20 and 79 years, the ability to answer a self-administered questionnaire without assistance, and functional status of class 3 or better by the criteria of the American College of Rheumatology (ACR) (see below) [26]. The baseline survey consisted of a self-administered questionnaire for the patients and a clinical data sheet completed by their rheumatologist. The questionnaire inquired about a variety of factors including activity of daily living, quality of life, and lifestyle (smoking, alcohol-drinking, physical exercise, sleep, diet, etc.) and about psychosocial factors (major life events, stress/personality, etc.). The items for the clinical data sheet included factors relevant to the disease status of RA, such as progression of arthritis, functional status, extra-articular complications, and factors associated with medical treatment.

In 2002, two years after the baseline survey, eight of the 12 hospitals/clinics participated in the follow-up survey, with the 23 rheumatologists who were working at these eight hospitals/clinics completing the same clinical data sheet as that used for the baseline survey and handing each patient participant a follow-up questionnaire with items selected from the baseline questionnaire. The follow-up survey was completed before 2003, when the first biologic agent (infliximab) became available in Japan for the treatment of patients with RA [27], because an expected high effectiveness of the agents [1] could overshadow the associations of lifestyle and psychosocial factors with the prognosis of patients with RA and lower the statistical power to detect them. The patients who agreed to cooperate with the follow-up survey sent their completed questionnaires to a central office located at the Department of Preventive Medicine, Kyushu University Graduate School of Medical Sciences. The rheumatologists who evaluated the patients’ clinical status were blinded from the patients’ answers for both the baseline and follow-up questionnaires.

Measurements

The rheumatologists assessed their patients’ functional status based on the criteria for classification of functional status in RA defined by ACR (ACR class), which classifies patients with RA into one of four classes as follows: Class I: completely able to perform usual activities of daily living (self-care, vocational, and avocational); Class II: able to perform usual self-care and vocational activities, but limited in avocational activities; Class III: able to perform usual self-care activities, but limited in

vocational and avocational activities; and Class IV: limited in ability to perform usual self-care, vocational, and avocational activities [26]. They also assessed the status of joint damage based on the classification by Steinbrocker et al. (joint stage: Stage I, Early; Stage II, Moderate; Stage III, Severe; and Stage IV, Terminal) [28]; specified afflicted joints defined as those with either tenderness, swelling, or deformity; and listed extra-articular complications using the following options: cervical myelopathy, cardiac/ pericardial manifestations, pulmonary/pleural manifestations, ocular manifestations, peripheral nervous manifestations, hematological manifestations, and others.

Personality factors including antiemotionality were assessed using the "Stress Inventory (SI)" [29,30]. SI is a self-administered questionnaire consisting of 45 items that was developed to assess response styles to stressors principally related to an interpersonal relationship or to chronic stress posed by the response style, that are relevant to chronic diseases [31-33]. Of the 12 scales constituting the SI, the "rationalizing conflicts/frustrations (RCF)" scale measures an extreme tendency to rationalize one's interpersonal situations accompanied by conflicts or frustrations, and was developed to represent antiemotionality. This scale consists of five questions, such as "do you under all circumstances try to control your reasoning and avoid, as much as possible, being emotional?" (Cronbach $\alpha = 0.78$). The answers receive a 1 to 6 rating, where 1 and 6 respectively correspond to "yes" and "no" or to "almost always" and "rarely", and the scores are averaged for the scale score, thus a higher score represents a higher tendency.

Analysis

The ACR class at follow-up was dichotomized (poorer function, Class 3 or 4 vs. better function, Class 1 or 2) and used as the outcome variable representing the functional prognosis of the patients with RA. The association between antiemotionality and the functional prognosis was examined using a multiple logistic regression model that included the dichotomized ACR class at follow-up as a dependent variable, the RCF score (1–6) as an independent variable, and baseline ACR class (Class I, II, III, IV = 1, 2, 3, 4, respectively) as a covariate. The association was then examined after controlling for known or potential confounding factors including sex; age; factors relevant to disease progression or activity: joint stage (Stage I, II, III, IV = 1, 2, 3, 4, respectively), afflicted joints count (1–45), number of extra-articular complications (0–5), C-reactive protein with log-transformation; medical treatments: methotrexate, corticosteroids, other DMARDs (yes = 1, no = 0); and socioeconomic status: education level (junior high school = 1, high school = 2, junior college = 3, college = 4). Finally, potential intervening effects of lifestyle factors were examined by adding the

following variables into this multivariate model: smoking (current smoker = 1, non-/ex-smoker = 0), alcohol drinking (current drinker = 1, non-/ex-drinker = 2), exercise habit (none = 0, 1–3 times/w = 1, 4+ times/w = 2), sleep habit (7–8 hrs/night = 1, others = 0), green-yellow vegetable intake (not eat = 1, once/w = 2, 2–4/w = 3, 5–6/w = 4, every day = 5) and fruit intake (same as green-yellow vegetables).

Results

At the time of the follow-up survey, seven of the 532 patients had died, 13 were visiting another hospital/clinic, and 33 had quit visiting for unknown reasons. The doctors in charge handed the follow-up questionnaire to the 479 patients who continued visiting, and 460 of them sent a completed questionnaire to the central office. Table 1 summarizes the demographics and baseline characteristics

Table 1 Demographic and baseline clinical characteristics of patients with rheumatoid arthritis (N = 460)

Characteristics	N	(%)
Age, yrs, mean (SD)	56.1	(9.6)
Female	406	(88.3)
Education completed		
Junior high school	114	(24.8)
High school	251	(54.6)
Junior college or higher	95	(20.6)
Disease duration, yrs, mean (SD)	11.4	(9.5)
ACR class		
1	76	(16.5)
2	322	(70.0)
3	62	(13.5)
Joint stage*		
1	55	(12.0)
2	98	(21.3)
3	113	(24.6)
4	194	(42.2)
Afflicted joint* count (0–49), mean (SD)	10.0	(8.8)
No. of extra-articular complications*		
0	351	(76.3)
1	83	(18.0)
2+	26	(5.7)
C-reactive protein, mg/dl, mean (SD)	1.63	(2.11)
Medical treatments†		
Methotrexate use	194	(42.2)
Corticosteroids use	205	(44.6)
Other DMARDs use	246	(53.5)

SD: standard deviation, ACR: American College of Rheumatology, DMARDs: disease modifying anti-rheumatic drugs. *See text for explanation. †Based on a yes/no choice. Values are N (%) unless otherwise stated.

of the subjects available for analysis. The characteristics of the patients who were lost to follow-up (visiting another hospital/clinic or having quit visiting for unknown reasons, N = 46) were not statistically different from those of the subjects included in the analysis (see Additional file 1: Table A1). Table 2 shows the baseline characteristics of the subjects according to the degree of antiemotionality. Greater antiemotionality was associated with female sex, older age, and greater number of afflicted joints.

The ACR class distribution at the follow-up was as follows: Class 1, 65 (14.1%); Class 2, 327 (71.0%); Class 3, 59 (12.8%); and Class 4, 9 (2.0%); thus, global functional status had improved for 56 (12.2%), was unchanged for 328 (71.3%), and had deteriorated for 76 (16.5%) over the two-year study period. Figure 1 shows the change in ACR class over the period according to the RCF score.

Table 3 shows the results of the multiple logistic regression analysis done to examine the association between antiemotionality and the functional prognosis. Greater antiemotionality (a higher RCF score) was associated with a poorer ACR class two-years after the baseline survey in the model that only included baseline ACR class as a covariate (Model 1). When multiple factors relevant to demographics, disease activity/progression, treatments, and socioeconomic status were additionally adjusted, the association between antiemotionality and the

Table 2 Association at baseline between antiemotionality and the demographic and clinical characteristics of patients with rheumatoid arthritis (N = 460)

Characteristics	Antiemotionality (RCF score*)			P†
	Low N = 145	Moderate N = 157	High N = 158	
Female	82.8	86.6	95.0	.001
Age, yrs, mean (SD)	54.9 (10.0)	56.0 (10.3)	57.4 (8.5)	.033
Education, college or higher	22.8	20.4	19.0	.85
Duration, yrs, mean (SD)	11.6 (9.8)	9.9 (8.5)	12.7 (10.0)	.41
ACR class > = 3	13.8	11.5	15.2	.75
Joint stage‡ > = 3	65.5	66.2	68.4	.45
Afflicted joints‡ count, mean (SD)	9.9 (9.4)	9.0 (7.8)	11.1 (9.0)	.028
No. extra-articular complications‡ > = 1	23.5	23.0	24.7	.49
CRP, mg/dl, mean (SD)	1.49 (1.87)	1.76 (2.50)	1.63 (1.90)	.12
Methotrexate use	38.6	43.4	44.3	.48
Corticosteroids use	44.8	42.0	46.8	.86
Other DMARDs use	54.5	54.1	51.9	.95

RCF: rationalizing conflicts/frustrations. ACR: American College of Rheumatology, CRP: C-reactive protein, DMARDs: disease modifying anti-rheumatic drugs. *The score of the rationalizing conflicts/frustrations scale of the Stress Inventory; the degrees of antiemotionality, "low", "moderate", and "high", were based on the tertiles of the RCF score, 3.8 and 4.6. †Based on Spearman's rank correlation. ‡See text for explanation. Values are % unless otherwise stated.




Figure 1 Changes in the functional status of patients with rheumatoid arthritis according to the degree of antiemotionality.

RCF score: the score of the rationalizing conflicts/frustrations scale of the Stress Inventory; "low", "moderate" and "high" represent the RCF score of < 3.8, 3.8 to < 4.6, and 4.6+, respectively. ACR: American College of Rheumatology. BL: at the time of the baseline survey, 2Y: two years after the baseline survey.

functional prognosis did not materially change (Model 2). A one point increment in the baseline RCF score was associated with a nearly 50% higher chance of having a poorer functional status rather than a better status at the follow-up. The possible intervening effect of lifestyle factors, including smoking, alcohol consumption, exercise, sleep hours, green/yellow vegetable intake, and fruit intake, was then examined. Additional adjustment for these factors, however, did not appreciably change the association between antiemotionality and the functional prognosis (OR = 1.60, 95% CI = 1.16-2.22, P = 0.005).

Discussion

This Japanese, multicenter, cohort study supports the hypothesis that a strong tendency toward antiemotionality, previously reported as a risk factor for the onset of RA [20], is also associated with the functional prognosis of patients with RA. This is, to the authors' knowledge, the first prospective study that supports the classical notion that a personality characterized by emotional suppression

Table 3 Multiple logistic regression analysis of the association between antiemotionality at baseline and the functional prognosis (ACR class at follow-up) of patients with rheumatoid arthritis

Baseline variables	Model 1			Model 2		
	OR	(95% CI)	P	OR	(95% CI)	P
Antiemotionality (RCF score*)	1.40	(1.05-1.87)	.023	1.47	(1.07-2.00)	.016
ACR class	10.52	(5.87-18.9)	<.001	6.13	(3.17-11.8)	<.001
Female sex				1.92	(0.53-6.98)	.32
Age				1.00	(0.96-1.04)	.99
Joint stage*				1.20	(0.80-1.78)	.38
Afflicted joints* count				1.03	(1.00-1.07)	.06
No. extra-articular complications*				1.40	(0.96-2.02)	.08
Log. C-reactive protein				1.30	(0.99-1.71)	.06
Methotrexate use				1.39	(0.67-2.88)	.38
Corticosteroids use				0.43	(0.20-0.94)	.034
Other DMARDs use				1.12	(0.54-2.32)	.76
Education level				0.80	(0.52-1.23)	.30

ACR: American College of Rheumatology. OR: odds ratio of a poorer function (Class 3 or 4) vs. a better function (Class 1 or 2) at follow-up. CI: confidence interval.
RCF: rationalizing conflicts/frustrations. *See text for explanation.

has a particular connection with the course of RA. The RCF score was differently correlated with specific aspects of negative emotionality in a study of its psychometrical validity [30]. Its positive and negative correlations respectively with “anger-in” and “anger-out”, concepts by Spielberger [34], support earlier cross-sectional findings of the inability to express aggressive feelings associated with a poorer course of this disease [10-12]. That the RCF score was not correlated with trait-anxiety or depression is consistent with the null association between “neurotic” personality and the course of RA in previous longitudinal studies [15,16].

Although this study was of a prospective design, the observed association will need to be further scrutinized for possible confounding effects. Factors representing disease activity, chronic symptoms and functional disability, and medical treatments that potentially affect the central nervous system may have affected the baseline RCF scores. Socioeconomic status, known as a possible prognostic factor for patients with RA [35], may also be correlated with RCF. However, the prospective association of antiemotionality with functional disability at follow-up was independent of all of these baseline factors.

Except for smoking’s exacerbating effect, no lifestyle factor has been established as affecting the disease course of RA [1,36]. However, lifestyle factors should be considered major pathways that potentially link psychosocial variables and physical outcomes. Various factors considered to be generally favorable or unfavorable for health, such as smoking, alcohol-drinking, physical exercise, sleep, and diet were examined for their potential intervening effects, but they did not explain the antiemotionality-prognosis association.

Another pathway to be considered is the psycho-physiological system. The hypothalamus is the center of the autonomic nervous system (ANS) and of the hypothalamus-pituitary-adrenal (HPA) axis, and it has been progressively elucidated that the ANS, HPA axis, and immune system are closely connected and that they interact with each other [5-7]. When the cerebral cortex, the center of reason, chronically and overly interferes with the limbic system, which is responsible for emotional responses, this may affect the function of the hypothalamus, which is closely connected to the limbic system, and in turn the function of the immune system. Such behaviors by which a person attempts to maintain imperturbability or to understand another person when confronting a frustrating interpersonal situation will be, when appropriately employed, harmless or rather preferable. However, when such behaviors are extremely employed, i.e., negative emotions are always rationalized and their expression is blocked, the emotions that lost outlet may damage the functioning of biological homeostasis. Ishii et al. found that patients with RA who were easily moved to tears as a response to psychological stress showed a better response to treatment and a better general prognosis than those who did not show such emotional responses [37].

This study has several limitations. First, no biomarkers, such as neuroendocrinological or immunological markers were measured that can be related to the above discussion of the psycho-physiological pathway. Second, although the outcome measure used, ACR class, is convenient for assessment and represents the overall functional status of patients with RA, its sensitivity to detect temporal changes

of the status is limited [38]. The assessment depends completely on the subjective evaluation of the rheumatologist, while other commonly used scoring systems, such as the ACR core set and the Disease Activity Score, integrate objective markers and subjective evaluations by both doctors and patients [39,40]. Third, the present data only included a small number of male patients, and thus could not address possible sex-related difference in the antiemotionality-prognosis association. Nevertheless, this study adds new findings to the currently insufficient prospective data concerning specific personalities and relevant properties that contribute to the progression of RA. The present results were based on a relatively large sample that was collected at multiple hospitals across Japan and can be generalized to Japanese patients with RA.

Conclusions

Antiemotionality, a personality trait characterized by strong suppression of becoming emotional and by overly appealing to reason in stressful interpersonal situations, is associated with a poorer functional prognosis for patients with RA. This finding sheds light on a seemingly forgotten issue in the care of patients with RA.

Additional file

Additional file 1: Table A1. Demographic and baseline clinical characteristics of patients with rheumatoid arthritis who were lost to follow-up (N = 46).

Abbreviations

ACR: American College of Rheumatology; AISIMS: Assessment and improvement of the system for interdisciplinary medical services for rheumatoid arthritis; ANS: Autonomic nervous system; Antiemotionality: Rational and antiemotional behavior; DMARD: Disease-modifying antirheumatic drug; HPA: Hypothalamus-pituitary-adrenal; RA: Rheumatoid arthritis; RCF: Rationalizing conflicts/frustrations; SI: The Stress Inventory.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

YN, TM, and JN conceived of the study. All authors designed the study and collected the data. JN and TM analyzed the data. JN prepared the draft of the manuscript, and MY and YN helped draft the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The hospitals/clinics that participated in the present cohort study were the Japanese Red Cross Kitami Hospital (Hokkaido prefecture), Yokohama Minami Kyousai Hospital (Kanagawa), Hot Spring of Rehabilitation Nakaizu Hospital (Shizuoka), National Nagoya Hospital (Aichi), Yukioka Hospital (Osaka), Miki Sanyo Hospital (Hyogo), Dohgo Spa Hospital (Ehime), and Kondo Rheumatism and Orthopedics Clinic (Fukuoka). This work was supported by a Health Sciences Research Grant from the Ministry of Health and Welfare of Japan and a Grant-in-Aid for Scientific Research (21590765) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Author details

¹Faculty of Arts and Science, Kyushu University, 6-1 Kasuga Park, Kasuga, Fukuoka 816-8580, Japan. ²Department of Preventive Medicine, Kyushu University Graduate School of Medical Sciences, Fukuoka, Japan. ³Hiroshima

University Graduate School of Biomedical & Health Sciences, Hiroshima, Japan. ⁴Taneichi Rheumatism Clinic, Sapporo, Japan. ⁵Yokohama Minami Kyousai Hospital, Yokohama, Japan. ⁶Hot Spring of Rehabilitation Nakaizu Hospital, Izu, Japan. ⁷Asai Rheumatism and Orthopedics Clinic, Nagoya, Japan. ⁸Yukioka Hospital, Osaka, Japan. ⁹Dohgo Spa Hospital, Matsuyama, Japan. ¹⁰Kondo Rheumatism and Orthopedics Clinic, Fukuoka, Japan. ¹¹Gratia Hospital, Minoh, Japan.

Received: 13 December 2013 Accepted: 20 February 2014

Published: 24 February 2014

References

1. Scott DL, Wolfe F, Huizinga TW: *Rheumatoid arthritis*. *Lancet* 2010, **376**(9746):1094–1108.
2. Scharloo M, Kaptein AA, Weinman JA, Hazes JMW, Breedveld FC, Rooijmans HGM: Predicting functional status in patients with rheumatoid arthritis. *J Rheumatol* 1999, **26**(8):1686–1693.
3. Evers AW, Kraaimaat FW, Geenen R, Jacobs JW, Bijlsma JW: Stress-vulnerability factors as long-term predictors of disease activity in early rheumatoid arthritis. *J Psychosom Res* 2003, **55**(4):293–302.
4. Stojanovich L, Marisavljevic D: Stress as a trigger of autoimmune disease. *Autoimmun Rev* 2008, **7**(3):209–213.
5. Jessop DS, Richards LJ, Harbuz MS: Effects of stress on inflammatory autoimmune disease: destructive or protective? *Stress* 2004, **7**(4):261–266.
6. Cutolo M, Straub RH: Stress as a risk factor in the pathogenesis of rheumatoid arthritis. *Neuroimmunomodulation* 2006, **13**(5–6):277–282.
7. de Brouwer SJ, Kraaimaat FW, Sweep FC, Creemers MC, Radstake TR, van Laarhoven Al, van Riel PL, Evers AW: Experimental stress in inflammatory rheumatic diseases: a review of psychophysiological stress responses. *Arthritis Res Ther* 2010, **12**(3):R89.
8. Alexander F: *Psychosomatic Medicine. Its Principles and Applications*. New York: W. W. Norton & Co. Press; 1950.
9. Rogers M, Brooks E: Psychosocial influences, immune function, and the progression of autoimmune disease. In *Psychoneuroimmunology. Volume 2*. 3rd edition. Edited by Ader R, Felten D, Cohen N. San Diego, San Francisco, New York, Boston, London, Sydney, Tokyo: Academic press; 2001:399–419.
10. King SH: Psychosocial factors associated with rheumatoid arthritis; an evaluation of the literature. *J Chronic Dis* 1955, **2**(3):287–302.
11. Cobb S: Contained hostility in rheumatoid arthritis. *Arthritis Rheum* 1959, **2**:419–426.
12. Moos RH: Personality factors associated with rheumatoid arthritis: a review. *J Chronic Dis* 1964, **17**:41–55.
13. Ward DJ: Rheumatoid arthritis and personality: a controlled study. *Br Med J* 1971, **2**(577):297–299.
14. Creed F: Psychological disorders in rheumatoid arthritis: a growing consensus? *Ann Rheum Dis* 1990, **49**(10):808–812.
15. Robinson H, Kirk RF Jr, Frye RF, Robertson JT: A psychological study of patients with rheumatoid arthritis and other painful diseases. *J Psychosom Res* 1972, **16**(1):53–56.
16. Crown S, Crown JM, Fleming A: Aspects of the psychology and epidemiology of rheumatoid disease. *Psychol Med* 1975, **5**(3):291–299.
17. Pincus T, Callahan LF, Bradley LA, Vaughn WK, Wolfe F: Elevated MMPI scores for hypochondriasis, depression, and hysteria in patients with rheumatoid arthritis reflect disease rather than psychological status. *Arthritis Rheum* 1986, **29**(12):1456–1466.
18. Grossarth-Maticek R: Synergetic effects of cigarette smoking, systolic blood pressure, and psychosocial risk factors for lung cancer, cardiac infarct and apoplexy cerebri. *Psychother Psychosom* 1980, **34**(4):267–272.
19. Grossarth-Maticek R, Kanazir DT, Schmidt P, Vetter H: Psychosocial and organic variables as predictors of lung cancer, cardiac infarct and apoplexy: some differential predictors. *Pers Individ Dif* 1985, **6**:313–321.
20. Grossarth-Maticek R, Eysenck HJ: Personality, stress and disease: description and validation of a new inventory. *Psychol Rep* 1990, **66**(2):355–373.
21. Grossarth-Maticek R, Eysenck HJ: Creative novation behaviour therapy as a prophylactic treatment for cancer and coronary heart disease: part I-description of treatment. *Behav Res Ther* 1991, **29**(1):1–16.
22. Eysenck HJ, Grossarth-Maticek R: Creative novation behaviour therapy as a prophylactic treatment for cancer and coronary heart disease: part II-effects of treatment. *Behav Res Ther* 1991, **29**(1):17–31.
23. Grossarth-Maticek R: *Autonomietraining: Gesundheit und Problemlösung durch Anregung der Selbstregulation*. New York, Berlin: de Gruyter; 2000.

24. Nagano J, Ichinose Y, Asoh H, Ikeda J, Ohshima A, Sudo N, Kubo C: A prospective Japanese study of the association between personality and the progression of lung cancer. *Intern Med* 2006, **45**(2):57–63.
25. Nishibayashi Y: Assessment and improvement of the system for interdisciplinary medical service for RA. In *Report for Research on Immunological and Allergic disease, Health Sciences Research Grant 2001, Volume 1, edn*. Edited by Kudo H. Tokyo: Bureau of the Research on Immunological and Allergic disease, Japan Ministry of Health and Welfare; 2002:1–9.
26. Hochberg MC, Chang RW, Dwosch I, Lindsey S, Pincus T, Wolfe F: The American College of Rheumatology 1991 revised criteria for the classification of global functional status in rheumatoid arthritis. *Arthritis Rheum* 1992, **35**(5):498–502.
27. Koike R, Takeuchi T, Eguchi K, Miyasaka N: Update on the Japanese guidelines for the use of infliximab and etanercept in rheumatoid arthritis. *Mod Rheumatol* 2007, **17**(6):451–458.
28. Steinbrocker O, Traeger CH, Batterman RC: Therapeutic criteria in rheumatoid arthritis. *J Am Med Assoc* 1949, **140**(8):659–662.
29. Nagano J, Sudo N: Development of a self-administered questionnaire to assess disease-prone personalities: Item construction and content validity (in Japanese). *Kenko Kagaku* 2001, **23**:41–52. English translation: <http://hdl.handle.net/2324/724>.
30. Nagano J, Sudo N, Kaihara C, Shimura M, Kubo C: Validity and reliability of the Stress Inventory: self-administered questionnaire to assess disease-prone personalities (in Japanese). *Kenko Shien* 2001, **3**(2):107–119. English translation: <http://hdl.handle.net/2324/15513>.
31. Nagano J, Nagase S, Sudo N, Kubo C: Psychosocial stress, personality, and the severity of chronic hepatitis C. *Psychosomatics* 2004, **45**(2):100–106.
32. Nagano J, Kono S, Toyomura K, Mizoue T, Yin G, Mibu R, Tanaka M, Kakeji Y, Maebara Y, Okamura T, Ikejiri K, Futami K, Yasunami Y, Maekawa T, Takenaka K, Ichimiya H, Imaizumi N: Personality and colorectal cancer: the Fukuoka colorectal cancer study. *Jpn J Clin Oncol* 2008, **38**(8):553–561.
33. Nagano J, Kakuta C, Motomura C, Odajima H, Sudo N, Nishima S, Kubo C: The parenting attitudes and the stress of mothers predict the asthmatic severity of their children: a prospective study. *BioPsychoSocial Medicine* 2010, **4**:12.
34. Spielberger CD: *Manual for the State-Trait Anger Expression Inventory*. Psychological Assessment Resources: Odessa; 1988.
35. McCollum L, Pincus T: A biopsychosocial model to complement a biomedical model: patient questionnaire data and socioeconomic status usually are more significant than laboratory tests and imaging studies in prognosis of rheumatoid arthritis. *Rheum Dis Clin N Am* 2009, **35**(4):699–712.
36. Kremer J: Nutrition and rheumatic diseases. In *Kelley's textbook of rheumatology. Volume 1*. 6th edition. Edited by Ruddy S, Harris EJ, Sledge C. Philadelphia: Saunders; 2001:713–727.
37. Ishii H, Nagashima M, Tanno M, Nakajima A, Yoshino S: Does being easily moved to tears as a response to psychological stress reflect response to treatment and the general prognosis in patients with rheumatoid arthritis? *Clin Exp Rheumatol* 2003, **21**(5):611–616.
38. Pincus T, Sokka T: Complexities in the quantitative assessment of patients with rheumatic diseases in clinical trials and clinical care. *Clin Exp Rheumatol* 2005, **23**(5 Suppl 39):S1–S9.
39. Aletaha D, Neogi T, Silman AJ, Funovits J, Felson DT, Bingham CO 3rd, Birnbaum NS, Burmester GR, Bykerk VP, Cohen MD, Combe B, Costenbader KH, Dougados M, Emery P, Ferraccioli G, Hazes JM, Hobbs K, Huizinga TW, Kavanaugh A, Kay J, Kvien TK, Laing T, Mease P, Menard HA, Moreland LW, Naden RL, Pincus T, Smolen JS, Stanislawska-Biernat E, Symmons D: 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 2010, **62**(9):2569–2581.
40. Prevoo ML, van 't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, van Riel PL: Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995, **38**(1):44–48.

doi:10.1186/1751-0759-8-8

Cite this article as: Nagano et al.: Rational/antiemotional behaviors in interpersonal relationships and the functional prognosis of patients with rheumatoid arthritis: a Japanese multicenter, longitudinal study. *BioPsychoSocial Medicine* 2014 **8**:8.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

