

REVIEW

Open Access

Function and mechanism of tumor suppressor gene LRRC4/NGL-2

Peiyao Li^{1,2,3,4}, Gang Xu^{1,2,3,4,5}, Guiyuan Li^{1,2,3,4} and Minghua Wu^{1,2,3,4*}

Abstract

LRRC4/NGL-2 (Leucine rich repeat containing 4/Netrin-G ligand-2), a relatively specific expressed gene in brain tissue, is a member of the LRRC4/ NGL (netrin-G ligand) family and belongs to the superfamily of LRR proteins. LRRC4/NGL-2 regulates neurite outgrowth and lamina-specific dendritic segmentation, suggesting that LRRC4/NGL-2 is important for the development of the nervous system. In addition, LRRC4/NGL-2 has been identified as a tumor suppressor gene. The overexpression of LRRC4/NGL-2 suppresses glioma cell growth, angiogenesis and invasion through complicated signaling regulation networks. LRRC4/NGL-2 also has the ability to form multiphase loops with miRNA, transcription factors and gene methylation modification; the loss of LRRC4/NGL-2 function may be an important event in multiple biological processes in gliomas. In summary, LRRC4/NGL-2 is a critical gene in the normal development and tumorigenesis of the nervous system.

Keywords: Leucine-rich repeat, Netrin, Netrin-G ligand, miRNA, Methylation, Glioma

Introduction

The leucine-rich repeat (LRR) superfamily is composed of a notably heterogeneous group of proteins containing leucine-rich domains (LRR domain) known to mediate highly specific protein-protein interaction or cell adhesion. Many of LRR-containing proteins are involved in the differentiation and development of normal nervous tissues [1-6].

The LRRC4 (GenBank accession No. AF196976) gene was cloned and characterized from human chromosome 7q31-32 using a computer-assisted positional cloning strategy combining 5'-RACE [7]. Lin et al. (2003) determined that there is a similarity of phenotype and sequence between LRRC4 and the netrin-G1 ligand [8]. Afterwards, Kim et al. demonstrated that LRRC4 directly interacts with netrin-G2, indicating that it is a ligand for netrin-G2; thus, LRRC4 is also named NGL-2 (netrin-G ligand-2) [9].

A genomic database analysis has since identified that LRRC4 is a member of the LRRC4 (NGL, netrin-G ligand) family and belongs to the superfamily of LRR proteins.

Moreover, there are three known members in the LRRC4 family; LRRC4C (NGL-1), LRRC4 (NGL-2) and LRRC4B (NGL-3) [10]. LRRC4/NGL-2 displays down-regulation or expression deletion in primary brain tumor biopsies and has the potential to suppress brain tumor growth [7].

Structure and distribution of LRRC4/NGL-2


LRRC4/NGL-2 is a member of the LRR superfamily. It contains two segments at its N-terminus and C-terminus for sequences representing a putative signal peptide and a transmembrane region, respectively. Following the signal peptide, the core LRR region consists of nine LRRs accompanied by typical amino-flanking (AF) and carboxy-flanking (CF) clusters [1]. The amino-terminal LRR-flanking domain contains cysteines in a C-X3-C-X-C-X8-C pattern, whereas the carboxy-terminal LRR-flanking domain conforms to a P-X2-C-X-C-X19-C-X2-C pattern. Cytosine clusters are known to enhance the stability of the central LRRs [11]. Adjacent to the CF region, one IgC2 domain is identified [8]. The transmembrane domain is followed by a cytoplasmic region, which ends with a PDZ domain-binding motif [12]. LRRC4C/NGL-1, LRRC4/NGL-2, and LRRC4B/NGL-3 share the same domain structure (Figure 1). The LRRC4 family displays a sequence identity of approximately 57-61% in the

* Correspondence: wuminghua554@aliyun.com

¹Hunan Cancer Hospital and the Affiliated Tumor Hospital of Xiangya Medical School Central South University, Changsha, Hunan, China

²Cancer Research Institute, Central South University, Changsha, Hunan 410078, P.R. China

Full list of author information is available at the end of the article



LRR and Ig domains, but their cytoplasmic domains show essentially no sequence identity, except in the PDZ-binding motif. Moreover, the extracellular region immediately preceding the transmembrane domain varies significantly among LRRC4s, suggesting that each LRRC4 may have a distinct function [10].

LRRC4/NGL-2 displayed a brain-specific expression pattern both in humans and mice. LRRC4/NGL-2 was detected in the human brain but not in other normal tissues including the heart, lung, liver and so on [13,14]. The mRNAs for LRRC4C/NGL-1 and LRRC4B/NGL-3 are mainly expressed in the brain, with minor expression in the liver and heart, respectively. The transcripts of the LRRC4 family members are distributed in distinct regions of embryonic and postnatal brains [9,8,13,15]. LRRC4/NGL-2 expression in both the human and mice brain was limited to the following areas: cerebellum, cerebral cortex, occipital pole, frontal lobe, temporal lobe and putamen [13]. The LRRC4C/NGL-1 and LRRC4B/NGL-3 proteins have been indicated to be mainly expressed in the brain but not in other tissues [9]. The absence of LRRC4C/NGL-1 and LRRC4B/NGL-3 protein expression in the liver and heart, respectively, is dissimilar to their mRNA expression patterns. The LRRC4C/NGL-1 protein is detected in the hippocampus, neocortex, and piriform cortex (Table 1). Within neurons, LRRC4/NGL-2 and LRRC4C/NGL-1 are selectively localized to distinct dendritic segments [15]. LRRC4/NGL family members are mainly localized to the postsynaptic side of excitatory synapses, and NGL-1 interacts with netrin-G1, NGL-2 with netrin-G2, and NGL-3 with LAR, which play

Table 1 The expression distribution of the LRRC4 family in brain tissues

LRRC4 family	Protein distribution
LRRC4/NGL-2	cerebellum, cerebral cortex, occipital pole, frontal lobe, temporal lobe, and putamen
LRRC4C/NGL-1	hippocampus, neocortex, and piriform cortex
LRRC4B/NGL-3	similar brain-specific

important roles in the development of axons, dendrites and synapses [10].

Role of LRRC4/NGL-2 in maintaining normal function in the central nervous system

A number of leucine-rich repeat proteins with LRRs and Ig regions, which seem to serve similar functions, have been found to be expressed predominantly in cells of nervous tissues [8,16,17], and are involved in the differentiation and development of normal nervous tissues [1,2]. LRRC4C/NGL-1, LRRC4/NGL-2 and LRRC4B/NGL-3 share the same domain structure. The structure character of LRRC4s implies that these genes might be involved in the development and differentiation of the nervous system. LRRC4C/NGL-1 was first identified as a ligand for netrin-G1 [8]. LRRC4C/NGL-1 has been found to interact with netrin-G1 but not with netrin-G2. In addition, LRRC4/NGL-2 interacts with netrin-G2 but not with netrin-G1 [9]. LRRC4B/NGL-3 does not interact with either netrin-G1 or netrin-G2 but was determined to interact with the receptor tyrosine phosphatase LAR [10,18]. The interaction between netrin-G1 and LRRC4C/NGL-1 has been implicated in the regulation of axonal outgrowth and migration [8]. The LRRC4C/NGL1-netrinG1 and LRRC4/NGL2-netrinG2 interactions have been implicated in the lamina-specific segmentation of dendrites [15]. In the early development of the brain, the expression of mLRRC4/NGL-2 increased with age, suggesting a possible role of LRRC4/NGL-2 in brain development [13]. LRRC4/NGL-2 regulates the formation of excitatory synapses through the recruitment of pre- and postsynaptic proteins [19], participates in the differentiation of neuron and glia cells, and promotes neurite outgrowth [20]. In addition, LRRC4/NGL-2 formed a complex with the NMDA receptor subunits involved in the regulation of NMDA(N-methyl-D-aspartate) receptor-mediated excitatory synaptic transmission [9]. LRRC4/NGL-2 is localized to the dendritic segment in the stratum radiatum of CA1(Cornu Ammonis 1 area) and functions as a key regulator of input-specific synapse development, which is critical for the functional integration of distinct inputs [21]. LRRC4/NGL-2 was found to localize selectively to the tips of HC(horizontal cell) axons, which form reciprocal connections with rods; moreover, LRRC4/NGL-2 works as a central component of pathway-specific development in the outer retina [22]. LRRC4B/NGL-3-LAR interaction is capable of mediating cell aggregation [23]. LRRC4B/NGL-3 might be involved in the maturation of excitatory synapses. A suggested function of the LRRC4B/NGL-3-LAR interaction is synapse formation [23]. The results from studies on the LRRC4 family suggest that LRRC4s play important roles in maintaining normal function in the central nervous system. In addition to

the functions in neuronal cell types, LRRC4/NGL proteins play a role in glial cells [10].

Aberrant expression of LRRC4/NGL-2 in tumors of the central nervous system

LRRC4/NGL-2 was found to be predominantly expressed in the normal brain tissues, though it was deleted or down-regulated in primary brain tumor biopsies (up to 87.5% in gliomas, 80.9% in meningiomas and 85.2% in pituitary and other brain tumors) [7]. LRRC4/NGL-2 was highly specific in brain tissue and grade I gliomas (WHO), but it was reduced or absent in grade II-III gliomas and absent in glioblastoma (WHO, grade IV) [24]. Thus, the loss of LRRC4/NGL-2 function may directly contribute to the increasing tumor grade and is a late event in the pathogenesis of gliomas. Furthermore, LRRC4/NGL-2 is downregulated expression in pituitary adenoma, polymorphisms or haplotypes in the LRRC4/NGL-2 and may have important research significance by predicting the risk of pituitary adenoma [25].

Role of LRRC4/NGL-2 as a tumor suppressor gene in glioma tumorigenesis

LRRC4/NGL-2 significantly inhibited glioma cell proliferation and greatly reduced its capacity to form colonies [24]. Furthermore, it reduced the growth and malignant grade of xenografts arising from glioma cells. The ultrastructure of glioma cell Tet-on-LRRC4/NGL-2 underwent a significant change, the nucleo-cytoplasmic ratio lessened, the nuclear shape became regular, heterochromatin in nuclei decreased while euchromatin increased, the volume of nucleoli lessened, rough ER increased significantly, the Golgi apparatus was well-developed and Golgi vesicles increased and were regularly arranged, most mitochondria were oval and their crista grew in number and were regularly arranged, and finally polyribosomes reduced while free ribosomes increased [26]. In addition, it has the ability to inhibit the invasion and angiogenesis of glioma cells.

LRRC4/NGL-2 suppressed glioma cell proliferation by delaying the cell cycle in late G1 and did not induce apoptosis of tumor cells [26]. Furthermore, LRRC4/NGL-2 inhibited the glioma tumor cell invasion through regulating the expression of the invasion-related molecules including CD44, MMP16, TB10 and annexin A2 [27]. In addition, LRRC4/NGL-2 might be a negative regulator of the RPTP-zeta receptor, contributing to the suppression of the invasion ability of glioma cells [28]. The reintroduction of LRRC4/NGL-2 might inhibit the expression of CXCR4 and SDF-1 α /CXCR4 axis-mediated cell invasion in vitro [29]. Moreover, LRRC4/NGL-2 inhibits glioblastoma cell proliferation, migration and angiogenesis by downregulating pleiotropic cytokine expression and response [30]. Together, these findings demonstrated

that LRRC4/NGL-2 can be identified as a tumor suppressor gene in the tumorigenesis of glioma. The inhibitory effect of LRRC4/NGL-2 on cell proliferation and invasion is dependent on its LRR cassette domain but not on IgC2 or Tm domain. In the LRR cassette domain, the third LRR motif of the core LRR is found to be indispensable for the function of LRRC4/NGL-2, and it plays a crucial role as a "proliferation-inhibition switch". The seventh LRR motif plays a crucial role as an "invasion-inhibition switch" [24].

Signaling transduction network of LRRC4/NGL-2 in glioma

To further analyze the mechanisms of LRRC4/NGL-2 on the suppression of tumorigenesis and progression in gliomas, Zhang et al. investigated alterations in gene expression related to the neurobiology of the Atlas array and found that 1) The overexpression of LRRC4/NGL-2 can elevate the expression levels of certain cell cycle progression regulators, such as RAP1GAP, ephrin-B3, somatostatin receptors, PTPN and NT-3, and regulate suppressing cell cycle progression by impacting on Raf/Rap/Ras pathways; 2) the overexpression of LRRC4/NGL-2 can down-regulate the expression of genes involved in tumor invasion and metastasis, including CD44, MMP16, Thymosin beta-10 and AnnexinA2; and 3) LRRC4/NGL-2 may act as a receptor for adhesion molecule Netrin-G2 by participating in the inhibition of tumorigenesis and suppression of glioma tumorigenesis by preventing the synthesis and release of toxic neurotransmitters [27]. LRRC4/NGL-2 can delay the cell cycle in late G1 by increasing the expression of cell cycle inhibitory molecules (p21, p27) and reducing the expression of cell cycle regulatory proteins (CyclinD1, CDK2, CyclinE, CDK4) via the down-regulation of growth factors (IGF, bFGF, EGF, PDGF, VEGF SDF-1 α) or receptors, the inhibition of K-Ras/c-Raf/ERK/MAPK, PI-3 K/AKT/NF- κ B, p70S6/PKC and STAT3, and the upregulation of the JNK2/c-Jun/mp53 (mutant p53) signaling pathway [24,27]. Moreover, the PMA-stimulated activation of PKC activates PI3K/Akt, but not pERK, in the presence of LRRC4/NGL-2, which is consistent with the fact that PKC is located downstream of ERK but upstream of PI3K/Akt [24]. The reintroduction of LRRC4/NGL-2 in U251 cells inhibits the expression of CXCR4 (CXC chemokine receptor 4) and SDF-1 (stromal cell-derived factor-1) α /CXCR4 axis-mediated downstream intracellular pathways such as ERK1/2 and Akt leading to proliferate, chemotactic and invasive effects (Figure 2) [29].

Epigenetic changes of LRRC4/NGL-2 in glioma

LRRC4/NGL-2 was a candidate tumor suppressor gene that may be involved in the pathogenesis of malignant glioma. However, no genetic alterations of the LRRC4/NGL-2 coding region were found in glioma [24]. The




Figure 2 Schematic diagram showing the signaling regulation network of LRRC4/NGL-2 in glioma. LRRC4/NGL-2 can delay the cell cycle in late G1 by increasing the expression of cell cycle inhibitory molecules (p21, p27) and reducing the expression of cell cycle regulatory proteins (CyclinD1, CDK2, CyclinE, CDK4) via the inhibition of K-Ras/c-Raf/ERK/MAPK, PI-3K/AKT/NF-κB, p70S6/PKC and STAT3, and the upregulation of the JNK2/c-Jun/mp53 signaling pathway. LRRC4/NGL-2 inhibits the expression of CXCR4 and SDF-1 alpha/CXCR4 axis-mediated downstream intracellular pathways leading to proliferate, chemotactic and invasive effects.

methylation-mediated inactivation of LRRC4/NGL-2 was found to be a frequent and glioma-specific event. Furthermore, the methylation of LRRC4/NGL-2 was detected in both the early and late stages of glioma, indicating that the inactivation of the LRRC4/NGL-2 gene might be essential in the early development of glioma and persist through the course of development [31]. Both DNA methylation and histone modification are not isolated epigenetic events; there are cross-talks between them. Trimethyl-H3 (Lys9) is correlated with LRRC4/NGL-2 promoter methylation, which suppresses LRRC4/NGL-2 expression and H3 acetylation, while trimethyl-H3 (lys4) is correlated with LRRC4/NGL-2 promoter demethylation, which activates LRRC4/NGL-2 expression [32].

The other dysregulation mechanisms of LRRC4/NGL-2 in glioma result from the miRNA-dependent regulatory network.

LRRC4/NGL-2 as the core of miRNA-dependent multiphase loops in glioma

MicroRNAs (miRNAs) are approximately 22 nt non-coding RNAs that interfere with the translation of coding mRNAs in a sequence-specific manner [33]. MiRNAs function as tumor suppressors or oncogenes to induce

cellular transformation and tumorigenesis [34-36]. *MiRNA-mediated gene regulation is now considered to be important in numerous biological processes and is also recognized as being crucial for regulating gene expression in tumors* [37,38]. Several key studies on miRNA expression patterns in gliomas have been completed to date. The abnormal expression of some miRNAs (miR-12b, miR-21, miR-221, miR-128, miR-181a, miR-181b and miR-181c et al.) were found in gliomas and might be involved in tumorigenesis and development [25,39-41]. A greater number of studies have been completed via the siRNA profiling of glioma; correlations between siRNA templates, their target genes, and candidate miRNA may help experimentally identify candidate miRNA targets in glioma [42]. *LRRC4/NGL-2* is a target gene of some miRNAs (such as miR-182 and miR-381); meanwhile, it is capable of regulating miRNAs as a tumor suppressor and forms multiphase loops with miRNAs, transcription factors and gene methylation modification in glioma [43-46].

The LRRC4/NGL-2-AP-2-miR-182- LRRC4/NGL-2 Loop

There is a set of cis-acting elements in the promoters of many miRNAs, which are regulated by a series of upstream transcription factors (TFs) [47]. Similar to the

transcription of the protein-coding gene, TFs regulate miRNA expression at the transcriptional level. Further, miRNA might influence the expression of its target genes and constitute a TF-miRNA-gene regulatory network [48,49]. Genes not only modulate the expression of miRNA target genes by influencing TF-mediated miRNAs expression but also regulate TFs through miRNAs [50,51]. In the regulatory processes where target genes are modulated by both miRNAs and transcription factors, TFs regulate miRNAs, and miRNAs suppress TFs. In this way, multiphase loops might be constituted at last [52-54].

LRRC4/NGL-2 is a target gene of miR-381 and miR182; meanwhile, the overexpression of *LRRC4/NGL-2* downregulated the expression of miR-381 and miR182 in glioma cells [43-45]. The interactions between miR-381/miR182 and *LRRC4/NGL-2* were identified to be involved in the pathological progression of astrocytoma [43]. BRD7, a transcriptional cofactor for p53, is highly expressed and negatively correlated with *LRRC4/NGL-2* expression in gliomas. Disturbing miR-182 and miR-381 inhibits the expression of BRD7, arrests glioma cells in the G0/G1 phase of the cell cycle, inhibits glioma cell growth and induces differentiation of glioma cells to astrocyte-like cell by upregulating *LRRC4/NGL-2* and suppressing the *LRRC4/NGL-2*-mediated binding of AP-2/SP1/E2F6/c-Myc to BRD7 in ERK/MAPK and PI-3 K/AKT signal pathways [43,45].

The transcription of miR-182 was induced by transcription factor AP-2; meanwhile, miR-182 can inhibit the expression of *LRRC4/NGL-2*, and *LRRC4/NGL-2* might inhibit the expression of AP-2 though negatively regulating the ERK/MAPK and PI-3 K/AKT signaling pathways. It was demonstrated that the *LRRC4*-AP-2-miR-182-*LRRC4* loop formed among *LRRC4/NGL-2*, miR-182 and AP-2 was involved in glioma development [43,45,46].

The *LRRC4/NGL-2*-miR-185/SP1-DNMT1-*LRRC4/NGL-2* Loop

MiRNAs regulate the epigenetic modification of genes through modulating the expression of DNMT (DNA methyltransferase), maintaining DNA methylation and mediating histone modifications [54-57]. During tumorigenesis, miRNAs not only mediate the methylation of DNAs by regulating DNMT but also can be methylated in its promoter. Thus, a miRNA-DNMT regulatory loop might be formed between miRNAs and DNMT [58-60].

The overexpression of *LRRC4/NGL-2* could increase the expression of miR-185, while miR-185 could regulate global methylation by inhibiting DNA methyltransferase DNMT1 and increasing the expression of such hypermethylation genes such as *LRRC4/NGL-2* in the end. It was indicated that the *LRRC4/NGL-2*-miR-185-DNMT1-*LRRC4/NGL-2* loop among *LRRC4/NGL-2*, miR185 and

DNMT1 participated in glioma development. In addition, DNMT1 was positively regulated by SP1, and it could increase the expression of *LRRC4/NGL-2*, while *LRRC4/NGL-2* could also inhibit SP1 by negatively regulating the ERK/MAPK and PI-3 K/AKT signal pathway. Therefore, the *LRRC4/NGL-2*-SP1-DNMT1-*LRRC4/NGL-2* loop formed among *LRRC4/NGL-2*, SP1 and DNMT1 took part in the glioma formation [44,46]. In addition, CDC42 and RhoA were the direct targets of miR-185. Furthermore, CDC42 and RhoA were inversely correlated with miR-185 expression in gliomas. *LRRC4/NGL-2* mediated its tumor suppressor by regulating miR-185 targets CDC42 and RhoA. *LRRC4/NGL-2* overexpression inhibited glioma cell growth and invasion through miR-185-mediated CDC42 and RhoA direct regulation and VEGFA indirect regulation [44].

Briefly, at the time of *LRRC4/NGL-2* regulating miRNAs as a tumor suppressor, those miRNAs were found to regulate the binding of transcription factors to DNA in their target-mediated signal pathways by directly targeting genes (such as *LRRC4/NGL-2*) or regulating the methylation and expression of such hypermethylation genes such as *LRRC4/NGL-2* by directly targeting DNA methyltransferase and controlling global methylation. Thus, multiphase loops, which had a core of *LRRC4/NGL-2*, were formed (Figure 3) [46]. They were *LRRC4/NGL-2*-AP-2-




Figure 3 Schematic diagram showing the multi-regulation loops of *LRRC4/NGL-2*, miRNAs, TFs and target genes in glioma.

LRRC4/NGL-2-AP-2-miR-182- *LRRC4/NGL-2* loop: *LRRC4/NGL-2* is a target of miR182; meanwhile, the overexpression of *LRRC4/NGL-2* downregulated the expression of miR182. The transcription of miR-182 was induced by AP-2; meanwhile, miR-182 can inhibit the expression of *LRRC4/NGL-2*, and *LRRC4/NGL-2* can inhibit the expression of AP-2 though negatively regulating the ERK/MAPK and PI-3K/AKT signaling pathways. The *LRRC4/NGL-2*-AP-2-miR-182- *LRRC4/NGL-2* loop was formed among *LRRC4/NGL-2*, miR-182 and AP-2. *LRRC4/NGL-2*-miR-185-DNMT1-*LRRC4/NGL-2* loop: The overexpression of *LRRC4/NGL-2* could increase the expression of miR-185, while miR-185 could regulate global methylation by targeting DNMT1 and increasing the expression of *LRRC4/NGL-2*. *LRRC4/NGL-2*-SP1-DNMT1-*LRRC4/NGL-2* loop: DNMT1 was positively regulated by SP1, and it could increase the expression of *LRRC4/NGL-2*, while *LRRC4/NGL-2* could also inhibit SP1 by negatively regulating the ERK/MAPK and PI-3K/AKT signal pathway.

miR-182-LRRC4/NGL-2, LRRC4/NGL-2-miR-185-DNMT1-LRRC4/NGL-2 and LRRC4/NGL-2-SP1-DNMT1-LRRC4/NGL-2. These loops were involved in glioma development with multiple positive feedback formations among them.

Conclusions

LRRC4/NGL-2, a member of the LRR superfamily thought to be involved in tumorigenesis and the development of the nervous tissues, has the potential to suppress gliomas cell growth, angiogenesis and invasion through complex regulation networks. In addition, LRRC4/NGL-2 has the ability to form multiphase loops with miRNA, transcription factors and target genes. These loops are involved in glioma development with multiple positive feedback formations among them. The results from studies on LRRC4/NGL-2 suggest that LRRC4/NGL-2 is not only a brain-specific gene, but it has been identified as a tumor suppressor gene for gliomas. Therefore, the loss of LRRC4/NGL-2 function may be an important event in multiple biological processes related to gliomas.

There are still many interesting questions that need to be addressed in future studies. Does the missing LRRC4/NGL-2 gene affect the development of the central nervous system? Does the lack of LRRC4/NGL-2 expression result in impaired learning and memory performance? To answer these questions, the role of LRRC4/NGL-2 in central nervous development and function should be verified through in vivo studies employing neuron-specific knockout mice.

Abbreviations

LRR: Leucine-rich repeat; LRRC4: Leucine rich repeat containing 4; NGL: Netrin-G ligand; AF: Amino-flanking; CF: Carboxy-flanking; Ig: Immunoglobulin; PDZ: PSD-95/Dlg/ZO-1; CXCR4: CXC chemokine receptor 4; SDF-1: Stromal cell-derived factor-1; TF: transcription factors; HC: Horizontal cell; CA1: Cornu Ammonis 1 area; NMDA: N-methyl-D-aspartate.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PYL collected the related paper and drafted the manuscript. MHW, GX and GYL participated in the design of the review and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by National Science Foundation of China (81171932).

Author details

¹Human Cancer Hospital and the Affiliated Tumor Hospital of Xiangya Medical School Central South University, Changsha, Hunan, China. ²Cancer Research Institute, Central South University, Changsha, Hunan 410078, P.R. China. ³Key Laboratory of Carcinogenesis and Cancer Invasion, Ministry of Education, Shanghai, China. ⁴Key Laboratory of Carcinogenesis, Ministry of Health, Changsha, Hunan, China. ⁵Medical College, University of South China, Hengyang, Hunan, China.

Received: 24 July 2014 Accepted: 15 December 2014
Published: 19 December 2014

References

1. Kobe B, Deisenhofer J: The leucine-rich repeat: a versatile binding motif. *Trends Biochem Sci* 1994, **19**:415–421.
2. Kobe B, Kajava AV: The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol* 2001, **11**:725–732.
3. Dolan J, Walshe K, Alsbury S, Hokamp K, O'Keefe S, Okafuji T, Miller SF, Tear G, Mitchell KJ: The extracellular leucine-rich repeat superfamily; a comparative survey and analysis of evolutionary relationships and expression patterns. *BMC Genomics* 2007, **8**:320.
4. Taguchi A, Wanaka A, Mori T, Matsumoto K, Imai Y, Tagaki T, Tohyama M: Molecular cloning of novel leucine-rich repeat proteins and their expression in the developing mouse nervous system. *Brain Res Mol Brain Res* 1996, **35**:31–40.
5. de Wit J, Ghosh A: Control of neural circuit formation by leucine-rich repeat proteins. *Trends Neurosci* 2014, **37**:539–550.
6. Bando T, Morikawa Y, Hisaoka T, Komori T, Miyajima A, Senba E: Dynamic expression pattern of leucine-rich repeat neuronal protein 4 in the mouse dorsal root ganglia during development. *Neurosci Lett* 2013, **548**:73–78.
7. Wang J, Qian J, Dong L, Li X, Tan C, Li J, Zhang B: Identification of LRRC4, a Novel Member of Leucine-rich Repeat (LRR) Superfamily, and Its Expression Analysis in Brain Tumor. *Prog Biochem Biophys* 2002, **29**:233–239.
8. Lin JC, Ho WH, Gurney A, Rosenthal A: The netrin-G1 ligand NGL-1 promotes the outgrowth of thalamocortical axons. *Nat Neurosci* 2003, **6**:1270–1276.
9. Kim S, Burette A, Chung HS, Kwon SK, Woo J, Lee HW, Kim K, Kim H, Weinberg RJ, Kim E: NGL family PSD-95-interacting adhesion molecules regulate excitatory synapse formation. *Nat Neurosci* 2006, **9**:1294–1301.
10. Woo J, Kwon SK, Kim E: The NGL family of leucine-rich repeat-containing synaptic adhesion molecules. *Mol Cell Neurosci*. 2009; **42**:1–10.
11. Gu W, Brodtkorb E, Steinlein OK: LG1 is mutated in familial temporal lobe epilepsy characterized by aphasic seizures. *Ann Neurol* 2002, **52**:364–367.
12. Ko J, Kim E: Leucine-rich repeat proteins of synapses. *J Neurosci Res* 2007, **85**:2824–2832.
13. Zhang Q, Wang J, Fan S, Wang L, Cao L, Tang K, Peng C, Li Z, Li W, Gan K, Liu Z, Li X, Shen S, Li G: Expression and functional characterization of LRRC4, a novel brain-specific member of the LRR superfamily. *FEBS Lett* 2005, **579**:3674–3682.
14. Dan L, Ming-hua W, Qiong C, He H, Chen H, Wei-song L, Xiao-ling L, Gui-yuan L: Preparation of anti-LRRC4 polyclonal antibody and its application in constructing expression profile of human gliomas with different pathological grades. *J Cent S Univ (Med Sci)* 2007, **32**:373–379.
15. Nishimura-Akiyoshi S, Niimi K, Nakashiba T, Itoharu S: Axonal netrin-Gs transneuronally determine lamina-specific subdendritic segments. *Proc Natl Acad Sci U S A* 2007, **104**:14801–14806.
16. Carim-Todd L, Escarceller M, Estivill X, Sumoy L: LRRN6A/LERN1 (leucine-rich repeat neuronal protein 1), a novel gene with enriched expression in limbic system and neocortex. *Eur J Neurosci* 2003, **18**:3167–3182.
17. Kuja-Panula J, Kiiltomaki M, Yamashiro T, Rouhianen A, Rauvala H: AMIGO, a transmembrane protein implicated in axon tract development, defines a novel protein family with leucine-rich repeats. *J Cell Biol* 2003, **160**:963–973.
18. Kwon SK, Woo J, Kim SY, Kim H, Kim E: Trans-synaptic adhesions between netrin-G ligand-3 (NGL-3) and receptor tyrosine phosphatases LAR, protein-tyrosine phosphatase delta (PTPdelta), and PTPsigma via specific domains regulate excitatory synapse formation. *J Biol Chem* 2010, **285**:13966–13978.
19. Biederer T: Hooking up new synapses. *Nat Neurosci* 2006, **9**:1203–1204.
20. Wu M, Huang H, Chen Q, Li D, Zheng Z, Xiong W, Zhou Y, Li X, Zhou M, Lu J, Shen S, Li G: Leucine-rich repeat C4 protein is involved in nervous tissue development and neurite outgrowth, and induction of glioma cell differentiation. *Acta Biochim Biophys Sin (Shanghai)* 2007, **39**:731–738.
21. DeNardo LA, de Wit J, Otto-Hitt S, Ghosh A: NGL-2 regulates input-specific synapse development in CA1 pyramidal neurons. *Neuron* 2012, **76**:762–775.
22. Soto F, Watkins KL, Johnson RE, Schottler F, Kerschensteiner D: NGL-2 regulates pathway-specific neurite growth and lamination, synapse formation, and signal transmission in the retina. *J Neurosci* 2013, **33**:11949–11959.
23. Woo J, Kwon SK, Choi S, Kim S, Lee JR, Dunah AW, Sheng M, Kim E: Trans-synaptic adhesion between NGL-3 and LAR regulates the formation of excitatory synapses. *Nat Neurosci* 2009, **12**:428–437.

24. Wu M, Huang C, Gan K, Huang H, Chen Q, Ouyang J, Tang Y, Li X, Yang Y, Zhou H, Zhou Y, Zeng Z, Xiao L, Li D, Tang K, Shen S, Li G: **LRRC4, a putative tumor suppressor gene, requires a functional leucine-rich repeat cassette domain to inhibit proliferation of glioma cells in vitro by modulating the extracellular signal-regulated kinase/protein kinase B/ nuclear factor-kappaB pathway.** *Mol Biol Cell* 2006, **17**:3534–3542.
25. Xiao L, Tu C, Chen S, Yu Z, Lei Q, Wang Z, Xu G, Wu M, Li G: **LRRC4 haplotypes are associated with pituitary adenoma in a Chinese population.** *Med Oncol* 2014, **31**:888.
26. Zhang QH, Wang LL, Cao L, Peng C, Li XL, Tang K, Li WF, Liao P, Wang JR, Li GY: **Study of a novel brain relatively specific gene LRRC4 involved in glioma tumorigenesis suppression using the Tet-on system.** *Acta Biochim Biophys Sin (Shanghai)* 2005, **37**:532–540.
27. Zhang QH, Wu MH, Wang LL, Cao L, Tang K, Peng C, Gan K, Li XL, Li GY: **Profiling of differentially expressed genes in LRRC4 overexpressed glioblastoma cells by cDNA array.** *Acta Biochim Biophys Sin (Shanghai)* 2005, **37**:680–687.
28. Wu M, Gan K, Huang C, Tang Y, Chen Q, Tang K, Li X, Shen S, Li G: **LRRC4 controls in vitro invasion of glioblastoma cells through inhibiting RPTP-zeta expression.** *J Neurooncol* 2006, **80**:133–142.
29. Wu M, Chen Q, Li D, Li X, Li X, Huang C, Tang Y, Zhou Y, Wang D, Tang K, Cao L, Shen S, Li G: **LRRC4 inhibits human glioblastoma cells proliferation, invasion, and proMMP-2 activation by reducing SDF-1 alpha/CXCR4-mediated ERK1/2 and Akt signaling pathways.** *J Cell Biochem* 2008, **103**:245–255.
30. Wu M, Huang C, Li X, Li X, Gan K, Chen Q, Tang Y, Tang K, Shen S, Li G: **LRRC4 inhibits glioblastoma cell proliferation, migration, and angiogenesis by downregulating pleiotropic cytokine expression and responses.** *J Cell Physiol* 2008, **214**:65–74.
31. Zhang Z, Li D, Wu M, Xiang B, Wang L, Zhou M, Chen P, Li X, Shen S, Li G: **Promoter hypermethylation-mediated inactivation of LRRC4 in gliomas.** *BMC Mol Biol* 2008, **9**:99.
32. Minghua W, Zuping Z, Xiaoping L, Hailin T: **Methylation Level of Genes and MiRNA-Mediated Methylation Modification Mechanism In Glioma.** In *Protein Purification and Analysis III - Methods and Applications*. 1st edition. Hong Kong: iConcept Press; 2014.
33. Mathupala SP, Mittal S, Guthikonda M, Sloan AE: **MicroRNA and brain tumors: a cause and a cure?** *DNA Cell Biol* 2007, **26**:301–10.
34. Zhang L, Liu T, Huang Y, Liu J: **microRNA-182 inhibits the proliferation and invasion of human lung adenocarcinoma cells through its effect on human cortical actin-associated protein.** *Int J Mol Med* 2011, **28**:381–88.
35. Bartel DP: **MicroRNAs: target recognition and regulatory functions.** *Cell* 2009, **136**:215–233.
36. Kusenda B, Mraz M, Mayer J, Pospisilova S: **MicroRNA biogenesis, functionality and cancer relevance.** *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2006, **150**:205–215.
37. Shi L, Cheng Z, Zhang J, Li R, Zhao P, Fu Z, You Y: **hsa-mir-181a and hsa-mir-181b function as tumor suppressors in human glioma cells.** *Brain Res* 2008, **1236**:185–193.
38. Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**:281–297.
39. Nelson PT, Baldwin DA, Kloosterman WP, Kauppinen S, Plasterk RH, Mourelatos Z: **RAKE and LNA-ISH reveal microRNA expression and localization in archival human brain.** *RNA* 2006, **12**:187–191.
40. Chan JA, Krichevsky AM, Kosik KS: **MicroRNA-21 is an antiapoptotic factor in human glioblastoma cells.** *Cancer Res* 2005, **65**:6029–6033.
41. Ciafre SA, Galardi S, Mangiola A, Ferracin M, Liu CG, Sabatino G, Negrini M, Maira G, Croce CM, Farace MG: **Extensive modulation of a set of microRNAs in primary glioblastoma.** *Biochem Biophys Res Commun* 2005, **334**:1351–1358.
42. Mathupala SP, Guthikonda M, Sloan AE: **RNAi based approaches to the treatment of malignant glioma.** *Technol Cancer Res Treat* 2006, **5**:261–269.
43. Tang H, Wang Z, Liu Q, Liu X, Wu M, Li G: **Disturbing miR-182 and -381 inhibits BRD7 transcription and glioma growth by directly targeting LRRC4.** *PLoS One* 2014, **9**:e84146.
44. Tang H, Wang Z, Liu X, Liu Q, Xu G, Li G, Wu M: **LRRC4 inhibits glioma cell growth and invasion through a miR-185-dependent pathway.** *Curr Cancer Drug Targets* 2012, **12**:1032–1042.
45. Tang H, Liu X, Wang Z, She X, Zeng X, Deng M, Liao Q, Guo X, Wang R, Li X, Zeng F, Wu M, Li G: **Interaction of hsa-miR-381 and glioma suppressor LRRC4 is involved in glioma growth.** *Brain Res* 2011, **1390**:21–32.
46. Minghua W: *Methylomes in Epigenetics and Epigenomics*. Rijeka: InTech; 2014.
47. Ozsolak F, Poling LL, Wang Z, Liu H, Liu XS, Roeder RG, Zhang X, Song JS, Fisher DE: **Chromatin structure analyses identify miRNA promoters.** *Genes Dev* 2008, **22**:3172–3183.
48. Ma L, Young J, Prabhala H, Pan E, Mestdagh P, Muth D, Teruya-Feldstein J, Reinhardt F, Onder TT, Valastyan S, Westermann F, Speleman F, Vandesompele J, Weinberg RA: **miR-9, a MYC/MYCN-activated microRNA, regulates E-cadherin and cancer metastasis.** *Nat Cell Biol* 2010, **12**:247–256.
49. Schulte JH, Horn S, Otto T, Samans B, Heukamp LC, Eilers UC, Krause M, Astrahantseff K, Klein-Hitpass L, Buettner R, Schramm A, Christiansen H, Eilers M, Eggert A, Berwanger B: **MYCN regulates oncogenic MicroRNAs in neuroblastoma.** *Int J Cancer* 2008, **122**:699–704.
50. Gehrke S, Imai Y, Sokol N, Lu B: **Pathogenic LRRK2 negatively regulates microRNA-mediated translational repression.** *Nature* 2010, **466**:637–641.
51. Brock M, Trenkmann M, Gay RE, Michel BA, Gay S, Fischler M, Ulrich S, Speich R, Huber LC: **Interleukin-6 modulates the expression of the bone morphogenetic protein receptor type II through a novel STAT3-microRNA cluster 17/92 pathway.** *Circ Res* 2009, **104**:1184–1191.
52. Rosa A, Brivanlou AH: **A regulatory circuitry comprised of miR-302 and the transcription factors OCT4 and NR2F2 regulates human embryonic stem cell differentiation.** *EMBO J* 2011, **30**:237–248.
53. Brabletz S, Bajdak K, Meidhof S, Burk U, Niedermann G, Firat E, Wellner U, Dimmler A, Faller G, Schubert J, Brabletz T: **The ZEB1/miR-200 feedback loop controls Notch signalling in cancer cells.** *EMBO J* 2011, **30**:770–782.
54. Liu S, Wu LC, Pang J, Santhanam R, Schwind S, Wu YZ, Hickey CJ, Yu J, Becker H, Maharry K, Radmacher MD, Li C, Whitman SP, Mishra A, Stauffer N, Eiring AM, Briesewitz R, Baiocchi RA, Chan KK, Paschka P, Caligiuri MA, Byrd JC, Croce CM, Bloomfield CD, Perrotti D, Garzon R, Marcucci G: **Sp1/ NFKappaB/HDAC/miR-29b regulatory network in KIT-driven myeloid leukemia.** *Cancer Cell* 2010, **17**:333–347.
55. Das S, Foley N, Bryan K, Watters KM, Bray I, Murphy DM, Buckley PG, Stallings RL: **MicroRNA mediates DNA demethylation events triggered by retinoic acid during neuroblastoma cell differentiation.** *Cancer Res* 2010, **70**:7874–7881.
56. Pan W, Zhu S, Yuan M, Cui H, Wang L, Luo X, Li J, Zhou H, Tang Y, Shen N: **MicroRNA-21 and microRNA-148a contribute to DNA hypomethylation in lupus CD4+ T cells by directly and indirectly targeting DNA methyltransferase 1.** *J Immunol* 2010, **184**:6773–6781.
57. Garzon R, Liu S, Fabbri M, Liu Z, Heaphy CE, Callegari E, Schwind S, Pang J, Yu J, Muthusamy N, Havelange V, Volinia S, Blum W, Rush LJ, Perrotti D, Andreeff M, Bloomfield CD, Byrd JC, Chan K, Wu LC, Croce CM, Marcucci G: **MicroRNA-29b induces global DNA hypomethylation and tumor suppressor gene reexpression in acute myeloid leukemia by targeting directly DNMT3A and 3B and indirectly DNMT1.** *Blood* 2009, **113**:6411–6418.
58. Liang R, Bates DJ, Wang E: **Epigenetic Control of MicroRNA Expression and Aging.** *Curr Genomics* 2009, **10**:184–193.
59. Grady WM, Parkin RK, Mitchell PS, Lee JH, Kim YH, Tsuchiya KD, Washington MK, Paraskeva C, Willson JK, Kaz AM, Kroh EM, Allen A, Fritz BR, Markowitz SD, Tewari M: **Epigenetic silencing of the intronic microRNA hsa-miR-342 and its host gene EVL in colorectal cancer.** *Oncogene* 2008, **27**:3880–3888.
60. Lujambio A, Calin GA, Villanueva A, Ropero S, Sanchez-Cespedes M, Blanco D, Montuenga LM, Rossi S, Nicoloso MS, Faller WJ, Gallagher WM, Eccles SA, Croce CM, Esteller M: **A microRNA DNA methylation signature for human cancer metastasis.** *Proc Natl Acad Sci U S A* 2008, **105**:13556–13561.

doi:10.1186/1476-4598-13-266

Cite this article as: Li et al.: Function and mechanism of tumor suppressor gene LRRC4/NGL-2. *Molecular Cancer* 2014 **13**:266.