




Pituitary Adenoma Coexistent with Sellar Clear Cell Meningioma Unattached to the Dura: Case Report and Treatment Considerations

Grégoire P. Chatain¹ Keanu Chee¹  Meghan Driscoll² B.K. Kleinschmidt-DeMasters^{1,2}
Kevin O. Lillehei¹

¹ Department of Neurosurgery, University of Colorado School of Medicine, Aurora, Colorado, United States

² Department of Pathology, University of Colorado, School of Medicine, Aurora, Colorado, United States

Address for correspondence Kevin O. Lillehei, MD, Department of Neurosurgery, University of Colorado School of Medicine, 12605 E 16th Ave, Aurora, CO, 80045, United States
(e-mail: kevin.lillehei@cuanschutz.edu).

J Neurol Surg Rep 2024;85:e1–e10.

Abstract

Collision tumors involving the sella are rare. Intrасellar collision tumors are most commonly composed of a combination of pituitary adenomas and pituitary neuroendocrine tumors; however, collision tumors consisting of a pituitary adenoma and intrасellar meningioma are exceedingly rare. The authors present the case of a 47-year-old man who presented with progressive right eye vision loss. Magnetic resonance imaging showed a large, heterogeneously enhancing sellar mass with suprasellar extension. Using a transcranial approach with a right subfrontal craniotomy, near-total resection of the mass was achieved. Histologic analysis confirmed a diagnosis of a gonadotroph adenoma with concomitant clear cell meningioma (CCM). This patient was discharged with improvement in visual acuity and no signs of diabetes insipidus. Given the indistinguishable radiographic characteristics of pituitary adenoma and CCM, a preoperative diagnosis of a collision tumor was difficult. This case was uniquely challenging since the CCM component lacked the classic dural attachment that is associated with meningiomas on neuroimaging. CCMs are classified as central nervous system (CNS) World Health Organization (WHO) grade 2 tumors and tend to behave more aggressively, therefore warranting close surveillance for signs of tumor recurrence. This is the first case to report a collision tumor consisting of pituitary adenoma and CCM.

Keywords

- ▶ collision tumors
- ▶ meningioma
- ▶ pituitary adenoma
- ▶ sella

Introduction

Coexistent tumors of the sellar region are rare. A recent study that reviewed 16,283 autopsy and surgical pathology cases found that double or triple pituitary adenomas/pituitary neuroendocrine tumors (PitNETs) were the most frequent combination, comprising 40/232 (17.2%) of cases presenting

with dual lesions.¹ The combination of meningioma and PitNET was one of the rarest combinations (6/232 [2.6%]), far behind PitNET + gangliocytoma (34/232 [14.7%]) or PitNET + sellar metastasis (12/232 [5.2%]).¹ Individually, both PitNETs and sellar meningiomas can attain large sizes and cause similar symptoms due to compromise of the optic apparatus,

received
August 28, 2023
accepted after revision
November 5, 2023

DOI <https://doi.org/10.1055/s-0043-1777792>.
ISSN 2193-6358.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

making the diagnosis and treatment of coexisting sellar tumors, or “collision tumors,” difficult.

Herein we present the case of a 47-year-old man who underwent a right frontal craniotomy for resection of a large sellar/suprasellar mass, found only after histological examination to be a dual PitNET and suprasellar meningioma. A transcranial surgical approach was performed due to the extent of suprasellar extension of the tumor and the relative narrowing hourglass deformity seen on magnetic resonance imaging (MRI) at the level of the diaphragma sella. It was only intraoperatively that two separate tumors were identified. Interestingly, intraoperative recognition of the suprasellar lesion as a meningioma was challenging due to the lack of typical dural attachment or vascular pedicle, as is seen for most meningiomas. Histological examination revealed two noncontiguous tumors, a clear cell meningioma, central nervous system (CNS) World Health Organization (WHO) grade 2, and a gonadotroph adenoma. The nondural attachment characteristic of the clear cell meningioma, as well as its occurrence in an atypical anatomical site for this meningioma type, is discussed in conjunction with review of the literature on meningioma and PitNET.

Case Description

History and Examination

The patient is a 47-year-old otherwise healthy man who presented to the optometrist with a 2-year history of progressive vision loss of the right eye. Formal visual acuity testing revealed sensitivity to hand-waving only, and an OS 20/60 with visual fields testing showing complete loss of vision in all four quadrants and sparing of the inferior nasal quadrant only in the left eye. An urgent MRI brain demonstrated a large homogeneously enhancing sellar mass with suprasellar extension (►Fig. 1). It was noted to cause significant mass effect on adjacent neural structures including the optic chiasm/nerves with significant cavernous sinus invasion. Following admission to the neurosurgery service, a complete pituitary workup was obtained. The data at the time were the following: cortisol, 8 µg/dL; adrenocorticotropic hormone (ACTH), 11.0 pg/mL; follicle stimulating hormone (FSH), 9.1 mIU/mL; luteinizing hormone (LH), 2.1 mIU/mL; prolactin (PRL), 11.1 ng/mL; free T4, 1.55 ng/dL; thyroid-stimulating hormone (TSH), 1.41 mIU/L; insulinlike growth factor 1 (IGF-1), 208 ng/mL.

Operation and Postoperative Course

Due to the radiographical evidence of significant suprasellar extension of the tumor toward the tuberculum sella along with a relative narrowing of the tumor at the level of the diaphragma sella, a transcranial approach was chosen over a transsphenoidal surgery. Although experienced with both the microscopic and endoscopic transnasal transsphenoidal approaches, we were concerned with our ability to safely access the suprasellar portion of the tumor, as well as the portion extending laterally over the right internal carotid artery. Therefore, the patient underwent a right frontal craniotomy and the dissection was carried out subfrontally




Fig. 1 (A,C) Coronal and (B,D) sagittal T1-weighted postcontrast brain magnetic resonance imaging (MRI). (A,B) Preoperative images demonstrating a heterogeneously enhancing mass (3.8 × 5.1 × 3.5 cm) expanding the sella turcica and extending superiorly into the suprasellar cisterns. Compression of optic apparatus is noted along with encasement of the paraclinoid internal carotid artery (ICA). A thin hyperintense layer of signal is seen between the intrasellar and suprasellar compartment. (C,D) Postoperative images illustrate the near total resection of the sellar and suprasellar mass with minimal residual enhancing tissue along the floor of the sella turcica.

to access the tumor. Intraoperative gross pathological findings revealed a well-circumscribed encapsulated suprasellar tumor that was dissected from of the brain parenchyma without obvious dural attachments. A frozen specimen section was suggestive of meningioma. As expected from preoperative imaging, it was significantly displacing the optic nerves and chiasm laterally and posteriorly (►Fig. 2). Following resection of the suprasellar lesion, the neurosurgeons were able to clearly visualize the pituitary stalk displaced to the left leading to the diaphragma sella. At this point, as no tumoral connection was seen between the resected suprasellar tumor and the diaphragma sella, a decision was made to open the diaphragma sella adjacent to the pituitary stalk on the right (►Fig. 2). At that point, intrasellar tumor was encountered, which was classic for a pituitary adenoma. Tumor was removed from within the sella using pituitary curettes through a transdiaphragmatic approach. Two separate tumor samples were sent for permanent pathology.

Histopathology

Microscopically, there were two distinct tumors. The meningioma showed numerous bands of collagen, with cells manifesting clear cytoplasm (►Fig. 3A). Immunohistochemical (IHC) staining for somatostatin receptor confirmed a meningothelial tumor (►Fig. 3B), while histochemical staining for periodic acid–Schiff (PAS) highlighted the glycogen-rich cytoplasm (►Fig. 3C, left). Diastase digestion proved the presence of glycogen by the absence of PAS staining after treatment, pathognomonic for clear cell meningioma (►Fig. 3C, right). The other, separate component of the




Fig. 2 Intraoperative image of the transcranial subfrontal approach illustrating the suprasellar region during tumor resection and the surrounding anatomy. Note the sella opening through which the pituitary adenoma was resected using ring curettes. CN II, cranial nerve II (optic nerve); ICA, internal carotid artery; OC, optic chiasm

resection showed pituitary adenoma/PitNET with patternless sheets of cells (**►Fig. 3D**), diffuse nuclear immunostaining for steroidogenic factor 1 (SF-1; **►Fig. 3E**), and the




Fig. 3 Microscopically, there were two distinct tumors. (A) The meningioma showed numerous bands of collagen, with cells manifesting clear cytoplasm. (B) Immunohistochemical (IHC) staining for somatostatin receptor confirmed a meningotheial tumor, (C) while IHC staining for periodic acid–Schiff (PAS) highlighted the glycogen-rich cytoplasm. Diastase digestion proved the presence of glycogen by absence of PAS staining after digestion, which is pathognomonic for clear cell meningioma (C). (D) The other separate component of the resection showed pituitary adenoma/pituitary neuroendocrine tumors (PitNET) with patternless sheets of cells, (E) diffuse nuclear immunostaining for steroidogenic factor 1, and (F) the typical patchy immunostaining for follicle stimulating hormone, which is indicative of a gonadotroph adenoma.

typical patchy immunostaining for FSH (**►Fig. 3F**) indicative of gonadotroph adenoma/PitNET.

Literature Search Strategy

A literature review was conducted using PubMed to identify available case reports and case series on collision tumors occurring in the sellar or suprasellar region of the brain from 1986 to 2022, linking “collision tumor” and “sella” or “suprasellar” or “parasellar” or “pituitary.” The search regarding clear cell meningioma utilized the search terms “clear cell meningioma” and “without dural attachment.” After independent evaluation for content and relevance, we identified a total of 30 articles published from the years 1986 to 2022 that describe collision tumors comprising PitNETs and various co-occurring tumors, and 6 articles published from 2009 to 2021 describing the diagnosis of clear cell meningioma without a dural attachment. The results of this literature review are shown in **►Tables 1–1–30** and **2. 31–37**

Discussion

Collision tumors affecting the sellar and suprasellar regions are rare. Due to indistinguishable radiographical characteristics, the preoperative diagnosis of a sellar collision tumor is difficult to differentiate from an isolated tumor type and is often preemptively diagnosed as a pituitary adenoma. Dual PitNETs are the most frequent type of sellar collision tumor, some of which are separate and some of which are contiguous.^{1,38} However, the second most common combination of tumor types varies significantly. Secondary tumors that have been described as co-occurring with pituitary adenomas include gangliocytomas, pilocytic astrocytomas, craniopharyngiomas, spindle cell astrocytomas, pituicytomas, granular cell tumors, and Rathke’s cleft cysts.^{19–21,25} Collision tumors consisting of a pituitary adenoma and parasellar meningioma are exceedingly rare. There are several case reports and series describing the coexistence of a pituitary adenoma and sellar/suprasellar meningioma, although none have reported a specific diagnosis of clear cell meningioma.^{1,2,4,11,12,14,17,18,23,26,28,30} Therefore, this case report represents the first description of a collision tumor consisting of a gonadotroph adenoma with a coexisting clear cell meningioma without a dural tail, the latter of which represented a significant diagnostic challenge.

Sellar or suprasellar meningiomas can closely mimic pituitary adenomas on neuroimaging.³⁹ Therefore, a definitive preoperative diagnosis of a pituitary adenoma and parasellar meningioma is not often possible on MRI. In considering our patient’s preoperative MRI, in retrospect it may have been possible to appreciate a small layer of hyperintense signal on T1 postcontrast sequences between the intrasellar and suprasellar tumor components, which may have delineated the distinct border of both the adenoma and meningioma. However, this layer, representing the diaphragma sella, was not found to be completely contiguous on the sagittal MRI, and was therefore considered an unreliable measure for confirming the radiographical appearance of a collision tumor.

Definitive diagnosis of a collision tumor requires histologic evaluation of the resected tumor specimen by the

Table 1 Literature review of case reports and case series on sellar and suprasellar collision tumors consisting of pituitary adenomas with various secondary neoplasm

Study	Patient age (y)	Sex	Clinical presentation	Tumor type 1	Tumor type 2	Surgical approach	Additional comment
Yamada et al ²	52	F	Headache with disturbances in visual acuity and galactorrhea	PA	Meningioma	Frontal osteoplastic craniotomy	Complete relief of headache and visual disturbance, with galactorrhea controlled with bromocriptine
Tajika et al ³	56	F	Mild acromegaly	GH-secreting PA	Gangliocytoma	Transsphenoidal excision	N/A
Prevedello et al ⁴	52	F	Continuous headache with right eye temporal visual field loss	PA	Meningioma	Endoscopic transsphenoidal excision	Postoperative resolution of headache, complete resolution of visual loss, and preservation of preoperative pituitary function
Karavitaki et al ^{5,a}	50 54	M M	Headache, sleep difficulties, decreased libido, hot flashes Proximal muscle weakness, central weight gain, back/hip pain, easy bruising, hypertension, decreased libido, erectile dysfunction, insomnia, and nocturia	Gonadotropic PA Corticotropic PA	Adamantinomatous craniopharyngioma Rathke's cleft cyst	Endoscopic transsphenoidal excision Endoscopic transsphenoidal excision	Postoperative hypogonadism requiring testosterone replacement therapy Postoperative diabetes insipidus and panhypopituitarism
Moshkin et al ⁶	12	M	Incidental finding	Silent PA subtype 3	Adamantinomatous craniopharyngioma	Right craniotomy	N/A
Koutouroussiou et al ^{7,a}	42 60 47 38 52 49 76 46	F M F M F F M F	Cushing's disease Hypogonadism and hyperprolactinemia Acromegaly and hypopituitarism Acromegaly, headache, and decreased libido Acromegaly, amenorrhea, and hyperprolactinemia Hypopituitarism Acromegaly and headache	ACTH-secreting PA Nonfunctioning PA GH-secreting PA GH-secreting PA GH-secreting PA GH-secreting PA GH-secreting PA Nonfunctioning PA GH-secreting PA	Rathke's cleft cyst Neurosarcoidosis Gangliocytoma Schwannoma Gangliocytoma Prolactinoma Rathke's cleft cyst Gangliocytoma	Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision Transsphenoidal excision	Postoperative hypocortisolism Persistent hypogonadism and tumor recurrence requiring reoperation Persistent hypopituitarism N/A Tumor recurrence requiring somatostatin analog treatment and radiotherapy Required dopamine agonist for persistent hyperprolactinemia Required hormone

Table 1 (Continued)

Study	Patient age (y)	Sex	Clinical presentation	Tumor type 1	Tumor type 2	Surgical approach	Additional comment
							replacement therapy for persistent panhypopituitarism N/A
Rivera et al ⁸	58	M	Rapid-onset diplopia	Prolactinoma	Plasmacytoma (multiple myeloma)	Stereotactic radiosurgery + cabergoline (prolactinoma) Nonoperative treatment regimen (plasmacytoma): Thalidomide Dexamethasone Pamidronate Bone marrow transplant	Postoperative secondary hypogonadism and hypothyroidism
Sahli et al ⁹	74	M	Progressive neuro-ophthalmologic deterioration	Gonadotrophic PA	Osteochondroma	Endoscopic transsphenoidal excision with adjuvant radiotherapy	Partial residual pituitary insufficiency with persistent FSH level elevation
Jin et al ¹⁰	37	F	Intermittent left eye blurring and headache	PA	Craniopharyngioma	Transsphenoidal excision (primary tumor) Interhemispheric transcallosal approach (residual tumor)	Transient postoperative diabetes insipidus and hyponatremia
Mahyash et al ¹¹	36	F	Frontal headache with visual field disturbance in the right eye	PA	Meningioma	Endoscopic transsphenoidal excision	Gross total resection with sufficient decompression of the optic apparatus
Karsy et al ¹²	70	F	Altered mental status, mutism, and incontinence	PA	Meningioma	Endoscopic transsphenoidal excision	Discharged home with permanent ventriculoperitoneal shunt and no significant neurological deficits
Matyja et al ^{13,a}	51 59 58 63	F M F F	Acromegaly and menstrual irregularities Headache and visual disturbances Headache and diplopia Acromegaly, headache, and sleep apnea syndrome	Somatotroph PA Somatotroph PA Somatotroph PA Somatotroph PA	Gangliocytoma Gangliocytoma Gangliocytoma Gangliocytoma	Transsphenoidal excision with adjuvant radiosurgery Endoscopic transsphenoidal excision Transsphenoidal excision Transsphenoidal excision	Persistent postoperative acromegaly with hypopituitarism N/A N/A N/A
Lim et al ¹⁴	65	F	Visual disturbances and vertigo	Nonfunctioning PA	Meningioma	Endoscopic transsphenoidal excision	Improvement in visual symptoms with normal postoperative hormonal studies

(Continued)

Table 1 (Continued)

Study	Patient age (y)	Sex	Clinical presentation	Tumor type 1	Tumor type 2	Surgical approach	Additional comment
Ban et al ¹⁵	74	M	Bilateral retro-orbital pain, left-sided ptosis, diplopia, headache, and nausea	FSH-secreting PA	DLBCL	Endoscopic transsphenoidal excision (PA) Chemotherapy (DLBCL)	N/A
Heng et al ¹⁶	46	F	Headache and decrease in visual acuity	Nonfunctioning PA	Gangliocytoma	Endoscopic transsphenoidal excision	Required Gamma Knife radiosurgery due to tumor recurrence
Zhao et al ^{17,a}	58 58	F F	Acromegaly Acromegaly	GH-secreting PA GH-secreting PA	Meningioma Meningioma	Transsphenoidal excision (PA) Craniotomy (meningioma) Transsphenoidal excision (PA) Craniotomy (meningioma)	N/A N/A
Amirjamshidi et al ^{18,a}	37 42	F M	Oligomenorrhea, headache, diplopia, and progressive visual impairment Acromegaly, decreased visual acuity with bitemporal hemianopsia	Prolactinoma GH-secreting PA	Meningioma Meningioma	Right pterional craniotomy with tumor resection (meningioma) Cabergoline (PA) Transsphenoidal excision (failed) with subsequent right pterional craniotomy with tumor resection	Continuous postoperative treatment with cabergoline Postoperative improvement in vision with no evidence of tumor recurrence
Levitus and Charitou ¹⁹	44	F	Incidental finding following head injury	GH-secreting PA	Gangliocytoma	Endoscopic transsphenoidal excision	Transient postoperative central adrenal insufficiency and permanent diabetes insipidus No tumor recurrence
Malli et al ²⁰	64	M	Bitemporal hemianopsia	Prolactinoma	Pilocytic astrocytoma	Endoscopic transsphenoidal excision with subfrontal craniotomy	N/A
Miyazaki et al ²¹	48	M	Memory disturbance, depression, and hemiplegia	PA	Adamantinomatous craniopharyngioma	Transsphenoidal excision (PA) Left frontoparietal craniotomy with tumor resection and cyst drainage (craniopharyngioma)	Improvement in all symptoms
Snyder et al ²²	49	F	Headache, dizziness, blurred vision, and nausea	Corticotrophic PA	Craniopharyngioma	Endoscopic transsphenoidal excision (primary tumor) Bifrontal inter-hemispheric	Postoperative CSF leak

Table 1 (Continued)

Study	Patient age (y)	Sex	Clinical presentation	Tumor type 1	Tumor type 2	Surgical approach	Additional comment
						transcallosal approach (residual tumor)	
de Vries et al ²³	75	F	Depression, fatigue, unintended weight loss	Nonfunctioning PA	Meningothelial meningioma	Extended endoscopic transplanum excision	Improvement of symptoms
Bteich et al ²⁴	35	M	Headache, progressive visual disturbance	Nonfunctioning PA	Papillary craniopharyngioma	Endoscopic transsphenoidal excision	N/A
de Almeida Verdolin et al ^{25,a}	Median = 60 (33–78)	3 F 2 M	Progressive visual field loss and/or headache	PA	Rathke's cleft cyst	Endoscopic transsphenoidal excision	N/A
Gezer et al ²⁶	34	F	Menstrual irregularities, proximal muscle weakness, and rapid weight gain	Corticotrophic PA	Meningioma	Endoscopic transsphenoidal excision	Postoperative resolution of weight gain, menstrual irregularities, and proximal muscle weakness
Shareef et al ²⁷	60	M	Prior history of PA with resection, nonremitting bitemporal visual deficit	PA	Adamantinomatous craniopharyngioma	Endoscopic transsphenoidal excision	Transient postoperative diabetes insipidus
Bao et al ^{28,a}	62 56	F F	Progressive visual loss in left eye Headache with progressive bilateral visual loss	Nonfunctioning PA Nonfunctioning PA	Meningioma Meningioma	Endoscopic transsphenoidal excision Transmaxillary-transpterygoid approach	Improvement in visual acuity postoperatively
Ren et al ²⁹	41	M	Intermittent headache	Lactotroph PA	DLBCL	Endoscopic transsphenoidal excision (PA) Chemotherapy (DLBCL)	No tumor recurrence
Schöning et al ^{1,a}	Mean = 53.8 ± 18.5	N/A	N/A	Double PitNET (n = 38) Triple PitNET (n = 2) PitNET (n = 34) PitNET (n = 6) PitNET (n = 5) PitNET (n = 12)	Gangliocytoma Meningioma Posterior lobe tumor Metastasis	N/A	N/A
Lu et al ³⁰	61	F	Progressive decline of binocular vision	PA	Meningioma	Endoscopic transsphenoidal excision	Stable visual acuity without tumor recurrence

Abbreviations: ACTH, adrenocorticotropic hormone; DLBCL, diffuse large B-cell lymphoma; F, female; FSH, follicle stimulating hormone; GH, growth hormone; M, male; PA, pituitary adenoma; PitNET, pituitary neuroendocrine tumor.

^aCase series

Table 2 Literature review of case reports on clear cell meningiomas without dural attachment

Study	Patient age (y)	Sex	Clinical presentation	Tumor location	Surgical approach	Additional comment
Miranda et al ³¹	10	F	Neck pain, progressive right-sided hemiparesis, gait instability, somnolence, bilateral nystagmus, voice changes, lower cranial nerve dysfunction, left tongue/uvula deviation, (+) Hoffman's sign, and extensor plantar reflex	Craniocervical junction	Posterior fossa craniotomy, C1–C2 laminectomy with gross total resection	Complete recovery without signs of residual tumor
Ko et al ³²	34	F	Leg/hip pain with voiding difficulties	Cauda equina (L2–L3)	L1–L3 laminectomy with gross total resection	No evidence of tumor recurrence at 2 y
Yin et al ³³	55	M	Intermittent pulsatile headache, right eye vision loss + diplopia	Sella	Transsphenoidal approach with gross total resection	Local tumor recurrence at 4-mo follow-up requiring stereotactic radiosurgery
Gupta et al ³⁴	19	M	Back pain radiating to bilateral leg with diminished knee and ankle reflexes bilaterally	Lumbosacral spine (L5–S2)	L4–S2 laminectomy with gross total resection	Complete recovery without signs of residual tumor
Tsuchiya et al ³⁵	10	M	Worsening low back and bilateral lower extremity pain	Lumbar spine (L3)	L3 laminectomy with gross total resection	Complete recovery without signs of residual tumor
Zhang et al ³⁶	45	F	Episodic low back and bilateral upper leg pain	Lumbar spine (L3)	L3 laminectomy with gross total resection	Complete recovery without signs of residual tumor
Maamri et al ³⁷	58	F	Low back pain with bilateral sciatica	Lumbar spine (L3)	L2–L3 laminectomy with gross total resection	Complete recovery

pathologist, as was the case for our patient. While a diagnosis of a PitNET in the context of a sellar/suprasellar collision tumor in itself is relatively unsurprising, a concomitant diagnosis of clear cell meningioma represents a unique entity. Clear cell meningiomas have a proclivity for the cerebellopontine angle and spine, particularly in the cauda equina region, and tends to affect children and young adults, neither of which were the case in our patient (47-year-old, sellar region).⁴⁰ Clear cell meningiomas are unique in that both germline and somatic *SMARCE1* mutations are present, unlike other meningioma types. While the histopathologic diagnosis can be supported by a loss of nuclear *SMARCE1* expression by IHC, it is not required in the cases with archetypal histological features such as dense bands of collagen, glycogen-rich cytoplasm, and clear cell morphology, as in our case.⁴¹ Importantly, clear cell meningiomas tend to exhibit more aggressive behavior, with higher rates of recurrence and occasional seeding of cerebrospinal fluid in comparison to other meningioma subtypes. As such, clear cell meningiomas have been designated a CNS WHO grade 2, thus warranting careful surveillance of our patient.

Few reports have described the diagnosis of a parasellar clear cell meningioma, all of which occurred as a single tumor.^{33,42–46}

Although more common in young adults and children, sellar region clear cell meningioma was diagnosed in 11- to 79-year-old patients.^{33,42–46} Additionally, in all but one case, a dural attachment was seen,^{42–46} while a single case described the occurrence of an intrasellar clear cell meningioma without a dural attachment.³³ Therefore, given the rarity of clear cell meningioma arising as a single-region tumor, this further emphasizes the diagnostic conundrum seen in our case where a gonadotropic adenoma was simultaneously diagnosed with a suprasellar clear cell meningioma without the classic dural attachment.

The association between pituitary adenomas and intracranial meningiomas has been widely described and are thought to arise as a delayed complication following radiotherapy for pituitary lesions.⁴⁷ However, collision tumors composed of simultaneously occurring pituitary adenoma and meningioma are difficult to explain, as the underlying etiology is not understood. One theory suggests that in patients with a growth hormone (GH) secreting macroadenoma, GH excess can induce meningioma growth resulting in collision tumor formation, although this has never been confirmed.^{48,49} Nonetheless, the co-occurrence of a pituitary adenoma and parasellar meningioma is likely an incidental finding of two common lesions within the sellar region.

Conclusion

We describe the diagnosis and treatment of a collision tumor composed of a pituitary adenoma and sellar region meningioma in a 47-year-old-man. Collision tumors arising in the sellar/suprasellar region of the brain are exceedingly rare entities, currently with unclear etiologies. In the absence of reliable radiographic measures to diagnose collision tumors using neuroimaging, histological evaluation remains the gold standard. A multidisciplinary approach between neurosurgeons and neuropathologists is critical for the management of these patients.

Informed Consent

Informed consent was deemed unnecessary for this work by the Colorado Multiple Institutional Review Board.

Conflict of Interest

None declared.

References

- Schöning JV, Flitsch J, Lüdecke DK, et al. Multiple tumorous lesions of the pituitary gland. *Hormones (Athens)* 2022;21(04):653–663
- Yamada K, Hatayama T, Ohta M, Sakoda K, Uozumi T. Coincidental pituitary adenoma and parasellar meningioma: case report. *Neurosurgery* 1986;19(02):267–270
- Tajika Y, Kubo O, Takeshita M, Tajika T, Shimizu T, Kitamura K. An intracranial collision tumor composed of intrasellar gangliocytoma and pituitary adenoma. *No Shinkei Geka* 1989;17(12):1181–1186
- Prevedello DM, Thomas A, Gardner P, Snyderman CH, Carrau RL, Kassam AB. Endoscopic endonasal resection of a synchronous pituitary adenoma and a tuberculum sellae meningioma: technical case report. *Neurosurgery* 2007;60(4, Suppl 2):E401, discussion E401
- Karavitaki N, Scheithauer BW, Watt J, et al. Collision lesions of the sella: co-existence of craniopharyngioma with gonadotroph adenoma and of Rathke's cleft cyst with corticotroph adenoma. *Pituitary* 2008;11(03):317–323
- Moshkin O, Scheithauer BW, Syro LV, Velasquez A, Horvath E, Kovacs K. Collision tumors of the sella: craniopharyngioma and silent pituitary adenoma subtype 3: case report. *Endocr Pathol* 2009;20(01):50–55
- Koutourousiou M, Kontogeorgos G, Wesseling P, Grotenhuis AJ, Seretis A. Collision sellar lesions: experience with eight cases and review of the literature. *Pituitary* 2010;13(01):8–17
- Rivera J, Alves S, Bianchi CC, Al-Mutawa N, Guiot MC, Zeitouni A. An unusual collision tumor comprising a prolactinoma and a plasmocytoma originating from the sellar region. *Pituitary* 2010;13(02):189–193
- Sahli R, Christ E, Kuhlen D, Giger O, Vajtai I. Sellar collision tumor involving pituitary gonadotroph adenoma and chondroma: a potential clinical diagnosis. *Pituitary* 2011;14(04):405–408
- Jin G, Hao S, Xie J, Mi R, Liu F. Collision tumors of the sella: coexistence of pituitary adenoma and craniopharyngioma in the sellar region. *World J Surg Oncol* 2013;11:178
- Mahvash M, Igressa A, Pechlivanis I, Weber F, Charalampaki P. Endoscopic endonasal transsphenoidal approach for resection of a coexistent pituitary macroadenoma and a tuberculum sellae meningioma. *Asian J Neurosurg* 2014;9(04):236
- Karsy M, Sonnen J, Couldwell WT. Coincident pituitary adenoma and sellar meningioma. *Acta Neurochir (Wien)* 2015;157(02):231–233
- Matyja E, Maksymowicz M, Grajkowska W, et al. Ganglion cell tumours in the sella turcica in close morphological connection with pituitary adenomas. *Folia Neuropathol* 2015;53(03):203–218
- Lim KZ, Goldschlager T, Chandra RV, Hall J, Uren B, Pullar M. Co-occurrence of pituitary adenoma with suprasellar and olfactory groove meningiomas. *Basic Clin Neurosci* 2016;7(04):361–365
- Ban VS, Chaudhary BR, Allinson K, Santarius T, Kirillos RW. Concomitant primary CNS lymphoma and FSH-pituitary adenoma arising within the sella. entirely coincidental? *Neurosurgery* 2017;80(01):E170–E175
- Heng LJ, Jia D, Gong L, Zhang W, Ma J, Qu Y. Endoscopic endonasal resection of a mixed lesion of gangliocytoma and nonfunctioning pituitary adenoma. *World Neurosurg* 2017;106:1050.e1–1050.e6
- Zhao Y, Zhang H, Lian W, et al. Collision tumors composed of meningioma and growth hormone-secreting pituitary adenoma in the sellar region: case reports and a literature review. *Medicine (Baltimore)* 2017;96(50):e9139
- Amirjamshidi A, Mortazavi SA, Shirani M, Saeedinia S, Hanif H. Coexisting pituitary adenoma and suprasellar meningioma—a coincidence or causation effect: report of two cases and review of the literature. *J Surg Case Rep* 2017;2017(05):rjx039
- Levitus CF, Charitou MMAN. An incidental collision tumor of the sella turcica. *AACE Clin Case Rep* 2019;5(04):e247–e249
- Malli A, Melissaris S, Dimitriadi A, Choreftaki T, Georgakoulias N. A coexisting pilocytic astrocytoma and a prolactinoma: a case report of collision tumors and literature review. *Cureus* 2019;11(06):e4911
- Miyazaki T, Kowari K, Eda H, Kambara M, Maruyama R, Akiyama Y. Ten-year follow-up of collision tumors composed of craniopharyngioma and pituitary adenoma: a case report and literature review. *Case Rep Med* 2019;2019:8080163
- Snyder R, Fayed I, Dowlati E, Seager A, Mason RB. Pituitary adenoma and craniopharyngioma collision tumor: diagnostic, treatment considerations, and review of the literature. *World Neurosurg* 2019;121:211–216
- de Vries F, Lobatto DJ, Zamanipoor Najafabadi AH, et al. Unexpected concomitant pituitary adenoma and suprasellar meningioma: a case report and review of the literature. *Br J Neurosurg* 2023;37(04):677–681
- Bteich F, El Khoury L, Nohra G, Trak V, Yazbek S, Akiki M. Pituitary adenoma and papillary craniopharyngioma: a rare case of collision tumor and review of the literature. *World Neurosurg* 2020;139:63–69
- de Almeida Verdolin A, Lamback EB, Ventura N, et al. Collision sellar lesions: coexistence of pituitary adenoma and Rathke cleft cyst—a single-center experience. *Endocrine* 2020;68(01):174–181
- Gezer E, Cantürk Z, Selek A, et al. Cushing's disease due to a pituitary adenoma as a component of collision tumor: a case report and review of the literature. *J Med Case Rep* 2020;14(01):59
- Shareef Z, Kerndt C, Nessel T, Mistry D, Figueroa B. Collision tumor in the pituitary, concurrent pituitary adenoma, and craniopharyngioma. *Case Rep Otolaryngol* 2020;2020:9584090
- Bao YY, Wu X, Ding H, Hong T. Endoscopic endonasal resection of coexisting pituitary adenoma and meningioma: two cases' report and literature review. *Neurochirurgie* 2021;67(06):611–617
- Ren S, Lu Q, Xiao Y, et al. Coexistence of pituitary adenoma and primary pituitary lymphoma: a case report and review of the literature. *Front Surg* 2022;9:842830
- Lu W, Shengkai Y, Yu W, Aimin L, Shiwei Y, Kang X. Case report: clinical report of co-occurrence of pituitary adenoma and meningioma in the sellar region after meningioma treatment. *Front Neurol* 2022;13:1042106
- Miranda P, Simal JA, Vila M, Hernández M, Menor F, Alvarez-Garijo JA. Posterior fossa clear cell meningioma without dural attachment in a child. *Childs Nerv Syst* 2009;25(03):389–392

- 32 Ko JK, Choi BK, Cho WH, Choi CH. Non-dura based intraspinal clear cell meningioma. *J Korean Neurosurg Soc* 2011;49(01):71–74
- 33 Yin S, Zhou P, Li Q, Jiang S. Intracellular clear cell meningioma mimicking invasive pituitary adenoma: a case report and review of the literature. *Turk Neurosurg* 2015;25(06):976–979
- 34 Gupta SK, Nayak N, Gandhoke CS, et al. A giant nondural-based lumbosacral clear cell meningioma mimicking schwannoma: a case report and review of the literature. *Asian J Neurosurg* 2021;16(01):44–50
- 35 Tsuchiya T, Ikeda S, Isoo A, et al. Intraoperative anatomical findings in pediatric clear cell meningioma of the lumbar spine: case report and literature review. *NMC Case Rep J* 2021;8(01):519–527
- 36 Zhang X, Zhang P, Wang JJ, et al. Intraspinal clear cell meningioma without dural attachment: a case report and literature review. *Medicine (Baltimore)* 2021;100(11):e25167
- 37 Maamri K, Hadj Taieb MA, Trifa A, Elkahla G, Njima M, Darmoul M. Spinal clear cell meningioma without dural attachment: a case report and literature review. *Radiol Case Rep* 2022;17(05):1760–1764
- 38 Vasquez CA, Downes A, Kleinschmidt-DeMasters BK, Youssef AS. Functioning pituitary adenoma with xanthogranulomatous features: review of literature and case report. *J Neurol Surg B Skull Base* 2019;80(05):449–457
- 39 Spallone A. Meningioma as a sequel of radiotherapy for pituitary adenoma. *Neurochirurgia (Stuttg)* 1982;25(02):68–72
- 40 Zhang H, Ma L, Shu C, Dong LQ, Ma YQ, Zhou Y. Spinal clear cell meningiomas: clinical features and factors predicting recurrence. *World Neurosurg* 2020;134:e1062–e1076
- 41 Gerkes EH, Fock JM, den Dunnen WF, et al. A heritable form of SMARCE1-related meningiomas with important implications for follow-up and family screening. *Neurogenetics* 2016;17(02):83–89
- 42 Baxter DS, Smith P, Stewart K, Murphy M. Clear cell meningioma presenting as rapidly deteriorating visual field and acuity during pregnancy. *J Clin Neurosci* 2009;16(11):1502–1504
- 43 Chen H, Li XM, Chen YC, et al. Intracranial clear cell meningioma: a clinicopathologic study of 15 cases. *Acta Neurochir (Wien)* 2011;153(09):1769–1780
- 44 Jain D, Sharma MC, Sarkar C, et al. Clear cell meningioma, an uncommon variant of meningioma: a clinicopathologic study of nine cases. *J Neurooncol* 2007;81(03):315–321
- 45 Ohba S, Sasaki H, Kimura T, Ikeda E, Kawase T. Clear cell meningiomas: three case reports with genetic characterization and review of the literature. *Neurosurgery* 2010;67(03):E870–E871, discussion E871
- 46 Prayson RA, Chamberlain WA, Angelov L. Clear cell meningioma: a clinicopathologic study of 18 tumors and examination of the use of CD10, CA9, and RCC antibodies to distinguish between clear cell meningioma and metastatic clear cell renal cell carcinoma. *Appl Immunohistochem Mol Morphol* 2010;18(05):422–428
- 47 Abs R, Parizel PM, Willems PJ, et al. The association of meningioma and pituitary adenoma: report of seven cases and review of the literature. *Eur Neurol* 1993;33(06):416–422
- 48 Furtado SV, Dadlani R, Ghosal N, Mahadevan A, Shankar SK, Hegde AS. Co-existing thyrotropin secreting pituitary adenoma and low grade glioma: clinical considerations and literature review. *J Neurosurg Sci* 2009;53(02):71–75
- 49 Black PM, Carroll R, Glowacka D, Riley K, Dashner K. Platelet-derived growth factor expression and stimulation in human meningiomas. *J Neurosurg* 1994;81(03):388–393