

POSTER PRESENTATION

Open Access

Recombinant human proinsulin expression in *Pichia pastoris* using PGK promoter

Enedina Nogueira de Assunção^{1*}, Santos Maria Cristina dos², Torres Fernando Araripe³, Astolfi-Filho Spartaco¹

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

There are indications, according to WHO, that diabetes will reach 333 million people in 2025, and already is among one of the main death causes in the world. In 2030 in Latin America, diabetes mellitus will be the second cause of deaths [1]. The actions to increase the production of insulin are strategic to respond the high demands for what we have to face in the near future [2]. *P. pastoris* methylotrophic yeast is being very used in the last decades to express heterologous genes to biotech uses [3]. The main expression vectors to *Pichia pastoris* are based in the strong promoter from *AOX1* gene which codifies alcohol oxidize enzyme. However there is a search for new promoters than *AOX1* due to certain disadvantages as the use of methanol as inducer that is toxic and originates undesirable subproducts during its metabolism [4]. This work has the main objective in to express in high levels the human proinsulin in *Pichia pastoris* using PGK promoter.

Methods

The human proinsulin codifying sequence was chemically synthesized (Genone inc.) with *P. pastoris* optimized codons. The synthesized sequence was subcloned under PGK promoter control in expression vector together with the codifying region from *Saccharomyces cerevisiae* factor- α signal peptide in order to have proinsulin expression/secretion. After linearization the vector containing the expression cassette was introduced in *P. pastoris* GS115 by electroporation. To select clones, with multicopies of expression cassette, was realized a zeocin resistance test in plate [5]. Resistant clones to 100 µg/mL of zeocin were selected and submitted to crescent concentrations of the antibiotic: 500, 1000 and 2000 µg/mL. The recombinant clones that produced/secreted more proinsulin was

detected by imunodot procedure using antibody anti-insulin/proinsulin (Pierce-D3E7/5B6/6). Selected recombinant clones were cultivated in YPD medium for 24 hours in 30°C at 200rpm and the supernatants were clarified by centrifugation and analyzed by ELISA and Western-blotting.

Results and conclusion

From 200 zeocin (100 µg/mL) resistant *P. pastoris* clones 25 were able to grow in 2000 µg/mL which indicates that they have integrated several expression/secretion cassettes in their genomes. From those one was notable for its higher human proinsulin expression/secretion in both experiments (imunodot and ELISA). The proinsulin over-expression of this selected clone was confirmed by PAGE and Western-blotting with anti-insulin/anti-proinsulin antibody. Additional experiments are in process in order to standardize the fermentative process for expression and secretion of human proinsulin as well as confirm recombinant protein amino acid sequence.

Acknowledgements

This work was supported by CNPq, CAPES, FAPEAM and FAPDF.

Authors' details

¹Divisão de Biotecnologia, Centro de Apoio Multidisciplinar Universidade

Federal do Amazonas, UFAM Manaus, AM, Brazil. ²Instituto de Ciências

Biológicas, Universidade Federal do Amazonas, UFAM Manaus, AM, Brazil.

³Departamento de Biologia Celular, Universidade de Brasília, DF, Brazil.

Published: 1 October 2014

References

- WHO/NCD/NCS/99.2: Definition, diagnosis and classification of diabetes mellitus and its complications. World Health Organization 1999 [http://whqlibdoc.who.int/hq/1999/WHO_NCD_NCS_99.2.pdf], Accessed on 02.09.2012.
- Brand-Miller Jennie: A Nova Revolução da Glicose. Rio de Janeiro: Campus/Elsevier 2003.

¹Divisão de Biotecnologia, Centro de Apoio Multidisciplinar Universidade Federal do Amazonas, UFAM Manaus, AM, Brazil
Full list of author information is available at the end of the article

3. Fazenda S, McNeil ML, Harvey B, L M: Heterologous protein production using the *Pichia pastoris* expression system. *Yeast* 2005, **22**:249-270.
4. Almeida JRM, Moraes LMP, Torres FAG: Molecular characterization of the phosphoglycerate kinase (PGK1) from the methylotrophic yeast *Pichia pastoris*. *Yeast* 2005, **22**:725-737.
5. Vassileva A, Chugh DA, Swaminathan S, Khanna N: Expression of hepatitis B surface antigen in the methylotrophic yeast *Pichia pastoris* using the GAP promoter. *J Biotechnol* 2001, **88**:21-35, doi: 10.1016/S0168-1656(01)00254.

doi:10.1186/1753-6561-8-S4-P88

Cite this article as: de Assunção *et al.*: Recombinant human proinsulin expression in *Pichiapastoris* using PGK promoter. *BMC Proceedings* 2014 **8**(Suppl 4):P88.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

