


# Epidemiology of C2 fractures and determinants of surgical management: analysis of a national registry

Kristin Salottolo,<sup>1</sup> Alejandro Betancourt,<sup>2</sup> Kaysie L Banton,<sup>3</sup> David Acuna,<sup>4</sup> Ripun Panchal,<sup>5,6</sup> David Bar-Or ,<sup>1</sup> Carlos H Palacio<sup>2</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tsaco-2023-001094>).

<sup>1</sup>Trauma Research, Swedish Medical Center, Englewood, Colorado, USA

<sup>2</sup>Trauma Services, South Texas Health System McAllen, McAllen, TX, USA

<sup>3</sup>Trauma Services, Swedish Medical Center, Englewood, CO, USA

<sup>4</sup>Trauma Services, Wesley Medical Center, Wichita, Kansas, USA

<sup>5</sup>Neurosurgery, American Neurospine Institute, PLLC, Plano, Texas, USA

<sup>6</sup>Neurosurgery, Medical City Plano, Plano, Texas, USA

## Correspondence to

Dr David Bar-Or; [davidbme49@gmail.com](mailto:davidbme49@gmail.com)

Received 24 January 2023

Accepted 26 May 2023

## ABSTRACT

**Objective** Operative management of axis fractures (C2) usually depend on the stability and location of the break and individual patient characteristics. We sought to describe the epidemiology of C2 fractures and hypothesized that determinants for surgery would differ by fracture diagnosis.

**Methods** Patients with C2 fractures were identified from the US National Trauma Data Bank from January 1, 2017, to January 1, 2020. Patients were classified by C2 fracture diagnosis: odontoid type II, odontoid types I and III, and non-odontoid fracture (hangman's fracture or fractures through base of the axis). The primary comparison was C2 fracture surgery versus non-operative management. Multivariate logistic regression was used to identify independent associations with surgery. Decision tree-based models were developed to identify determinants for surgery.

**Results** There were 38 080 patients; 42.7% had an odontoid type II fracture; 16.5% had an odontoid type I/III fracture; and 40.8% had a non-odontoid fracture. All examined patient demographics, clinical characteristics, outcomes, and interventions differed by C2 fracture diagnosis. Overall, 5292 (13.9%) were surgically managed (17.5% odontoid type II, 11.0% odontoid type I/III, and 11.2% non-odontoid;  $p < 0.001$ ). The following covariates increased odds of surgery for all three fracture diagnoses: younger age, treatment at a level I trauma center, fracture displacement, cervical ligament sprain, and cervical subluxation. Determinants of surgery differed by fracture diagnosis: for odontoid type II, age  $\leq 80$  years, a displaced fracture, and cervical ligament sprain were determinants; for odontoid type I/III, age  $\leq 85$  years, a displaced fracture, and cervical subluxation were determinants; for non-odontoid fractures, cervical subluxation and cervical ligament sprain were the strongest determinants for surgery, by hierarchy.

**Conclusions** This is the largest published study of C2 fractures and current surgical management in the USA. Odontoid fractures, regardless of type, had age and fracture displacement as the strongest determinants for surgical management, whereas associated injuries were determinants of surgery for non-odontoid fractures.

**Level of evidence** III.

## BACKGROUND

Fractures of the second vertebrae (C2), or axis, are the most common cervical spine injuries and typically result from motor vehicle collisions or falls that cause the neck and head to snap forward or backward or twist suddenly. These axis fractures are usually characterized into odontoid fractures and

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Controversy exists in the management of fractures to the second vertebrae (C2 fractures), with wide variation in practice in part due to the location of the break and the frequency of these injuries among the elderly.

## WHAT THIS STUDY ADDS

⇒ There were significant differences in demographics, injury patterns, outcomes, and surgical management based on C2 fracture diagnosis. Decision tree modeling identified similar determinants of surgery for type I, II, and III odontoid fractures being age and fracture displacement, whereas determinants of surgery for non-odontoid fractures were associated injuries of cervical subluxation and cervical ligament sprain.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY


⇒ This is the largest published descriptive study of C2 fractures and surgical management in the USA to date. The findings from our models inform current management practices in the USA and may be used for additional study of outcomes based on determinants of surgery.

what is colloquially known as hangman's fractures. Odontoid fractures are further classified as type I (fractures through the tip), type II (fractures through the base), and type III (fractures through the body of the C2 vertebral body).<sup>1</sup> The overall incidence of C2 fractures has increased annually in the USA,<sup>2</sup> as well as in other countries.<sup>3</sup> This increase is driven largely by the aging population who are at increased risk of odontoid fractures specifically.<sup>4,5</sup>

Controversy exists in the management of C2 fractures. Historically, the overall gestalt of fracture management is that hangman's fractures and type I and III odontoid fractures are treated non-operatively with cervical orthoses, whereas there is wide variation in practice for the management of odontoid type II fractures<sup>5,6</sup> due to the frequency of these injuries among the elderly.<sup>7,8</sup> The degree to which the fracture is displaced and the age and frailty of the patient both play a role in whether surgical or non-operative management is the preferred approach. However, recent treatment guidelines are generally based on a review of the literature rather than taking a statistical approach for prediction and classification.<sup>5,9-11</sup>

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Salottolo K, Betancourt A, Banton KL, et al. *Trauma Surg Acute Care Open* 2023;**8**:e001094.



**Figure 1** Patient disposition. ED, emergency department; ICD-10, International Classification of Diseases, 10th Revision.

Our study objectives are twofold: (1) to provide a detailed epidemiological evaluation of C2 fractures using a large national registry and (2) to analyze associations with operative intervention and model determinants of surgery using decision tree analysis. We hypothesized that determinants of surgery would differ by C2 fracture diagnosis.

## METHODS

This national registry analysis used the American National Trauma Data Bank (NTDB), which is the largest aggregation of admissions due to traumatic injury.<sup>12</sup> Patients admitted for a C2 fracture from January 1, 2017, to January 1, 2020, were included (figure 1). Exclusions were patients who died in the ED (0.1% of C2 fracture admissions), subsequent, non-index admissions (0.7%), and penetrating injuries (1.9%).

Study groups were determined based on C2 fracture diagnosis using the International Classification of Diseases, 10th Revision, diagnosis code (figure 2): odontoid type II fracture (S12.11), odontoid type I or III fracture (S12.12), and other fractures (S12.19, used primarily for hangman's fracture but also includes fractures through the base of the axis; this group is defined as non-odontoid fractures in the table and text). Patients with other C2 diagnoses were excluded: traumatic spondylolisthesis without fracture (S12.13–S12.15) and unspecified fractures (S12.1 and S12.10; these fractures did not have sufficient information in the medical record to determine if it was an odontoid I, II, III, or non-odontoid fracture).


The dependent variable of interest was surgical intervention of the C2 fracture, identified using ICD-10 procedure codes for cervical fusion and internal fixation. ICD-10 diagnosis codes and procedure codes used throughout this study can be found in online supplemental table 1.

There were 26 covariates of interest, categorized as follows:

- Demographics: age in years (median and IQR; patients >89 years were coded as 90 years old); sex (male and female); race (non-Hispanic (NH) white, NH black, NH other, Hispanic, and unspecified); individual comorbidities with >10% incidence.
- Hospital characteristics: transferred in from another facility for definitive care (yes or no) and trauma level (level I or II or III/IV or unverified/not applicable), based on American College of Surgeons (ACS) verification; for hospitals

unverified by the ACS but state designated I or II, the state designation was used.

- Injury characteristics, in addition to fracture type: fracture displacement (displaced or non-displaced, based on ICD-10 diagnosis codes); cause of injury (ground-level fall, fall from height, motor vehicle crash or motorcycle crash (MVC), all other causes); abnormal emergency department (ED) vital signs of systolic blood pressure (SBP <90 mm Hg), heart rate (HR <60 or >120 beats/min), and respiratory rate (RR



**Figure 2** C2 fracture diagnoses. (A). Odontoid type II fracture (ICD-10 diagnosis: S12.11). (B) Odontoid type I/III fracture (ICD-10 diagnosis: S12.12). (C) Other, non-odontoid fracture (hangman's fracture and C2 base fracture, ICD-10 diagnosis: S12.19). ICD-10, International Classification of Diseases, 10th Revision.

<12 or >20 breaths/min); ED Glasgow Coma Scale (GCS) motor component (median and IQR, scores range from 1 (worst) to 6 (best); severe concomitant injury to the head/neck, face, thorax, abdomen/pelvis, or extremities (defined with the Abbreviated Injury Scale region and score  $\geq 3$ ); and associated injury diagnoses for cervical ligament sprain, vertebral subluxation and dislocation, and intracranial injury (defined by ICD-10 diagnosis codes).

In-hospital outcomes of mortality (%), intensive care unit (ICU) admission, hospital length of stay (LOS), ICU LOS, and development of a complication in-hospital (yes or no) are also reported.

### Statistical analysis

Analyses were performed with SAS V.9.4. A conservative alpha of <0.01 was used for statistical significance based on the large sample size.  $\chi^2$  tests (categorical variables, reported using %, n) and Wilcoxon rank-sum tests (continuous variables, reported using medians and IQR) were used to examine differences between study covariates and fracture diagnosis, as well as operative management. Collinearity was assessed with Spearman's rank correlation; values  $\geq 0.4$  had moderate-to-strong correlation. Intracranial injury and pre-existing hypertension were collinear with severe head/neck injury and age, respectively, and were not used in the models, but all remaining covariates were analyzed. For both regression models and decision trees, model fit was assessed with c-statistics.

Multivariate logistic regression with backward selection and exit criteria of p value of <0.01 determined independent associations with operative management for each fracture diagnosis. ORs and the associated 99% CIs were calculated from the regression models.

For decision trees, the high-performance split procedure (PROC HPSPLIT) was used to build a tree-based statistical model for classifying patients into operative versus non-operative management for each fracture diagnosis. Candidate splits for each node were based on criteria for impurity (entropy), which uses an entropy score from 0 to 1 to determine how to grow the tree based on how similar or impure the data were within each 'branch'. Cost complexity pruning was used to ensure the complexity (number of branches) was reduced to prevent overfitting.

### RESULTS

There were 38 080 patients; 42.7% had an odontoid type II fracture; 16.5% had an odontoid type I/III fracture; and 40.8% had a non-odontoid fracture (figure 1). Among the odontoid fractures, the majority (72.1%) were type II fractures. The population median age was 74 (58 to 85) years; ground-level falls and MVCs were the most common causes of injury (31.9% and 31.4%, respectively); patients frequently presented with severe injuries besides the C2 fracture, including head/neck injuries (29.7%), extremity injuries (25.2%), and thoracic injuries (22.7%), and nearly half of the patients (45.6%) were treated at a level I trauma center.

As shown in table 1, there were statistically significant differences in all demographics, comorbidities, and presenting clinical characteristics by fracture diagnosis ( $p < 0.001$ ), with the exception of similar rates of mental/personality disorder comorbidities ( $p = 0.63$ ). For example, compared with odontoid type I/III and non-odontoid fractures, patients with odontoid type II fractures were older (median age: 80 years vs. 75 years and 65 years, respectively), were injured in a ground-level fall (44.8% vs. 31.0% and 18.6%), and were more likely to present with

**Table 1** Characteristics of patients with C2 fracture by fracture diagnosis

Characteristic, % or median (IQR)	Odontoid type II (n=16 265)	Odontoid types I and III (n=6288)	Non-odontoid* (n=15 527)	P value
Age (years), median (IQR)	80 (69–88)	75 (59–85)	65 (47–79)	<0.001
Female sex	52.0	53.6	47.7	<0.001
Race/ethnicity				<0.001
NH white	83.7	80.7	74.9	
NH black	4.8	6.0	9.1	
Hispanic	4.1	5.1	7.7	
NH other	2.7	3.7	4.7	
Unspecified	4.8	4.6	4.1	
Comorbidities†				
Anticoagulant use preinjury	18.5	14.7	11.3	<0.001
Dementia	16.0	11.8	7.1	<0.001
Diabetes	19.3	17.1	16.0	<0.001
Functional dependence	19.8	15.4	9.9	<0.001
Hypertension	61.4	52.8	44.2	<0.001
Mental/personality disorder	10.9	11.1	10.6	0.63
Smoker	10.2	14.4	17.9	<0.001
Interfacility transfer	48.3	42.4	36.7	<0.001
Trauma level				<0.001
Level I verified/designated	54.4	58.1	58.2	
Level II verified/designated	35.8	32.7	33.9	
Level III/IV verification	5.5	4.9	4.1	
Unverified by ACS	4.2	4.3	3.7	
Cause of injury				<0.001
MVC	18.0	32.5	45.0	
Fall from height	31.4	28.3	24.8	
Ground-level fall	44.8	31.0	18.6	
Other cause	5.7	8.1	11.6	
GCS motor score, median (IQR)	6 (6–6)	6 (6–6)	6 (6–6)	<0.001
Abnormal ED vital signs				
SBP <90 mm Hg	2.3	3.2	3.8	<0.001
HR <60 or >120	8.9	9.0	10.0	0.004
RR <12 or >20	16.1%	19.0%	21.8%	<0.001
Severe concomitant injury, AIS score $\geq 3$				
Head/neck injury	21.8	30.0	37.7	<0.001
Facial injury	8.2	7.3	9.0	<0.001
Thoracic injury	13.9	23.0	21.7	<0.001
Abdomen/pelvic injury	3.4	6.0	8.7	<0.001
Extremity injury	17.9	26.6	32.4	<0.001
Other associated injuries				
Displaced C2 fracture	73.7	71.3	67.8	<0.001
Cervical ligament sprain	3.9	5.2	7.4	<0.001
Vertebral subluxation/dislocation	3.2	2.8	3.9	<0.001
Intracranial injury	13.1	17.2	21.5	<0.001

Bold values denotes statistical significance with alpha <0.01.  
 \*Non-odontoid fractures are other C2 fractures (S12.19) including hangman's fracture and fractures through the base of the axis.  
 †Comorbidities with at least 10% incidence are tabulated.  
 ACS, American College of Surgeons; AIS, Abbreviated Injury Scale; ED, emergency department; GCS, Glasgow Coma Scale; HR, heart rate; MVC, motor vehicle crash or motorcycle crash; NH, non-Hispanic; RR, respiratory rate; SBP, systolic blood pressure.

comorbidities but less likely to present with abnormal ED vital signs and concomitant injuries.

There were also significant differences in outcomes by fracture diagnosis (table 2). Compared with patients with odontoid

**Table 2** Procedures and outcomes of patients with C2 fracture

Variable, % or median (IQR)	Odontoid type II (n=16265)	Odontoid types I and III (n=6288)	Non-odontoid* (n=15527)	P value
Operative management	17.5	11.0	11.2	<0.001
Spinal fusion only	36.9	41.9	52.5	
Internal fixation without fusion	32.3	23.5	12.2	
Internal fixation with fusion	30.7	34.6	35.3	
Other procedures				
Cervical joint/disc	1.3	1.3	3.2	<0.001
Cervical spinal cord	0.6	0.6	1.0	<0.001
Neurosurgery	0.7	1.4	2.1	<0.001
Mechanical ventilation	9.2	11.2	13.0	<0.001
Tracheostomy	2.5	3.4	4.5	<0.001
In-hospital outcomes				
Complication	11.8	13.0	13.0	0.002
In-hospital mortality	7.8	7.6	6.5	<0.001
ICU admission	42.7	45.6	51.2	<0.001
ICU LOS	3 (2–6)	3 (2–7)	3 (2–7)	<0.001
Hospital LOS	5 (3–8)	5 (3–9)	5 (3–9)	0.01
Disposition home/home health	37.1	39.1	47.2	<0.001

\*Non-odontoid fractures are other C2 fractures (S12.19) including hangman's fracture and fractures through the base of the axis.  
ICU, intensive care unit; LOS, length of stay.

type II or type I/III fractures, patients with non-odontoid fractures had the lowest mortality rate (7.8%, 7.6%, and 6.5%, respectively;  $p < 0.001$ ) as well as the most favorable rate of hospital discharge disposition home (37.1%, 39.1%, and 47.2%,  $p < 0.001$ ), but the highest ICU admission rate (42.7%, 45.6%, and 51.2%,  $p < 0.001$ ).

### Operative management

Overall, 5292 (13.9%) were operatively managed. The rate of operative management of C2 fractures differed by fracture diagnosis and was 17.5% for patients with odontoid type II fractures, 11.0% for odontoid type I/III fractures, and 11.2% for non-odontoid fractures. Most patients (76%) who were operatively managed had cervical fusion, and 24% had internal fixation without fusion.

As shown in table 3, the operative management group was younger than non-operatively managed patients (median age: 68 years vs. 75 years,  $p < 0.001$ ), less likely to be female (42.5% vs. 51.8%,  $p < 0.001$ ), less likely to be white (77.7% vs. 79.9%,  $p < 0.001$ ), less likely to be injured from a ground-level fall (29.0% vs. 32.3%,  $p < 0.001$ ), more likely to be transferred in (46.3% vs. 42.0%,  $p < 0.001$ ), more likely to be treated at a level I trauma center (61.1% vs. 55.9%,  $p < 0.001$ ), more likely to have a displaced C2 fracture (86.5% vs. 68.4%,  $p < 0.001$ ), more likely to have associated injuries of cervical ligament sprain (13.3% vs. 4.3%,  $p < 0.001$ ) and vertebral subluxation (9.6% vs. 2.4%,  $p < 0.001$ ), and to have concomitant head/neck injuries (33.0% vs. 29.1%,  $p < 0.001$ ) but less likely to have concomitant extremity injuries (23.0% vs. 25.6%,  $p < 0.001$ ). There were no differences for the operative and non-operative management groups in abnormal ED vital signs of GCS, SBP, RR, and HR and concomitant injuries to the face, thorax, abdomen, and pelvis. Compared with the non-operative group, the operative group had lower mortality (3.9% vs. 7.8%) but otherwise had worse outcomes (table 3).

**Table 3** Characteristics of patients with C2 fracture by operative management

Characteristics, n (%) or median (IQR)	Nonoperative (32 788)	Operative (5292)	P value
Age, median (IQR)	75 (59–86)	68 (52–78)	<0.001
Female sex	51.8	42.5	<0.001
Race/ethnicity			<0.001
NH white	79.9	77.7	
NH black	6.5	8.2	
Hispanic	5.4	6.1	
NH other	3.6%	4.1%	
Unspecified	4.6%	3.8%	
Comorbidities*			
Anticoagulant use preinjury	15.6	10.7	<0.001
Dementia	12.6	5.9	<0.001
Diabetes	17.7	16.5	0.03
Functional dependence	15.9	9.9	<0.001
Hypertension	53.7	48.2	<0.001
Mental/personality disorder	10.6	12.1	0.002
Smoker	13.1	20.0	<0.001
Interfacility transfer	42.0	46.3	<0.001
Trauma level			
Level I verification/designation	55.9	61.1	<0.001
Level II verification/designation	34.7	33.3	
Level III/IV verification	5.2	3.1	
Unverified by ACS	4.3	2.6	
Cause of injury			<0.001
MVC	31.2	32.5	
Fall from height	28.3	27.9	
Ground-level fall	32.3	29.0	<0.001
Other cause	8.2	10.7	
GCS motor component score, median (IQR)	6 (6–6)	6 (6–6)	0.03
Abnormal ED vital signs			
SBP <90 mm Hg	3.1	2.6	0.04
HR <60 or >120	9.5	8.8	0.15
RR <12 or >20	18.9	18.9	0.87
Severe concomitant injury, AIS score $\geq 3$			
Head/neck injury	29.1	33.0	<0.001
Facial injury	8.2	9.2	0.02
Thoracic injury	22.7	22.5	0.80
Abdomen/pelvic injury	6.0	6.2	0.65
Extremity injury	25.6	23.0	<0.001
Other associated injuries			
Displaced C2 fracture	68.4	86.5	<0.001
Cervical ligament sprain	4.3	13.3	<0.001
Vertebral subluxation/dislocation	2.4	9.6	<0.001
Intracranial injury	17.4	15.8	0.004
C2 fracture diagnosis			
Odontoid type II	40.9	53.9	<0.001
Odontoid types I and III	17.1	13.1	
Non-odontoid† fracture	42.0	33.0	
In-hospital outcomes			
Complication	11.2	20.7	<0.001
In-hospital mortality	7.8	3.9	<0.001
ICU admission	43.0	68.9	<0.001
Median ICU LOS (days)	3 (2–6)	5 (3–10)	<0.001
Median hospital LOS (days)	5 (3–8)	9 (6–15)	<0.001
Disposition home/home health	42.0	38.7	<0.001

\*Comorbidities with at least 10% incidence are tabulated.

†Non-odontoid fractures are other C2 fractures (S12.19) including hangman's fracture and fractures through the base of the axis.

ACS, American College of Surgeons; AIS, Abbreviated Injury Scale; ED, emergency department; GCS, Glasgow Coma Scale; HR, heart rate; ICU, intensive care unit; LOS, length of stay; MVC, motor vehicle crash or motorcycle crash; NH, non-Hispanic; RR, respiratory rate; SBP, systolic blood pressure.

**Table 4** Multivariate logistic regression analysis of operative management for patients with C2 fracture

Covariate of interest	Odontoid type II (n=16 265) C-statistic: 0.72		Odontoid types I and III (n=6288) C-statistic: 0.71		Non-odontoid* (n=15 527) C-statistic: 0.73	
	AOR (99% CI)	P value	AOR (99% CI)	P value	AOR (99% CI)	P value
Age, 10-year increment	0.82 (0.78 to 0.85)	<0.001	0.85 (0.80 to 0.91)	<0.001	0.90 (0.87 to 0.93)	<0.001
Male vs. female sex	1.21 (1.07 to 1.36)	<0.001	–	–	1.27 (1.09 to 1.47)	<0.001
Fall, height vs. same-level fall	1.05 (0.92 to 1.21)	0.34	0.98 (0.72 to 1.33)	0.85	–	–
MVC vs. same-level fall	0.79 (0.65 to 0.97)	0.003	0.60 (0.42 to 0.84)	<0.001	–	–
Other cause vs. same-level fall	1.14 (0.89 to 1.47)	0.17	0.98 (0.63 to 1.52)	0.89	–	–
GCS motor score, 1-point increment	1.11 (1.05 to 1.18)	<0.001	1.14 (1.03 to 1.26)	<0.001	–	–
SBP ≤90 mm Hg vs. >90 mm Hg	0.62 (0.39 to 0.98)	0.008	–	–	–	–
Pulse >120 vs. ≤120 beats/min	–	–	–	–	0.78 (0.61 to 1.00)	0.01
Thoracic injury (AIS score ≥3) vs. no	0.80 (0.66 to 0.96)	0.002	–	–	–	–
Displaced vs. non-displaced	3.11 (2.63 to 3.68)	<0.001	2.48 (1.82 to 3.38)	<0.001	2.50 (2.07 to 3.00)	<0.001
Cervical ligament sprain vs. no	2.48 (1.95 to 3.17)	<0.001	3.07 (2.14 to 4.41)	<0.001	3.14 (2.58 to 3.83)	<0.001
Cervical subluxation vs. no	2.32 (1.77 to 3.05)	<0.001	3.74 (2.35 to 5.95)	<0.001	4.98 (3.88 to 6.39)	<0.001
Transferred in vs. no	1.19 (1.05 to 1.35)	<0.001	–	–	–	–
Level I TC vs. ACS non-verified	1.70 (1.20 to 2.39)	<0.001	2.29 (1.12 to 4.69)	0.003	2.11 (1.30 to 3.44)	<0.001
Level II TC vs. ACS non-verified	1.85 (1.31 to 2.62)	<0.001	2.05 (0.99 to 4.27)	0.012	1.88 (1.15 to 3.09)	0.001
Level III/IV TC vs. ACS non-verified	1.33 (0.86 to 2.06)	0.09	0.98 (0.36 to 2.67)	0.96	1.07 (0.54 to 2.12)	0.79
Comorbidities						
Anticoagulant vs. no	0.76 (0.64 to 0.90)	<0.001	–	–	0.72 (0.55 to 0.96)	0.003
Dementia vs. no	0.56 (0.45 to 0.69)	<0.001	–	–	0.44 (0.28 to 0.68)	<0.001
Functionally dependent vs. no	0.75 (0.63 to 0.90)	<0.001	0.51 (0.34 to 0.79)	<0.001	–	–
Mental/personality disorder vs. no	1.23 (1.03 to 1.47)	0.003	–	–	–	–
Smoker vs. no	1.33 (1.12 to 1.58)	<0.001	1.63 (1.23 to 2.17)	<0.001	–	–

Bolding denotes adjusted  $p \leq 0.01$ , statistically significant. The following covariates were not significantly associated with surgery in any of the three models after adjustment and are not presented in the table: race/ethnicity, diabetes comorbidity, abnormal respiratory rate, abdominal/pelvic injury (AIS score  $\geq 3$ ), head/neck injury (AIS score  $\geq 3$ ), facial injury (AIS score  $\geq 3$ ), and extremity injury (AIS score  $\geq 3$ ).

\*Non-odontoid fractures are other C2 fractures (S12.19) including hangman's fracture and fractures through the base of the axis.

ACS, American College of Surgeons; AIS, Abbreviated Injury Scale; AOR, adjusted OR; GCS, Glasgow Coma Scale; MVC, motor vehicle or motorcycle collision; SBP, systolic blood pressure; TC, trauma center.

### Multivariate models

The logistic regression models identified covariates that were independently associated with operative management by fracture diagnosis (table 4). For all three fracture diagnoses, the following covariates were independently associated with increased odds of operative management: younger age, admission to a level I trauma center (vs. unverified/not applicable trauma centers), a displaced C2 fracture, and associated injuries of cervical ligament sprain and vertebral subluxation. The following covariates were not significantly associated with surgery in any of the three models, by fracture diagnosis: race/ethnicity, diabetes comorbidity, abnormal RR, and severe injuries to the head/neck, abdomen/pelvis, face, or extremities.

For patients with odontoid type II fractures, the odds of surgery also increased with higher ED GCS motor scores, transfer in for definitive care, for men, and for comorbidities of mental/personality disorder and smoking; whereas there were decreased odds of surgery with severe thoracic injury, ED SBP of  $\leq 90$  mm Hg, MVC cause of injury, and comorbidities of chronic anticoagulant use, dementia, and functional dependence.

For patients with odontoid type I or III fractures, the odds of surgery also increased with higher ED GCS motor scores, for men, and for smokers, whereas there were decreased odds of surgery for patients with MVC cause of injury and patients who were functionally dependent.


For patients with non-odontoid fracture, the odds of surgery also decreased with comorbidities of chronic anticoagulant use

and dementia, for women, and for patients with an abnormal ED pulse of  $> 120$  beats/min.

The decision tree analysis identified different determinants for operative management based on fracture diagnosis. For patients with a type II odontoid fracture, age was the most important determinant of surgery; patients  $> 80$  years old were only estimated to be surgically managed in 9% of cases (figure 3). A displaced odontoid type II fracture and a cervical ligament sprain were the two remaining determinants of surgical management. The estimated surgery rate was 24.9% for patients  $\leq 80$  years old and increased to 29.3% if the fracture was displaced, and was 49.2% if there was also cervical ligament sprain.

The decision tree for odontoid type I/III fractures similarly identified age as the most important determinant for surgery (figure 4). Patients  $> 85$  years were rarely estimated to have surgery ( $< 4\%$ ). A displaced odontoid type II fracture and a cervical subluxation were the two remaining determinants of surgical management. The estimated surgery rate was 13.7% for patients  $\leq 85$  years old, increased slightly to 16.3% if the fracture was displaced, and was 46.7% if there was also cervical subluxation.

The decision tree for non-odontoid fractures had more nodes, but many of these nodes had small patient groups sizes of  $< 5\%$  of the total sample (figure 5). The presence of cervical subluxation was the most important determinant for surgery of non-odontoid fractures, although only 10% of patients had cervical subluxation with an estimated rate of surgery of 13.7%. Among



**Figure 3** Decision tree for operative intervention of odontoid type II fractures. Shading indicates largest (darkest) to smallest (lightest, <5% of total N) patient group size. Misclassification rate=17.1%, area under the receiver operating characteristic (AUROC) curve=0.68.

patients without cervical subluxation (90% of patients), determinants of surgery were cervical ligament sprain and age. The estimated rate of surgery was 8.8% in patients without a cervical ligament sprain and 29.7% for patients with cervical ligament sprain who were  $\leq 80$  years old. A displaced fracture was not a consideration for surgery in this fracture diagnosis.


The regression models had model fit statistics indicating fair fit, with area receiver operating characteristic (AUROC) values of 0.71 to 0.73. The decision tree models had slightly poorer fit with AUROC values of 0.68 for odontoid type II and type I/III fractures and 0.62 for non-odontoid fractures.

## DISCUSSION

This population-based study identified significant differences in demographics, injury patterns, and outcomes based on fracture diagnosis for patients with fractures to the second vertebrae. These variations in injuries and populations are indicative of differences in surgical management of the fracture. Our regression analysis identified between 9 and 16 variables that were independently associated with operative intervention. When considering decision tree modeling, only three covariates were determinants of surgical management for odontoid type II and odontoid I/III fractures, and these determinants were similar, by hierarchy: patient age, whether the fracture was displaced, and associated cervical injuries of either cervical subluxation or ligament sprain. The decision tree for non-odontoid fractures had more determinants for

surgery, but most patients did not have cervical subluxation and were differentiated by presence of cervical ligament sprain and age. Thus, the overarching determinants for surgery of C2 fractures were the patient's age and the cervical injuries themselves, and not more general injury characteristics such as the cause of injury, presenting vital signs, and severe concomitant injuries (polytrauma), nor patient demographics like sex, race, and comorbidities, nor hospital characteristics like trauma level and transfer status.

This large registry analysis illustrates the epidemiology of C2 fractures in the USA. Radovanovic *et al* previously described patterns of C2 fracture in geriatric patients treated in London, England.<sup>13</sup> Odontoid type II fractures were the majority of fractures (57%), which is higher than the 43% we reported, possibly due to their studies' older age inclusion criteria. Also contrary to our findings, they reported few differences between fracture types with respect to cause of injury, demographics, and outcomes. They also reported only 10.6% were injured in an MVC, which is lower than the 31% rate in our population and lower than a prior NTDB analysis of octogenarians where 17% were injured by MVC. Using the Swedish national registry, Robinson *et al* report that in geriatric patients, approximately 63% of C2 fractures are odontoid type II and 26% are odontoid type III, and in younger patients, approximately 34% are odontoid type II, 17% are odontoid type III, and 24% are hangman's fractures.<sup>3</sup> Our C2 population was 43% odontoid type II fractures, which were within the range of these Swedish studies.



**Figure 4** Decision tree for operative intervention of odontoid type I/III fractures. Shading indicates largest (darkest) to smallest (lightest, <5% of total N) patient group size. Misclassification rate=10.5%, area receiver operating characteristic=0.68.


Surgical rates of C2 fractures vary widely. Regarding age, in Sweden, the surgical rate was 22% among  $\geq 70$  years<sup>14</sup>; in England, it was 27% among  $\geq 65$  years<sup>13</sup>; in New Zealand, it was 11% among  $\geq 70$  years,<sup>15</sup> and in the NTDB, it was previously reported to be 10% for octogenarians.<sup>16</sup> The surgical rate among non-geriatric patients in Sweden was higher, at approximately 35%.<sup>17</sup> In our population, surgical management peaked in patients aged 50–69 years at 18% and was lowest in patients  $\geq 80$  years old (8% surgically managed).

Few studies have compared the surgical rate by fracture diagnosis. In the Swedish registry, the surgical rate was 40% across all C2 fractures but was 53% with odontoid type II injuries.<sup>3</sup> A Norwegian study of 336 patients reported surgical rates of 32% with odontoid type II fractures but only 4% with odontoid type III fractures.<sup>18</sup> The surgical rate in our study was low at 14%, and the disparity with our data compared with other regions may be driven by the lower rate of surgical management for patients with odontoid type II fractures, at only 17.5%.

We used regression analysis and decision tree modeling to determine surgical versus non-operative management, as opposed to other studies that describe preferred treatment guidelines based on a review of the literature. Carvalho *et al* recommend surgical treatment for type III fractures with  $>5$  mm displacement and type II fractures with  $>4$  mm to 6 mm displacement and who are non-geriatric.<sup>9</sup> Nourbakhsh and Hanson prefer conservative management for type I and III fractures regardless of age, and surgical management of type II fractures with  $>4$  mm

displacement.<sup>11</sup> Wagner *et al* take a more general approach and prefer surgical management of geriatric patients with odontoid type II fractures.<sup>5</sup> Rizvi *et al* examined compliance with their recommendation to operate on younger patients with displaced odontoid type II fractures, older patients with type II fractures regardless of displacement, and all displaced type III fractures regardless of age; the non-compliance rate of 36% was largely driven by age, and the authors conclude that age should play a larger consideration in decision trees for treatment choice.<sup>18</sup> Our findings agree with the aforementioned studies, that displacement of the odontoid fracture and the patients' age both play a role in whether surgical or non-operative management is preferred. The most favored age cut-off in our study was  $<80$  years old for odontoid type II fractures and  $<85$  years old for odontoid type I/III fractures, which is older than geriatric age in the aforementioned studies (often described as the 'old old' or octogenarians). Our analysis was more specific than prior studies because it modeled each fracture diagnosis separately and analyzed age continuously.

The gestalt that odontoid type II fractures are surgically managed, and hangman's fractures and odontoid type I/III fractures are non-operatively managed does not appear to be supported by our analysis, as seen in the overall surgical rates were 17.5% and 11.1%, respectively. Still, when comparing the decision trees for odontoid type II and type I/III fractures, the estimated surgical rates were approximately twofold higher for odontoid type II fractures compared with odontoid type I/III



**Figure 5** Decision tree for operative intervention of non-odontoid fractures. Shading indicates largest (darkest) to smallest (lightest, <5% of total N) patient group size. Misclassification rate=10.6%, area receiver operating characteristic=0.62. GCS, Glasgow Coma Scale.

fractures in the old old cohort, 9.1% versus 3.8% surgery, and in the younger cohort, 24.9% versus 13.7% surgery; as well as in the younger cohort who had a displaced fracture, 29.3% versus 16.3% surgery.

One of the studies' biggest limitations is that the models' findings should not be considered the standard of care or to guide optimal treatment decisions. Rather, these models demonstrate current practice patterns in the USA by C2 fracture diagnosis. This study is limited in determining whether patients were mismanaged or inappropriately selected for surgery, in part because there are no long-term (postdischarge) outcomes in the NTDB. Additional study is needed to compare outcomes by surgical management of C2 fractures based on the variables identified in this study (fracture diagnosis, fracture displacement, patients' age, and associated cervical injuries), but also to adjust for variables that would influence outcomes in trauma patients such as abnormal vital signs, severe concomitant injuries, and comorbidities.

Second, the degree of fracture displacement was not available. Other studies recommended surgery based on whether the fracture was displaced >4, 5, or 6 mm. Displacement was a significant determinant for classifying into operative management, and our model fit statistics would likely have been improved had we been able to examine the degree of displacement. However, it should be noted that displacement of C2 fractures can be difficult to assess because it may change with the patient's position and posture and even with respiration.<sup>9</sup> Third, nearly 20% of patients with C2 fracture did not have more detailed diagnoses and were considered 'unspecified', and these patients were excluded from our analysis comparing types of C2 fractures. Fourth, 3% of patients had more than one C2 fracture diagnosis; patients with an odontoid type II fracture were characterized

in that group even when there was also an odontoid type I/III or non-odontoid fracture, and patients with an odontoid type I/III fracture were characterized in that group even when they also had a non-odontoid fracture. We performed a sensitivity analysis to exclude the 3% of patients with more than one C2 fracture diagnosis, and there were no differences in the decision tree models (order and variables of nodes) or the model fit statistics (data not shown). Finally, the NTDB only includes data from contributing trauma centers, and these results might not be generalizable to non-participating hospitals. There were 183 of approximately 198 (93%) level I trauma centers contributing data and 206 of approximately 252 (82%) level II trauma centers contributing data, but far fewer III/IV and non-ACS verified centers contributed data.

## CONCLUSIONS

At nearly 40 000 patients, this is the largest published descriptive study of C2 fractures and current management in the USA to date. There were significant differences in demographics, injury patterns, surgical procedures, and outcomes based on C2 diagnosis. The hierarchy (order) that was identified in the decision trees suggests age is the predominant determinant of surgery for patients with an odontoid fracture, regardless of type, followed by fracture displacement. On the contrary, associated cervical diagnoses of subluxation/dislocation and cervical ligament sprain were the predominant determinants of operative intervention for patients with non-odontoid fractures in US trauma centers. Our findings reflect other authors' suggested criteria for operative intervention of odontoid fractures that they based on review of the literature, but our findings are based on statistical modeling of a large US database and provide more specific



age criterion. The model performance for surgical management was similarly adequate for decision tree modeling and logistic regression modeling. The findings from our analysis may be used for studying outcomes by surgical management, especially when examined by fracture displacement, fracture location, and age. These findings could also be used as a reference to inform current management practices in the USA.

**Contributors** All authors made substantial contributions to the manuscript as follows: KS is responsible for conceptualization, methodology, software, formal analysis, and drafting of the manuscript. AB, KLB, DA, RP, and CHP are responsible for verification and critical revisions of the manuscript. DB-O is responsible for project administration, writing the manuscript, and is responsible for the overall content as guarantor. All authors provided final approval of the submitted manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** National Trauma Database review is not applicable. This study did not require institutional review board approval because the analysis of secondary, deidentified, publicly available data does not constitute research involving human subjects under the federal Common Rule, 45 CFR Part 46.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. The datasets used and/or analyzed during the current study are publicly available. The NTDB remains the full and exclusive copyrighted property of the American College of Surgeons. The American College of Surgeons is not responsible for any claims arising from works based on the original Data, Text, Tables, or Figures. Data are available via <https://www.facs.org/quality-programs/trauma/tqp/center-programs/ntdb/datasets>. The authors did not have any special access privileges to this data.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

David Bar-Or <http://orcid.org/0000-0002-3685-314X>

#### REFERENCES

- 1 Anderson LD, D'Alonzo RT. Fractures of the odontoid process of the axis. *J Bone Joint Surg Am* 1974;56:1663–74.
- 2 Zusman NL, Ching AC, Hart RA, Yoo JU. Incidence of second cervical vertebral fractures far surpassed the rate predicted by the changing age distribution and growth among elderly persons in the United States. *Spine* 2013;38:752–6.
- 3 Robinson AL, Möller A, Robinson Y, Olerud C. C2 fracture subtypes, incidence, and treatment allocation change with age: a retrospective cohort study of 233 consecutive cases. *Biomed Res Int* 2017;2017:8321680.
- 4 Fehlings MG, Arun R, Vaccaro AR, Arnold PM, Chapman JR, Kopjar B. Predictors of treatment outcomes in geriatric patients with odontoid fractures: Aospine North America multi-centre prospective GOF study. *Spine (Phila Pa 1976)* 2013;38:881–6.
- 5 Wagner SC, Schroeder GD, Kepler CK, Schupper AJ, Kandziara F, Vialle EN, Oner C, Fehlings MG, Vaccaro AR. Controversies in the management of geriatric odontoid fractures. *J Orthop Trauma* 2017;31 Suppl 4:S44–8.
- 6 Oner C, Rajasekaran S, Chapman JR, Fehlings MG, Vaccaro AR, Schroeder GD, Sadiqi S, Harrop J. Spine trauma-what are the current controversies *J Orthop Trauma* 2017;31:S1–6.
- 7 Iyer S, Hurlbert RJ, Albert TJ. Management of odontoid fractures in the elderly: a review of the literature and an evidence-based treatment algorithm. *Neurosurgery* 2018;82:419–30.
- 8 Ryan MD, Henderson JJ. The epidemiology of fractures and fracture-dislocations of the cervical spine. *Injury* 1992;23:38–40.
- 9 Carvalho AD, Figueiredo J, Schroeder GD, Vaccaro AR, Rodrigues-Pinto R. Odontoid fractures: a critical review of current management and future directions. *Clin Spine Surg* 2019;32:313–23.
- 10 Gonschorek O, Vordemvenne T, Blattert T, Katscher S, Schnake KJ. Treatment of odontoid fractures: recommendations of the spine section of the German society for orthopaedics and trauma (DGOU). *Global Spine J* 2018;8:125–175.
- 11 Nourbakhsh A, Hanson ZC. Odontoid fractures: a standard review of current concepts and treatment recommendations. *J Am Acad Orthop Surg* 2022;30:e561–72.
- 12 American College of Surgeons Committee on Trauma. National trauma data Bank version 2017 - 2019. Chicago, IL, 2021.
- 13 Radovanovic I, Urquhart JC, Rasoulinejad P, Gurr KR, Siddiqi F, Bailey CS. Patterns of C-2 fracture in the elderly: comparison of etiology, treatment, and mortality among specific fracture types. *J Neurosurg Spine* 2017;27:494–500.
- 14 Robinson A-L, Olerud C, Robinson Y. Surgical treatment improves survival of elderly with axis fracture-a national population-based multiregistry cohort study. *Spine J* 2018;18:1853–60.
- 15 Chan H-YH, Segreto FA, Horn SR, Bortz C, Choy GG, Passias PG, Deverall HH, Baker JF. C2 fractures in the elderly: single-center evaluation of risk factors for mortality. *Asian Spine J* 2019;13:746–52.
- 16 Dhall SS, Yue JK, Winkler EA, Mummaneni PV, Manley GT, Tarapore PE. Morbidity and mortality associated with surgery of traumatic C2 fractures in octogenarians. *Neurosurgery* 2017;80:854–62.
- 17 Robinson AL, Olerud C, Robinson Y. Epidemiology of C2 fractures in the 21st century: a national registry cohort study of 6,370 patients from 1997 to 2014. *Adv Orthop* 2017;2017:6516893.
- 18 Rizvi SAM, Helseth E, Rønning P, Mirzamohammadi J, Harr ME, Brommeland T, Aarhus M, Høstmælingen CT, Ølstørn H, Rydning PNF, et al. Odontoid fractures: impact of age and comorbidities on surgical decision making. *BMC Surg* 2020;20:236.