

RESEARCH

Open Access

Novel gene expression responses in the ovine abomasal mucosa to infection with the gastric nematode *Teladorsagia circumcincta*

Pamela A Knight^{1*}, Susan E Griffith¹, Alan D Pemberton¹, Judith M Pate¹, Lauren Guarneri¹, Katherine Anderson¹, Richard T Talbot¹, Sarah Smith¹, David Waddington¹, Mark Fell¹, Alan L Archibald¹, Stewart TG Burgess², David W Smith², Hugh RP Miller¹ and Ivan W Morrison¹

Abstract

Infection of sheep with the gastric nematode *Teladorsagia circumcincta* results in distinct Th2-type changes in the mucosa, including mucous neck cell and mast cell hyperplasia, eosinophilia, recruitment of IgA/IgE producing cells and neutrophils, altered T-cell subsets and mucosal hypertrophy. To address the protective mechanisms generated in animals on previous exposure to this parasite, gene expression profiling was carried out using samples of abomasal mucosa collected pre- and post- challenge from animals of differing immune status, using an experimental model of *T. circumcincta* infection. Recently developed ovine cDNA arrays were used to compare the abomasal responses of sheep immunised by trickle infection with worm-naïve sheep, following a single oral challenge of 50 000 *T. circumcincta* L3. Key changes were validated using qRT-PCR techniques. Immune animals demonstrated highly significant increases in levels of transcripts normally associated with cytotoxicity such as granulysin and granzymes A, B and H, as well as mucous-cell derived transcripts, predominantly calcium-activated chloride channel 1 (*CLCA1*). Challenge infection also induced up-regulation of transcripts potentially involved in initiating or modulating the immune response, such as heat shock proteins, complement factors and the chemokine *CCL2*. In contrast, there was marked infection-associated down-regulation of gene expression of members of the gastric lysozyme family. The changes in gene expression levels described here may reflect roles in direct anti-parasitic effects, immuno-modulation or tissue repair. (Funding; DEFRA/SHEFC (VT0102) and the BBSRC (BB/E01867X/1)).

Introduction

Parasitic gastroenteritis (PGE), caused by trichostrongylid nematodes, is the most commonly diagnosed systemic disease of sheep in the U.K. The principal causative nematode is the abomasal parasite *Teladorsagia circumcincta*. Control currently depends on the use of anthelmintics, but is failing due to the rapid emergence of drug resistance in the target nematodes [1]. Immunity builds up slowly on repeated exposure to the parasite, indicating vaccination could be a feasible alternative, but vaccine development is hampered by a lack of knowledge of the host-parasite interaction to infective

larvae. This immunity can be replicated experimentally by giving animals a low level trickle infection over several weeks, which results in a significant level of protective immunity to *T. circumcincta* challenge, measurable by reduced worm burdens, stunting of the worms and increased levels of larval arrest [2-4]. Protective immunity includes both cellular and humoral components; previously exposed animals undergo a local blast cell response in the draining lymphatics which can convey protection to genetically identical naïve recipients [2,5], while local IgA/IgE responses have been associated with certain protective responses such as stunted growth and reduced fecundity of the worms [4,6,7]. *T. circumcincta* challenge in previously immunised sheep elicits local predominantly Th2 cytokine expression, compared to a more Th1-bias in naïve animals [8,9]. This response is

* Correspondence: pamaknight@gmail.com

¹The Roslin Institute and R(D)SVS, University of Edinburgh, Roslin, Midlothian, EH25 9RG, Scotland, UK

Full list of author information is available at the end of the article

accompanied by distinct Th2-type changes in the mucosa, such as mucous neck cell and mast cell hyperplasia, eosinophilia, recruitment of IgA/IgE producing cells and neutrophils, altered T-cell subsets and mucosal hypertrophy [10-14]. However, the molecular changes involved, and the relative contributions of these factors to both control of infection and the clinical symptoms of disease, are still poorly understood. The host immune responses may act concordantly to generate an unfavourable micro-environment [15], which could involve generation of specific antibodies to reduce worm fecundity or feeding [6,7], or promote rapid expulsion [16]. To identify the molecular changes generated in the abomasal mucosa in animals after previous exposure to this parasite, gene expression profiling was carried out using ovine cDNA microarrays on samples of abomasal mucosa collected pre- and post-challenge from animals of differing immune status. The significance of the key changes observed is discussed.

Materials and Methods

All experimental research described in this manuscript was carried out in accordance with Moredun Research Institute, Roslin Institute and R(D)SVS guidelines. All experimental protocols were approved by the Moredun Research Institute Experiments and Ethical Review Committee and authorised under the UK Animals (Scientific Procedures) Act 1986.

Infections and sample collection

A series of experimental trials were set up to compare the immune responses of “previously infected” yearling sheep immunised by an eight-week trickle infection (referred to subsequently as “immune” throughout this manuscript), with worm-free naïve yearling sheep, at different timepoints post-challenge with a single dose of 50 000 *T. circumcincta* L3. The details of these infection trials are summarised in Table 1. All sheep were housed

under worm-free conditions. Previously described work has established that there were significant differences in anti-parasite responses between the “naïve” and “immune” groups used in this study [4,11]. Samples of host material collected post-mortem included abomasal fold (anterior to the fundic region) for RNA extraction [17]. RNA was extracted as described previously [17] and assessed for quality and quantity using an Agilent 2100 Bioanalyzer (Agilent Technologies UK Ltd, Edinburgh, U.K.) according to ARK-Genomics standard protocols [18].

Hybridisations and statistical analysis

Ovine cDNA microarrays (ARK-Genomics *O. aries* 12 K v1.0) were generated at ARK-Genomics using PCR-amplified products from individual cDNA clones predominantly from the KN511 library; a normalised cDNA library generated from gastrointestinal tract and lymphoid tissues of worm-free and *T. circumcincta* infected sheep, supplemented with an existing sheep/brain library [19]. Expressed Sequence Tags (ESTs) for each cDNA clone have been deposited in the EMBL/Genbank public DNA sequence databases. The array was annotated on the basis of homologous bovine sequences using Uni-gene, DfCI Gene Index, Ensembl cDNA library and the International Protein Index.

In order to characterise responses at day 5 (Expt.1) and day 0/day 2 (Expt. 2) post-challenge (Table 1), RNA samples were fluorescently labelled and competitively hybridised to the ovine cDNA microarrays using ARK-Genomics standard protocols [18]. Samples were paired in a dye-balanced arrangement as indicated in Table 2. The groups of samples being paired will be termed throughout this manuscript as Nvd5/Nvd0 (i.e. naive day 5 post-challenge vs. day 0 (unchallenged)), Imd5/Nvd5 (i.e. immune vs. naive day 5 post-challenge), Imd0/Nvd0 (i.e. immune vs. naive day 0 (unchallenged)) and Nvd2/Nvd0 or Imd2/Imd0 (i.e. naive or immune

Table 1 Design of *T. circumcincta* infection trials in yearling sheep

Trial	Group	Treatment		Days post-challenge ^a ; no. of sheep				
		Trickle infn ^b	clearance ^c	0 ^d	2	5	10	21
Expt. 1	Naïve (Nv)	-	+	6	-	6	6	6
	Immune (Im) ^e	+	+	-	-	6	6	-
Expt. 2	Naïve (Nv)	-	+	6	6	-	-	-
	Immune (Im) ^e	+	+	6	6	-	-	-

The experimental model of infection used in this study is summarised. Details are given in the footnotes. Two experimental trials were carried out; Expt.1; addressing differences between immune (“previously infected”) and naïve sheep at days 5 and 10 post challenge, and Expt.2, addressing differences between immune and naïve and sheep without challenge (day 0) and early post-challenge (day 2).

^a1 × 50 000 *T. circumcincta* L3 on Day 0

^b2000 *T. circumcincta* L3 3 × per wk for eight weeks

^cLevamisole (7.5 mg/kg) 7 days before challenge

^dDay 0 = unchallenged

^epreviously infected by trickle infection

Table 2 Hybridisation design

Experiment	Hybridisations (d = days post-challenge)		
Expt. 1	Nvd0/Nvd5	Nvd5/Imd5	
Day 5 post-challenge	(n = 6)	(n = 6)	
Expt. 2	Nvd2/Nvd0	Imd0/Nvd0	Imd2/Imd0
Day 2 post-challenge	(n = 5 ^a)	(n = 5 ^a)	(n = 5 ^a)

Samples in both cDNA hybridisation experiments were paired as indicated with a dye-balanced arrangement (three cy3/cy5 & three cy5/cy3 hybridisations). Nv = "naive" yearlings, worm naïve prior to challenge; Im = "immune" yearlings, previously infected by trickle infection prior to challenge (see Table 1). Common samples linked the sets of paired hybridisations (Expt 1: Nvd5 samples, Expt. 2: Nvd0 & Imd0 samples). An indirect comparison between the results from all three hybridisations gave the (Nvd2/Nvd0)/(Imd2/Imd0) gene list. ^aOne chip failed hybridisation QC in Expt 2 and so was removed from the analysis, along with chips linked to it.

day 2 post-challenge vs. day 0 (unchallenged)), and as shown in Table 2. Scanning and data capture using Bluefuse feature extraction software (BlueGnome Limited, Cambridge, United Kingdom) was undertaken according to ARK-Genomics standard protocols [18]. Log₂ (intensity ratios) were subjected to spatial and intensity dependent normalisations to remove technical bias by spatial row and column averaging [20] and M-A lowess correction [21]. Means of normalized gene log₂ (intensity ratios) were compared for treatment effects and treatment-by-dye interactions by ANOVA followed by *t*-tests, using modifications of the Limma package [22]. Statistical analysis to generate gene lists was based on each sample coming from a different sheep, and fitting the two (Expt. 1) or three (Expt. 2) dye-balanced arrangements with a common error variance. *T*-tests were modified by the Limma eBayes correction [23]. Genes with Benjamini & Hochberg false discovery rate ≤ 0.05 [24] were considered to show significantly different expression levels. Normalisations and analyses were weighted by the Bluefuse "confidence" measure [25]. Subsets of the data from Expt. 2; genes whose expression levels were significantly changed (FDR ≤ 0.05) ≥ 1.5 fold from the Imd0/Nvd0 and Imd2/Imd0 gene lists (Table 1)); were analyzed through the use of Ingenuity Pathways Analysis (Ingenuity® Systems (Redwood City, CA, U.S.A) [26]) by inputting human gene symbols (HUGO) for their putative ovine orthologues.

RT-PCR and multiplex qRT-PCR analysis

Subsequent validation assays for selected groups of genes whose expression levels were significantly changed (FDR ≤ 0.05) ≥ 2 fold were carried out initially by semi-quantitative RT-PCR and sequence analysis of products to confirm identity. Methods for reverse transcription, polymerase chain reaction and purification of PCR products have been described previously [17,27]. Gene-specific primers were designed from the ovine sequences (ESTs) for the relevant KN511 cDNA clones, or

previously published sequences, using the Primer 3 program; [28]. Primers and PCR conditions for RT-PCR are shown in Additional file 1: Table S1. Three "housekeeping" genes found to show no significant change in any of the array analyses (*ATPase*, *RW1* and *TM57*; Table 3) were used as positive controls. Sequence analysis of PCR products was carried out using multiple alignment and sequence similarity search programs available on [29,30] and [31].

In order to provide quantitative data on gene expression for the transcripts investigated above, two multiplex competitive qRT-PCR assays were designed from published sequences as summarised in Table 3; which included a total of 21 study genes and 3 "housekeeping" genes. Multiplex competitive qRT-PCR assays were designed by Sequenom Inc. (San Diego, CA, USA) using QGE assay designer software, version 3.4 (Sequenom® MassARRAY® System; [32]). These assays were used to compare levels of transcripts for each gene in naïve sheep with immune sheep at days 0, 2, 5 and 10 and 21 post-challenge, by running the multiplex qRT-PCR assays with cDNA from all 60 RNA samples summarised in Table 1. The assays and data collection were carried out using Sequenom standard protocols for quantitative gene expression [33]. Significant differences were identified using the Mann-Whitney U-test for non-parametric data.

Results

Results of microarray analysis

The numbers of transcripts that exhibited significant changes in expression levels (Benjamini & Hochberg FDR ≤ 0.05) in Expts. 1 and 2 are summarised in the Venn diagrams in Figure 1, and in Additional file 1 Table S2. In both experiments, the highest numbers of transcripts showing significantly altered expression levels (approx > 2K) were associated with the challenged immune group (day 2 or day 5) compared to day 0 of the same group, or compared to naïve animals at the same timepoint post-challenge (Figure 1a and 1b). There were no significantly differentially expressed genes in the Nvd2/Nvd0 comparison from Expt. 2. Any transcripts showing significant (FDR ≤ 0.05) treatment-by-dye interactions (Additional file 1: Table S2; Expt. 2) were removed from the gene lists. This precautionary measure excluded pairs of treatment estimates (from cy5-cy3 or cy3-cy5 dye orientations) with conflicting signs and some pairs of treatment estimates with the largest proportionate differences in magnitude (graphs not presented). The complete gene lists of significantly differentially expressed genes, with significant treatment-by-dye interactions excluded, are available under Array Express [34] or by contacting the authors. Data from Expt. 2 were analysed through the use of


Table 3 Details of gene sequences, and likely function, selected for further analysis

Functional Category	Gene Name	Clone/probe ID on KN511 cDNA array	EMBL acc. no.	Assay
Cytotoxicity	GNLY	C0009264H11.P1KAM13F	FE028690	W1
	CatC	sh1-ao011_10.s2	DY479012	W2
	GZMA	KN511_9486f02	FE037399	W2
	GZMB	KN511_9253n09	FE021442	W1
	GZMH	KN511_9260m05	FE026184	W2
Mucus composition	CLCA1	020502OAPP1012080HT OAPP	EE748540	W1
	ITLN1	- ^a	AM087961	W2
	ITLN2	KN511_9264a24.p1kaM13F	EF521881	W2
Heat shock response	ITLN3	- ^a	AM888394	W2
	HSPA8	KN511_9479o03	DY491193	W1
	HSPCA	KN511_9253c04	FE020983	W2
Proinflammatory response	ST1	KN511_9486e08	FE037366	W1
	PLA2G2A	KN511_9482g22	FE034658	W1
	CF1	KN511_9257m10	FE024140	W1
Tissue remodelling	CCL2	KN511_9477e07	FE031007	W1
	MMP13	KN511_9487m11.p1kM13F	AY091604	W1
	MMP23	sh1-bjd04_10.s2	DY483383	W2
Digestion	CST3	KN511_9476g20	FE030675	W1
	LZM1A	KN511_9487n22.p1kM13F ^b	M32492	W1
Unknown	LZM4A	KN511_9487n22.p1kM13F ^b	M32497	W1
	MALAT	KN511_9484i14	FE036272	W1
"Housekeeping" genes	ATPase	KN511_9488b23.p1kM13F	X02813	W1/W2
	RW1	KN511_9261h17	FE026635	W1/W2
	TM57	KN511_9488i21	FE030125	W1/W2

The EMBL accession no. for the sequences used to design multiplex competitive qRT-PCR assays (Sequenom® MassARRAY® System) is given, together with the representative probe on the ovine cDNA array. The two multiplex competitive qRT-PCR assays were designated W1 and W2 as indicated.

^aOnly *ITLN2* was represented on the arrays.


^bThese cDNA probes would not be able to distinguish between *LZM1A*, 3A and 4A.



Ingenuity Pathways Analysis (Ingenuity® Systems, [26]). Data were analysed from two datasets of genes whose expression levels were significantly changed ($FDR \leq 0.05$) ≥ 1.5 fold in Expt. 2 (see Table 2): Imd0/Nvd0; to highlight changes induced by trickle infection immunisation, and Imd2/Imd0; to highlight early (<48 h) changes induced following challenge of immunised animals. For each differentially expressed ovine transcript of interest the gene identifier (HUGO) for the putative orthologous human gene and the transcript's expression value were uploaded into the application. The output of the top 20 most significant (Fisher's exact test) biological functions are shown in Figure 2. Significant networks ($P \leq 10^{-10}$; Fischer's exact test) identified from the Imd0/Nvd0 and Imd2/Imd0 datasets from Expt. 2 are shown in Additional file 2 Table S3. Pathways were identified from the Ingenuity Pathways Analysis (Ingenuity® Systems, [26]) library of canonical pathways that were most significant to the data set. The top 20 most significant canonical pathways from the Imd0/Nvd0 and

Imd2/Imd0 datasets from Expt. 2 are shown in Additional file 2 Table S4.

The biological function associated with the most significant changes (Figure 2) in the Imd0/Nvd0 comparison was "Cell Cycle", which also features as the most significant network (Additional file 2 Table S3) probably reflecting the increased cellular turnover occurring in the immune versus naïve abomasum. "Cancer" was the most significant function from the Imd2/Imd0 comparison, which is likely to relate to increased transcription factor activity following challenge of the immunised sheep. It is notable that the functions "Inflammatory Disease", "Immunological Disease", "Infectious Disease" and "Cell-mediated Immune Response" were identified in the Imd2/Imd0 comparison only, which is indicative of increased innate and adaptive immune responses within 48 h following challenge. The most significant network from the Imd2/Imd0 comparison is "Hematological Disease, Small Molecule Biochemistry, Cellular Compromise" (Additional file 2 Table S3); further



pathway analysis of this subset (data not shown) showed the most significant pathways within this network were the “Complement System”, “Granzyme B signalling” (i.e. Granzyme B pathway), “Huntington’s Disease signalling” and “NRF2-mediated oxidative stress response”; the latter two pathways predominantly involve heat shock proteins. It is notable that NFkB signalling predominated in many of the pathways within this network. The 20 most significant pathways from the whole Imd0/Nvd0 and Imd2/Imd0 datasets are shown in Additional file 2 Table S4.

The 20 most significantly ($FDR \leq 0.05$) up- and down-regulated transcripts from both experiments are shown in Tables 4 (up-regulated transcripts) and 5 (down-regulated transcripts). Fold changes of a selection of these transcripts, grouped under possible function, are illustrated diagrammatically in Figure 3. Transcripts of particular biological interest, listed in Table 3, were selected for further investigation, initially by semi-quantitative RT-PCR (Additional file 1 Table S1) and sequence analysis to validate expression in the abomasum and sequence identity (data not shown) and subsequently, by multiplex competitive qRT-PCR analyses (Table 3). The aims of these assays were to validate the key findings from the array analysis (Figure 3), and to examine temporal changes in response to challenge infection in both the naïve and immune groups, using samples collected at post-mortem on days 0-21 post-challenge as summarised in Table 1. The results of the multiplex qRT-PCR analyses are shown in Figures 4, 5, 6 and 7. The results are discussed below in the context of transcripts specific to particular cell types and/or likely to share a similar function.

Transcripts normally associated with cells exhibiting cytotoxicity

Transcripts for granzysin were identified as the most highly up-regulated in immune sheep (+94 fold), compared with naïve at day 5 post-infection (p.i.) (Table 4; Figure 3), and were confirmed to show significantly ($FDR \leq 0.05$) highly up-regulated expression in immune animals at all timepoints post-challenge by multiplex qRT-PCR (Figure 4). We also observed a +2-5 fold up-regulation of transcripts for granzymes A, B and H in immune animals on day 5 post-challenge compared with naïve, and a >7 fold up-regulation of cathepsin C transcripts in immune animals on day 2 after challenge compared to naïve or to day 0 (Figure 3a). Results from the multiplex qRT-PCR analysis showed significantly higher numbers of transcripts for granzymes A and H in immune abomasal samples compared to naïve at day 5 post-challenge, with elevated levels persisting at day 10, but expression of granzyme B was highly variable and not significant (Figure 4). Transcripts for cathepsin C

were generally 10-fold more abundant in challenged animals at days 5-21 compared to day 0, although there were no significant differences between the immune and naïve groups (Figure 4). The 20 most significant pathways identified from the Imd2/Imd0 comparison included “Crosstalk between Dendritic cells and NK cells”, “Granzyme B Signalling” and “Leukocyte Extravasation Signalling”, indicative of an increased cell-mediated immune response (Additional file 2 Table S4).

Mucous-cell associated transcripts

The mucous-cell associated transcript calcium-activated chloride channel 1 (*CLCA1*) was one of the most up-regulated transcripts seen in immune sheep in both experiments. *CLCA1* was the second most up-regulated transcript in immune versus naïve sheep day 5 post-challenge (Expt. 1) and was also among the most up-regulated transcripts in the immune versus naïve sheep at day 0 and day 2 post-challenge (Expt 2) (Table 4, Figure 3b)). Significant ($FDR \leq 0.05$) up-regulation of *CLCA1* transcripts in immune versus naïve sheep was confirmed in the multiplex qRT-PCR analysis (day 5 post-challenge, Figure 5) and there was a general up-regulation of expression in response to challenge (i.e. compared to day 0) in both groups, significant ($P \leq 0.01$) in naïve animals on days 5-21. We also observed significant up-regulation of transcripts for the mucous cell product trefoil factor 3 (*TFF3*) in immune animals compared to naïve (Table 4; Figure 3b)), while trefoil factor 2 (*TFF2*) was highly down-regulated (Table 5).

While members of the intelectin family, galactose-binding lectins expressed by mucous neck cells [27], did not feature in the lists of significantly up-regulated genes from the microarray analysis (only *ITLN2* (KN511_9264a24.p1kaM13F) was represented on the array), specific assays for all three of these genes were included in the multiplex qRT-PCR analysis as a positive control, since *ITLNs* 1, 2 and 3 have been shown to be up-regulated in abomasal mucosa in response to *T. circumcincta* [35]. There was a significant up regulation in response to challenge in the naïve group for all 3 intelectins, with earlier up-regulation of *ITLN-3* than *ITLN-1* and -2, although these assays did not show significant differences between naïve and immune sheep (Figure 5). We also observed infection-associated up-regulation of transcripts for two galectins, which belong to the C-type lectin family; galectin 1 (+1.5 fold), and galectin 4 (+1.6 fold).

Mast -cell associated transcripts

Significantly ($FDR \leq 0.05$) up-regulated gene expression of the high affinity IgE receptor in immune animals was detected on day 2 (+1.5 fold) and day 5 (+1.8 fold) post-challenge, compared to the naïve group at the same

Table 4 The 20 most up-regulated transcripts detected in Expts. 1 and 2 (FDR ≤ 0.05)

Sequence.ID	TIGR_TC	TC. description	Gene symbol	Fold Change
EXPT 1. Comparison = Nvd5/Nvd0				
KN511_9480n02.p1kaM13F	TC385850	—	—	5.44
C0009213B13	—	—	—	1.81
KN511_9261j08.p1kaM13F	—	—	—	1.77
KN511_9255d18.p1kaM13F	TC307335	laminin B1	LAMB1	1.64
KN511_9486f02.p1kaM13F	TC305120	UP GRAA_BOVIN (Q7YRZ7) Granzyme A precursor, complete	gzmA	1.59
KN511_9488i01.p1kaM13F	TC311345	homologue to UP Q96IL1_HUMAN (Q96IL1) DIAPH1 protein (Fragment), partial (43%)	LOC786565	1.55
KN511_9483h09.p1kaM13F	TC302202	UP Q3T119_BOVIN (Q3T119) FCGRT protein, complete	FCGRT	1.53
CN822687	TC381381	similar to emb X79547.1 MIECCOMP Equus caballus mitochondrial DNA complete sequence, partial (4%)	—	1.52
KN511_9257n17.p1kaM13F	—	—	—	1.52
KN511_9479d15.p1kaM13F	TC361081	similar to UP O97916_BOVIN (O97916) Reverse transcriptase-like, partial (43%)	—	1.51
KN511_9486m17.p1kaM13F	TC303503	UP C1QA_BOVIN (Q5E9E3) Complement C1q subcomponent subunit A precursor, complete	C1QA	1.50
KN511_9257l13.p1kaM13F	—	—	—	1.50
KN511_9479o01.p1kaM13F	TC355858	GB AAI02065.1 74353831 BC102064 CCL5 protein {Bos taurus} (exp = -1; wgp = 0; cg = 0), complete	CCL5	1.50
CN823601	TC342786	—	—	1.50
KN511_9486d22.p1kaM13F	TC337098	UP Q3SZV9_BOVIN (Q3SZV9) Heat shock 10kDa protein 1 (Chaperonin 10), complete	HSPE1	1.50
KN511_9477j24.p1kaM13F	TC301808	UP Q3ZC91_BOVIN (Q3ZC91) Sphingomyelin phosphodiesterase, acid-like 3A, complete	LOC781963	1.48
KN511_9485e22.p1kaM13F	—	—	—	1.48
KN511_9253n18.p1kaM13F	TC317864	UP Q3KZ51_SCHJA (Q3KZ51) SJCHGC03568 protein (Fragment), partial (6%)	MGC139367	1.48
KN511_9265g17.p1kaM13F	TC319596	homologue to UP TCEA3_HUMAN (O75764) Transcription elongation factor A protein 3 (Transcription elongation factor S-II protein 3) (Transcription elongation factor TFIIIS.h), complete	MGC137536	1.48
KN511_9265b23.p1kaM13F	—	—	—	1.48
EXPT 1 Comparison = Imd5/Nvd5				
KN511_9264h11.p1kaM13F	TC331976	homologue to UP Q864L7_BOVIN (Q864L7) Granulysin/NK-lysin-like protein (Fragment), complete	GNLY	94.0
KN511_9257k12.p1kaM13F	—	—	—	5.91
KN511_9263m23.p1kaM13F	TC306994	similar to UP Q9TUB5_PIG (Q9TUB5) Epithelial chloride channel protein, partial (95%)	CLCA1	5.15
KN511_9253n09.p1kaM13F	TC328399	similar to UP Q67BC3_HUMAN (Q67BC3) Endogenous granzyme B, partial (90%)	LOC508646	5.14
KN511_9265g07.p1kaM13F	TC351879	—	LOC614719	4.70
KN511_9260m05.p1kaM13F	—	—	—	4.64
KN511_9260b08.p1kaM13F	TC330838	—	—	4.27
CN823175	—	—	—	3.14
CN823115	TC329726	homologue to GB CAA44699.1 440 BTIGG1HCX anti-testosterone antibody {Bos taurus} (exp = -1; wgp = 0; cg = 0), partial (70%)	IGHG1	2.72
KN511_9480o15.p1kaM13F	—	—	—	2.66
KN511_9476e05.p1kaM13F	TC330547	similar to UP NBR1_PONPY (Q5RC94) Next to BRCA1 gene 1 protein (Neighbor of BRCA1 gene 1 protein), partial (23%)	LOC515032	2.64
KN511_9486a03.p1kaM13F	TC350940	similar to UP CAD26_MOUSE (P59862) Cadherin-like protein 26 precursor, partial (11%)	LOC617096	2.47
CO202749	TC366134	similar to UP Q48MR5_PSE14 (Q48MR5) Uncharacterized protein family UPF0016, partial (58%)	—	2.45
KN511_9486o13.p1kaM13F	TC322636	weakly similar to UP Q9R0M7_MOUSE (Q9R0M7) Aldo-keto reductase AKR1C12, partial (91%)	LOC507734	2.42
CN824147	—	—	—	2.38
CN823337	TC313767	UP Q2FLD4_METHJ (Q2FLD4) ABC transporter related, partial (6%)	—	2.37
CO203065	TC321552	—	—	2.34
KN511_9487e05.p1kaM13F	TC302266	GB AAC67307.1 1930063 BTU92535 neuronal axonal membrane protein {Bos taurus} (exp = -1; wgp = 0; cg = 0), complete	BASP1	2.30

Table 4 The 20 most up-regulated transcripts detected in Expts. 1 and 2 (FDR \leq 0.05) (Continued)

KN511_9261f23.p1kaM13F	—	—	—	2.29
KN511_9478a01.p1kaM13F	TC347423	similar to UP GRAH_HUMAN (P20718) Granzyme H precursor (Cytotoxic T-lymphocyte proteinase) (Cathepsin G-like 2) (CTSG2) (CCP-X) (Cytotoxic serine protease C) (CSP-C), partial (93%)	LOC617313	2.28
KN511_9486l10.p1kaM13F	TC315163	similar to GB AAH12303.1 15126763 BC012303 PDZK1 interacting protein 1 {Homo sapiens} (exp = -1; wgp = 0; cg = 0), complete	PDZK1IP1	2.21
CN824077	—	—	—	2.21
CN824594	TC359921	homologue to UP Q4K9Y4_PSEF5 (Q4K9Y4) BNR/Asp-box repeat protein, partial (41%)	—	2.14
KN511_9260m24.p1kaM13F	—	—	—	2.13
EXPT 2 Comparison = Imd0/Nvd0				
KN511_9484k05.p1kaM13F	TC323139	similar to UP Q4TA31_TETNG (Q4TA31) Chromosome undetermined SCAF7459, whole genome shotgun sequence, partial (10%)	—	3.59
KN511_9263m23.p1kaM13F	TC306994	similar to UP Q9TUB5_PIG (Q9TUB5) Epithelial chloride channel protein, partial (95%)	CLCA1	2.99
KN511_9484p11.p1kaM13F	—	—	—	2.08
KN511_9485e08.p1kaM13F	TC312631	—	—	1.61
KN511_9478a06.p1kaM13F	TC312631	—	—	1.49
KN511_9481c23.p1kaM13F	TC314684	—	—	1.46
CN823114	TC301217	homologue to UP NDRG1_HUMAN (Q92597) Protein NDRG1 (N-myc downstream-regulated gene 1 protein) (Differentiation-related gene 1 protein) (DRG-1) (Reducing agents and tunicamycin-responsive protein) (RTP) (Nickel-specific induction protein Cap43) (Rit42), partial (97%)	NDRG1	1.37
KN511_9257n17.p1kaM13F	—	—	—	1.35
CN823685	TC315004	similar to UP Q3TT48_MOUSE (Q3TT48) Adult male pituitary gland cDNA, RIKEN full-length enriched library, clone:5330433L19 product:proprotein convertase subtilisin/kexin type 2, full insert sequence. (Fragment), partial (3%)	LOC615685	1.33
KN511_9476g20.p1kaM13F	TC337413	UP CYTC_BOVIN (P01035) Cystatin C precursor (Colostrum thiol proteinase inhibitor), complete	CST3	1.31
—	—	—	—	1.30
KN511_9259b18.p1kaM13F	TC371167	—	—	1.29
KN511_9479a08.p1kaM13F	TC318244	UP CATA_BOVIN (P00432) Catalase, partial (5%)	MGC128112	1.29
CN823587	—	—	—	1.27
CO202780	TC357183	UP Q71V68_MOUSE (Q71V68) Ldb1a, complete	LOC526472	1.26
KN511_9262a18.p1kaM13F	TC308953	UP Q3T013_BOVIN (Q3T013) BCL2/adenovirus E1B 19kDa interacting protein 3-like, partial (84%)	BNIP3L	1.26
CO202648	TC301621	homologue to UP Q9HB23_HUMAN (Q9HB23) Lysyl-tRNA synthetase, complete	MGC127504	1.25
KN511_9254i12.p1kaM13F	TC381609	homologue to UP Q963G4_PLAFA (Q963G4) MB2 (Fragment), partial (3%)	LOC508133	1.25
KN511_9261e18.p1kaM13F	—	—	—	1.24
EXPT 2 Comparison = Imd2/Imd0				
KN511_9481e19.p1kaM13F	TC318483	—	LRIG1	10.2
KN511_9256h20.p1kaM13F	TC309152	similar to UP O97916_BOVIN (O97916) Reverse transcriptase-like, partial (19%)	—	10.0
KN511_9257i19.p1kaM13F	TC375873	—	—	9.71
KN511_9481f19.p1kaM13F	TC345827	homologue to GB CAI11042.1 55859631 AL353665 match: proteins: Q92802 {Homo sapiens} (exp = 0; wgp = 1; cg = 0), partial (35%)	LOC541201	9.17
KN511_9480f18.p1kaM13F	—	—	—	8.94
CN822322	TC309241	nucleolin [Bos taurus]	NCL	8.03
KN511_9483a22.p1kaM13F	TC316336	UP Q3ZCJ8_BOVIN (Q3ZCJ8) Cathepsin C, complete	CTSC	7.52
KN511_9263m23.p1kaM13F	TC306994	similar to UP Q9TUB5_PIG (Q9TUB5) Epithelial chloride channel protein, partial (95%)	CLCA1	7.33
KN511_9483l13.p1kaM13F	TC340529	homologue to UP Q9BTX0_HUMAN (Q9BTX0) RNA binding motif protein 10, isoform 2, partial (69%)	RBM10	6.73
KN511_9488g12.p1kaM13F	TC304325	similar to GB CAJ18388.1 71059689 CT010180 {Mus musculus} (exp = -1; wgp = 0; cg = 0), complete	MGC127625	6.46
KN511_9263i18.p1kaM13F	TC332274	UP Q7ZU59_BRARE (Q7ZU59) H2AV protein (Fragment), partial (87%)	H2AFZ	6.45
KN511_9253j02.p1kaM13F	TC331773	homologue to UP HSP76_PIG (Q04967) Heat shock 70 kDa protein 6 (Heat shock 70 kDa protein B'), complete	HSPA6	6.11

Table 4 The 20 most up-regulated transcripts detected in Expts. 1 and 2 (FDR \leq 0.05) (Continued)

KN511_9487k02.p1kaM13F	TC385703	UP VPS28_MOUSE (Q9D1C8) VPS28 protein homolog (Caspase-activated DNase inhibitor that interacts with ASK1) (CIIA), complete	VPS28	5.97
KN511_9265g07.p1kaM13F	TC351879	—	LOC614719	5.84
KN511_9476g04.p1kaM13F	TC371801	homologue to SP O60506 HNRPQ_HUMAN Heterogeneous nuclear ribonucleoprotein Q (hnRNP Q) (hnRNP-Q)(Synaptotagmin binding, cytoplasmic RNA interacting protein) (Glycine-and tyrosine-rich RNA binding protein) (GRY-RBP) (NS1-associatedprotein 1). {Homo sapiens} (exp = -1; wgp = -1; cg = -1), partial (51%)	SYNCRIP	5.58
KN511_9263f17.p1kaM13F	TC355165	similar to UP Q9XSA0_SHEEP (Q9XSA0) Pulmonary surfactant-associated protein B (Fragment), partial (16%)	—	4.64
KN511_9476p04.p1kaM13F	TC329988	UP Q5E973_BOVIN (Q5E973) Ribosomal protein L18, complete	RPL18	4.47
KN511_9480l18.p1kaM13F	TC364833	—	—	4.38
KN511_9265h18.p1kaM13F	TC303880	homologue to UP O35328_MOUSE (O35328) Proline-rich protein 9-1 (Fragment), partial (6%)	—	4.18
KN511_9260b08.p1kaM13F	TC330838	—	—	4.15

The experiment and the samples being compared are given as subheadings at the top of each section. Nv = "naïve" yearlings, worm naïve prior to challenge; Im = "immune" yearlings, previously infected by trickle infection prior to challenge, d = days post-challenge. Details of the experimental trials and hybridisation design are given in Tables 1 and 2.

timepoints. Up-regulated expression of cathepsin C, another potential mast cell transcript, has already been mentioned in the previous section. *SMCP-1* and trypsinase, which are expressed by abomasal mucosal mast cells during nematode infection [36,37], did not feature on this array.

Heat shock proteins

Significant up-regulation of a number of transcripts for heat shock proteins was detected in both microarray experiments. Transcripts for the heat shock proteins *HSP1A* (HSP70 protein 1A) and *HSPA6* (HSP 70 protein 6) feature among some of the most significantly (FDR \leq 0.05) highly up-regulated transcripts detected in immune sheep, up-regulated (+3-7 fold) by day 2 post-challenge in this group in comparison to naïve (Figure 3c) and Table 4). The *STIP1* (HSP70/90 organising protein; HOP) and *HSPA8* (HSP70 protein 8) genes also appear to show significantly up-regulated expression in immune animals compared to naïve (+1.5-2 fold) (Figure 3c) and Table 4). Transcripts for *HSPCA* (HSP90a) appear to show significantly (FDR \leq 0.05) increased levels on day 2 post-challenge in immune animals, although the expression pattern varies across the different comparisons (Figure 3c) and Table 4). The most significant network identified in the Imd2/Imd0 comparison (Network 1;"Cellular Compromise"; Additional file 2 Table S3) links *STIP1* with a number of heat shock proteins in the *HSP70/90* family. Multiplex qRT-PCR analysis showed that transcripts for *STIP1* were significantly ($P \leq$ 0.01-0.05) increased day 5-21 post-challenge in the naïve group, with a similar trend in immune sheep, although there were no immune day 0 animals within the same trial (Figure 6). Multiplex qRT-PCR analysis, however, did not support all the

findings for *HSPA8* and *HSPCA*, although there was a significant ($P \leq$ 0.05) challenge-associated increase in levels of *HSPA8* transcripts in the naïve group by day 10 (Figure 6).

Transcripts associated with pro-inflammatory responses

The microarray analyses demonstrated significant (FDR \leq 0.05) up-regulation of a number of transcripts associated with eicosanoid metabolism. Transcripts for *PLA2G2A*, which liberates arachidonic acid from phospholipids, were significantly up-regulated in response to challenge (+3 fold), and in immune animals compared to naïve at day 2 and day 5 post-challenge (+1.5-2 fold) (Figure 3d). The multiplex qRT-PCR analysis showed significant ($P \leq$ 0.05) up-regulation of *PLA2G2A* transcripts at day 5 post-challenge, but did not show significant differences between the naïve and immune groups (Figure 6). Transcripts for prostaglandin E synthase (*CPGES*), another member of this pathway, appear to be significantly (FDR \leq 0.05) up-regulated (+1.6 fold) in immune animals by day 2 post-challenge compared to naïve animals. "Arachidonic Acid Metabolism" features among the significant pathways from the Imd0/Nvd0 comparison (Additional file 2 Table S4).

Transcripts for the chemokine *CCL2* (MCP-1) were significantly up-regulated >2 fold on day 2 post-challenge in immune sheep, compared to naïve or unchallenged sheep (Figure 3d). Multiplex qRT-PCR analysis showed that transcripts for *CCL2* were significantly ($P \leq$ 0.01-0.05) increased day 10-21 post-challenge in the naïve group in Expt. 1, with a similar trend in the immune group, but did not show significant differences between the two groups (Figure 6). Transcripts for the chemokine *CCL5* (RANTES) were among the most up-regulated in naïve sheep on day 5 post-challenge (+1.5

Table 5 The 20 most down-regulated transcripts detected in Expts 1 and 2 (FDR ≤ 0.05)

Sequence.ID	TIGR_TC	TC.description	Gene symbol	Fold Change
EXPT 1 Comparison = d5Nv_d0Nv				
KN511_9487m11.p1kaM13F	TC336019	UP MMP13_BOVIN (O77656) Collagenase 3 precursor (Matrix metalloproteinase-13) (MMP-13), complete	MMP13	-2.68
KN511_9481c03.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-2.37
KN511_9476d13.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-2.33
KN511_9485a08.p1kaM13F	TC302650	milk lysozyme [Bos taurus]	LYZ1	-2.30
KN511_9485g08.p1kaM13F	TC346615	similar to UP O97916_BOVIN (O97916) Reverse transcriptase-like, partial (32%)	—	-2.23
KN511_9257d12.p1kaM13F	TC322737	similar to UP Q3VD89_9SPHN (Q3VD89) ABC-2, partial (5%)	—	-2.16
KN511_9257a18.p1kaM13F	TC308928	similar to GB AAK18773.1 13324523 F272846531 Fanconi anemia complementation group D2 protein, isoform 2 {Homo sapiens} (exp = -1; wgp = 0; cg = 0), partial (18%)	LOC515845	-2.14
KN511_9262d01.p1kaM13F	—	—	—	-2.12
KN511_9261n23.p1kaM13F	TC362037	homologue to UP IFRD1_PIG (Q551U6) Interferon-related developmental regulator 1, complete	IFRD1	-2.12
KN511_9476d14.p1kaM13F	TC349311	similar to UP NEK4_HUMAN (P51957) Serine/threonine-protein kinase Nek4 (NimA-related protein kinase 4) (Serine/threonine-protein kinase 2) (Serine/threonine-protein kinase NRK2), partial (33%)	MGC159441	-2.08
KN511_9261o08.p1kaM13F	TC305060	homologue to UP Q96BA7_HUMAN (Q96BA7) HNRPU protein, partial (93%)	MGC142835	-2.07
KN511_9258d23.p1kaM13F	—	—	—	-2.07
KN511_9487p21.q1kaM13R	TC342358	homologue to UP DC1L2_HUMAN (O43237) Cytoplasmic dynein 1 light intermediate chain 2 (Dynein light intermediate chain 2, cytosolic) (LIC53/55) (LIC-2), complete	LOC519789	-2.05
KN511_9262k18.p1kaM13F	TC375209	similar to UP ZSWM3_HUMAN (Q96MP5) Zinc finger SWIM domain-containing protein 3, partial (27%)	LOC512244	-2.05
KN511_9485d23.q1kaM13R	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-2.04
KN511_9254g13.p1kaM13F	TC319944	—	—	-2.02
KN511_9487h17.p1kaM13F	TC327406	UP Q5DTN6_MOUSE (Q5DTN6) MKIAA4095 protein (Fragment), partial (8%)	LOC614823	-1.96
CO202908	TC360504	UP Q3MHM4_BOVIN (Q3MHM4) Heat shock 70 kDa protein 8, complete	HSPA8	-1.87
KN511_9482g13.p1kaM13F	TC331274	homologue to UP TENA_PIG (Q29116) Tenascin precursor (TN) (Hexabrachion) (Cytotactin) (Neuronectin) (GMEM) (JI) (Miotendinous antigen) (Glioma-associated-extracellular matrix antigen) (GP 150-225) (Tenascin-C) (TN-C) (P230), partial (10%)	MGC140517	-1.87
KN511_9482f05.p1kaM13F	TC377013	—	—	-1.86
EXPT 1 Comparison = d5Im_d5Nv				
KN511_9256g17.p1kaM13F	TC314535	homologue to UP Q94513_DROME (Q94513) Boundary element associated factor (LD44361p) (CG10159-PB, isoform B), partial (5%)	—	-6.51
KN511_9256g17.p1kaM13F	TC314535	homologue to UP Q94513_DROME (Q94513) Boundary element associated factor (LD44361p) (CG10159-PB, isoform B), partial (5%)	—	-5.97
KN511_9254i23.p1kaM13F	TC354719	similar to UP TFF2_HUMAN (Q03403) Trefoil factor 2 precursor (Spasmodytic polypeptide) (SP) (Spasmodysin), complete	MGC139191	-5.54
KN511_9257i15.p1kaM13F	TC354719	similar to UP TFF2_HUMAN (Q03403) Trefoil factor 2 precursor (Spasmodytic polypeptide) (SP) (Spasmodysin), complete	MGC139191	-5.42
KN511_9260b10.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.92
KN511_9260m19.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.79
KN511_9483j03.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.52
KN511_9265j02.p1kaM13F	CB220717	—	—	-3.45
KN511_9264e04.p1kaM13F	TC314849	UP Q6SJV6_BOVIN (Q6SJV6) Foveolin precursor, complete	FOV	-3.33
KN511_9262k17.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.25
KN511_9476c18.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.25

Table 5 The 20 most down-regulated transcripts detected in Expts 1 and 2 (FDR \leq 0.05) (Continued)

KN511_9476d13.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.17
KN511_9488a15.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.16
KN511_9259k01.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.14
KN511_9265m20.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.10
KN511_9263j17.p1kaM13F	TC323852	similar to UP Q86XP6_HUMAN (Q86XP6) GDDR (Blottin precursor) (TFIZ1 protein precursor), complete	LOC512001	-3.10
KN511_9487e19.p1kaM13F	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.08
KN511_9485d23.q1kaM13R	TC365645	UP LYSC2_BOVIN (Q06283) Lysozyme C-2 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-3.05
KN511_9486h17.p1kaM13F	TC314849	UP Q65JV6_BOVIN (Q65JV6) Foveolin precursor, complete	FOV	-3.04
KN511_9265h22.p1kaM13F	TC304583	UP LYSC1_BOVIN (Q06285) Lysozyme C-1 precursor (1,4-beta-N-acetylmuramidase C), complete	LYZ2	-2.99
EXPT 2 Comparison = Day 0 Im vs Day 0 Nv				
KN511_9484p13.p1kaM13F	TC376600	similar to UP CENPA_MOUSE (O35216) Centromere protein A (CENP-A) (Centromere autoantigen A), partial (15%)	CENP-A	-1.75
KN511_9479i16.p1kaM13F	—	—	—	-1.75
KN511_9253l11.p1kaM13F	TC303509	homologue to UP RIR2_HUMAN (P31350) Ribonucleoside-diphosphate reductase M2 subunit (Ribonucleotide reductase small subunit) (Ribonucleotide reductase small chain), partial (88%)	RRM2	-1.73
KN511_9488d17.p1kaM13F	TC314207	similar to UP TOP2A_PIG (O46374) DNA topoisomerase 2-alpha (DNA topoisomerase II, alpha isozyme), partial (17%)	TOP2A	-1.71
KN511_9265l09.p1kaM13F	TC340393	UP CCNA2_BOVIN (P30274) Cyclin-A2 (Cyclin-A) (Fragment), complete	CCNA2	-1.70
KN511_9488f20.p1kaM13F	TC355225	homologue to UP BIRC5_CANFA (Q81009) Baculoviral IAP repeat-containing protein 5 (Apoptosis inhibitor survivin), complete	BIRC5	-1.66
KN511_9487m04.p1kaM13F	TC355225	homologue to UP BIRC5_CANFA (Q81009) Baculoviral IAP repeat-containing protein 5 (Apoptosis inhibitor survivin), complete	BIRC5	-1.65
KN511_9253i15.p1kaM13F	TC340307	UP CP51A_BOVIN (Q4PJW3) Cytochrome P450 51A1 (CYPLI) (P450LI) (Sterol 14-alpha demethylase) (Lanosterol 14-alpha demethylase) (LDM) (P450-14DM) (P45014DM), complete	CYP51	-1.63
KN511_9265a16.p1kaM13F	TC303827	—	KIF11	-1.58
KN511_9264b09.p1kaM13F	—	—	—	-1.56
KN511_9256j19.p1kaM13F	TC355225	homologue to UP BIRC5_CANFA (Q81009) Baculoviral IAP repeat-containing protein 5 (Apoptosis inhibitor survivin), complete	BIRC5	-1.53
KN511_9261d22.p1kaM13F	—	—	—	-1.50
KN511_9481l06.p1kaM13F	TC303509	homologue to UP RIR2_HUMAN (P31350) Ribonucleoside-diphosphate reductase M2 subunit (Ribonucleotide reductase small subunit) (Ribonucleotide reductase small chain), partial (88%)	LOC508167	-1.49
KN511_9476j14.p1kaM13F	TC333525	homologue to GB AAH08056.2 39644830 BC008056 PLEK2 protein {Homo sapiens} (exp = -1; wgp = 0; cg = 0), partial (21%)	MGC142714	-1.48
KN511_9482e10.p1kaM13F	TC330132	similar to UP CV106_HUMAN (Q6P0N0) Protein C14orf106 (P243), partial (33%)	LOC507661	-1.47
KN511_9257p16.p1kaM13F	TC340307	UP CP51A_BOVIN (Q4PJW3) Cytochrome P450 51A1 (CYPLI) (P450LI) (Sterol 14-alpha demethylase) (Lanosterol 14-alpha demethylase) (LDM) (P450-14DM) (P45014DM), complete	CYP51	-1.46
KN511_9256f06.p1kaM13F	TC332145	similar to UP Q9BDH4_PIG (Q9BDH4) Amphiregulin long form, complete	MGC152310	-1.45
CN824006	TC306385	homologue to UP Q4S6A0_TETNG (Q4S6A0) Chromosome 9 SCAF14729, whole genome shotgun sequence, partial (59%)	LOC404053	-1.42
KN511_9257l02.p1kaM13F	TC327524	homologue to UP Q9BW51_HUMAN (Q9BW51) TTK protein kinase, partial (44%)	LOC524925	-1.41
KN511_9261e22.p1kaM13F	TC301804	homologue to UP IMA2_HUMAN (P52292) Importin alpha-2 subunit (Karyopherin alpha-2 subunit) (SRP1-alpha) (RAG cohort protein 1), complete	KPNA2	-1.41
EXPT 2 Comparison = Day 2 Im vs Day 0 Im				
KN511_9479k18.p1kaM13F	CX950602	—	—	-3.53
KN511_9488b02.p1kaM13F	DY126632	—	—	-3.24

Table 5 The 20 most down-regulated transcripts detected in Expts 1 and 2 (FDR \leq 0.05) (Continued)

KN511_9257115.p1kaM13F	TC354719	similar to UP TFF2_HUMAN (Q03403) Trefoil factor 2 precursor (Spasmodic polypeptide) (SP) (Spasmodic), complete	TFF2	-3.14
KN511_9254123.p1kaM13F	TC354719	similar to UP TFF2_HUMAN (Q03403) Trefoil factor 2 precursor (Spasmodic polypeptide) (SP) (Spasmodic), complete	TFF2	-3.03
KN511_9264e04.p1kaM13F	TC314849	UP Q65JV6_BOVIN (Q65JV6) Foveolin precursor, complete	FOV	-2.81
KN511_9263j17.p1kaM13F	TC323852	similar to UP Q86XP6_HUMAN (Q86XP6) GDDR (Blottin precursor) (TFIZ1 protein precursor), complete	GKN2	-2.45
KN511_9486h17.p1kaM13F	TC314849	UP Q65JV6_BOVIN (Q65JV6) Foveolin precursor, complete	FOV	-2.09
KN511_9261k22.p1kaM13F	TC327109	similar to UP O97916_BOVIN (O97916) Reverse transcriptase-like, partial (21%)	—	-1.93
KN511_9486b13.p1kaM13F	TC341767	—	—	-1.90
KN511_9484h13.p1kaM13F	TC350450	similar to UP UDBS_MOUSE (P17717) UDP-glucuronosyltransferase 2B5 precursor (UDPGT) (M-1), partial (49%)	UGT2B17	-1.81
KN511_9263c22.p1kaM13F	—	—	—	-1.80
KN511_9264a19.p1kaM13F	—	—	—	-1.77
KN511_9484i14.p1kaM13F	TC362885	similar to UP MALAT_HUMAN (Q9UH22) Metastasis-associated lung adenocarcinoma transcript 1, partial (72%)	MALAT1	-1.75
KN511_9480l06.p1kaM13F	TC362885	similar to UP MALAT_HUMAN (Q9UH22) Metastasis-associated lung adenocarcinoma transcript 1, partial (72%)	MALAT1	-1.74
KN511_9481o23.p1kaM13F	TC350450	similar to UP UDBS_MOUSE (P17717) UDP-glucuronosyltransferase 2B5 precursor (UDPGT) (M-1), partial (49%)	UGT2B17	-1.73
CN824758	—	—	—	-1.72
KN511_9488a17.p1kaM13F	TC318925	—	—	-1.71
KN511_9480o03.p1kaM13F	—	—	—	-1.70
KN511_9263o20.p1kaM13F	TC380127	similar to UP UDBS_MOUSE (P17717) UDP-glucuronosyltransferase 2B5 precursor (UDPGT) (M-1), partial (42%)	UGT2B17	-1.70
KN511_9484i14.p1kaM13F	TC362885	similar to UP MALAT_HUMAN (Q9UH22) Metastasis-associated lung adenocarcinoma transcript 1, partial (72%)	MALAT1	-1.67

The experiment and the samples being compared are given as subheadings at the top of each section. Nv = "naïve" yearlings, worm naïve prior to challenge; Im = "immune" yearlings, previously infected by trickle infection prior to challenge, d = days post-challenge. Details of the experimental trials and hybridisation design are given in Tables 1 and 2.

fold; Figure 3d)), and were up-regulated in immune sheep compared to naïve at this timepoint (+1.4 fold).

Gene expression of several members of the complement cascade also appear to be up-regulated in response to *T. circumcincta* challenge. Complement factor 1 (CF1), and other members of the classical C1-complex, were significantly (FDR \leq 0.05) up-regulated in immune sheep in response to challenge by day 2 (+2-3 fold), but not in naïve animals (Table 4 and Figure 3d)). The "Complement System" was the third most significant pathway identified from the Imd2/Imd0 comparison (Figure 2), and the most significant pathway within the Network 1 subset from this comparison (Additional file 2 Table S3). However, the multiplex qRT-PCR analysis specific for complement factor 1 did not show the same pattern of expression detected in the arrays, with a lot of sheep-to-sheep variation, although there was a clear trend towards up-regulated gene expression in response to challenge in both groups in Expt. 1 (Figure 6).


Matrix metalloproteinases and protease inhibitors

Transcripts for the matrix metalloproteinase *MMP-23B* were significantly (FDR \leq 0/05) up-regulated (+1.7 fold) in immune versus naïve sheep on day 5 pi in the

microarray analysis (Figure 3e), while *MMP-13* was down-regulated (-2.3 fold) in the immune group compared to naïve at day 5 post-challenge (Figure 3e and Table 5). There was also significant up-regulation of the protease inhibitor cystatin C (*CST3*) in immune sheep compared to naïve at day 0 and day 5 post-challenge (+1.4/+1.5 fold) (Figure 3e and Table 5). The multiplex qRT-PCR analysis confirmed transcripts for *MMP-13* were expressed at significantly ($P \leq$ 0.05) lower levels in immune versus naïve sheep at day 0, while *MMP23B* gene expression showed a similar pattern at day 2 pi ($P \leq$ 0.05; immune vs. naïve) (Figure 6). Transcripts for cystatin C appeared to be expressed at high but fairly consistent levels in both groups, although significantly higher levels were detected in naïve sheep than immune sheep at day 2 post-challenge (Figure 6).

Immunoglobulins


Genes featuring the "immunoglobulin" descriptor occur 31 times among the significantly (FDR \leq 0.05) up-regulated transcripts in immune animals on day 5 post-challenge compared to day 5 naïve, and 13 times in the day 2 immune versus day 0 immune group (see [34] for complete dataset).



Gastric lysozyme family

Members of the gastric lysozyme family [38] were the most consistently down-regulated transcripts ($FDR \leq 0.05$) detected in the Expt. 1 microarray analyses (day 5 post-challenge) (Figure 3f and Table 5). RT-PCR and sequence analysis have previously indicated infection-associated down-regulation of transcripts for the ovine

gastric lysozyme genes *1A*, *2A*, *3A* and *4A*, and confirmed these transcripts all exhibited some level of polymorphism (Anderson and Knight; data not shown). Gastric lysozymes *1A* and *4A* were selected for further analysis in the multiplex qRT-PCR assay (Table 3). This analysis confirmed that transcripts for lysozymes *1A* and *4A* were both highly expressed in the naïve abomasum



and were significantly ($P \leq 0.05-0.005$) down-regulated in immune versus naïve sheep at days 2 and 5 post-challenge, although these differences became less apparent from day 10 onwards as levels in the naïve animals reduced (Figure 7).

MALAT-1

Metastasis-associated lung adenocarcinoma transcript 1 (*MALAT-1*) was identified as among the most significantly ($FDR \leq 0.05$) down-regulated transcripts in immune sheep on day 2 post-challenge (-1.8-2 fold), compared to naïve animals or day 0 (Table 4 and Figure 3g). This observation from the array data was confirmed by the multiplex qRT-PCR analysis, with significantly reduced levels of *MALAT-1* transcripts in immune versus naïve animals at day 0 and day 2 post-challenge (Figure 7). *MALAT-1* is understood to be a novel non-coding RNA which is highly conserved across species and up-regulated during metastasis, but its function is unknown to date [39]. We also observed transcripts for "boundary element associated factor" as the most highly down-regulated transcript in immune versus naïve animals at day 5 post-challenge (-6.5 fold, Table 5); again the function of this gene product is not clear but it

appears to be involved in transcriptional regulation by compartmentalization of the genome [40].

Discussion

The molecular mechanisms that contribute to protection from gastrointestinal nematodes have been extensively investigated in rodent models (e.g. reviews by [41-43]), but progress in understanding the responses to natural infections in ruminants, especially sheep, has been slower partly due to the limited range of appropriate species-specific reagents and genomics tools. Here we summarise findings using a novel ovine cDNA microarray, which is the first global transcriptomic analysis of ovine immune responses to *Teladorsagia circumcincta* challenge. Our data presented here are consistent with the observed increased adaptive and innate immune response taking place by days 2 and 5 post-challenge in the abomasal mucosa of sheep previously exposed to the parasite using this experimental protocol [11]. As discussed below, many of our findings highlight the importance of genes which have been investigated in rodent models of gastrointestinal nematode infection and previously highlighted in other ruminant studies, such as the expression changes in immunoglobulin





Figure 5 Results of competitive multiplex qRT-PCR analysis to investigate temporal changes in the expression levels of mucous-cell associated transcripts; *CLCA1* and *ITLNs* 1-3, as indicated, throughout the experiment trials summarised in Table 1. Open circles represent data from naïve sheep; closed triangles represent data from immune ("previously infected") sheep. Significant difference between Nv and Im at same timepoint; * $P \leq 0.05$, ** $P \leq 0.005$, *** $P \leq 0.005$; Significant difference between timepoint and day 0 of same group (where available); + $P \leq 0.05$, ++ $P \leq 0.005$, +++ $P \leq 0.005$ (Mann-Whitney U-test for non-parametric data).

transcripts, mucus and mast cell products, and members of the arachidonic acid pathway, and point to common mechanisms operating in response to nematode parasitism of both the gastric and intestinal mucosa. However, we have also highlighted novel changes such as altered expression levels of granulysin, gastric lysozymes, members of the matrix-metalloprotease family, and *STIP1*, which indicate responses that are unique to this model or have not been identified to date.

As reported previously, in all the trials (Table 1), sheep immunised by previous infection ("immune") had significantly ($P \leq 0.01$) lower worm burdens than naïve animals at the same timepoint, even at 48 h post-challenge [4,11]. Similarly, worms recovered from the immunised sheep showed higher levels of stunting [4]. These observations confirmed there were significantly increased anti-parasite responses in the immunised groups compared to naïve, and that these responses manifested themselves within 48 h of challenge. Previous work using materials from these infection trials indicated that there were clear histological and/or histochemical phenotypic differences elicited in the host

response between the "naïve" and "immune" groups, such as mucosal hypertrophy, mast cell and eosinophil counts, and in mucous composition ([8,44] and Craig et al., unpublished observations). In order to characterise these differences at a molecular level we examined changes in gene expression at the transcript level in the same experimental animals.

There were some discrepancies in this study between the findings from the cDNA arrays and the multiplex qRT-PCR results, which is likely to be due, in part, to the inability of some of the cDNA probes (+400bp) to distinguish between groups of highly similar genes. This is particularly obvious in the case of the highly similar family of ruminant gastric lysozyme genes [38]; only the specific RT-PCR/competitive qRT-PCR techniques would have been able to reliably distinguish between transcripts for lysozymes *1A* and *4A*, which are 92% identical at the DNA level. A similar explanation is likely to account for inconsistencies between the microarray and qRT-PCR findings in identifying changes in transcripts for members of the C1 complement family, and for the *ITLN* genes 1, 2 and 3, which were unlikely




to be distinguished by the *ITLN2* cDNA probe on the array; there were also temporal differences in expression between the three genes (Figure 5). These findings highlight the importance of validation of cDNA array data by more sensitive techniques, such as quantitative PCR-based analyses.

As immunity to *T. circumcincta* is associated with local IgA and IgE responses typical of gastrointestinal nematode infections [4,6] it is unsurprising that genes featuring the "immunoglobulin" descriptor were highly represented among the significantly up-regulated

transcripts in immune animals. Our findings for immunoglobulin transcripts are consistent both with an increased adaptive response in sheep previously exposed to the parasite, and with observations from other global analyses of responses to gastrointestinal nematode infection [45-47].

Our findings highlight the universal importance of mucus components in mammalian responses to parasitic nematode infection, such as *CLCA1*, members of the *ITLN* family and trefoil factor 3, which potentially contribute to expulsion by altering mucus composition and



making the environment of the parasite inhospitable. A similar pattern of gene expression has been identified in response of the abomasal mucosa to *Haemonchus contortus* infection [46]. There is a clear association between intestinal goblet cell hyperplasia, release of their effector molecules and altered mucus composition with the trapping and expulsion of gastrointestinal nematodes, mostly based on murine studies (reviewed by [43,48,49]). Sheep given trickle infection of *T. circumcincta* are known to exhibit hyperplasia of mucous-neck cells [14], which are phenotypically similar to goblet cells in the intestine, but this occurs as a later event in naïve sheep in response to challenge [10]. Previous work [44] has also demonstrated differences in PAS staining in naïve versus immune sheep ($P < 0.05$, Kruskal-Wallis at day 10 post-challenge) from the experimental samples used in this study, indicating altered abomasal mucous composition between the two groups.

CLCA1 (the putative orthologue of *CLCA3* in mice), which featured as one of the most consistently up-regulated transcripts in immune sheep, is secreted by goblet/mucous-neck cells in association with Th2-type inflammatory responses, such as murine models of pulmonary inflammation and gastro-intestinal nematode infection [50]. We have confirmed that *CLCA1* is up-regulated by IL-4 and IL-13 in a human goblet cell line [43], and by IL-4 in ovine gastric epithelial cells (Knight et al., in preparation). The significant up-regulation of transcripts for the mucous cell product trefoil factor 3 (*TFF3*) in immune animals compared to naïve was in contrast to *TFF2*, which is highly down-regulated (Tables 4 and 5). *TFF3* is IL-4/IL-13-regulated and interacts with the mucin MUC2 (not on array) to alter mucus viscosity, with which it co-localises in human intestinal goblet cells, and up-regulation of both these mucus components has been associated with responses to nematode infection in mice (reviewed by [43]). Up-regulation of

TFF3 and down-regulation of *TFF2* transcripts has also been demonstrated in the abomasal mucosa of sheep during the response to *H. contortus* [46,51,52], thus switching to a more "intestinal" phenotype in terms of TFF expression.

The multiplex qRT-PCR analyses showed the galactose-binding lectins *ITLNs* 1, 2 and 3 were significantly up-regulated in response to challenge in naïve sheep, with earlier up-regulation of *ITLN3* than *ITLNs* -1 and -2 (Figure 5). *ITLN* protein is localised to mucous neck cells in the abomasal mucosa, and highly up-regulated in infected compared to worm-free naïve sheep [27,35]. We also observed infection-associated up-regulation of transcripts for two other galactose-binding lectins; galectin 1, which is thought to bind gastrointestinal mucins [53], and galectin 4, which has been shown to be localised to gastric mucous cells in mice [54]. There is a clear temporal association between expression of members of the intelectin family by murine intestinal goblet cells and resistance to gastrointestinal nematode infection in mice [55,56]. The *ITLN* family, like *CLCA1*, are highly regulated by Th2 cytokines in murine and human goblet cells [43], and this has also been confirmed in ovine tracheal goblet cells [57] and gastric epithelial cultures (Knight et al, in preparation). Both *CLCA1* and intelectin protein were up-regulated in mucosal washings from previously infected, but not naïve, animals [58] indicating they act concordantly as part of an increased Th2-type response to alter mucus composition. It is possible that members of both the intelectin and galectin family are involved in interaction with mucins to change the rheological properties or adhesiveness of the mucus, and/or adhere to the worms to exert effector function or target them for immune clearance.

Significant up-regulation of the high affinity IgE receptor in immune animals day 2 and day 5 post-challenge,

along with up-regulation of cathepsin C, is consistent with an increased mast cell response in this group. T-cell mediated mucosal mast cell hyperplasia, accompanied by release of mast-cell mediators, is a characteristic feature of gastrointestinal nematode infections in mammals (reviewed by [43,59]). While numbers of mast cells (globule leucocytes) in the ovine abomasal mucosa are normally very low in uninfected sheep, they markedly increase in response to *T. circumcincta* infection in sensitised sheep, accompanied by release of sheep mast cell protease (SMCP) [13,60]. An assessment of the samples used in this study confirmed that the immune group had significantly higher numbers of mast cells than the naïve group at days 0, 2, 5 and 10 post-challenge, with the naïve animals showing very little increase in mucosal mast cell numbers in response to challenge ([8] and Craig et al., unpublished data).

We see evidence of up-regulation of a number of transcripts associated with eicosanoid metabolism, in particular *PLA2G2A*, which can initiate and regulate inflammation [61]. Increased synthesis of phospholipase and other members of the arachidonic acid cascade have been associated with intestinal nematode infection in rodents, cattle and swine [62-66], and so are likely to play a common key role in the induction and/or regulation of nematode-induced allergic inflammation; we would postulate a similar set of events occurring in the ovine abomasal mucosa in response to parasite challenge. Members of the complement cascade also appear to be up-regulated in response to *T. circumcincta* challenge; up-regulation of the complement C1 and C4 has similarly been linked to resistance to intestinal nematode infection in cattle [65]. Complement activation serves to initiate and propagate pro-inflammatory responses, although the role of complement in immunity to parasitic helminths is unclear [67]. Proteomic analysis has indicated complement and other plasma proteins are up-regulated in the mucosa of *T. circumcincta* infected versus naïve sheep, although these are likely to be derived from plasma rather than local expression [58].

We observed significant up-regulation of the chemokines *CCL2* and *CCL5* in response to larval challenge (Table 4, Figures 3c) and 6). *CCL2*, which is produced by a range of cell types including mast cells [68] has been shown to be released locally in response to gastrointestinal nematode infection in mice, and has been implicated in resistance to *Trichuris muris* by steering towards a Th2-type response [69]. *CCL5* is chemotactic for T cells, eosinophils, and basophils and implicated in a wide range of inflammatory diseases [70]. A previous RT-PCR-based analysis has shown that the abomasal mucosa is a source for a range of chemokine transcripts, and that *CCL5* transcripts are up-regulated in response

to *T. circumcincta* challenge (Griffith et al., unpublished observations).

One of the most surprising findings from this study, which has not been evidenced from rodent models, is the striking up-regulation of transcripts for granulysin in immune animals, which is normally associated with antimicrobial activity [71], along with other transcripts (granzymes, cathepsin C) normally associated with lymphocytes exhibiting cytotoxicity. Granulysin and NK-lysin are secreted, antimicrobial lipid-binding proteins belonging to the saposin family [71]. They have been identified in human, swine and bovine cytotoxic T lymphocytes and natural killer (NK) cells [72,73], but no murine counterpart has been identified to date. Granulysin is active against a broad range of intracellular and extracellular microbes [74], and granulysin activity has also been associated with apoptosis and necrosis of keratinocytes [75]. Granzymes A, B and H, like granulysin, are also normally associated with the activity of cytotoxic T-cells and NK cells [76], while cathepsin C is required for the processing and correct functioning of granzymes A and B and mast cell proteinases [77]. Interestingly, the pattern of expression of granulysin transcripts did not closely parallel that of the granzymes and cathepsin C (Figure 4), which suggests a different cellular source in the abomasal mucosa. This is the first time this molecule has been associated with gastrointestinal nematode infection, and the first association of this molecule with ovine disease. We have recently confirmed increased levels of granulysin protein in the abomasal mucosa of immune, but not naïve, sheep, and the potential biological role of granulysin in nematode infection is under further investigation (Griffith et al, in preparation).

Expression levels of transcripts for a number of heat shock proteins were significantly altered in the microarray analyses; in particular *STIP1*, which facilitates the association of the HSP70/90 complex which is implicated in the folding/regulation of a range of signalling proteins [78], was significantly increased in response to challenge (Figure 6). Host heat shock proteins are expressed constitutively in all cells but synthesis is increased in response to certain stressors or infection [79]. There has been little evidence to identify a direct role for heat shock proteins in gastrointestinal nematode infection, although *T. spiralis* has been shown to elicit host heat shock protein production during muscle migration [80]. The up-regulation and/or release of heat shock proteins is likely to be solely the effect of cellular damage caused to the mucosa by the parasite, but their release may also activate immune cells or facilitate antigen presentation [79,81].

Both the microarray and multiplex qRT-PCR analysis demonstrated significant down-regulation of transcripts

for *MMP-13*, as well as apparently high levels of cystatin C transcript expression. In a separate study (Knight et al., unpublished data), we have detected markedly decreased levels of transcripts for *MMP-7* and *TIMP-1* (tissue inhibitor of metalloproteinases) in immune versus naïve sheep exposed to *T. circumcincta* (*MMP-7* and *TIMP-1* were not this on array). Matrix metalloproteinases are implicated in a wide range of processes including regulation of inflammatory responses such as modulating response to cytokines and activation of β -defensins, as well as tissue dissolution/remodelling [82,83] and have been associated with T-cell mediated tissue damage to the gut mucosa [84]. The apparently higher levels of these transcripts in naïve compared to immune sheep may reflect increased tissue damage that is generally observed in the abomasa of animals that have not built up a level of immunity to the parasite. It is also possible that they have a role in inflammatory processes/tissue remodelling that is “replaced” in previously exposed sheep by mast cell proteases, which may have a similar function [85], following expansion of the mucosal mast cell population. The altered profile in matrix metalloproteinases may also have implications in interaction with incoming larvae, as helminth parasites are understood to produce proteinase inhibitors to protect themselves from degradation by host proteinases [86], and parasites of the gastrointestinal tract can interact with host proteolytic pathways with immunomodulatory effects [87,88]. Furthermore, *T. circumcincta* larvae abundantly produce the cysteine protease cathepsin F [89], which raises the possibility that high local levels of cystatin C could serve a protective function. The potential interaction of these matrix-metalloproteinases and protease inhibitors with proteases or inhibitors produced by incoming larvae would merit further investigation.

The most consistently down-regulated transcripts in immune versus naïve sheep detected at day 5 post-challenge were members of the ruminant gastric lysozyme family. It should be noted that the array contained more than 50 spots representing members of this family which were down-regulated in the day 5 immune vs. naïve comparison, and 5 spots in the day 5 naïve vs. day 0 naïve comparison. The large numbers of significant spots may be partly due to the overrepresentation of the lysozyme family on the cDNA chip, having a high proportion of sequences in the KN511 library (2.5%) [90]. There are four highly similar ruminant gastric lysozyme genes, which are thought to have evolved from the lysozyme C gene family by gene-duplication events [38]. They are highly expressed in the ovine abomasum and are thought to act as a major digestive enzyme, functional at low pH, for the large amounts of bacteria entering from the

rumen, by breaking down peptidoglycan cell walls which cannot be hydrolysed by conventional digestive enzymes. This analysis confirmed that transcripts for lysozymes *1A* and *4A* were both highly expressed in the naïve abomasum and significantly down-regulated in immune versus naïve sheep at days 2 and 5 post-challenge, although these differences became less apparent from day 10 onwards as levels in the naïve animals reduced (Figure 7). This pattern is in contrast to a parallel proteomics study, where lysozyme *4A* increased in previously infected animals compared to naïve at day 0 and day 2 post-challenge (Brown, Pemberton et al., in preparation) and members of the lysozyme family were found to be up-regulated in abomasal epithelial extracts and mucosal washes from previously infected versus naïve sheep [58]. This disparity is likely to be due to differing rates of transcription, processing and storage of the gastric lysozymes prior to secretion. Sheep parasitized by *T. circumcincta* show diminution of parietal cell number, raised abomasal pH and hypergastrinaemia [14,91], so we could speculate that the alterations in lysozyme production, and the possible inability of lysozymes to function in the raised pH of the abomasum, both reflect these changes and may contribute to the resultant nutritional loss seen in infected animals.

In summary, the up-regulated expression of mucous-cell related transcripts such as *CLCA1*, *TFE3* and *ITLN*s, as well as transcripts for the high affinity IgE receptor, *CCL2*, and members of the arachidonic acid pathway in response to ovine gastric nematode infection, parallel findings from mouse models of intestinal nematode infection by ourselves and others [43,56,92]. Many of these changes reflect a common Th2-driven immune response occurring across species and in both areas of the gastrointestinal tract. However, other findings point to novel changes, such as the distinct local up-regulation of granulysin in immune sheep, a gene which is present in humans, cattle and pigs but not mice, and the alteration in the gastric lysozyme profile which is unique to ruminants.

As can be seen from the pattern of expression of the gastric lysozymes discussed previously, it needs to be borne in mind that changes in the transcriptome may not directly reflect contemporaneous changes in local levels of the corresponding protein, many of which are stored before subsequent release into the local microenvironment. Many of the changes described here are likely to reflect the pronounced cellular changes in the abomasal mucosa induced by exposure to *T. circumcincta* larvae. Nevertheless, the data presented here gives valuable insights into some of the molecular mechanisms that may be operating at the ovine host-parasite interface to control gastric nematode infection.

Additional material

Additional file 1: Table S1 Details of primers and PCR conditions used for RT-PCR analysis. Details of primers and PCR conditions used for RT-PCR analysis, and sequences used to design PCR probes. **Table S2** Total numbers of genes whose expression levels were significantly altered detected in each hybridisation experiment Total numbers of genes whose expression levels were significantly altered (Benjamini & Hochberg FDR ≤ 0.05) detected in each hybridisation experiment; (Table S2a: Expt.1; day 5 post-challenge and Table S2b: Expt. 2; day 2 post-challenge) including dye interactions.

Additional file 2: Table S3 Significantly highly represented networks identified using Ingenuity Pathways Analysis software (Ingenuity® Systems, [26]). Networks that were significantly highly represented ($P \leq 10^{-10}$; Fischer's exact test) identified from the lmd0/Nvd0 and lmd2/lmd0 datasets from Expt. 2 (see Table II), using Ingenuity Pathways Analysis software (Ingenuity® Systems, [26]). **Table S4** Canonical pathways identified using Ingenuity Pathways Analysis software (Ingenuity® Systems, <http://www.ingenuity.com>). The top 20 most significant canonical pathways from the lmd0/Nvd0 and lmd2/lmd0 datasets from Expt. 2 (See Table II), identified using Ingenuity Pathways Analysis software (Ingenuity® Systems, [26]), are shown.

Acknowledgements

We thank Steve Smith and Aileen Halliday (MRI) for assistance with and access to tissue samples collected at post-mortem and Sequenom Inc, San Diego, CA for running the multiplex qRT-PCR assays (Sequenom® MassARRAY® System). This work was funded by DEFRA/SFC (VT0102) and the BBSRC (BB/E01867X/1).

Author details

¹The Roslin Institute and R(D)SVS, University of Edinburgh, Roslin, Midlothian, EH25 9RG, Scotland, UK. ²Moredu Research Institute, Pentlands Science Park, Bush Loan, Penicuik, Midlothian, EH26 0PZ, Scotland, UK.

Authors' contributions

PK planned and supervised the study and drafted the manuscript, SG carried out the sequencing and RT-PCR analysis of granulysin, JP assisted with post-mortem sample collection and carried out RNA extractions/QC checks, AP assisted with interpretation of data and participated in manuscript preparation, LG carried out the sequence analysis of MMPs and inhibitors, KA carried out the sequence analysis of the lysozyme family, RT supervised the microarray experiments, SS carried out the hybridisations, DW carried out statistical analysis of the microarray experiments, MF carried out annotation of the ARK-Genomics *O. aries* 12K v1.0 array and assisted with data processing, AA supervised the microarray experiments and advised on manuscript preparation, SB carried out the Ingenuity analysis and assisted with presentation of the output, DS supervised the experimental infections and advised on manuscript preparation, HM and IM conceived and obtained the funding for the VTRI overall project and HM advised on manuscript preparation. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 4 October 2010 Accepted: 17 June 2011

Published: 17 June 2011

References

1. Bartley DJ, Jackson F, Jackson E, Sargison N: Characterisation of two triple resistant field isolates of *Teladorsagia* from Scottish lowland sheep farms. *Vet Parasitol* 2004, **123**:189-199.
2. Smith WD, Jackson F, Jackson E, Williams J: Local immunity and *Ostertagia circumcincta*: changes in the gastric lymph of immune sheep after a challenge infection. *J Comp Pathol* 1983, **93**:479-488.
3. Smith WD, Jackson F, Jackson E, Williams J, Miller HR: Manifestations of resistance to ovine ostertagiasis associated with immunological responses in the gastric lymph. *J Comp Pathol* 1984, **94**:591-601.
4. Halliday AM, Routledge CM, Smith SK, Matthews JB, Smith WD: Parasite loss and inhibited development of *Teladorsagia circumcincta* in relation to the kinetics of the local IgA response in sheep. *Parasite Immunol* 2007, **29**:425-434.
5. Smith WD, Jackson F, Jackson E, Graham R, Williams J, Willadsen SM, Fehilly CB: Transfer of immunity to *Ostertagia circumcincta* and IgA memory between identical sheep by lymphocytes collected from gastric lymph. *Res Vet Sci* 1986, **41**:300-306.
6. Stear MJ, Strain S, Bishop SC: How lambs control infection with *Ostertagia circumcincta*. *Vet Immunol Immunopathol* 1999, **72**:213-218.
7. Huntley JF, Schallig HD, Kooyman FN, Mackellar A, Jackson F, Smith WD: IgE antibody during infection with the ovine abomasal nematode, *Teladorsagia circumcincta*: primary and secondary responses in serum and gastric lymph of sheep. *Parasite Immunol* 1998, **20**:565-571.
8. Craig NM: Cytokine gene expression in naive and previously infected sheep and lambs after challenge with the abomasal nematode *Teladorsagia circumcincta*. PhD thesis University of Edinburgh, UK; 2010.
9. Craig NM, Miller HR, Smith WD, Knight PA: Cytokine expression in naive and previously infected lambs after challenge with *Teladorsagia circumcincta*. *Vet Immunol Immunopathol* 2007, **120**:47-54.
10. Scott I, Khalaf S, Simcock DC, Knight CG, Reynolds GW, Pomroy WE, Simpson HV: A sequential study of the pathology associated with the infection of sheep with adult and larval *Ostertagia circumcincta*. *Vet Parasitol* 2000, **89**:79-94.
11. Halliday AM, Morrison WI, Smith WD: Kinetics of the local cellular response in the gastric lymph of immune and susceptible sheep to infection with *Teladorsagia circumcincta*. *Parasite Immunol* 2009, **31**:402-411.
12. Huntley JF, Jackson F, Coop RL, Macalodow C, Houdijk JG, Familton AS, Xieh HL, Stankiewicz M, Sykes AR: The sequential analysis of local inflammatory cells during abomasal nematode infection in periparturient sheep. *Vet Immunol Immunopathol* 2004, **97**:163-176.
13. Balic A, Bowles VM, Liu YS, Meeusen EN: Local immune responses in sensitized sheep following challenge infection with *Teladorsagia circumcincta*. *Parasite Immunol* 2003, **25**:375-381.
14. Scott I, Hodgkinson SM, Khalaf S, Lawton DE, Collett MG, Reynolds GW, Pomroy WE, Simpson HV: Infection of sheep with adult and larval *Ostertagia circumcincta*: abomasal morphology. *Int J Parasitol* 1998, **28**:1383-1392.
15. Miller HR, Jackson F, Newlands G, Appleyard WT: Immune exclusion, a mechanism of protection against the ovine nematode *Haemonchus contortus*. *Res Vet Sci* 1983, **35**:357-363.
16. Miller HRP, Jackson F, Newlands GFJ, Huntley JF: Rapid expulsion of gastrointestinal nematodes in sheep: a role for immediate hypersensitive reactions in the mucosa. In *Immunology of the Sheep*. Edited by: Morris B, Myasaka M. Roche, Basle; 1986:460-482.
17. Knight PA, Pate J, Smith WD, Miller HR: An ovine chitinase-like molecule, chitinase-3 like-1 (YKL-40), is up-regulated in the abomasum in response to challenge with the gastrointestinal nematode, *Teladorsagia circumcincta*. *Vet Immunol Immunopathol* 2007, **120**:55-60.
18. ARK-Genomics: Centre for Comparative & Functional Genomics website. [<http://www.ark-genomics.org>].
19. Smith S, et al: ARK-Genomics Conference 2008: 3rd International Symposium on Animal Functional Genomics, ISAFG-P71. [<http://www.ark-genomics.org/downloads/ISAFG-2008-Abstracts.pdf>].
20. Baird D, Johnstone P, Wilson T: Normalization of microarray data using a spatial mixed model analysis which includes splines. *Bioinformatics* 2004, **20**:3196-3205.
21. Smyth GK, Speed T: Normalization of cDNA microarray data. *Methods* 2003, **31**:265-273.
22. Smyth GK: Limma: linear models for microarray data. In *Bioinformatics and Computational Biology Solutions using R and Bioconductor Gentleman R*. Edited by: Carey V, Dudoit S, Irizarry R, Huber W. Springer, New York; 2005:397-420.
23. Smyth GK: Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004, **3**:Article3.
24. Benjamini Y, Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Statist Soc B* 1995, **57**:289-300.
25. Jaffrezic F, de Koning DJ, Boettcher PJ, Bonnet A, Buitenhuis B, Closset R, Dejean S, Delmas C, Detilleux JC, Dovc P, Duval M, Foulley JL, Hedegaard J,

- Hornshøj H, Hulsegge I, Janss L, Jensen K, Jiang L, Lavric M, Le Cao KA, Lund MS, Malinverni R, Marot G, Nie H, Petzl W, Pool MH, Robert-Granie C, San Cristobal M, van Schothorst EM, Schubert HJ, Sorensen P, Stella A, Tosser-Klopp G, Waddington D, Watson M, Yang W, Zerbe H, Seyfert HM: **Analysis of the real EADGENE data set: comparison of methods and guidelines for data normalisation and selection of differentially expressed genes (open access publication).** *Genet Sel Evol* 2007, **39**:633-650.
26. Ingenuity Systems, Inc. website. [<http://www.ingenuity.com>].
27. French AT, Knight PA, Smith WD, Brown JK, Craig NM, Pate JA, Miller HR, Pemberton AD: **Up-regulation of intelectin in sheep after infection with *Teladorsagia circumcincta*.** *Int J Parasitol* 2008, **38**:467-475.
28. The National Center for Biotechnology Information primer designing tool Primer3 and BLAST. [<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>].
29. European Bioinformatics Institute Tools for Sequence Analysis. [<http://www.ebi.ac.uk/Tools/sequence.html>].
30. The National Center for Biotechnology Information website. [<http://www.ncbi.nlm.nih.gov>].
31. The Ensembl Genome Browser. [<http://www.ensembl.org/>].
32. The Sequenom® Gene Expression Analysis website. [<http://www.sequenom.com/Genetic-Analysis/Applications/QGE-Gen-Expression>].
33. Andell Y, *et al*: **Multiplexed Gene Expression Analysis Using Competitive PCR and MassARRAY™.** [http://www.sequenom.com/getdoc/564717df-0437-4701-8f60-78ec61d95d09/Multiplexed_Gene_Expression_Analysis].
34. The ArrayExpress Archive. [<http://www.ebi.ac.uk/arrayexpress/>], - Accession numbers: Expt. 1: E-TABM-1149; Expt. 2: E-TABM-1150.
35. French AT, Knight PA, Smith WD, Pate JA, Miller HR, Pemberton AD: **Expression of three intelectins in sheep and response to a Th2 environment.** *Vet Res* 2009, **40**:53.
36. McAleese SM, Pemberton AD, McGrath ME, Huntley JF, Miller HR: **Sheep mast-cell proteinases-1 and -3: cDNA cloning, primary structure and molecular modelling of the enzymes and further studies on substrate specificity.** *Biochem J* 1998, **333**:801-809.
37. Pemberton AD, McAleese SM, Huntley JF, Collie DD, Scudamore CL, McEuen AR, Walls AF, Miller HR: **cDNA sequence of two sheep mast cell tryptases and the differential expression of tryptase and sheep mast cell proteinase-1 in lung, dermis and gastrointestinal tract.** *Clin Exp Allergy* 2000, **30**:818-832.
38. Wen Y, Irwin DM: **Mosaic evolution of ruminant stomach lysozyme genes.** *Mol Phylogenet Evol* 1999, **13**:474-482.
39. Ji P, Diederichs S, Wang W, Boing S, Metzger R, Schneider PM, Tidow N, Brandt B, Buerger H, Bulk E, Thomas M, Berdel WE, Serve H, Muller-Tidow C: **MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer.** *Oncogene* 2003, **22**:8031-8041.
40. Pathak RU, Rangaraj N, Kallappagoudar S, Mishra K, Mishra RK: **Boundary element-associated factor 32B connects chromatin domains to the nuclear matrix.** *Mol Cell Biol* 2007, **27**:4796-4806.
41. Artis D, Grencis RK: **The intestinal epithelium: sensors to effectors in nematode infection.** *Mucosal Immunol* 2008, **1**:252-264.
42. Nair MG, Guild KJ, Artis D: **Novel effector molecules in type 2 inflammation: lessons drawn from helminth infection and allergy.** *J Immunol* 2006, **177**:1393-1399.
43. Knight PA, Brown JK, Pemberton AD: **Innate immune response mechanisms in the intestinal epithelium: potential roles for mast cells and goblet cells in the expulsion of adult *Trichinella spiralis*.** *Parasitology* 2008, **135**:655-670.
44. French AT: **The co-regulation of the mucus associated molecules intelectin, resistin like molecule beta and beta galactoside alpha 2,3 sialyl transferase in a T helper celltype 2 response.** *PhD thesis* University of Edinburgh, UK; 2008.
45. Diez-Tascon C, Keane OM, Wilson T, Zadissa A, Hyndman DL, Baird DB, McEwan JC, Crawford AM: **Microarray analysis of selection lines from outbred populations to identify genes involved with nematode parasite resistance in sheep.** *Physiol Genomics* 2005, **21**:59-69.
46. Rowe A, Gondro C, Emery D, Sangster N: **Sequential microarray to identify timing of molecular responses to *Haemonchus contortus* infection in sheep.** *Vet Parasitol* 2009, **161**:76-87.
47. Andronicos N, Hunt P, Windon R: **Expression of genes in gastrointestinal and lymphatic tissues during parasite infection in sheep genetically resistant or susceptible to *Trichostrongylus colubriformis* and *Haemonchus contortus*.** *Int J Parasitol* 2010, **40**:417-429.
48. de Veer MJ, Kemp JM, Meeusen EN: **The innate host defence against nematode parasites.** *Parasite Immunol* 2007, **29**:1-9.
49. Miller HRP, Knight PA, Pemberton AD: **Mucus; modulation by the TH2 response to enhance an innate defensive barrier against gut nematodes.** *Parasite Immunol* 2006, **28**:259-262.
50. Gibson A, Lewis AP, Affleck K, Aitken AJ, Meldrum E, Thompson N: **hCLCA1 and mCLCA3 are secreted non-integral membrane proteins and therefore are not ion channels.** *J Biol Chem* 2005, **280**:27205-27212.
51. Dunphy JL, Balic A, Barcham GJ, Horvath AJ, Nash AD, Meeusen EN: **Isolation and characterization of a novel inducible mammalian galectin.** *J Biol Chem* 2000, **275**:32106-32113.
52. Ingham A, Reverter A, Windon R, Hunt P, Menzies M: **Gastrointestinal nematode challenge induces some conserved gene expression changes in the gut mucosa of genetically resistant sheep.** *Int J Parasitol* 2008, **38**:431-442.
53. Wasano K, Hirakawa Y: **Recombinant galectin-1 recognizes mucin and epithelial cell surface glycolycales of gastrointestinal tract.** *J Histochem Cytochem* 1997, **45**:275-283.
54. Nio-Kobayashi J, Takahashi-Iwanaga H, Iwanaga T: **Immunohistochemical localization of six galectin subtypes in the mouse digestive tract.** *J Histochem Cytochem* 2009, **57**:41-50.
55. Pemberton AD, Knight PA, Gamble J, Colledge WH, Lee JK, Pierce M, Miller HR: **Innate BALB/c enteric epithelial responses to *Trichinella spiralis*: inducible expression of a novel goblet cell lectin, intelectin-2, and its natural deletion in C57BL/10 mice.** *J Immunol* 2004, **173**:1894-1901.
56. Datta R, deSchoolmeester ML, Hedeler C, Paton NW, Brass AM, Else KJ: **Identification of novel genes in intestinal tissue that are regulated after infection with an intestinal nematode parasite.** *Infect Immun* 2005, **73**:4025-4033.
57. French AT, Bethune JA, Knight PA, McNeilly TN, Wattegedera S, Rhind S, Miller HR, Pemberton AD: **The expression of intelectin in sheep goblet cells and up-regulation by interleukin-4.** *Vet Immunol Immunopathol* 2007, **120**:41-46.
58. Athanasiadou S, Pemberton A, Jackson F, Inglis N, Miller HR, Thevenod F, Mackellar A, Huntley JF: **Proteomic approach to identify candidate effector molecules during the *in vitro* immune exclusion of infective *Teladorsagia circumcincta* in the abomasum of sheep.** *Vet Res* 2008, **39**:58.
59. Miller HR: **Mucosal mast cells and the allergic response against nematode parasites.** *Vet Immunol Immunopathol* 1996, **54**:331-336.
60. Huntley JF, Gibson S, Brown D, Smith WD, Jackson F, Miller HR: **Systemic release of a mast cell proteinase following nematode infections in sheep.** *Parasite Immunol* 1987, **9**:603-614.
61. Gilroy DW, Newson J, Sawmynaden P, Willoughby DA, Croxtall JD: **A novel role for phospholipase A2 isoforms in the checkpoint control of acute inflammation.** *FASEB J* 2004, **18**:489-498.
62. Larsh JE Jr: ***Trichinella spiralis*: phospholipase in sensitized mice after challenge.** *Exp Parasitol* 1975, **37**:233-238.
63. Brown JK, Knight PA, Thornton EM, Pate JA, Coonrod S, Miller HR, Pemberton AD: ***Trichinella spiralis* induces de novo expression of group IVC phospholipase A2 in the intestinal epithelium.** *Int J Parasitol* 2008, **38**:143-147.
64. Goven AJ, DeBuysscher EV: **Intestinal phospholipase B activity in swine inoculated with *Trichinella spiralis*.** *Am J Vet Res* 1979, **40**:1469-1471.
65. Araujo RN, Padilha T, Zarlenga D, Sonstegard T, Connor EE, Van Tassel C, Lima WS, Nascimento E, Gasbarre LC: **Use of a candidate gene array to delineate gene expression patterns in cattle selected for resistance or susceptibility to intestinal nematodes.** *Vet Parasitol* 2009, **162**:106-115.
66. Li RW, Gasbarre LC: **A temporal shift in regulatory networks and pathways in the bovine small intestine during *Cooperia oncophora* infection.** *Int J Parasitol* 2009, **39**:813-824.
67. Giacomin PR, Gordon DL, Botto M, Daha MR, Sanderson SD, Taylor SM, Dent LA: **The role of complement in innate, adaptive and eosinophil-dependent immunity to the nematode *Nippostrongylus brasiliensis*.** *Mol Immunol* 2008, **45**:446-455.
68. Brown JK, Knight PA, Wright SH, Thornton EM, Miller HR: **Constitutive secretion of the granule chymase mouse mast cell protease-1 and the chemokine, CCL2, by mucosal mast cell homologues.** *Clin Exp Allergy* 2003, **33**:132-146.

69. de Schoolmeester ML, Little MC, Rollins BJ, Else KJ: **Absence of CC chemokine ligand 2 results in an altered Th1/Th2 cytokine balance and failure to expel *Trichuris muris* infection.** *J Immunol* 2003, **170**:4693-4700.
70. Navratilova Z: **Polymorphisms in CCL2&CCL5 chemokines/chemokine receptors genes and their association with diseases.** *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2006, **150**:191-204.
71. Bruhn H: **A short guided tour through functional and structural features of saposin-like proteins.** *Biochem J* 2005, **389**:249-257.
72. Endsley JJ, Furrer JL, Endsley MA, McIntosh MA, Maue AC, Waters WR, Lee DR, Estes DM: **Characterization of bovine homologues of granulysin and NK-lysin.** *J Immunol* 2004, **173**:2607-2614.
73. Endsley JJ, Furrer JL, McIntosh MA, Estes DM: **Antimicrobial activity of a bovine granulysin homologue.** *Faseb J* 2003, **17**:C171-C171.
74. Stenger S, Hanson DA, Teitelbaum R, Dewan P, Niazi KR, Froelich CJ, Ganz T, Thoma-Uszynski S, Melian A, Bogdan C, Porcelli SA, Bloom BR, Krensky AM, Modlin RL: **An antimicrobial activity of cytolytic T cells mediated by granulysin.** *Science* 1998, **282**:121-125.
75. Chung WH, Hung SI, Yang JY, Su SC, Huang SP, Wei CY, Chin SW, Chiou CC, Chu SC, Ho HC, Yang CH, Lu CF, Wu JY, Liao YD, Chen YT: **Granulysin is a key mediator for disseminated keratinocyte death in Stevens-Johnson syndrome and toxic epidermal necrolysis.** *Nat Med* 2008, **14**:1343-1350.
76. Chowdhury D, Lieberman J: **Death by a thousand cuts: granzyme pathways of programmed cell death.** *Annu Rev Immunol* 2008, **26**:389-420.
77. Meade JL, de Wynter EA, Brett P, Sharif SM, Woods CG, Markham AF, Cook GP: **A family with Papillon-Lefevre syndrome reveals a requirement for cathepsin C in granzyme B activation and NK cell cytolytic activity.** *Blood* 2006, **107**:3665-3668.
78. Daniel S, Bradley G, Longshaw VM, Soti C, Csermely P, Blatch GL: **Nuclear translocation of the phosphoprotein Hop (Hsp70/Hsp90 organizing protein) occurs under heat shock, and its proposed nuclear localization signal is involved in Hsp90 binding.** *Biochim Biophys Acta* 2008, **1783**:1003-1014.
79. Wallin RP, Lundqvist A, More SH, von Bonin A, Kiessling R, Ljunggren HG: **Heat-shock proteins as activators of the innate immune system.** *Trends Immunol* 2002, **23**:130-135.
80. Martinez J, Perez Serrano J, Bernadina WE, Rodriguez-Caabeiro F: **Influence of parasitization by *Trichinella spiralis* on the levels of heat shock proteins in rat liver and muscle.** *Parasitology* 1999, **118**:201-209.
81. Stewart GR, Young DB: **Heat-shock proteins and the host-pathogen interaction during bacterial infection.** *Curr Opin Immunol* 2004, **16**:506-510.
82. Burke B: **The role of matrix metalloproteinase 7 in innate immunity.** *Immunobiology* 2004, **209**:51-56.
83. Parks WC, Wilson CL, López-Boado YS: **Matrix metalloproteinases as modulators of inflammation and innate immunity.** *Nat Rev Immunol* 2004, **4**:617-629.
84. Di Sabatino A, Pickard KM, Rampton D, Kruidenier L, Rovedatti L, Leakey NA, Corazza GR, Monteleone G, MacDonald TT: **Blockade of transforming growth factor beta upregulates T-box transcription factor T-bet, and increases T helper cell type 1 cytokine and matrix metalloproteinase-3 production in the human gut mucosa.** *Gut* 2008, **57**:605-612.
85. Miller HR, Pemberton AD: **Tissue-specific expression of mast cell granule serine proteinases and their role in inflammation in the lung and gut.** *Immunology* 2002, **105**:375-390.
86. Knox DP: **Proteinase inhibitors and helminth parasite infection.** *Parasite Immunol* 2007, **29**:57-71.
87. Antalis TM, Shea-Donohue T, Vogel SN, Sears C, Fasano A: **Mechanisms of disease: protease functions in intestinal mucosal pathobiology.** *Nat Clin Pract Gastroenterol Hepatol* 2007, **4**:393-402.
88. Hewitson JP, Grainger JR, Maizels RM: **Helminth immunoregulation: the role of parasite secreted proteins in modulating host immunity.** *Mol Biochem Parasitol* 2009, **167**:1-11.
89. Redmond DL, Smith SK, Halliday A, Smith WD, Jackson F, Knox DP, Matthews JB: **An immunogenic cathepsin F secreted by the parasitic stages of *Teladorsagia circumcincta*.** *Int J Parasitol* 2006, **36**:277-286.
90. The National Center for Biotechnology Information Library browser: Ovis aries. [<http://www.ncbi.nlm.nih.gov/UniGene/lbrowse2.cgi?TAXID=9940>].
91. Simpson HV: **Pathophysiology of abomasal parasitism: is the host or parasite responsible?** *Vet J* 2000, **160**:177-191.
92. Knight PA, Pemberton AD, Robertson KA, Roy DJ, Wright SH, Miller HR: **Expression profiling reveals novel innate and inflammatory responses in the jejunal epithelial compartment during infection with *Trichinella spiralis*.** *Infect Immun* 2004, **72**:6076-6086.

doi:10.1186/1297-9716-42-78

Cite this article as: Knight *et al.*: Novel gene expression responses in the ovine abomasal mucosa to infection with the gastric nematode *Teladorsagia circumcincta*. *Veterinary Research* 2011 **42**:78.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

