

Review Article

Male sexual health and dysfunction

pISSN: 2287-4208 / eISSN: 2287-4690
World J Mens Health 2021 Jan 39(1): 31-37
<https://doi.org/10.5534/wjmh.190151>

The World Journal of
MEN'S HEALTH



Vitamin D and Male Erectile Function: An Updated Review

Onder Canguven^{1,2}, Ahmad H. Al Malki^{1,3}

¹Department of Urology, Hamad General Hospital, Hamad Medical Corporation, Doha, Qatar, ²Department of Urology, Weill Cornell Medicine, New York, NY, USA, ³Department of Urology, Qatar University, Doha, Qatar

Literature support that vitamin-D is important for different systems of the human body including, but not limited to endocrine and immune systems, vasculature and endothelial function of the body. Male erectile function depends on many factors and can be perceived as a health indicator of the body. Epidemiological data have shown that vitamin-D deficiency is also associated with erectile dysfunction. In this review, our aim is to interpret the mechanisms by which vitamin-D might regulate anatomy and physiology of penis. Evidence showed that vitamin-D is needed for an adequate erectile function. Briefly, vitamin-D is crucial for a better healthy body and sexual activity.

Keywords: Andrology; Endothelium; Erectile dysfunction; Hormones; Male; Vascular endothelium

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Human sexual activity is the manner in which humans experience and express their sexuality. Male penile erection is a physiological phenomenon in which the penis becomes firm, engorged, and enlarged in response to sexual arousal during sexual activity [1]. Penile erection is the result of a complex interaction of psychological, neural, vascular, and endocrine factors in the presence of a normal penis anatomy [2]. Penile erection is initiated by nitric oxide (NO), a potent vasodilator, released from the nerve endings, further production of NO depends on increased blood flow which is also known as shear stress [3]. In most vascular beds, the NO is released by a stimulus that increases the shear stress over the endothelium. Endothelial

dysfunction is a key variable in the pathogenesis of atherosclerosis and its complications, including erectile dysfunction (ED).

ED is defined as the inability to attain or maintain erection sufficient for satisfactory sexual activity [2]. Over the past 20 years, there have been numerous reports on the epidemiology of ED all over the world [4-7]. In all studies, it was demonstrated that the prevalence of ED increases with age [4,7]. In other words, higher age is strongly associated with higher incidence of ED [7,8]. In men under the aged 40 to 49 years, the prevalence of ED was relatively low (~40%) and in men aged 70 to 79 years, the prevalence of ED was significantly higher (~90%) [6,7,9]. The etiology of ED has been studied comprehensively and found to be multifactorial. ED is frequently found in vascular syndromes, such

Received: Nov 28, 2019 **Revised:** Dec 17, 2019 **Accepted:** Dec 24, 2019 **Published online** Jan 16, 2020

Correspondence to: Onder Canguven <https://orcid.org/0000-0002-1010-0609>

Department of Urology, Hamad General Hospital, Hamad Medical Corporation, 3050, Doha, Qatar.

Tel: +974-44391864, **Fax:** +974-44391842, **E-mail:** ocanguven@yahoo.com

as atherosclerotic cardiovascular (ASCV) diseases, hypertension, cerebrovascular disease, peripheral arterial disease and diabetes mellitus (DM) [8,10-12]. In fact, the penis is a highly vascularized organ and erections are primarily vascular events. Both ED and ASCV diseases have frequently identical functional and morphologic basis [3,12]. Because of its vascularity, ED is even accepted as a marker of coronary endothelial dysfunction and atherosclerosis [10,12].

In the last decades, several studies have correlated ED risk factors with vitamin-D (VD) deficiency (VDD) [5,13,14]. Observational and interventional studies published in literature have established a strong relationship between VD levels and ED. This important knowledge turned scientists to go in deep for the effect of VD on erectile function. Shortly, it is now well known that VD is a part of well-functioning penis starting from the early days of our lives.

VITAMIN-D

VD, which is actually a steroid hormone, has much more assorted role in human body for a vitamin than previously recognized just for calcium metabolism [15]. Since it was shown that VD-receptor (VDR) is expressed in all tissues of human body including penis, VD regulates cellular differentiation and function across many cell types [1,16-23]. VD effects are facilitated through binding to the VDR and stimulating not only genomic but also non-genomic effects [19,22]. Upon VD binding, VDR translocates from the plasma membrane to the nucleus where it transcriptionally activates genes *via* the VD response element, thereby affecting transcription of other genes [19]. Human Genome Project study results showed that a human DNA has between 20,000 and 25,000 genes [24], and more important to us, surprisingly, over 3,000 genes are responsive to VD [25].

Unfortunately, the estimated worldwide prevalence of VDD is really high and common among the elderly. VDD is about 50% at older ages and this result underlines the importance of VDD for public health [26,27]. If we consider that VD effect many tissues, VDD becomes more and more important for a healthy sexual life, in other words for erectile function.

1. Essentials of vitamin-D

Vitamins D2 (ergocalciferol) and D3 (cholecalciferol)

are two dominant forms of VD. Vitamin D2 is synthesized by invertebrates and plants after exposure to ultraviolet radiation [1]. Vitamin D3 is naturally present in a small range of foods (such as oily fish, egg and fortified dairy products) and is also made endogenously in the skin. Dietary VD typically comprises only 10% to 20% of circulating levels of VD [28]. If we look at the VD physiology, one can see that Vitamin D2 and D3 are biologically inactive and subjected to double hydroxylation, first mainly in the liver and then in the kidney, to produce the biologically active compound calcitriol [1,14,21,29,30]. Although $1,25(\text{OH})_2\text{D}$ is the most active form of VD and undertakes the major biological function, it has a shorter half-life (~0.3 and 25 days, respectively) and a 1,000-fold lower concentration than $25(\text{OH})\text{D}$ [31]. More importantly, serum $25(\text{OH})\text{D}$ is relatively stable and not directly influenced by diet (*e.g.*, calcium intake) and life style (*e.g.*, mobility) [31]. Hence, total-body VD stores are best reflected by $25(\text{OH})\text{D}$ [31]. Earlier, it was proved that availability of ultraviolet-B exposure, and its resultant VD production in skin, is highest in late spring through early fall and lowest from late fall through early spring [26]. In addition to seasonal variations, there are many other factors that can affect the prevalence of VDD. These can be summarized as region and latitude, diet and supplement use, clothing, obesity, smoking, concerns about sun damage, and the nature of built environment [32].

VITAMIN-D AND ERECTILE DYSFUNCTION

A general population study showed that low serum VD was associated with higher prevalence of peripheral arterial disease [33]. Likewise, a very recent cross-sectional analyses (3,390 men aged >20 years, free of ASCV disease) also reported that VDD was associated with an increased ED prevalence [14]. In support, deficient levels of VD <20 ng/mL were associated with increased ED risk, where a decreased prevalence of ED was associated with VD levels >35 ng/mL [14]. When researchers restricted their analysis to the 562 men (among 3,390 men) adjusted for sex hormone levels, the association of VD with ED became even stronger [14]. The biological link between VDD and ED exhibits several interlaced mechanisms that could suggest that the link of VD with ED appears to be independent of sex hormones [14].

1. Vitamin-D and architecture of the penis

There are many risk factors in the etiology of ED which might be directly arising from VDD. Since the latest literature showed the importance of VD on erectile function, Fernandes-Lima et al [34] investigated the effects of VD restricted diet on Wistar rats offspring penis morphology. Fernandes-Lima et al [34] demonstrated that VD restriction during perinatal and post-natal periods induced metabolic and structural changes and represented important risk factors for ED in the penis of the adult offspring. The latter findings suggest that VD is an important micronutrient in maintaining the cytoarchitecture of the penis [34]. In other words, human body requires VD for the proper anatomical development of the penis during embryonal life.

2. Vitamin-D and endothelial function

The pathophysiology of ED is multifactorial, but common point is a vascular disorder related to decreased endothelial function. Indeed, one of the most significant mechanism of VD on erectile function seems to work *via* endothelial integrity [30,35]. In molecular levels, VD stabilizes the quiescent endothelium, regulates certain stages of endothelial activation, and is involved in the repair of the damaged endothelium *in vitro* and *in vivo* models. A recent 12 cross-sectional studies, including 2,086 subjects of varying ethnic groups, showed an association between endothelial dysfunction and VDD [36]. Also, VD may directly protect endothelial cells against oxidative stress, and hence, VDD may contribute to ED through inflammation [30].

Many ED patients are VD deficient, particularly patients with arteriogenic ED [5]. Epidemiological data in humans have shown that VDD is associated with hypertension, left ventricular hypertrophy, increased arterial stiffness, and endothelial dysfunction in normal subjects [30]. Men with ED have an increased prevalence of endothelial dysfunction, and VD may improve endothelial function [14]. A placebo-controlled randomized trial demonstrated that even a single large dose of VD improves endothelial function in patients with type 2 diabetes and VDD [37]. Since endothelial function improvement is the cornerstone of the treatment of ED, VD supplementation may be beneficial in the treatment of ED patients.

3. Vitamin-D and nitric oxide production

Another important mechanism of action of VD seems

to be *via* NO mediated vascular dilation. NO synthases (NOS) are a family of enzymes that catalyze the production of NO from L-arginine. NO-pathway is a well-known physiologic signal essential to penile erection. Shortly, sexual stimulation releases neurotransmitters from the corpus cavernosa as well as NO (a relaxing factor) from the endothelial cells of the penis [28]. Any kind of disorders that reduce NO synthesis or release in the erectile tissue are commonly associated with ED. Activated VD stimulates the production of NO in endothelial cells and NOS [38], and is a key to vascular dilation and thereby critical to preventing ED. Molinari et al [38] demonstrated that VD is able to stimulate NO production in human umbilical vein endothelial cells through endothelial NOS (eNOS) activation.

The finding of an involvement of VD in NO production by endothelial cells is quite relevant. Recently, Andrukhova et al [30] reported that VDR mutant mice are characterized by lower bioavailability of the vasodilator NO due to reduced expression of the key NO synthesizing enzyme *i.e.*, eNOS. Reduction in eNOS ends with endothelial dysfunction, increased arterial stiffness, increased aortic impedance, structural remodeling of the aorta, and impaired systolic and diastolic heart function at later ages, independent of changes in the renin-angiotensin system [30]. This may also clarify why endothelium derived; NO-evoked dilation is halved in arteries from VD deficient male rats [35]. Under the light of recent scientific researches, unsurprisingly, there exists a higher prevalence of ED among VD deficient patients compared to those with optimal levels [14].

4. Vitamin-D and atherosclerotic cardiovascular diseases

Penile erection predominantly depends on vascular event and according to several studies there is a strong association between ED and ASCV diseases [39]. Unfortunately, VDD is associated with atherogenic dyslipidemia, DM, and reduced serum testosterone (T) levels all of which are associated with endothelial dysfunction and are classic risk factors for the onset of ED [13,14,18].

Recently, Barassi et al [5] demonstrated a higher presence of VDD in arterial ED patients compared with non-arterial-ED patients and a lower serum VD levels in more severe ED patients. Melamed et al [33] has shown that low serum VD levels are associated with a higher prevalence of peripheral arterial disease in the general population. Additionally, an earlier

study done with more than 7,000 VD deficient patients showed an association between VDD and several cardiovascular disease (CVD) states such as hypertension, coronary artery disease as well as ASCV risk factors (DM, dyslipidemia *etc.*) [18]. However, a recent cross-sectional analysis of 3,390 men aged >20 years, found that VDD in individuals free of ASCV diseases was associated with an increased prevalence of ED [14].

The available literature evaluating VD supplementation and its relationship to ASCV diseases are limited in number. However, in an observational retrospective study involving more than 10,000 patients Vacek et al [18] showed that VD supplement use improved survival in deficient subjects, supporting the potential benefit of this intervention. Later, in a randomized placebo-controlled preliminary study, Al-Dujaili et al [40] suggested that daily VD supplementation may ameliorate CVD risk factors.

5. Vitamin-D and endocrine system

T modulates nearly every component involved in erectile function and its deficiency associated with ED [41]. In an ageing male population, hypogonadism is common (30% prevalence in men >60 years), and is associated with ASCV risks (*e.g.*, atherogenic lipid profile, insulin resistance and obesity) [42]. The Longitudinal Aging Study Amsterdam, showed serum VD was positively associated with total and bioavailable T levels [43]. In another study, with large population group (2,854 men), investigators revealed that a lower VD level is associated with a higher prevalence of hypogonadism [44]. The latter result was supported by another study of 652 men over 40 years of age [45]. VD was significantly and positively associated with T levels before and after adjustment for age and ethnicity [46].

VD is positively associated with T, exhibits a concordant seasonal changeability [47], elevates when T was supplemented in hypogonadism [48]. Surprisingly, the reverse situation is also true, suggesting that VD supplementation might increase T levels [49]. In a clinical randomized controlled trial, which is the first on this topic in literature, Pilz et al [49] investigated the effect of VD supplementation on androgens in men. The results were significant and the researchers observed that overweight men with VDD had a clinically meaningful increase in serum T levels after VD supplementation for 1 year [49]. Recently, it was

also demonstrated that VD supplementation improves T levels, metabolic syndrome and erectile function in middle-aged VD deficient men [8]. Canguven et al [8] supported earlier research where serum total-T effectively increased after VD treatment [49,50]. Likewise, Wehr et al [47] demonstrated that androgen levels were associated with VD levels in 2,299 older men; and also a cross-sectional analysis of men in a national survey found that increasing VD ends with higher total-T levels [51]. Certainly, it has been advocated that the most advantageous serum VD concentrations begin at 30 ng/mL, and the best are between 36 to 40 ng/mL [52].

6. Vitamin-D and androgen receptors

An extra mechanism of VD on erectile function seems to function *via* binding to T receptors. Computer (*in silico*) modeling shows that besides activating the VDR, 1,25-D displays high affinity for some of the body's other nuclear receptors. This suggests that when 1,25-D increases above its normal range, it binds the α/β thyroid, the glucocorticoid, and the T receptors, displacing their native ligands [53]. Marshall [54] showed the symmetry with which endogenous ligands exhibited very similar affinities across some members of the type 1 nuclear receptor family [54]. For example, 1,25-D docked into the VDR with a (nanomolar) Kd of 8.48, but also exhibited a Kd of 8.05 into the T receptor.

7. Vitamin-D and immune system

ED is associated with an incremental inflammatory activation and inflammation plays an important pathophysiological role in both ED and ASCV diseases [12,39,55]. It has been extensively debated that inflammation can exert a detrimental effect on the ASCV system *via* two pathways: chronic, low-grade inflammation and an acute systemic inflammatory response. The former has been implicated in atherosclerotic processes [56], while the latter accounts for adverse ASCV events following severe inflammatory stimulation. Sildenafil, one of the phosphodiesterase inhibitors used as first line treatment in ED, induces a significant acute decrease in levels of pro-inflammatory markers/mediators [55]. The anti-inflammatory action might play an important role in addition to relaxation of penile smooth muscles.

Ding et al [50] revealed that VD supplement may protect the cells through suppressing inflammation factors and alleviating cell apoptotic death, as well to

reproduction and T synthesis. Ding et al' study [50] indicated that VD played a protective role of testes and testicular damage induced by diabetes, and the possible mechanism might be regulating attenuating inflammation and inactivating caspase cascade. Emerging data suggests that VD has a potential role in regulating inflammation. In research, it was shown that VD inhibits the expression of inflammatory cytokines in monocytes, including interleukins-1, 6, 8, 12, and tumor necrosis factor- α [57,58]. In summary, VD may directly defend endothelial cells against oxidative stress; and VDD may contribute to ED through inflammation.

CONCLUSIONS

Based on the evidence from bench and bed, it was shown that VD is important for erectile function. Measurement of VD in ED patients is very crucial with supplementation as required. VD supplementation potentially represents a low-cost, low-risk method to treat and prevent ED. In summary, we need VD starting from the early embryonal days to the end of our lives for a better, healthy and sexually active life.

Conflict of Interest

The authors have nothing to disclose.

Author Contribution

Conceptualization: OC. Data curation: OC. Formal analysis: OC, AHAM. Funding acquisition: OC. Investigation: OC. Methodology: OC. Project administration: OC. Resources: OC, AHAM. Software: OC. Supervision: OC. Validation: OC, AHAM. Visualization: OC. Writing – original draft: OC. Writing – review & editing: OC, AHAM.

REFERENCES

1. Talib RA, Khalafalla K, Cangüven Ö. The role of vitamin D supplementation on erectile function. *Turk J Urol* 2017;43:105-11.
2. Khanzada U, Khan SA, Hussain M, Adel H, Masood K, Adil SO, et al. Evaluation of the causes of erectile dysfunction in patients undergoing penile Doppler ultrasonography in Pakistan. *World J Mens Health* 2017;35:22-7.
3. Burnett AL. Novel nitric oxide signaling mechanisms regulate the erectile response. *Int J Impot Res* 2004;16 Suppl 1:S15-9.
4. Ayta IA, McKinlay JB, Krane RJ. The likely worldwide increase in erectile dysfunction between 1995 and 2025 and some possible policy consequences. *BJU Int* 1999;84:50-6.
5. Barassi A, Pezzilli R, Colpi GM, Corsi Romanelli MM, Melzi d'Eril GV. Vitamin D and erectile dysfunction. *J Sex Med* 2014;11:2792-800.
6. Safarinejad MR. Prevalence and risk factors for erectile dysfunction in a population-based study in Iran. *Int J Impot Res* 2003;15:246-52.
7. Corona G, Lee DM, Forti G, O'Connor DB, Maggi M, O'Neill TW, et al.; EMAS Study Group. Age-related changes in general and sexual health in middle-aged and older men: results from the European Male Ageing Study (EMAS). *J Sex Med* 2010;7:1362-80.
8. Canguven O, Talib RA, El Ansari W, Yassin DJ, Al Naimi A. Vitamin D treatment improves levels of sexual hormones, metabolic parameters and erectile function in middle-aged vitamin D deficient men. *Aging Male* 2017;20:9-16.
9. Feldman HA, Johannes CB, Derby CA, Kleinman KP, Mohr BA, Araujo AB, et al. Erectile dysfunction and coronary risk factors: prospective results from the Massachusetts male aging study. *Prev Med* 2000;30:328-38.
10. Montorsi P, Ravagnani PM, Galli S, Rotatori F, Briganti A, Salonia A, et al. Common grounds for erectile dysfunction and coronary artery disease. *Curr Opin Urol* 2004;14:361-5.
11. Pegge NC, Twomey AM, Vaughton K, Gravenor MB, Ramsey MW, Price DE. The role of endothelial dysfunction in the pathophysiology of erectile dysfunction in diabetes and in determining response to treatment. *Diabet Med* 2006;23:873-8.
12. Meluzín J, Vasků A, Kincl V, Panovský R, Srámková T. Association of coronary artery disease, erectile dysfunction, and endothelial nitric oxide synthase polymorphisms. *Heart Vessels* 2009;24:157-63.
13. Caretta N, de Kreutzenberg SV, Valente U, Guarneri G, Ferlin A, Avogaro A, et al. Hypovitaminosis D is associated with erectile dysfunction in type 2 diabetes. *Endocrine* 2016;53:831-8.
14. Farag YMK, Guallar E, Zhao D, Kalyani RR, Blaha MJ, Feldman DI, et al. Vitamin D deficiency is independently associated with greater prevalence of erectile dysfunction: the National Health and Nutrition Examination Survey (NHANES) 2001-2004. *Atherosclerosis* 2016;252:61-7.
15. McCollum EV, Simmonds N, Becker JE, Shipley PG. An experimental demonstration of the existence of a vitamin which promotes calcium deposition. *J Biol Chem* 1922;53:293-312.
16. Blomberg Jensen M, Nielsen JE, Jørgensen A, Rajpert-De Meyts E, Kristensen DM, Jørgensen N, et al. Vitamin D recep-

- tor and vitamin D metabolizing enzymes are expressed in the human male reproductive tract. *Hum Reprod* 2010;25:1303-11.
17. Haussler MR, Haussler CA, Whitfield GK, Hsieh JC, Thompson PD, Barthel TK, et al. The nuclear vitamin D receptor controls the expression of genes encoding factors which feed the "Fountain of Youth" to mediate healthful aging. *J Steroid Biochem Mol Biol* 2010;121:88-97.
 18. Vacek JL, Vanga SR, Good M, Lai SM, Lakkireddy D, Howard PA. Vitamin D deficiency and supplementation and relation to cardiovascular health. *Am J Cardiol* 2012;109:359-63.
 19. Deeb KK, Trump DL, Johnson CS. Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat Rev Cancer* 2007;7:684-700.
 20. Niruban SJ, Alagiakrishnan K, Beach J, Senthilselvan A. Association between vitamin D and respiratory outcomes in Canadian adolescents and adults. *J Asthma* 2015;52:653-61.
 21. Kerley CP, Elnazir B, Faul J, Cormican L. Vitamin D as an adjunctive therapy in asthma. Part 2: a review of human studies. *Pulm Pharmacol Ther* 2015;32:75-92.
 22. Nagpal S, Na S, Rathnachalam R. Noncalcemic actions of vitamin D receptor ligands. *Endocr Rev* 2005;26:662-87.
 23. Griffin MD, Lutz W, Phan VA, Bachman LA, McKean DJ, Kumar R. Dendritic cell modulation by 1 α ,25 dihydroxyvitamin D₃ and its analogs: a vitamin D receptor-dependent pathway that promotes a persistent state of immaturity in vitro and in vivo. *Proc Natl Acad Sci U S A* 2001;98:6800-5.
 24. Collins FS, Morgan M, Patrinos A. The Human Genome Project: lessons from large-scale biology. *Science* 2003;300:286-90.
 25. Bossé Y, Lemire M, Poon AH, Daley D, He JQ, Sandford A, et al. Asthma and genes encoding components of the vitamin D pathway. *Respir Res* 2009;10:98.
 26. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266-81.
 27. Ginde AA, Liu MC, Camargo CA Jr. Demographic differences and trends of vitamin D insufficiency in the US population, 1988-2004. *Arch Intern Med* 2009;169:626-32.
 28. Sorenson M, Grant WB. Does vitamin D deficiency contribute to erectile dysfunction? *Dermatoendocrinol* 2012;4:128-36.
 29. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al.; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96:1911-30.
 30. Andrukhova O, Slavic S, Zeitz U, Riesen SC, Heppelmann MS, Ambrisko TD, et al. Vitamin D is a regulator of endothelial nitric oxide synthase and arterial stiffness in mice. *Mol Endocrinol* 2014;28:53-64.
 31. Lips P. Relative value of 25(OH)D and 1,25(OH)₂D measurements. *J Bone Miner Res* 2007;22:1668-71.
 32. Munns CF, Simm PJ, Rodda CP, Garnett SP, Zacharin MR, Ward LM, et al.; APSU Vitamin D Study Group. Incidence of vitamin D deficiency rickets among Australian children: an Australian Paediatric Surveillance Unit study. *Med J Aust* 2012;196:466-8.
 33. Melamed ML, Muntner P, Michos ED, Uribarri J, Weber C, Sharma J, et al. Serum 25-hydroxyvitamin D levels and the prevalence of peripheral arterial disease: results from NHANES 2001 to 2004. *Arterioscler Thromb Vasc Biol* 2008;28:1179-85.
 34. Fernandes-Lima F, Gregório BM, Nascimento FAM, Costa WS, Gallo CBM, Sampaio FJB. Effects of vitamin D restricted diet administered during perinatal and postnatal periods on the penis of Wistar rats. *Biomed Res Int* 2018;2018:6030646.
 35. Tare M, Emmett SJ, Coleman HA, Skordilis C, Eyles DW, Morley R, et al. Vitamin D insufficiency is associated with impaired vascular endothelial and smooth muscle function and hypertension in young rats. *J Physiol* 2011;589:4777-86.
 36. Dalan R, Liew H, Tan WKA, Chew DEK, Leow MKS. Vitamin D and the endothelium: basic, translational and clinical research updates. *IJC Metab Endocr* 2014;4:4-17.
 37. Sugden JA, Davies JI, Witham MD, Morris AD, Struthers AD. Vitamin D improves endothelial function in patients with Type 2 diabetes mellitus and low vitamin D levels. *Diabet Med* 2008;25:320-5.
 38. Molinari C, Uberti F, Grossini E, Vacca G, Carda S, Invernizzi M, et al. 1 α ,25-dihydroxycholecalciferol induces nitric oxide production in cultured endothelial cells. *Cell Physiol Biochem* 2011;27:661-8.
 39. Tostes RC, Carneiro FS, Lee AJ, Giachini FR, Leite R, Osawa Y, et al. Cigarette smoking and erectile dysfunction: focus on NO bioavailability and ROS generation. *J Sex Med* 2008;5:1284-95.
 40. Al-Dujaili EA, Munir N, Iniesta RR. Effect of vitamin D supplementation on cardiovascular disease risk factors and exercise performance in healthy participants: a randomized placebo-controlled preliminary study. *Ther Adv Endocrinol Metab* 2016;7:153-65.
 41. Isidori AM, Buvat J, Corona G, Goldstein I, Jannini EA, Lenzi A, et al. A critical analysis of the role of testosterone in erectile function: from pathophysiology to treatment-a systematic review. *Eur Urol* 2014;65:99-112.
 42. Kapoor D, Clarke S, Stanworth R, Channer KS, Jones TH. The effect of testosterone replacement therapy on adipocyto-

- kines and C-reactive protein in hypogonadal men with type 2 diabetes. *Eur J Endocrinol* 2007;156:595-602.
43. Rafiq R, van Schoor NM, Sohl E, Zillikens MC, Oosterwerff MM, Schaap L, et al. Associations of vitamin D status and vitamin D-related polymorphisms with sex hormones in older men. *J Steroid Biochem Mol Biol* 2016;164:11-7.
 44. Wang N, Han B, Li Q, Chen Y, Chen Y, Xia F, et al. Vitamin D is associated with testosterone and hypogonadism in Chinese men: results from a cross-sectional SPECT-China study. *Reprod Biol Endocrinol* 2015;13:74.
 45. Tak YJ, Lee JG, Kim YJ, Park NC, Kim SS, Lee S, et al. Serum 25-hydroxyvitamin D levels and testosterone deficiency in middle-aged Korean men: a cross-sectional study. *Asian J Androl* 2015;17:324-8.
 46. Chin KY, Ima-Nirwana S, Wan Ngah WZ. Vitamin D is significantly associated with total testosterone and sex hormone-binding globulin in Malaysian men. *Aging Male* 2015;18:175-9.
 47. Wehr E, Pilz S, Boehm BO, März W, Obermayer-Pietsch B. Association of vitamin D status with serum androgen levels in men. *Clin Endocrinol (Oxf)* 2010;73:243-8.
 48. Francis RM, Peacock M, Aaron JE, Selby PL, Taylor GA, Thompson J, et al. Osteoporosis in hypogonadal men: role of decreased plasma 1,25-dihydroxyvitamin D, calcium malabsorption, and low bone formation. *Bone* 1986;7:261-8.
 49. Pilz S, Frisch S, Koertke H, Kuhn J, Dreier J, Obermayer-Pietsch B, et al. Effect of vitamin D supplementation on testosterone levels in men. *Horm Metab Res* 2011;43:223-5.
 50. Ding C, Wang Q, Hao Y, Ma X, Wu L, du M, et al. Vitamin D supplement improved testicular function in diabetic rats. *Biochem Biophys Res Commun* 2016;473:161-7.
 51. Anic GM, Albanes D, Rohrmann S, Kanarek N, Nelson WG, Bradwin G, et al. Association between serum 25-hydroxyvitamin D and serum sex steroid hormones among men in NHANES. *Clin Endocrinol (Oxf)* 2016;85:258-66.
 52. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr* 2006;84:18-28.
 53. Proal AD, Albert PJ, Marshall TG. Dysregulation of the vitamin D nuclear receptor may contribute to the higher prevalence of some autoimmune diseases in women. *Ann N Y Acad Sci* 2009;1173:252-9.
 54. Marshall TG. Vitamin D discovery outpaces FDA decision making. *Bioessays* 2008;30:173-82.
 55. Vlachopoulos C, Ioakeimidis N, Rokkas K, Angelis A, Terentes-Printzios D, Stefanadis C, et al. Acute effect of sildenafil on inflammatory markers/mediators in patients with vasculogenic erectile dysfunction. *Int J Cardiol* 2015;182:98-101.
 56. Libby P, Ridker PM, Hansson GK; Leducq Transatlantic Network on Atherothrombosis. Inflammation in atherosclerosis: from pathophysiology to practice. *J Am Coll Cardiol* 2009;54:2129-38.
 57. Almerighi C, Sinistro A, Cavazza A, Ciaprin C, Rocchi G, Bergamini A. 1 α ,25-dihydroxyvitamin D₃ inhibits CD40L-induced pro-inflammatory and immunomodulatory activity in human monocytes. *Cytokine* 2009;45:190-7.
 58. D'Ambrosio D, Cippitelli M, Cocciolo MG, Mazzeo D, Di Lucia P, Lang R, et al. Inhibition of IL-12 production by 1,25-dihydroxyvitamin D₃. Involvement of NF-kappaB downregulation in transcriptional repression of the p40 gene. *J Clin Invest* 1998;101:252-62.