



Physiological and gene transcription assays to assess responses of mussels to environmental changes

Katrina L. Counihan¹, Lizabeth Bowen², Brenda Ballachey³, Heather Coletti⁴, Tuula Hollmen⁵, Benjamin Pister⁶ and Tammy L. Wilson^{4,7}

¹Alaska SeaLife Center, Seward, AK, United States of America

²US Geological Survey, Western Ecological Research Center, Davis, CA, United States of America

³US Geological Survey, Alaska Science Center, Anchorage, AK, United States of America

⁴Inventory and Monitoring Program, Southwest Alaska Network, National Park Service, Anchorage, AK, United States of America

⁵College of Fisheries and Ocean Sciences, University of Alaska—Fairbanks and Alaska SeaLife Center, Seward, AK, United States of America

⁶Ocean Alaska Science and Learning Center, National Park Service, Anchorage, AK, United States of America

⁷Department of Natural Resource Management, South Dakota State University, Brookings, SD, United States of America

ABSTRACT

Coastal regions worldwide face increasing management concerns due to natural and anthropogenic forces that have the potential to significantly degrade nearshore marine resources. The goal of our study was to develop and test a monitoring strategy for nearshore marine ecosystems in remote areas that are not readily accessible for sampling. Mussel species have been used extensively to assess ecosystem vulnerability to multiple, interacting stressors. We sampled bay mussels (*Mytilus trossulus*) in 2015 and 2016 from six intertidal sites in Lake Clark and Katmai National Parks and Preserves, in south-central Alaska. Reference ranges for physiological assays and gene transcription were determined for use in future assessment efforts. Both techniques identified differences among sites, suggesting influences of both large-scale and local environmental factors and underscoring the value of this combined approach to ecosystem health monitoring.

Submitted 5 April 2019
Accepted 31 August 2019
Published 4 October 2019

Corresponding author
Katrina L. Counihan,
katrinac@alaskasealife.org

Academic editor
Carlos De Rezende

Additional Information and
Declarations can be found on
page 23

DOI 10.7717/peerj.7800

© Copyright
2019 Counihan et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Conservation Biology, Ecosystem Science, Marine Biology

Keywords *Mytilus trossulus*, Gene transcription, Biomarker, Nearshore marine ecosystem, Alaska, Ecosystem management, Monitoring

INTRODUCTION

As a result of rapid changes to the global climate and increases in anthropogenic stressors, challenges are mounting along coastal regions around the world. These pressures have the potential to significantly degrade nearshore marine resources and are a threat not only in developed areas, but also in remote regions, including coastal areas of Alaska (Breitburg & Riedel, 2005). Alaska has tens of thousands of miles of coastline that are only accessible by boat or plane and during short periods of the summer due to remoteness and weather conditions. Recognizing that change is inevitable, technologies for monitoring the status of nearshore marine ecosystems in remote areas are necessary for managing and maintaining

healthy coastal communities. One approach for assessing ecosystem status is the use of indicator species, including intertidal bivalves ([Beyer et al., 2017](#)).

Intertidal bivalve communities are spatially and temporally variable ([Bodkin et al., 2017](#)) and are vulnerable to a wide range of influences and disturbances ([Denny & Wetthey, 2001](#); [Menge & Branch, 2001](#)). Species composition, abundance, and physiology are influenced by a complex variety of stressors, the cumulative effects of which are generally not well understood ([Gunderson, Armstrong & Stillman, 2016](#)). In Alaska, nearshore communities have been affected by earthquakes, including the major 1964 Alaska earthquake ([Baxter, 1971](#)), volcanic eruptions ([DeGange et al., 2010](#); [Walker et al., 2013](#); [Jewett & Drew, 2014](#); [Zimmermann et al., 2018](#)), and hydrocarbon contamination from the 1989 Exxon Valdez oil spill ([Armstrong et al., 1995](#); [Driskell et al., 1996](#); [Lees, Houghton & Driskell, 1996](#); [Fukuyama, Shigenaka & Hoff, 2000](#)). Further stressors will be forthcoming with increases in ocean temperature, decreases in pH ([Lesser et al., 2010](#); [Hoegh-Guldberg & Bruno, 2010](#); [Gaylord et al., 2011](#); [Bijma et al., 2013](#)), contaminants ([Franzellitti et al., 2010](#)), and human harvest of marine species ([Jamieson, 1993](#)).

Intertidal invertebrates are important members of nearshore communities, and in the Gulf of Alaska are a primary food source for a variety of marine and terrestrial vertebrate and invertebrate predators including brown bears ([Smith & Partridge, 2004](#)), sea stars ([Paul & Feder, 1975](#); [Fukuyama & Oliver, 1985](#)), shorebirds ([Gill Jr & Handel, 1990](#)), sea ducks ([Lewis, Esler & Boyd, 2007](#)), sea otters ([Calkins, 1978](#); [Doroff & DeGange, 1994](#); [Coletti et al., 2016](#)) and human subsistence users ([Fall & Field, 1996](#)). Bay mussels (*Mytilus trossulus*) are ubiquitous throughout nearshore communities in the Gulf of Alaska and northeast Pacific. Changes in bay mussel populations due to abiotic or biotic factors may result in alterations of the entire intertidal community structure ([Harley, 2011](#); [Navarrete, Menge & Daley, 2000](#)). For example, a reduction in mussel abundance due to predation or thermal stress decreased overall species diversity in rocky intertidal communities ([Harley, 2011](#)).

Nearshore species can serve as a focal point for understanding variables that influence the nearshore ecosystem, as they integrate marine and terrestrial stressors into their behavior and condition and reflect these drivers in their abundance and population trends over time. In our work, bay mussels were selected as an indicator species because they are abundant, an important component of the nearshore food web and, as prodigious filter feeders, will consume and sequester contaminants. Studies using mussels have included assessment of contaminants and changing climate conditions ([Akaishi et al., 2007](#); [Beyer et al., 2017](#); [Coray, St.Jean & Bard, 2007](#); [Gagné et al., 2007](#); [Halldórsson, Svavarsson & Granmo, 2005](#); [Halldórsson et al., 2008](#); [Shaw et al., 2011](#)), which are frequent goals of long-term monitoring.

In recent years, there have been major advances in assays to evaluate the physiological condition of mussels using biomarkers (biological metrics that quantify a physiological response including detection of proteins or their activity) and gene transcription (measurement of alterations in transcription of specific genes) to elucidate changes at the molecular level in response to environmental changes and contaminant exposure ([Livingstone et al., 2000](#); [Evans & Hofmann, 2012](#); [Hüning et al., 2013](#); [Bolognesi & Cirillo, 2014](#)). Gene transcription is the process by which information from the DNA template of a

particular gene is transcribed into messenger RNA (mRNA) and eventually translated into a functional protein. The amount of a particular gene that is expressed is physiologically dictated by a number of intrinsic and extrinsic factors, and analysis of mRNA and protein levels can provide information about dynamic changes in the functional state of an organism. A holistic approach that combines biomarker and gene transcription is gaining momentum in other regions, with researchers evaluating and applying these methods to mussels and other invertebrates for environmental monitoring to improve sensitivity and efficiency (Banni et al., 2017; Beyer et al., 2017; Carella et al., 2018; Sforzini et al., 2018). An advantage of measuring gene transcription is that mRNA production is the earliest observable sign an organism is responding to a stressor, but mRNA is generally not as stable as proteins (De Sousa Abreu et al., 2009; McLoughlin et al., 2006). While protein levels respond more slowly than mRNA, their often greater stability makes assessing protein biomarkers advantageous. Thus, combining the two approaches can be complementary as increases in gene transcripts and proteins are often detected concurrently, but they can also provide information that may be missed when using one method alone.

Often, large scale investigations into populations and ecosystems have been driven by catastrophic changes such as oil spills (Wells, Butler & Hughes, 1995; Peterson et al., 2003), mortality events (Lessios, 1988) and ecological extinction (Jackson et al., 2001). However, investigations 'after the fact' may be constrained by a lack of baseline data, limiting insight into ecological pathways and causes of change and recovery (Breitburg & Riedel, 2005). The goal of this study was to work toward an approach that combines biomarker and genetic methods to assess the condition of nearshore ecosystems, including those in areas that are remote and logistically difficult to access. The specific objectives to accomplish this goal included: (1) acquire baseline physiological and gene transcription data for the bay mussel across sites in the Gulf of Alaska, (2) assess the relationship between physiological and gene transcription assays for validation and support of both techniques, and (3) determine if site-level differences were present. Mussels were sampled in 2015 and 2016 from six intertidal sites in two national parks. The sites were expected to reflect a range of representative habitat types to generate reference baseline data applicable to mussels in this region. We hypothesized that we would be able to establish baseline data for the sites that were sampled and that the biomarker and gene transcription assays would provide complementary results. Further, we hypothesized that biomarker and gene transcription results would not be significantly different among sites.

MATERIALS & METHODS

Study organisms and area

Mussels (*Mytilus trossulus*) were collected during late morning on the rising tide in July of 2015 and 2016 at three sites within each of two national parks in southcentral Alaska: Lake Clark National Park and Preserve (LACL; sites: Fossil Point, Silver Salmon, and Chinitna Bay) and Katmai National Park and Preserve (KATM; sites: Kukak, Kafia, & Takli) (Fig. 1) (Alaska Department of Fish and Game permit # CF-15-088 for KATM and LACL, 2015, and # CF-16-089 for KATM and LACL, 2016). Due to the remote nature of these parks, all




Figure 1 Mussel sampling sites in Lake Clark National Park and Preserve (NPP) and Katmai National Park and Preserve (NPP). Sites were sampled during June of 2015 and 2016.

Full-size [DOI: 10.7717/peerj.7800/fig-1](https://doi.org/10.7717/peerj.7800/fig-1)

sites are only accessible by boat or plane during the summer months. Twenty mussels were collected from each site in both 2015 and 2016 (total of 120 each year), with 10 designated for physiological assays and 10 for gene transcription; we therefore collected a total of 240 mussels, and used 120 of them for each methodology. Mussels collected in KATM were from sites that were randomly selected as part of a long-term nearshore monitoring program (Dean, Bodkin & Coletti, 2014). These sites were established on mixed-sediment beaches in protected to semi-protected areas (Weitzman et al., 2017). Currently, no long-term monitoring of coastal bivalve resources exists along the LACL coastline, so sampling locations in LACL were selected based on the presence of mussels. Mussels were present on and collected from bedrock outcroppings. Overall, our intent was to capture the range of natural variation across a set of sites that are representative of coastline habitats in lower Cook Inlet and the Gulf of Alaska.

The mussels were kept submerged in seawater and processed as soon as possible following collection, generally within 1–4 h. For biomarker assays, hemolymph was extracted from the posterior adductor muscle with a tuberculin needle. The hemolymph and mussel were frozen in liquid nitrogen for transport to the lab where they were transferred to a -80°C freezer until processing.

For gene transcription assays, gill tissue was removed, placed in cryovials with RNAlater[®], and frozen at -20°C until analyses.

Table 1 Biomarkers selected for this study and their primary functions and interactions.

Biomarker	Biological Process	Environmental Interaction	References
Condition factor	Growth	Ocean acidification Nutrient availability Contaminants Temperature	<i>Carmichael, Shriver & Valiela (2004); Gagné et al. (2007)</i>
Shell Thickness	Growth	Ocean acidification Nutrient availability Predation Density	<i>Blanchard & Feder (2000); Freeman (2007); Gaylord et al. (2011); Xavier, Branch & Wieters (2007)</i>
Hemocyte Count	Immune function	Contaminants Pathogens	<i>Galloway & Depledge (2001)</i>
Hydrogen Peroxide	Immune function	Contaminants Pathogens	<i>Galloway & Depledge (2001)</i>
RNA:DNA	Metabolic condition	Nutrient availability	<i>Caldarone et al. (2001), Lesser et al. (2010)</i>
Cytochrome P450	Detoxification	Contaminants	<i>Rewitz et al. (2006)</i>
Heat Shock Protein 40	Thermal stress	Temperature Stress	<i>Cruz-Rodriguez & Chu (2002)</i>

Invertebrate physiology

Seven biomarkers were used to assess the physiological status of the mussels and are summarized in [Table 1](#).

Morphometrics, condition factor and age

The length, width, height, and total wet weight of mussels were measured prior to dissection. The posterior adductor muscle and digestive gland were excised for assays. The condition factor was calculated by dividing the total mussel weight by the shell length. Age was determined by counting shell growth rings. The median and range of morphometrics and ages by site are summarized in [Table 2](#).

Shell thickness

A micrometer was used to measure the shell thickness at 5 regularly spaced points around the shell approximately one mm from the edge (*Versteegh & Hansson, 2012*).

Hemocyte count

A 0.01 ml sample of hemolymph was diluted 1:1 with tris-buffered saline (TBS) and the number of cells were counted using a hemocytometer in three replicates (*Akaishi et al., 2007*).

Hydrogen peroxide production

Hemolymph samples were diluted 1:1 with tris-buffered saline (TBS). Samples were tested in triplicate by pipetting 0.05 ml of the 1:1 hemolymph:TBS solution into a 96-well plate. After incubating the plate in the dark for 1 h, 0.05 ml of phenol red solution (phosphate buffered saline pH 7.4, 5.5 mM dextrose, 0.56 mM phenol red, 8.5 U ml⁻¹ horseradish peroxidase, type II) was added to each well and incubated for another 30 min in the dark. The reaction was stopped by adding 0.01 ml of 1 N NaOH, and the plate was read on a

Table 2 Medians and ranges of morphometrics and ages of mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves during 2015 and 2016 ($n = 10$ mussels per site per year).

Park	Site	Year	Length (mm)		Width (mm)		Height (mm)		Weight (g)		Age (years)	
			Median	Range	Median	Range	Median	Range	Median	Range	Median	Range
Katmai	Takli	2015	41.81	22.27-56.28	21.63	19.98-25.81	16.93	15.05-20.74	6.15	4.48-11.34	6	5-10
		2016	41.72	33.36-44.92	19.29	17.09-22.83	16.01	13.47-18.70	5.91	3.99-9.19	7	5-9
	Kukak	2015	38.73	29.91-48.79	19.75	15.90-24.03	15.51	12.03-18.74	4.08	1.68-7.78	6.5	4-8
		2016	34.69	30.76-44.22	18.06	16.18-21.01	13.56	11.32-16.26	4.05	2.24-7.36	7	6-11
	Kafliia	2015	48.94	44.20-57.96	23.39	21.17-26.36	19.47	17.09-22.82	8.11	6.56-13.91	9	7-12
		2016	37.24	30.67-45.80	17.12	15.97-19.87	14.11	12.16-17.45	4.28	2.11-7.65	7	5-10
Fossil Point	2015	60.26	41.14-78.55	25.94	20.23-32.73	25.06	16.03-30.06	16.18	4.57-24.19	8	5-11	
	2016	54.73	42.11-63.74	24.99	23.13-29.44	22.51	18.66-25.90	10.53	5.90-19.12	9	6-12	
Lake Clark	Silver	2015	48.02	37.14-53.83	22.54	19.16-25.74	22.36	15.94-23.80	9.06	4.16-12.72	7	5-12
	Salmon	2016	45.88	41.96-55.39	21.46	20.02-26.19	18.55	14.51-24.95	6.12	4.73-14.89	8	5-12
	Chinitna Bay	2015	42.76	37.73-57.16	20.48	17.06-23.85	20.44	16.34-25.45	9.02	5.32-14.68	7	5-8
		2016	41.69	33.61-46.19	20.07	15.43-22.45	20.23	14.87-24.31	9.43	4.65-14.05	7.5	6-12

Molecular Devices SpectraMax Plus microplate reader (Sunnyvale, CA, USA) at 620 nm (*Akaishi et al., 2007*).

RNA:DNA ratio

Half of the posterior adductor muscle was homogenized and 0.15 ml of 1% sarcosyl tris-EDTA (STEB) was added to the homogenate. The mixture was vortexed for 60 min and then 1.35 ml of tris-EDTA (TE) buffer was added. The sample was centrifuged 15 min at 14,000 x g at room temperature and the supernatant saved for testing. The samples were diluted 1:20 and 0.075 ml of each sample was added to a 96-well plate in duplicate. Genomic, unshered DNA from calf thymus (Sigma-Aldrich, St. Louis, MO) was used to prepare a DNA standard curve (0.1 $\mu\text{g ml}^{-1}$, 0.2 $\mu\text{g ml}^{-1}$, 0.4 $\mu\text{g ml}^{-1}$, 0.8 $\mu\text{g ml}^{-1}$, 1.6 $\mu\text{g ml}^{-1}$, 3.2 $\mu\text{g ml}^{-1}$, 6.4 $\mu\text{g ml}^{-1}$, 10.0 $\mu\text{g ml}^{-1}$). RNA from bovine pancreas (Sigma-Aldrich) was used for a RNA standard curve (0.4 $\mu\text{g ml}^{-1}$, 0.8 $\mu\text{g ml}^{-1}$, 1.6 $\mu\text{g ml}^{-1}$, 3.0 $\mu\text{g ml}^{-1}$, 6.0 $\mu\text{g ml}^{-1}$, 8.0 $\mu\text{g ml}^{-1}$, 12.0 $\mu\text{g ml}^{-1}$, 16.0 $\mu\text{g ml}^{-1}$). The wells had 0.075 ml of ethidium bromide solution (2 $\mu\text{g ml}^{-1}$) added to them and the microplate was shaken for 15 min. The plate was read in a SpectraMax Gemini EM fluorescent microplate reader (Molecular Devices) with 525 nm excitation, 600 nm emission to determine the total nucleic acid reading. Each well had 0.0075 ml of RNase solution (20 U ml^{-1}) added to it and the plate was shaken for 20 min and read on the microplate reader again with the same settings. The second reading was the DNA only reading. The RNA content was determined by subtracting the second reading from the first (*Caldarone et al., 2001*).

Cytochrome P450 activity

The digestive gland was homogenized in buffer (25 mM Hepes, 125 mM NaCl, 0.1 mM EDTA, 0.1 mM dithiothreitol) at a 1:5 weight:volume ratio. The mixture was centrifuged at 1,500 x g for 10 min at 2 °C and the supernatant transferred to a clean tube. The supernatant was centrifuged at 10,000 x g for 20 min at 2 °C. The supernatant was discarded and the pellet resuspended in 0.15 ml of microsome buffer (25 mM Hepes, 140 mM NaCl, 1 mM KH_2PO_4). A 96-well plate was inoculated with 0.05 ml of each microsome sample in triplicate. Each well had 0.05 ml of 50 μM BFC solution (7-benzyl-4(trifluoromethyl) coumarin in phenol red-free Dulbecco's Modified Eagle Medium) added and the plate was incubated for 4 h at room temperature. The plate had 0.04 ml of stop solution (80% CH_3CN , 20% 0.5 M Tris base) added to each well and it was read on a SpectraMax Gemini EM fluorescent microplate reader (Molecular Devices) at 410 excitation, 530 emission (*Mensah-Osman et al., 2007*).

Heat shock protein

Half of the posterior adductor muscle was homogenized and lysed in buffer (150 mM NaCl, 1% Triton-X 100, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulphate, 50 mM Tris, 1 mM phenylmethylsulfonyl fluoride). The homogenate was centrifuged at 12,000 rpm for 20 min at 4 °C and the supernatant collected. The protein concentration was determined using a Bradford assay. The sample was mixed 1:1 with 2x Laemmli buffer, boiled at 100 °C for 5 min and loaded on a SDS-PAGE gel. Each gel was run with a positive heat shock protein control and a molecular weight marker. The gel was electrophoretically transferred

Table 3 Genes selected for the transcription panel and their primary functions and interactions.

Gene	Biological process	Environmental interaction	References
Calmodulin (CaM)	Metabolism, shell formation	Ocean acidification Temperature Dissolved oxygen	<i>Chen et al. (2012); Li et al. (2004)</i>
Caspase 8 (Casp 8)	Apoptosis, necrosis, inflammation	Pathogens Contaminants	<i>Romero et al. (2011)</i>
Macrophage migration inhibitory factor (MIF)	Innate immunity	Pathogens	<i>Parisi et al. (2012), Philipp et al. (2012)</i>
Calponin (CNN)	Hypoxia	Ocean acidification Dissolved oxygen	<i>Hüning et al. (2013), Li et al. (2016)</i>
Chitinase (CHI)	Metabolism, hypoxia	Ocean acidification Dissolved oxygen	<i>Banni et al. (2011), Hüning et al. (2013)</i>
Cytochrome C Oxidase IV (CCOIV)	Hypoxia	Dissolved oxygen	<i>Fukuda et al. (2007)</i>
Heat shock protein 70 (HSP70)	Thermal stress, molecular chaperone	Temperature Pathogens Contaminants Hypoxia	<i>De Maio (1999), Iwama et al. (1999), Tsan & Gao (2004)</i>
Heat shock protein 90 (HSP90)	Thermal stress, molecular chaperone	Temperature Pathogens Contaminants Hypoxia	<i>De Maio (1999), Iwama et al. (1999), Tsan & Gao (2004)</i>
Hypoxia-inducible factor alpha (HIFa)	Hypoxia	Dissolved oxygen	<i>Wu (2002)</i>
Myticin B (MytB)	Innate immunity	Pathogens	<i>Balseiro et al. (2011)</i>
Mytilin (Myt)	Innate immunity	Pathogens Ocean acidification	<i>Balseiro et al. (2011), Mitta et al. (2000)</i>
Metallothionein 20 (MT20)	Detoxification	Contaminants—metals	<i>Banni et al. (2007)</i>
Cytochrome P450, family 3 (Cyp3)	Detoxification	Contaminants	<i>Giuliani et al. (2013)</i>
Tumor protein 53 (p53)	Apoptosis	Contaminants—PAHs	<i>Goodson et al. (2006), Banni et al. (2009)</i>

onto a polyvinyl difluoride (PVDF) membrane. Total protein was visualized by staining the gel with Pierce reversible protein stain (Thermo Scientific, Pittsburgh, PA USA). The stain was removed and the membrane was probed with a mouse anti-HSP 40 primary antibody (Abcam, Cambridge, MA USA). The membrane was washed and incubated with a secondary alkaline phosphatase labeled anti-mouse antibody (Abcam). The membrane was washed again and alkaline phosphatase substrate added. The membrane was photographed to document bands and analyzed with Image Studio Lite software, version 5.2.5 (Li-Core, Lincoln, NE, USA).

Invertebrate gene transcription

Transcription of 14 genes was used to assess the physiologic status of the mussels; genes used are summarized in [Table 3](#).

Stability of the proposed reference gene (18S) was determined using the web-based analysis tool RefFinder (<http://www.leonxie.com/referencegene.php>) (*Xie et al., 2012*). Sufficient stability was found, thus cycle threshold crossing values (C_T) for the genes

of interest were normalized to 18S. Although sex can potentially have impacts on gene transcription, most studies investigating gene transcription in the specific genes identified in our panel do not differentiate between sexes (*Zeng et al., 2012; Woo et al., 2011; Lacroix et al., 2014; Balseiro et al., 2011; Hüning et al., 2013; Woo et al., 2013; Cellura et al., 2007; Châtel et al., 2015; Franzellitti et al., 2010; Place, O'Donnell & Hofmann, 2008; Li et al., 2010; Giannetto et al., 2015; Dondero et al., 2006; Sureda et al., 2011; Zanette et al., 2013; Giuliani et al., 2013; Di et al., 2011*), and we did not differentiate by sex in this study.

Tissue collection and RNA extraction

Gill tissue was collected from each mussel and placed immediately into RNAlater[®] (Ambion/Life Technologies, Grand Island, New York). All tissue samples were stored at -20°C . Total RNA was extracted from pulverized gill tissue using the RNeasy Lipid Tissue Mini Kit (Qiagen; <http://www.qiagen.com>). To remove contaminating genomic (g)DNA, the spin columns were treated with $10\text{ U } \mu\text{l}^{-1}$ of RNase-free DNase I (DNase, Amersham Pharmacia Biotech Inc.; <http://www.apbiotech.com>) at 20°C for 15 min. RNA was then stored at -80°C pending further analyses.

cDNA synthesis

A standard cDNA synthesis was performed on $2\text{ }\mu\text{g}$ of RNA template from each mussel. Reaction conditions included 4 units reverse transcriptase (Omniscript, Qiagen, Valencia, CA, USA), $1\text{ }\mu\text{M}$ random hexamers, 0.5 mM each dNTP, and 10 units RNase inhibitor, in RT buffer (Qiagen, Valencia, CA, USA). Reactions were incubated for 60 min at 37°C , followed by an enzyme inactivation step of 5 min at 93°C , and then stored at -30°C until further analysis.

Primer design

Degenerate primers were designed based upon multi-species alignments (GenBank). Briefly, degenerate primer pairs developed for the mussel were used on cDNA from three randomly selected mussel samples. Degenerate primer pairs were designed to amplify 14 genes of interest and one ribosomal housekeeping gene (Table 4). The PCR amplifications using these primers were performed on 20 ng of each cDNA sample in $50\text{ }\mu\text{l}$ volumes containing 20–60 pmol of each primer, 40 mM Tris-KOH (pH 8.3), 15 mM KOAc, 3.5 mM Mg (OAc)₂, $3.75\text{ }\mu\text{g/ml}$ bovine serum albumin (BSA), 0.005% Tween-20, 0.005% Nonidet-P40, $200\text{ }\mu\text{M}$ each dNTP, and 5U of Advantage[®] 2 Taq polymerase (Clontech, Palo Alto, CA, USA). The PCR was performed on an MJ Research PTC-200 thermal cycler (MJ Research, Watertown, MA, USA) and consisted of 1 cycle at 94°C for 3 min, and then 40 cycles at 94°C for 30 s, at 60°C for 30 s, and 72°C for 2 min, with a final extension step of 72°C for 10 min. The products of these reactions were electrophoresed on 1.5% agarose gels and resulting bands visualized by ethidium bromide staining. Definitive bands representing PCR products of a predicted base pair size of the targeted gene were excised from the gel, and extracted and purified using a commercially available nucleic acid-binding resin (Qiaex II Gel extraction kit; Qiagen, Valencia, CA, USA).

Table 4 *Mytilus trossulus*-specific quantitative real-time polymerase chain reaction primers used in the analysis of mussels.

Gene	Forward primer sequence (5'→3')	Reverse primer sequence (5'→3')
<i>CaM</i>	TCTGTTGACAAAGATGGCG	GCATCTACTTCGTTAATCATGT
<i>Casp 8</i>	CCCAACCAGTAGTAACACCAGAC	GTATGAACCATGCCCTATATCA
<i>MIF</i>	TACACCCAGACCAAATGATG	TTCTCCTAATGCTCCAATACTG
<i>CNN</i>	ATACTCCGGCGGAGACAGT	TCTTCTTCGGAATCTCTTGT
<i>CHI</i>	ATATCATCTACTCATTCCGCA	AGTGATAGTTTCAAGGCTG
<i>CCOIV</i>	GATGTAGTGGCTCTCAAGGAT	AGATCTGTTTCCATTACCTGT
<i>HSP70</i>	GGTGGTGAAGACTTTGACAACAG	CTAGTTTGGCATCACGTAGAGC
<i>HSP90</i>	GATCTCCAACCTCATCTGATGC	GTGTGTTGTTATCCTTGTCTG
<i>HIFa</i>	ATACCTTGGCATCTCACAGAT	GACTTCTTCTTGTGGTGGTC
<i>MytB</i>	AATGTCTTCGTTGTTCCAG	AATGCCAGTTTCCACCTTG
<i>Myt</i>	GTTATTCTGGCTATCGCTCTTG	GTATAATGTCAAACAGAACGGGTC
<i>MT20</i>	GATCTACTAAGCAGACCAGC	TACATCCGGAACATCCACAG
<i>Cyp3</i>	AGTTACAGTACTTGGACAGATTCGT	TGCCTCAAGTAATGCCAGCCTCA
<i>P53</i>	TGTGTAGACTGAGGGATTCATTGG	TCACCTTCTTCATCAGTTTGTTTTT
<i>18s</i>	GTGCTCTTGACTGAGTGCTCG	CGAGGTCCTATTCCATTATCC

Real-time PCR

Real-time PCR reactions for the individual genes of interest and the housekeeping gene (18S) were run in separate wells (Bowen *et al.*, 2018). Briefly, 1 μ l of cDNA was added to a mix containing 12.5 μ l of QuantiTect Fast SYBR Green[®] Master Mix (5 mM Mg 2+) (Qiagen, Valencia, CA, USA), 0.5 μ l each of forward and reverse sequence specific primers (Invitrogen, Carlsbad, CA, USA), and 10.5 μ l of RNase-free water; total reaction mixture was 25 μ l. The reaction mixture cDNA samples for each gene of interest and 18S were loaded into Fast 96 well plates in duplicate and sealed with optical sealing tape (Applied Biosystems, Foster City, CA, USA). Reaction mixtures that contained water but no cDNA were used as negative controls.

Statistical analysis

Analysis of qPCR data was conducted using normalized C_T values (housekeeping gene threshold crossing subtracted from the gene of interest threshold crossing); the lower the normalized value, the more transcripts are present. A change in normalized value of 2 is approximately equivalent to a 4-fold change in the amount of the transcript. A normalized value of 34 or higher indicated a quantity less than the detection limit for transcription of that gene. Two samples contained values that were less than the detection limit for transcription and were thus omitted from statistical analyses because they were influential outliers that affected within and among site means.

For all data, gene transcription and biomarker assays, medians, 2.5% and 97.5% percentiles, and ranges were calculated (NCSS, Statistical and Power Analysis Software). As mussel age (based on ring counts) and size were not found to have significant effects on biomarkers or gene transcription, they were omitted from further analyses. We used generalized linear mixed effects models (GLMM) to estimate site means for each gene

transcription factor and biomarker parameter. We included sampling year as a random effect to account for variances introduced by collecting sample units over the course of 2 years. Input data were examined for normality prior to analysis; a log linear model was used for hemocyte count, RNA:DNA ratio, P450, and HSP40 biomarkers to normalize variances, and reported means were back-transformed into the original data scale. We fit separate models for each gene and biomarker with maximum likelihood estimation using the lme4 package in R 3.5.0 (*R Development Core Team, 2012*), which accounts for the unbalanced data using Satterthwaite's method. We conducted post-hoc Tukey tests of site level differences. To obtain site means for each year, we fit a Gaussian generalized linear model (GLM) with a site by year interaction. The resulting site means were used to test for correlations between genes and biomarker parameters. Relationships between gene transcript and biomarker data, as well as within gene transcript and biomarker data, were assessed in R using Pearson correlations. We considered correlations that were >0.30 (or <-0.30) to be of possible biological significance. To further evaluate relationships between and within responses, we used R to conduct principal components analysis (PCA). As for the Pearson correlations, we evaluated the relationships within gene transcript and biomarker datasets using all available data records, and the relationships between gene transcripts and biomarkers using site by year means obtained from the GLM interaction model. We visualized multivariate interactions by plotting the contributions of each variable to PC1 and PC2, and by plotting the quality of representation of each metric to all principal components that could be fit. We showed data from all dimensions that could be fit to better represent the exploratory nature of our analysis.

RESULTS

Biomarker and gene transcript medians, 2.5% and 97.5% percentiles, and ranges were determined for use in future monitoring efforts ([Table 5](#)). Site-specific medians and ranges for the biomarker and gene transcription assays for each year are provided as [Files S1](#) and [S2](#).

The Principal Components Analysis for the biomarkers, gene transcripts, and mix of biomarkers and gene transcripts revealed that three, five, and six principal components had eigenvalues >1 , respectively. The first three dimensions of the biomarker PCA explained 65% of the variance in the data, and 27% was attributed to the first principal component. The first five dimensions of the gene transcript PCA explained 69% of the variation in the data, with 31% going to the first component. Two-thirds of the variation was explained by six dimensions for the combined PCA, with 33% explained by the first principal. Only 12 dimensions could be fit for the combined dataset because of the sample size restrictions caused by using site means for each year.

Within the biomarker assays we found correlations between condition factor and shell thickness ($R_{\text{pearson}} = 0.48$), and condition factor and HSP40 ($R_{\text{pearson}} = 0.24$, [Fig. 2](#)). Condition factor, shell thickness and HSP40 also contributed the most to PC1 when the physiological markers were analyzed by PCA ([Fig. 3](#)), with shell thickness having the highest quality of representation on this axis ([Fig. 4](#)). Hydrogen peroxide production, P450 activity

Table 5 Medians, 2.5 and 97.5 percentiles and ranges obtained from 120 mussels collected at 6 sites in Lake Clark and Katmai National Parks and Preserves for all variables across all sites and years. For gene transcription, higher numbers indicate less transcription, and a value of 34 indicates a quantity less than detection limits.

Condition Factor			Shell Thickness (mm)			Hemocyte Count (cells mL ⁻¹)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
0.16	0.07–0.31	0.06–0.35	0.67	0.27–1.69	0.24–2.22	241,667	13,333–1,540,000	3,333–1,740,000
Hydrogen Peroxide (OD ₆₂₀)			RNA:DNA (ratio g ⁻¹)			P450 (activity mg ⁻¹ protein)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
0.05	0.01–0.10	0.01–0.14	69.49	4.09–663.70	1.74–833.49	27.64	9.48–127.94	7.62–190.33
HSP40 (Arbitrary Units)			CaM (C _T)			Casp8 (C _T)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
2.88	0.32–24.98	0.18–40.03	16.19	13.91–19.42	13.57–21.46	10.48	8.26–12.91	7.74–13.82
MIF (C _T)			CNN (C _T)			CHI (C _T)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
17.88	12.37–22.17	11.92–23.22	25.12	21.67–27.93	17.70–34	18.66	16.01–23.97	13.30–34
CCOIV (C _T)			HSP70 (C _T)			HSP90 (C _T)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
18.38	12.73–22.29	11.56–23.14	11.31	7.99–14.65	6.63–16.58	13.54	9.66–16.66	9.31–17.89
HIFa (C _T)			MytB (C _T)			Myt (C _T)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
14.13	12.24–16.43	11.86–17.02	12.6	5.80–18.69	4.67–19.72	15.91	12.02–19.24	10.69–22.11
MT20 (C _T)			Cyp3 (C _T)			P53 (C _T)		
Median	2.5–97.5%	Range	Median	2.5–97.5%	Range	Median	2.5–97.5%	Range
10.14	5.48–15.22	4.43–20.73	14.67	12.01–17.18	11.11–18.10	12.79	10.31–15.77	9.39–34

and RNA:DNA contributed to PC2, with RNA:DNA representing an opposite effect from the other two variables (Fig. 3). PC3 corresponded to hemocyte count. PC1-3 contributed 27.5%, 22.1% and 14.5%, respectively, to variation in the physiological data and 64.1% of overall variation.

Strong correlations were noted between many of the gene transcripts (Fig. 5). No strong negative correlations in levels of transcription were noted among the genes. Analysis of gene transcription by PCA indicated no gene transcripts were anticorrelated, matching the Pearson correlation results. PC1 corresponded to CaM, Casp8, CNN, HSP70, Myt, Cyp3 and p53 (Fig. 3) with Casp8 providing the highest quality of representation on this axis (Fig. 4). This was similar to the inference obtained by the Pearson correlations where PC1 maps to variables that have strong correlations (>0.30) with both CaM and Casp8. CHI, HSP70, HSP90 and HIFa contributed the most to PC2 (Fig. 3). MT20 contributed most substantially to PC3. PC1-3 corresponded to 31.3%, 13.6% and 8.9%, respectively, of variation in gene transcription data and 53.8% of variation overall.

Increased P450 activity was associated with increased transcription of 4 genes (most notably CCOIV) and decreased transcription of MT20. A higher condition factor was associated with increased gene transcription: nine of 14 genes had a negative correlation with condition factor that was >0.30. In contrast, we observed an inverse relation between




Figure 2 Pearson correlations between biomarkers obtained from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. The numbers are Pearson correlations between biomarkers. Bold red numbers are statistically significant correlations ($P < 0.05$). The graphs depict actual observations (black circles) and the best fit line (red) between paired biomarkers.

Full-size [DOI: 10.7717/peerj.7800/fig-2](https://doi.org/10.7717/peerj.7800/fig-2)

RNA:DNA and gene transcription, with nine of 14 genes having a correlation > -0.30 (Table 6).

PCA analysis showed that PC1 was comprised primarily of the gene transcripts: CaM, Casp8, CNN, HSP70, MytB, Myt, Cyp3, p53, and the biomarkers RNA:DNA and condition factor, with condition factor having an opposite relationship than the other measures (Fig. 3). PC2 corresponded to MIF, MT20, hydrogen peroxide production, P450, shell thickness and hemocyte count, with shell thickness and hemocyte count opposing the other variables (Fig. 3). CHI, HSP70, HSP90 and HIFa, with Myt, hydrogen peroxide production and P450 activity opposing those measures, contributed the most to PC3. PC1-3 corresponded to 33.3%, 19.4% and 12.5%, respectively, of the variance in the metrics and 65.3% of variance overall.

DISCUSSION

The overall goal of our research was to use biomarker and genetic methods to assess the condition of nearshore species in remote areas of Alaska, as nearshore ecosystems, particularly at higher latitudes, are vulnerable to changing environmental conditions (Taylor et al., 2017). Our approach included establishment of reference ranges for




Figure 3 Correlation plots based on Principle Components Analysis of variables measured from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. (A) Gene transcripts, (B) biomarkers, (C) all metrics. The arrows correspond to each measured variable and maps the contribution of that variable to the first and second principle components (Dim 1 and 2). Arrows that are close together are correlated, and those that are opposite are anti-correlated. Those orthogonal to one another are not correlated. Variables that contribute strongly to either the first or second axis are nearer the axes than variables that do not contribute strongly. The length and color of the arrow illustrates the strength of the contribution of each variable, with the longer arrows rendered in warm colors contributing more than those that are shorter and rendered in cool colors.

Full-size DOI: [10.7717/peerj.7800/fig-3](https://doi.org/10.7717/peerj.7800/fig-3)




Figure 4 Quality of representation plots based on Principle Components Analyses of variables measured from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. (A) gene transcripts, (B) biomarkers, (C) all metrics. The charts depict all metrics and all Principle Components that could be calculated for each analysis. The size and shade of the circle denotes how well the variable is represented by each principle component. The large and dark circles denote good representation, while the small and light circles denote poor representation. This plot has the advantage of showing potential variable correlations in multiple dimensions.

Full-size  DOI: [10.7717/peerj.7800/fig-4](https://doi.org/10.7717/peerj.7800/fig-4)




Figure 5 Pearson correlations between gene transcripts obtained from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. The numbers are Pearson correlations between gene transcripts. Red numbers are statistically significant correlations ($P < 0.05$). The graphs depict all observations (black circles) and the best fit line (red) between paired gene transcripts.

Full-size [DOI: 10.7717/peerj.7800/fig-5](https://doi.org/10.7717/peerj.7800/fig-5)

biomarker and gene transcription assays of mussels collected at sites in LACL and KATM and identification of correlations between the assays. Due to the remote nature of the parks and other logistical constraints, sampling opportunities were limited to a one week window in July 2015 and again in July 2016. Other environmental monitoring studies generally have been conducted in areas with known contamination concerns, allowing for a clear contrast among experimental groups (Akaishi et al., 2007; Banni et al., 2017; Carella et al., 2018; Gagné et al., 2007; Halldórsson et al., 2008; Sforzini et al., 2018; Shaw et al., 2011). However, our goal was to develop a method to monitor sites for changes, rather than assess sites already impacted. Therefore, our collection sites were not known to be compromised by anthropogenic activity and were considered relatively pristine. Any differences in biomarker or gene transcription results were expected to be due to natural variation and to provide a range of representative values which could act as a reference for continuing studies.

Table 6 Results of the Pearson correlations between gene transcripts and biomarkers obtained from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. Statistically significant correlations ($P < 0.05$) are bolded. Values reflect the numerical correlation; however, the biological correlation will be the inverse, as lower C_T values reflect higher levels of gene transcription.

	Condition factor	P450 activity	RNA:DNA
CaM	-0.40	0.04	0.59
Casp8	-0.49	-0.18	0.83
MIF	-0.33	0.26	0.11
CNN	-0.21	0.04	0.45
CHI	0.22	0.06	-0.25
CCOIV	-0.24	-0.71	0.39
HSP70	-0.47	-0.20	0.46
HSP90	-0.16	-0.34	0.18
HiFa	-0.54	-0.05	0.16
Myt B	-0.60	-0.34	0.57
Myt	-0.49	0.10	0.65
MT20	-0.08	0.62	-0.09
Cyp3	-0.71	-0.12	0.52

In this light, although differences in physiological patterns among sites are of interest, the more important goal is to identify the range of values that can be expected under what we assume to be “normal” conditions. Biomarker and gene transcript levels may differ between two sites, but both sites, nevertheless, may fall within a range considered to be normal. Many factors can cause natural variation in biomarker and gene transcription levels, including predators (*Reimer & Tedengren, 1996*), temperature, dissolved oxygen (*Abele & Puntarulo, 2004*) and wave exposure (*Dahlhoff & Menge, 1996*). There may have been differences in these and other physical (e.g., substrate type) or biological factors across the sites that contributed to normal background variation.

Several considerations must be taken into account when interpreting gene transcription data. First, within an individual, there are tradeoffs in the allocation of limited resources among competing physiological functions (*Vera-Massieu et al., 2015*). Increased physiological activity and gene transcription in response to stressors can present a metabolic challenge for any species, including bivalves (*Graham et al., 2010*), and may result in reduction of fitness evidenced by decreased reproductive capability, increased susceptibility to disease, or disadvantageous behavioral changes (*Graham et al., 2010; Martin et al., 2010*). Additionally, biological processes identified by our panel are influenced by multiple genes, and each gene may contribute to multiple functions. For example, although several genes on our panel respond to ocean acidification, including calmodulin (CaM), calponin (CNN), and chitinase (CHI), each plays a different role in the overall mechanism (*Li et al., 2004; Banni et al., 2011; Chen et al., 2012; Hüning et al., 2013; Li et al., 2016*). Thus, transcript levels of genes with similar endpoint functions may not necessarily correlate with each other. Identifying patterns of variation for each gene measured (among and within sites) is an important step in determining their value for monitoring studies.

Significant differences were observed among sites for biomarker as well as gene transcription parameters. For example, five out of seven biomarkers varied significantly among sites (condition factor, shell thickness, RNA:DNA ratio, HSP40, hemocyte count) (Fig. 6), and nine of 14 genes varied significantly among sites (CNN, CaM, CHI, Cyp3, HIFa, HSP70, HSP90, MT20, and Myt) (Fig. 7). Given that we had no reason to expect differences among sites due to obvious impairments, we might assume that the differences we detected were part of the natural variation we would observe in these populations when sites are compared across a large spatial scale.

Condition factor and shell thickness were higher at Chinitna Bay, Fossil Point, and Silver Salmon (all in LACL) as compared to Kafia, Kukak, and Takli (all in KATM). Elevated mussel condition factor has been associated with the presence of higher quality and/or quantity of nutrients (Carmichael, Shriver & Valiela, 2004), suggesting that nutrient availability varied between parks. Water originating from Upper Cook Inlet flows along the LACL coastline, eventually merging with the Alaska Coastal Current (Nagorski et al., 2008). The KATM coastline is dominated by the Alaska Coastal Current, which carries a high amount of freshwater to the region (Nagorski et al., 2007). Differences in oceanographic processes between LACL and KATM likely influence nutrient availability along the coast.

Shell thickness can be influenced by changes in predation pressure, mussel density or abiotic factors. Studies conducted with *M. galloprovincialis* have demonstrated that mussels living at higher densities are smaller with thicker shells (Xavier, Branch & Wieters, 2007). Predation can induce mussels to thicken their shells as a defense mechanism (Freeman, 2007). Abiotic factors such as temperature, salinity and wave action can influence shell thickness, as well (Akester & Martel, 2000; Blanchard & Feder, 2000). Based on observations, Chinitna Bay is more exposed than the other sites and mussels at that location had the thickest shells, potentially as a result of experiencing more wave action. However, mussel density, predators and other abiotic factors were not quantified during this study.

Mussel hemocyte count was the most variable biomarker within sites, and other studies have observed similar variability (Akaishi et al., 2007; Coray, St.Jean & Bard, 2007; Duchemin et al., 2008). Mussels at Kafia had a significantly lower hemocyte count than mussels at the other sites. This result indicates that despite relatively high variability, differences in hemocyte count can be identified. Wild mussels are constantly exposed to antigens that may stimulate an immune response and elevate hemocyte count with high variability between individuals (Galloway & Depledge, 2001). Less variability was observed in the hydrogen peroxide assay suggesting it may be a more suitable biomarker for monitoring immune activity than hemocyte count.

Variability among sites was also observed in RNA:DNA and HSP40. Mussels at Kafia and Kukak had significantly higher RNA:DNA compared to those from Silver Salmon, indicating differences in protein production among sites. HSP40 levels were higher at Kukak, Chinitna Bay and Fossil Point as compared to Kafia suggesting an elevated response to an unknown stressor at those three sites.

Relationships within the biomarker assays, within the gene transcription panel, and between the biomarkers and genes were determined using Pearson correlations and PCA, and the results of both analyses were complementary. A positive Pearson correlation was




Figure 6 Boxplots of biomarker data obtained from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. Random effects model results are denoted by red diamonds (mean) and red arrows (95% confidence intervals). Sites sharing a lowercase letter did not differ statistically based on post-hoc testing ($P < 0.05$). (A) condition factor, (B) shell thickness, (C) hemocyte count, (D) hydrogen peroxide, (E) RNA:DNA, (F) P450 activity, (G) HSP40.

Full-size DOI: 10.7717/peerj.7800/fig-6




Figure 7 Boxplots of gene transcription data (normalized C_T values) obtained from 120 mussels collected at six sites in Lake Clark and Katmai National Parks and Preserves. Random effects model results are denoted by red diamonds (mean) and red arrows (95% confidence intervals). Sites sharing a low-ercase letter did not differ statistically based on post-hoc Tukey testing ($P < 0.05$). (A) CaM, (B) Casp8, (C) MIF, (D) CNN, (E) CHI, (F) CCOIV, (G) HSP70, (H) HSP90, (I) HIFa, (J) MytB, (K) Myt, (L) MT20, (M) Cyp3, (N) p53.

Full-size DOI: 10.7717/peerj.7800/fig-7

found between condition factor and shell thickness and between shell thickness and HSP40. Additionally, in the PCA analysis, condition factor, shell thickness and HSP40 contributed to PC1 with high quality of representation on that axis. Condition factor indicates the nutritional status of the mussels, and it was not surprising that shell thickness would relate to overall condition as mussels with more nutrients might be expected to allocate more nutrients to shell formation. Mussels with nutrient deficits have been shown to metabolize their shell (*Masthanamma, Purushotham & Ramamurthi, 1984*). Shell thickness influences body temperature (*Caddy-Retalic, Benkendorff & Fairweather, 2011*), and possibly mussels with thicker shells absorb more heat, resulting in HSP40 production. Additional research is needed to understand the association between shell formation and HSP40 production.

Given the multiple functions of many of the genes in our transcript panel, as well as the interconnectedness of genes in general, we expected and found numerous correlations among the genes (*Fig. 5*). For example, CaM, CNN, and CHI are all sensitive to changes in ocean acidification, and Casp8, MIF, MytB, Myt, HSP70, and HSP90 can all be influenced by pathogen exposure. Contaminant exposure can influence the transcription of Casp8, HSP70, HSP90, MT20, Cyp3, and p53, and changes in dissolved oxygen can alter the transcription of CaM, CNN, CHI, CCOIV, and HIFa. As expected, there was a Pearson correlation between CNN and CaM, but not with CHI, as we would have anticipated. The strong correlations observed among CNN, Casp8 and Myt suggest potential links among processes associated with shell formation and pathogen presence. In addition, Casp8 was strongly correlated with Cyp3, MytB, and p53, suggesting a link between pathogen and contaminant exposures. CaM, Casp8, CNN, HSP70, Myt, Cyp3 and p53 contributed substantially to PC1 according to the PCA.

An objective of our study was the comparison of the two methodologies, gene transcription and biomarker assays. Several of the biomarkers and genes were associated with similar physiological functions, and we anticipated correlations would arise. Although we did not expect complete agreement, we expected the methods to support one another. Several correlations between the biomarker and gene transcription assays were identified. CaM, Casp8, CNN, HSP70, MytB, Myt, Cyp3, p53, RNA:DNA and condition factor contributed to PC1 in the PCA analysis of all metrics, with condition factor having an opposing relationship to the other variables. A negative Pearson correlation between condition factor and MytB, Cyp3 and p53 was identified, while a positive correlation between RNA:DNA and CaM, Casp8, Myt and p53 was present. The statistical analysis used the C_T values for gene transcription, and high C_T values indicate low gene expression. Therefore, the relationship between high gene expression and high condition factor makes biological sense, as a good condition factor implies ample resources, which translates into an overall ability to increase transcription of genes needed to combat stressors. RNA:DNA was associated with lower transcription of several genes, which was not an expected finding. However, the small number of genes in this study (14) may not be representative of patterns in the overall transcriptome, which numbers many thousands of genes.

Higher activity of P450 was strongly associated with lower MT20 transcription and increased CCOIV transcription in Pearson correlations. In PCA analysis, MT20 and P450 contributed to PC2. MT20 and P450 are responsible for detoxifying metals and xenobiotics,

respectively. Contaminants can contain complex chemical mixtures that induce different detoxification pathways (Islam & Tanaka, 2004). CCOIV is transcribed during hypoxic conditions, and contaminants can damage or interfere with gill function inhibiting respiration and resulting in hypoxia (Sokolova & Lannig, 2008). We expected P450 activity to correlate with Cyp3 transcription, but this was not observed. The cytochrome P450 family, which includes Cyp3, contains numerous proteins (Rewitz et al., 2006). The gene transcription assay is specific for Cyp3, but the P450 assay detects the enzyme activity of multiple cytochrome isoforms, not only Cyp3 (Kobayashi et al., 2002). Therefore, overall P450 enzyme activity may be increased while Cyp3 transcription is not.

The results of this study examined the strength of using both gene transcription and biomarker assays to evaluate coastal ecosystems. This approach has been implemented for mussel biomonitoring in other coastal regions including the Mediterranean (Carella et al., 2018; Sforzini et al., 2018). In both of these studies, site-level differences were detected using genetic techniques and biomarker analyses. The results correlated with the presence of known contamination (Carella et al., 2018; Sforzini et al., 2018). In our study, differences between sites also were identified, but without known sources of contamination. A limitation of our findings is that biomarker and gene transcription assays were not run on the same individual mussels (i.e., ten mussels were collected for each set of assays at each site in each year). Therefore, our correlations are computed on a site basis, and we are not able to evaluate how metrics from the two sets of assays relate on an individual mussel basis.

At the beginning of this study, we made assumptions about the pristine nature of the intertidal sites (Weeks, 1999; Weeks, 2003). The sites are along Alaskan national park coastlines, with very little direct anthropogenic stressors. However, variations in biomarkers and genes suggest some differences exist among sites. Most notably, mussels from LACL were in better condition with thicker shells, perhaps associated with quantity and/or quality of nutrients in the vicinity of those sites. Developing reference biomarker and gene transcription levels in mussels is important to properly differentiate among changes due to climate or anthropogenic activity and natural variation. The capability to discern local and large-scale changes is beneficial for monitoring remote locations where sampling opportunities are limited. Effects of some environmental impacts, such as oil spills, may be readily observed, but subtle changes in the marine environment can be difficult to detect. The sensitivity of biomarker and gene transcription assays supports the identification of subtle changes. Additionally, this approach enables the measurement of cumulative effects of several stressors. However, further studies comparing the biomarker and gene transcription assays, under controlled conditions and in the field, with both sets of assays measured on the same individual mussels, are needed to provide a more thorough understanding of how the various metrics respond and relate.

In addition to the need for controlled exposure studies, our results highlight an important reality pertaining to ecosystem monitoring: although combined technologies provide better resolution of potential causative factors, an approach using longitudinal monitoring to continually assess populations for subtle yet significant changes is necessary to provide insight into ecosystem health. In the future, we will improve our understanding of

ecosystems at risk if we take a proactive approach to monitoring prior to occurrence of population level effects (Bennett, Thompson & Mortenson, 2006).

CONCLUSIONS

Intertidal communities are important in marine and terrestrial food webs and understanding the condition of these intertidal resources will support management in the maintenance of healthy coastal ecosystems. This study generated baseline gene transcription and biomarker data that will be useful for monitoring these remote areas of the Alaskan coastline for environmental change. The gene transcription and biomarker assays were advantageous because they provided a significant amount of information regarding physiological responses of mussels to environmental conditions from small amounts of tissue, which is necessary for assessing remote intertidal communities where sampling opportunities and environmental data are constrained by logistics. Additionally, the assays were sensitive enough to detect differences between sites with no obvious impacts. Results between the gene transcription and biomarker assays were often complementary, but some differences were noted, likely due to variations in the rate of production and turnover of mRNA and proteins. Integrating gene transcription and biomarker assays provided a more comprehensive assessment of mussel condition than either approach alone. Additional controlled experiments will strengthen our understanding of the value of this approach. We anticipate implementing this holistic approach to evaluate stressors affecting intertidal communities and changes occurring within those communities over time.

ACKNOWLEDGEMENTS

We would like to thank the NPS LACL staff, Jim Bodkin (USGS), Carissa Turner (NPS), Bob “Bunyan” Peterson (KATM), A.K. Miles (USGS), Shannon Waters (USGS), Jennifer Goetsch (ASLC) and Emily Johnson (ASLC) for assistance with this project. Gulf Watch Alaska supported field efforts in Katmai National Park and Preserve. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the National Park Service Park Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Park Service Park Foundation.

Competing Interests

Katrina Counihan is employed by the Alaska SeaLife Center.

Author Contributions

- Katrina L. Counihan, Lizabeth Bowen, Brenda Ballachey, Heather Coletti, Tuula Hollmen and Benjamin Pister conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Tammy L. Wilson analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Field collections were approved by the National Park Service. Field collection permit numbers are Alaska Department of Fish and Game permit # CF-15-088 for Katmai and Lake Clark National Parks and Preserves, 2015, and # CF-16-089 for Katmai and Lake Clark National Parks and Preserves, 2016.

Data Availability

The following information was supplied regarding data availability:

The raw measurements and code used in the statistical analyses are available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7800#supplemental-information>.

REFERENCES

- Abele D, Puntarulo S. 2004.** Formation of reactive species and induction of antioxidant defense systems in polar and temperate marine invertebrates and fish. *Comparative Biochemistry and Physiology Part A Molecular & Integrative Physiology* **138**:405–415 DOI [10.1016/j.cbpb.2004.05.013](https://doi.org/10.1016/j.cbpb.2004.05.013).
- Akaishi F, St-Jean S, Bishay F, Clarke J, Rabitto I, De Oliveira Ribeiro C. 2007.** Immunological responses, histopathological finding and disease resistance of blue mussel (*Mytilus edulis*) exposed to treated and untreated municipal wastewater. *Aquatic Toxicology* **82**:1–14 DOI [10.1016/j.aquatox.2007.01.008](https://doi.org/10.1016/j.aquatox.2007.01.008).
- Akester R, Martel A. 2000.** Shell shape, dysodont tooth morphology and hinge-ligament thickness in the bay mussel *Mytilus trossulus* correlate with wave exposure. *Canadian Journal of Zoology* **78**:240–253 DOI [10.1139/z99-215](https://doi.org/10.1139/z99-215).
- Armstrong DA, Dinnel PA, Orensanz JM, Armstrong JL, McDonald TL, Cusimano RF, Nemeth RS, Landolt ML, Skalski JR, Lee RF, Huggett RJ. 1995.** Status of selected bottomfish and crustacean species in Prince William Sound following the Exxon Valdez oil spill. In: Wells PG, Butler JN, Hughes JS, eds. *Exxon Valdez oil spill: fate and effects in Alaskan waters*. Philadelphia: American Society for Testing and Materials, 485–547.

- Balseiro P, Falcó A, Romero A, Dios S, Martínez-López A, Figueras A, Estepa A, Novoa B. 2011.** *Mytilus galloprovincialis* Myticin C: a chemotactic molecular with antiviral activity and immunoregulatory properties. *PLOS ONE* 6:e23140 DOI 10.1371/journal.pone.0023140.
- Banni M, Dondero F, Jebali J, Guerbej H, Boussetta H, Viarengo A. 2007.** Assessment of heavy metal contamination using real-time PCR analysis of mussel metallothionein mt10 and mt20 expression: a validation along the Tunisian coast. *Biomarkers* 12:369–383 DOI 10.1080/13547500701217061.
- Banni M, Negri A, Mignone F, Boussetta H, Viarengo A, Dondero F. 2011.** Gene expression rhythms in the mussel *Mytilus galloprovincialis* (Lam.) across an annual cycle. *PLOS ONE* 6:e18904 DOI 10.1371/journal.pone.0018904.
- Banni M, Negri A, Rebelo M, Rapallo F, Boussetta H, Viarengo A, Dondero F. 2009.** Expression analysis of the molluscan p53 protein family mRNA in mussels (*Mytilus* spp.) exposed to organic contaminants. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 149:414–418 DOI 10.1016/j.cbpc.2008.09.017.
- Banni M, Sforzini S, Arlt VM, Barranger A, Dallas LJ, Oliveri C, Aminot Y, Pacchioni B, Millino C, Lanfranchi G, Readman JW, Moore MN, Viarengo A, Jha AN. 2017.** Assessing the impact of Benzo [a] pyrene on marine mussels: application of a novel targeted low density microarray complementing classical biomarker responses. *PLOS ONE* 12:e0178460 DOI 10.1371/journal.pone.0178460.
- Baxter RE. 1971.** Earthquake effects on clams of Prince William Sound. In: *The great Alaska earthquake of 1964*. Washington, D.C: National Academy of Sciences, 238–245.
- Bennett AJ, Thompson WL, Mortenson DC. 2006.** *Vital signs monitoring plan, Southwest Alaska Network, inventory and monitoring program*. Anchorage: National Park Service, Southwest Alaska Network.
- Beyer J, Green NW, Brooks S, Allen IJ, Ruus A, Gomes T, Brate ILN, Schoyen M. 2017.** Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: a review. *Marine Environmental Research* 130:338–365 DOI 10.1016/j.marenvres.2017.07.024.
- Bijma J, Pörtner HO, Yesson C, Rogers AD. 2013.** Climate change and the oceans—what does the future hold? *Marine Pollution Bulletin* 74:495–505 DOI 10.1016/j.marpolbul.2013.07.022.
- Blanchard A, Feder H. 2000.** Shell growth of *Mytilus trossulus* Gould, 1850, in Port Valdez, Alaska. *The Veliger* 43:34–42.
- Bodkin JL, Coletti HA, Ballachey BE, Monson D, Esler D, Dean TA. 2017.** Variation in abundance of Pacific blue mussel (*Mytilus trossulus*) in the Northern Gulf of Alaska, 2006–2015. *Deep Sea Research Part II: Tropical Studies in Oceanography* 147:87–97 DOI 10.1016/j.dsr2.2017.04.008.
- Bolognesi C, Cirillo S. 2014.** Genotoxicity biomarkers in aquatic bioindicators. *Current Zoology* 60:273–284 DOI 10.1093/czoolo/60.2.273.
- Bowen L, Miles AK, Ballachey B, Waters S, Bodkin J, Lindeberg M, Esler D. 2018.** Gene transcription patterns in response to low level petroleum contaminants in *Mytilus*

- trossulus* from field sites and harbors in southcentral Alaska. *Deep Sea Research Part II* 147:27–35 DOI 10.1016/j.dsr2.2017.08.007.
- Breitburg D, Riedel G. 2005.** Multiple stressors in marine systems. In: Norse E, Crowder L, eds. *Marine conservation biology: the science of maintaining the sea's biodiversity*. Washington, D.C.: Island Press, 167–182.
- Caddy-Retalic S, Benkendorff K, Fairweather P. 2011.** Visualizing hotspots: applying thermal imaging to monitor internal temperatures in intertidal gastropods. *Molluscan Research* 31:106–113.
- Caldarone EM, Wagner M, Onge-Burns JS, Buckley LJ. 2001.** Protocol and guide for estimating nucleic acids in larval fish using a fluorescence microplate reader. In: *Northeast Fisheries Science Center Reference Document 01–11*. Woods Hole: National Marine Fisheries Service, 22.
- Calkins DG. 1978.** Feeding behavior and major prey species of the sea otter, *Enhydra lutris*, in Montague Strait, Prince William Sound, Alaska. *Fishery Bulletin* 76:125–131.
- Carella F, Aceto S, Mangoni O, Mollica MP, Cavaliere G, Trinchese G, Aniello F, De Vico G. 2018.** Assessment of the health status of mussels *Mytilus galloprovincialis* along the Campania coastal areas: a multidisciplinary approach. *Frontiers in Physiology* 9:683 DOI 10.3389/fphys.2018.00683.
- Carmichael RH, Shriver AC, Valiela I. 2004.** Changes in shell and soft tissue growth, tissue composition, and survival of quahogs, *Mercenaria*, and softshell clams, *Mya arenaria*, in response to eutrophic-driven changes in food supply and habitat. *Journal of Experimental Marine Biology and Ecology* 313:75–104 DOI 10.1016/j.jembe.2004.08.006.
- Cellura C, Toubiana M, Parrinello N, Roch P. 2007.** Specific expression of antimicrobial peptide and HSP70 genes in response to heat-shock and several bacterial challenges in mussels. *Fish & Shellfish Immunology* 22:340–350 DOI 10.1016/j.fsi.2006.06.007.
- Châtel A, Faucet-Marquis V, Gourlay-Francé C, Pfohl-Leszkowicz A, Vincent-Hubert F. 2015.** Genotoxicity and activation of cellular defenses in transplanted zebra mussels *Dreissena polymorpha* along the Seine river. *Ecotoxicology and Environmental Safety* 114:241–249 DOI 10.1016/j.ecoenv.2014.03.023.
- Chen ZF, Wang H, Matsumura K, Qian PY. 2012.** Expression of calmodulin and myosin light chain kinase during larval settlement of the barnacle *Balanus amphitrite*. *PLOS ONE* 7:e31337 DOI 10.1371/journal.pone.0031337.
- Coletti HA, Bodkin JL, Monson DH, Ballachey BE, Dean TA. 2016.** Detecting and inferring cause of change in an Alaska nearshore marine ecosystem. *Ecosphere* 7:e01489 DOI 10.1002/ecs2.1489.
- Coray C, St. Jean S, Bard S. 2007.** Immunomodulation in blue mussels (*Mytilus edulis* and *Mytilus trossulus*) exposed to municipal effluent in Eastern Canada. *Water Quality Research Journal Canada* 42:194–201 DOI 10.2166/wqrj.2007.023.
- Cruz-Rodriguez L, Chu F. 2002.** Heat-shock protein (HSP70) response in the eastern oyster, *Crassostrea virginica*, exposed to PAHs sorbed to suspended artificial clay

- particles and to suspended field contaminated sediments. *Aquatic Toxicology* **60**:157–168 DOI [10.1016/S0166-445X\(02\)00008-5](https://doi.org/10.1016/S0166-445X(02)00008-5).
- Dahlhoff E, Menge B. 1996.** Influence of phytoplankton concentration and wave exposure on the ecophysiology of *Mytilus californianus*. *Marine Ecology Progress Series* **144**:97–107 DOI [10.3354/meps144097](https://doi.org/10.3354/meps144097).
- De Maio AM. 1999.** Heat shock proteins: facts, thoughts, and dreams. *Shock* **11**:1–2.
- De Sousa Abreu R, Penalva L, Marcotte E, Vogel C. 2009.** Global signatures of protein and mRNA expression levels. *Molecular BioSystems* **5**(12):1512–1526 DOI [10.1039/b908315d](https://doi.org/10.1039/b908315d).
- Dean TA, Bodkin JL, Coletti HA. 2014.** Protocol narrative for nearshore marine ecosystem monitoring in the Gulf of Alaska, version 1.1. Natural resource report NPS/SWAN/NRR-2014/756. National Parks Service, Fort Collins, Colorado.
- DeGange AR, Byrd GV, Walker LR, Waythomas CF. 2010.** Introduction—the impacts of the 2008 eruption of Kasatochi Volcano on the terrestrial and marine ecosystems in the Aleutian Islands, Alaska. *Arctic, Antarctic, and Alpine Research* **42**:245–249 DOI [10.1657/1938-4246-42.3.245](https://doi.org/10.1657/1938-4246-42.3.245).
- Denny M, Wetthey D. 2001.** Physical processes that generate patterns in marine communities. In: Bertness M, Hay M, Gaines S, eds. *Marine community ecology*. Sunderland: Sinauer Press, 3–37.
- Di Y, Schroeder DC, Highfield A, Readman JW, Jha AN. 2011.** Tissue-specific expression of p53 and ras genes in response to the environmental genotoxicant benzo (α) pyrene in marine mussels. *Environmental Science & Technology* **45**:8974–8981 DOI [10.1021/es201547x](https://doi.org/10.1021/es201547x).
- Dondero F, Dagnino A, Jonsson H, Capri F, Gastaldi L, Viarengo A. 2006.** Assessing the occurrence of a stress syndrome in mussels (*Mytilus edulis*) using a combined biomarker/gene expression approach. *Aquatic Toxicology* **78**:S13–S24 DOI [10.1016/j.aquatox.2006.02.025](https://doi.org/10.1016/j.aquatox.2006.02.025).
- Doroff AM, DeGange AR. 1994.** Sea otter, *Enhydra lutris*, prey composition and foraging success in the northern Kodiak Archipelago. *Fishery Bulletin* **92**:704–710.
- Driskell WB, Fukuyama AK, Houghton JA, Lees DC, Mearns AJ, Shigenaka G. 1996.** Recovery of Prince William Sound intertidal infauna from Exxon Valdez oiling and shoreline treatments, 1989 through 1992. *American Fisheries Society Symposium* **18**:362–378.
- Duchemin M, Auffret M, Wessel N, Fortier M, Morin Y, Pellerin J, Fournier M. 2008.** Multiple experimental approaches of immunotoxic effects of mercury chloride in the blue mussel, *Mytilus edulis*, through *in vivo*, *in tubo* and *in vitro* exposures. *Environmental Pollution* **153**:416–423 DOI [10.1016/j.envpol.2007.08.015](https://doi.org/10.1016/j.envpol.2007.08.015).
- Evans TG, Hofmann GE. 2012.** Defining the limits of physiological plasticity: how gene expression can assess and predict the consequences of ocean change. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **367**:1733–1745 DOI [10.1098/rstb.2012.0019](https://doi.org/10.1098/rstb.2012.0019).
- Fall JA, Field LJ. 1996.** Subsistence uses of fish and wildlife before and after the Exxon Valdez oil spill. In: Rice SD, Spies RB, Wolfe DA, Wright BA, eds. *Proceedings of*

- the Exxon valdez oil spill symposium*, vol. 18. Bethesda: American Fisheries Society Symposium, 819–836.
- Franzellitti S, Buratti S, Donnini F, Fabbri E. 2010.** Exposure of mussels to a polluted environment: insights into the stress syndrome development. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **152**:24–33 DOI [10.1016/j.cbpc.2010.02.010](https://doi.org/10.1016/j.cbpc.2010.02.010).
- Freeman A. 2007.** Specificity of induced defenses in *Mytilus edulis* and asymmetrical predator deterrence. *Marine Ecology Progress Series* **334**:145–153 DOI [10.3354/meps334145](https://doi.org/10.3354/meps334145).
- Fukuda R, Zhang H, Kim JW, Shimoda L, Dang CV, Semenza GL. 2007.** HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells. *Cell* **129**:111–122 DOI [10.1016/j.cell.2007.01.047](https://doi.org/10.1016/j.cell.2007.01.047).
- Fukuyama AK, Oliver JS. 1985.** Sea star and walrus predation on bivalves in Norton Sound, Bering Sea, Alaska. *Ophelia* **24**:17–36 DOI [10.1080/00785236.1985.10426616](https://doi.org/10.1080/00785236.1985.10426616).
- Fukuyama AK, Shigenaka G, Hoff RZ. 2000.** Effects of residual Exxon Valdez oil on intertidal *Protothaca staminea*: mortality, growth, and bioaccumulation of hydrocarbons in transplanted clams. *Marine Pollution Bulletin* **40**:1042–1050 DOI [10.1016/S0025-326X\(00\)00055-2](https://doi.org/10.1016/S0025-326X(00)00055-2).
- Gagné F, Blaise C, André C, Gagnon C, Salazar M. 2007.** Neuroendocrine disruption and health effects in *Elliptio complanata* mussels exposed to aeration lagoons for wastewater treatment. *Chemosphere* **68**:731–743 DOI [10.1016/j.chemosphere.2006.12.101](https://doi.org/10.1016/j.chemosphere.2006.12.101).
- Galloway TS, Depledge MH. 2001.** Immunotoxicity in invertebrates: measurement and ecotoxicological relevance. *Ecotoxicology* **10**:5–23 DOI [10.1023/A:1008939520263](https://doi.org/10.1023/A:1008939520263).
- Gaylord B, Hill T, Sanford E, Lenz E, Jacobs L, Sato K, Russell A, Hettinger A. 2011.** Functional impacts of ocean acidification in an ecologically critical foundation species. *The Journal of Experimental Biology* **214**:2586–2594 DOI [10.1242/jeb.055939](https://doi.org/10.1242/jeb.055939).
- Giannetto A, Maisano M, Cappello T, Oliva S, Parrino V, Natalotto A, De Marco G, Barberi C, Romeo O, Mauceri A, Fasulo S. 2015.** Hypoxia-inducible factor α and Hif-prolyl hydroxylase characterization and gene expression in short-time air-exposed *Mytilus galloprovincialis*. *Marine Biotechnology* **17**:768–781 DOI [10.1007/s10126-015-9655-7](https://doi.org/10.1007/s10126-015-9655-7).
- Gill Jr RE, Handel CM. 1990.** The importance of subarctic intertidal habitats to shorebirds: a study of the central Yukon-Kuskokwim Delta, Alaska. *The Condor* **92**:709–725 DOI [10.2307/1368690](https://doi.org/10.2307/1368690).
- Giuliani ME, Benedetti M, Arukwe A, Regoli F. 2013.** Transcriptional and catalytic responses of antioxidant and biotransformation pathways in mussels, *Mytilus galloprovincialis*, exposed to chemical mixtures. *Aquatic Toxicology* **134–135**:120–127 DOI [10.1016/j.aquatox.2013.03.012](https://doi.org/10.1016/j.aquatox.2013.03.012).
- Goodson MS, Crookes-Goodson WJ, Kimbell JR, McFall-Ngai MJ. 2006.** Characterization and role of p53 family members in the symbiont-induced morphogenesis of the *Euprymna scolopes* light organ. *The Biological Bulletin* **211**:7–17 DOI [10.2307/4134573](https://doi.org/10.2307/4134573).

- Graham AL, Shuker DM, Pollitt LC, Auld SKJR, Wilson AJ, Little TJ. 2010.** Fitness consequences of immune responses: strengthening the empirical framework for ecoimmunology. *Functional Ecology* 25:1–13 DOI [10.1111/j.1365-2435.2010.01777.x](https://doi.org/10.1111/j.1365-2435.2010.01777.x).
- Gunderson AR, Armstrong EJ, Stillman JH. 2016.** Multiple stressors in a changing world: the need for an improved perspective on physiological responses to the dynamic marine environment. *Annual Review of Marine Science* 8:357–378 DOI [10.1146/annurev-marine-122414-033953](https://doi.org/10.1146/annurev-marine-122414-033953).
- Halldórsson HP, De Pirro M, Romano C, Svavarsson J, Sarà G. 2008.** Immediate biomarker responses to benzo[a]pyrene in polluted and unpolluted population of the blue mussel (*Mytilus edulis* L.) at high-latitudes. *Environmental International* 34:483–489 DOI [10.1016/j.envint.2007.11.002](https://doi.org/10.1016/j.envint.2007.11.002).
- Halldórsson HP, Svavarsson J, Granmo Å. 2005.** The effect of pollution on scope for growth of the mussel (*Mytilus edulis* L.) in Iceland. *Marine Environmental Research* 59:47–64 DOI [10.1016/j.marenvres.2004.02.001](https://doi.org/10.1016/j.marenvres.2004.02.001).
- Harley C. 2011.** Climate change, keystone predation and biodiversity loss. *Science* 334:1124–1127 DOI [10.1126/science.1210199](https://doi.org/10.1126/science.1210199).
- Hoegh-Guldberg O, Bruno JF. 2010.** The impact of climate change on the world's marine ecosystems. *Science* 328:1523–1528 DOI [10.1126/science.1189930](https://doi.org/10.1126/science.1189930).
- Hüning AK, Melzner F, Thomsen J, Gutowska MA, Kramer L, Frickenhaus S, Rosenstiel P, Pörtner HO, Philipp EER, Lucassen M. 2013.** Impacts of seawater acidification on mantle gene expression patterns of the Baltic Sea blue mussel: implications for shell formation and energy metabolism. *Marine Biology* 160:1845–1861 DOI [10.1007/s00227-012-1930-9](https://doi.org/10.1007/s00227-012-1930-9).
- Islam MS, Tanaka M. 2004.** Impacts of pollution on coastal and marine ecosystems including coastal and marine fisheries and approach for management: a review and synthesis. *Marine Pollution Bulletin* 48:624–649 DOI [10.1016/j.marpolbul.2003.12.004](https://doi.org/10.1016/j.marpolbul.2003.12.004).
- Iwama GK, Mathilakath MV, Forsyth RB, Ackerman PA. 1999.** Heat shock proteins and physiological stress in fish. *American Zoology* 39:901–909 DOI [10.1093/icb/39.6.901](https://doi.org/10.1093/icb/39.6.901).
- Jackson J, Kirby M, Berger W, Bjorndal K, Botsford L, Bourque B, Bradbury R, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan HS, Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ, Warner RR. 2001.** Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293:629–638 DOI [10.1126/science.1059199](https://doi.org/10.1126/science.1059199).
- Jamieson G. 1993.** Marine invertebrate conservation: evaluation of fisheries over-exploitation concerns. *American Zoology* 33:551–567 DOI [10.1093/icb/33.6.551](https://doi.org/10.1093/icb/33.6.551).
- Jewett SC, Drew GS. 2014.** Recolonization of the intertidal and shallow subtidal community following the 2008 eruption of Alaska's Kasatochi Volcano. *Biogeosciences Discussions* 11:3799–3836 DOI [10.5194/bgd-11-3799-2014](https://doi.org/10.5194/bgd-11-3799-2014).
- Kobayashi K, Urashima K, Shimada N, Chiba K. 2002.** Substrate specificity for rat cytochrome P450 (CYP) isoforms: screening with cDNA-expressed systems of the rat. *Biochemical Pharmacology* 63:889–896 DOI [10.1016/S0006-2952\(01\)00843-7](https://doi.org/10.1016/S0006-2952(01)00843-7).
- Lacroix C, Coquillé V, Guyomarch J, Auffret M, Moraga D. 2014.** A selection of reference genes and early-warning mRNA biomarkers for environmental monitoring

- using *Mytilus* spp. as sentinel species. *Marine Pollution Bulletin* **86**:304–313
[DOI 10.1016/j.marpolbul.2014.06.049](https://doi.org/10.1016/j.marpolbul.2014.06.049).
- Lees D, Houghton JP, Driskell WB. 1996.** Short-term effects of several types of shoreline treatment on rocky intertidal biota in Prince William Sound. In: Rice SD, Spies RB, Wolfe DA, Wright BA, eds. *Exxon Valdez oil spill symposium proceedings*, vol. 18. Bethesda: American Fisheries Society Symposium, 329–348.
- Lesser M, Bailey M, Merselis D, Morrison J. 2010.** Physiological response of the blue mussel *Mytilus edulis* to differences in food and temperature in the Gulf of Maine. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **156**:541–551 [DOI 10.1016/j.cbpa.2010.04.012](https://doi.org/10.1016/j.cbpa.2010.04.012).
- Lessios H. 1988.** Mass mortality of *Diadema antillarum* in the Caribbean: what have we learned? *Annual Review of Ecology and Systematics* **19**:371–393
[DOI 10.1146/annurev.es.19.110188.002103](https://doi.org/10.1146/annurev.es.19.110188.002103).
- Lewis TL, Esler D, Boyd WS. 2007.** Effects of predation by sea ducks on clam abundance in soft-bottom intertidal habitats. *Marine Ecology Progress Series* **329**:131–144
[DOI 10.3354/meps329131](https://doi.org/10.3354/meps329131).
- Li H, Venier P, Prado-Alvarez M, Gestal C, Toubiana M, Quartesan R, Borghesan F, Novoa B, Figueras A, Roch P. 2010.** Expression of *Mytilus* immune genes in response to experimental challenges varied according to the site of collection. *Fish & Shellfish Immunology* **28**:640–648 [DOI 10.1016/j.fsi.2009.12.022](https://doi.org/10.1016/j.fsi.2009.12.022).
- Li S, Huang J, Liu C, Liu Y, Zheng G, Xie L, Zhang R. 2016.** Interactive effects of seawater acidification and elevated temperature on the transcriptome and biomineralization in the pearl oyster *Pinctada fucata*. *Environmental Science & Technology* **50**:1157–1165 [DOI 10.1021/acs.est.5b05107](https://doi.org/10.1021/acs.est.5b05107).
- Li S, Xie L, Zhang C, Zhang Y, Gu M, Zhang R. 2004.** Cloning and expression of a pivotal calcium metabolism regulator: calmodulin involved in shell formation from pearl oyster (*Pinctada fucata*). *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **138**:235–243 [DOI 10.1016/j.cbpc.2004.03.012](https://doi.org/10.1016/j.cbpc.2004.03.012).
- Livingstone DR, Chipman JK, Lowe DM, Minier C, Pipe RK. 2000.** Development of biomarkers to detect the effects of organic pollution on aquatic invertebrates: recent molecular, genotoxic, cellular and immunological studies on the common mussel (*Mytilus edulis* L.) and other mytilids. *International Journal of Environment and Pollution* **13**:56–91 [DOI 10.1504/IJEP.2000.002311](https://doi.org/10.1504/IJEP.2000.002311).
- Martin LB, Hopkins WA, Mydlarz LD, Rohr JR. 2010.** The effects of anthropogenic global changes on immune functions and disease resistance. *Annals of the New York Academy of Sciences* **1195**:129–148 [DOI 10.1111/j.1749-6632.2010.05454.x](https://doi.org/10.1111/j.1749-6632.2010.05454.x).
- Masthanamma P, Purushotham K, Ramamurthi R. 1984.** Metabolism of *Lamellidens marginalis* (Lamarck) in relation to starvation stress. I. Mortality, body weight, and oxygen consumption. *Indian Journal of Comparative Animal Physiology* **2**:60–63.
- McLoughlin K, Turteltaub K, Bankaitis-Davis D, Gerren R, Siconolfi L, Storm K, Cheronis J, Trollinger D, Macejak D, Tryon V, Bevilacqua M. 2006.** Limited dynamic range of immune response gene expression observed in healthy blood donors using RT-PCR. *Molecular Medicine* **12**:185–195 [DOI 10.2119/2006-00018.McLoughlin](https://doi.org/10.2119/2006-00018.McLoughlin).

- Menge BA, Branch GM. 2001.** Rocky intertidal communities. In: Bertness MD, Gaines SD, Hay ME, eds. *Marine community ecology*. Sunderland: Sinaur Associates, Inc., 3–37.
- Mensah-Osman E, Thomas D, Tabb M, Larios J, Hughes D, Giordano T, Lizyness M, Rae JM, Blumberg B, Hollenberg PF, Baker LH. 2007.** Expression levels and activation of a PXR variant are directly related to drug resistance in osteosarcoma cell lines. *Cancer* **109**:957–965 DOI [10.1002/cncr.22479](https://doi.org/10.1002/cncr.22479).
- Mitta G, Vandenbulcke F, Hubert F, Salzet M, Roch P. 2000.** Involvement of mytilins in mussel antimicrobial defense. *Journal of Biological Chemistry* **275**:12954–1262 DOI [10.1074/jbc.275.17.12954](https://doi.org/10.1074/jbc.275.17.12954).
- Nagorski S, Eckert G, Hood E, Pyare S. 2007.** Assessment of coastal water resources and watershed conditions at Katmai National Park and Preserve, Alaska. Natural resource technical report NPS/NRWRD/NRTR—2007/372. National Park Service, Fort Collins, Colorado.
- Nagorski S, Hood E, Eckert G, Pyare S. 2008.** Assessment of coastal water resources and watershed conditions: Lake Clark National Park and Preserve. Natural Resource Technical Report NPS/NRPC/WRD/NRTR—2008/144. National Park Service, Fort Collins, Colorado.
- Navarrete S, Menge B, Daley B. 2000.** Species interactions in intertidal food webs: prey or predation regulation of intermediate predators. *Ecology* **81**:2264–2277 DOI [10.1890/0012-9658\(2000\)081\[2264:SIIFW\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[2264:SIIFW]2.0.CO;2).
- Parisi MG, Toubiana M, Mangano V, Parrinello N, Cammarata M, Roch P. 2012.** MIF from mussel: coding sequence, phylogeny, polymorphism, 3D model and regulation of expression. *Developmental & Comparative Immunology* **36**:688–696 DOI [10.1016/j.dci.2011.10.014](https://doi.org/10.1016/j.dci.2011.10.014).
- Paul AJ, Feder HM. 1975.** The food of the sea star *Pycnopodia helianthoides* (Brandt) in Prince William Sound, Alaska. *Ophelia* **14**:15–22 DOI [10.1080/00785236.1975.10421968](https://doi.org/10.1080/00785236.1975.10421968).
- Peterson CH, Rice SD, Short JW, Esler D, Bodkin JL, Ballachey BE, Irons DB. 2003.** Long-term ecosystem response to the Exxon Valdez oil spill. *Science* **302**:2082–2086 DOI [10.1126/science.1084282](https://doi.org/10.1126/science.1084282).
- Philipp EER, Kraemer L, Melzner F, Poustka AJ, Thieme S, Findeisen U, Schreiber S, Rosenstiel P. 2012.** Massively parallel RNA sequencing identifies a complex immune gene repertoire in the lophotrochozoan *Mytilus edulis*. *PLOS ONE* **7**:e33091 DOI [10.1371/journal.pone.0033091](https://doi.org/10.1371/journal.pone.0033091).
- Place SP, O'Donnell MJ, Hofmann GE. 2008.** Gene expression in the intertidal mussel *Mytilus californianus*: physiological response to environmental factors on a biogeographic scale. *Marine Ecology Progress Series* **356**:1–14 DOI [10.3354/meps07354](https://doi.org/10.3354/meps07354).
- R Development Core Team. 2012.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <http://www.R-project.org/> (accessed on 1 May 2017).

- Reimer O, Tedengren M. 1996.** Phenotypical improvement of morphological defenses in the mussel *Mytilus edulis* induced by exposure to the predator *Asterias rubens*. *OIKOS* 75:383–390 DOI [10.2307/3545878](https://doi.org/10.2307/3545878).
- Rewitz KF, Styrihave B, Lobner-Olesen A, Andersen O. 2006.** Marine invertebrate cytochrome P450: emerging insights from vertebrate and insect analogies. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 143:363–381 DOI [10.1016/j.cbpc.2006.04.001](https://doi.org/10.1016/j.cbpc.2006.04.001).
- Romero A, Estévez-Calvar N, Dios S, Figueras A, Novoa B. 2011.** New insights into the apoptotic process in mollusks: characterization of caspase genes in *Mytilus galloprovincialis*. *PLOS ONE* 6:e17003 DOI [10.1371/journal.pone.0017003](https://doi.org/10.1371/journal.pone.0017003).
- Sforzini S, Oliveri C, Orru A, Chessa G, Pacchioni B, Millino C, Jha AN, Viarengo A, Banni M. 2018.** Application of a new targeted low-density microarray and conventional biomarkers to evaluate the health status of marine mussels: a field study in Sardinian coast, Italy. *Science of the Total Environment* 628–629:319–328 DOI [10.1016/j.scitotenv.2018.01.293](https://doi.org/10.1016/j.scitotenv.2018.01.293).
- Shaw JP, Dondero F, Moore MN, Negri A, Dagnino A, Readman JW, Lowe DR, Frickers PE, Beesley A, Thain JE, Viarengo A. 2011.** Integration of biochemical, histochemical and toxicogenomic indices for the assessment of health status of mussels from the Tamar Estuary, UK. *Marine Environmental Research* 72:13–24 DOI [10.1016/j.marenvres.2011.05.003](https://doi.org/10.1016/j.marenvres.2011.05.003).
- Smith T, Partridge S. 2004.** Dynamics of intertidal foraging by coastal brown bears in southwestern Alaska. *Journal of Wildlife Management* 68:233–240 DOI [10.2193/0022-541X\(2004\)068\[0233:DOIFBC\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2004)068[0233:DOIFBC]2.0.CO;2).
- Sokolova IM, Lannig G. 2008.** Interactive effects of metal pollution and temperature on metabolism in aquatic ectotherms: implications of global climate change. *Climate Research* 37:181–201 DOI [10.3354/cr00764](https://doi.org/10.3354/cr00764).
- Sureda A, Box A, Tejada S, Blanco A, Caixach J, Deudero S. 2011.** Biochemical responses of *Mytilus galloprovincialis* as biomarkers of acute environmental pollution caused by the Don Pedro oil spill (Eivissa Island, Spain). *Aquatic Toxicology* 101:540–549 DOI [10.1016/j.aquatox.2010.12.011](https://doi.org/10.1016/j.aquatox.2010.12.011).
- Taylor P, Maslowski W, Perwitz J, Wuebbles D. 2017.** *Arctic changes and their effects on Alaska and the rest of the United States*. Publications, Agencies and Staff of the US Department of Commerce 582. Available at <https://digitalcommons.unl.edu/usdeptcommercepub/582/>.
- Tsan MF, Gao B. 2004.** Cytokine function of heat shock proteins. *American Journal of Physiology-Cell Physiology* 286:C739–C744 DOI [10.1152/ajpcell.00364.2003](https://doi.org/10.1152/ajpcell.00364.2003).
- Vera-Massieu C, Brock PM, Godinez-Reyes CR, Acevedo-Whitehouse K. 2015.** Activation of an inflammatory response is context-dependent during early development of the California sea lion. *Royal Society Open Science* 2(4):150108 DOI [10.1098/rsos.150108](https://doi.org/10.1098/rsos.150108).
- Versteegh L, Hansson T. 2012.** Assessment of shell thickness in *Mytilus edulis* as a biomarker for establishment of its health and thiamine status. Available at <http://www.raysforexcellence.se/> (accessed on 10 May 2012).

- Walker LR, Sikes DS, DeGange AR, Jewett SC, Michaelson G, Talbot SL, Talbot SS, Wang B, Williams JC. 2013.** Biological legacies: direct early ecosystem recovery and food web reorganization after a volcanic eruption in Alaska. *Ecoscience* **20**:240–251 DOI [10.2980/20-3-3603](https://doi.org/10.2980/20-3-3603).
- Weeks DP. 1999.** Katmai National Park and Preserve Alaska water resources scoping report. Technical report NPS/NRWRD/NRTR-99/226. United States Department of the Interior, Washington D.C.
- Weeks DP. 2003.** Lake Clark National Park and Preserve Alaska water resources scoping report. Technical report NPS/NRWRD/NRTR-2003. United States Department of the Interior, Washington DC.
- Weitzman BP, Bodkin JL, Kloecker KA, Coletti HA. 2017.** SOP for monitoring intertidal bivalves on mixed-sediment beaches—version 2.0: Southwest Alaska Inventory and Monitoring Network. Natural resource report NPS/SWAN/NRR—2017/1443. National Park Service, Fort Collins, Colorado.
- Wells P, Butler J, Hughes J. 1995.** *Exxon Valdez oil spill: fate and effects in Alaskan waters*. Philadelphia: ASTM International.
- Woo S, Denis V, Won H, Shin K, Lee G, Lee TK, Yum S. 2013.** Expressions of oxidative stress-related genes and antioxidant enzyme activities in *Mytilus galloprovincialis* (Bivalvia, Mollusca) exposed to hypoxia. *Zoological Studies* **52**:15 DOI [10.1186/1810-522X-52-15](https://doi.org/10.1186/1810-522X-52-15).
- Woo S, Jeon HY, Kim SR, Yum S. 2011.** Differentially displayed genes with oxygen depletion stress and transcriptional responses in the marine mussel, *Mytilus galloprovincialis*. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **6**:348–356 DOI [10.1016/j.cbd.2011.07.003](https://doi.org/10.1016/j.cbd.2011.07.003).
- Wu R. 2002.** Hypoxia: from molecular responses to ecosystem responses. *Marine Pollution Bulletin* **45**:35–45 DOI [10.1016/S0025-326X\(02\)00061-9](https://doi.org/10.1016/S0025-326X(02)00061-9).
- Xavier B, Branch G, Wieters E. 2007.** Abundance, growth and recruitment of *Mytilus galloprovincialis* on the west coast of South Africa in relation to upwelling. *Marine Ecology Progress Series* **346**:189–201 DOI [10.3354/meps07007](https://doi.org/10.3354/meps07007).
- Xie F, Xiao P, Chen D, Xu L, Zhang B. 2012.** miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Molecular Biology* **80**:75–84 DOI [10.1007/s11103-012-9885-2](https://doi.org/10.1007/s11103-012-9885-2).
- Zanette J, Jenny MJ, Goldstone JV, Parente T, Woodin BR, Bainy AC, Stegeman JJ. 2013.** Identification and expression of multiple CYP1-like and CYP3-like genes in the bivalve mollusk *Mytilus edulis*. *Aquatic Toxicology* **128**:101–112 DOI [10.1016/j.aquatox.2012.11.017](https://doi.org/10.1016/j.aquatox.2012.11.017).
- Zeng LG, Wang JH, Li YJ, Sheng JQ, Gu Q, Hong YJ. 2012.** Molecular characteristics and expression of calmodulin cDNA from the freshwater pearl mussel, *Hyriopsis schlegelii*. *Genetics and Molecular Research* **11**:42–52 DOI [10.4238/2012.January.9.5](https://doi.org/10.4238/2012.January.9.5).
- Zimmermann M, Ruggerone GT, Freymueller JT, Kinsman N, Ward DH, Hogrefe KR. 2018.** Volcanic ash deposition, eelgrass beds, and inshore habitat loss from the 1920s to the 1990s at Chignik, Alaska. *Estuarine, Coastal and Shelf Science* **202**:69–86 DOI [10.1016/j.ecss.2017.12.001](https://doi.org/10.1016/j.ecss.2017.12.001).