


*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1637

Introduction: There is a growing literature suggesting disgust plays a major role in religiosity. Asceticism is a personality trait characterized by abstinence from sensual pleasures, often for the purpose of pursuing spiritual goals. Since few studies suggested that higher disgust levels may lead to greater fear of sin, contamination disgust may serve as effective mechanisms for inflated scrupulosity.

Objectives: We aimed to investigate the role of contamination disgust on a specific religious belief in two clinical groups: eating disorders and anxiety disorders.

Methods: We enrolled 84 patients admitted in the Psychiatric Unit of Careggi with diagnosis of Anxiety Disorders (AD) and Eating Disorders (ED). We administered them: Disgust Scale Revised (DS-R) and Eating Disorders Inventory 2 (EDI-2)

Results: A t-test for independent samples between AD patients and ED patients showed no difference in DS-R contamination disgust subscale ($t=1.437$, $p=0.153$), while significantly higher EDI-2 asceticism scores were detected in ED patients ($t=2.452$, $p=0.010$). An ANCOVA model having EDI2 Asceticism subscale as dependent variable, and DS contamination disgust subscale, diagnosis, and interaction between contamination disgust and diagnosis was estimated. The overall model was significant ($F(1,136)=4.854$, $p=0.003$) and accounted for 9.7% of variance of EDI2 Asceticism subscale ($R^2=0.097$). The effect of contamination disgust was positive and significant ($\beta=0.302$, $t=2.781$, $p=0.006$), and accounted for 5.4% of variance of EDI2 Asceticism (partial $\eta^2=0.054$). No effect was detected for diagnosis ($F(1,136)=0.012$, $p=0.912$) or interaction between contamination disgust and diagnosis ($F(1,136)=1.346$, $p=0.248$).



Conclusions: Contamination disgust may have a trans-diagnostic effect on asceticism and may be a possible driver for specific religious behaviors.

Disclosure: No significant relationships.

Keywords: Asceticism; anxiety disorders; eating disorders; disgust

EPV0021

Attention deficit hyperactivity disorder in adulthood and the heritability of this condition

J. Borges¹, R. Almeida Leite¹, M. Almeida^{1*}, S. Morais² and N. Madeira²

¹Departamento De Psiquiatria E Saúde Mental, Centro Hospitalar do Baixo Vouga, Aveiro, Portugal and ²Centro De Responsabilidade Integrado De Psiquiatria, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal

*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1638

Introduction: Attention Deficit Hyperactivity Disorder (ADHD) is a common neurodevelopmental disorder characterized by inattention and/or hyperactivity-impulsivity resulting from the interaction of genetic and environmental risk factors. Family studies shows that persistent ADHD is very familial.

Objectives: We aim to review the literature on this condition and its heritability and describe the implications that a possible misdiagnosis can have during life.

Methods: Bibliography review was performed using the databases PubMed and Cochrane, using the following keywords: “ADHD”; “Adults”; “Heritability”; “Family” and “Rater effect”.

Results: Childhood ADHD persists into adolescence and adulthood substantially, identified in some studies, as going up to 78%. The prevalence of ADHD in children and adults is between 2.5% and 5% worldwide. Family studies have shown that children of adults with ADHD are at higher risk of having ADHD. Some large-scale twin studies of adult ADHD, used self-report assessments of ADHD symptoms and estimated the heritability of this condition to be between 30 to 40%, which differs from other studies that analyse parents and teachers responses and estimates heritability to be between 60 and 90%.

Conclusions: Since there is a direct influence of the evaluators in estimating the extent of ADHD heritability, future studies need to clarify and describe in detail all the related characteristics of the raters. Although ADHD is widely studied, there is still a lot to learn about its etiology. The diagnosis of ADHD is clinical and complex and must be considered both in childhood and adolescence and in adulthood, with special emphasis on the family antecedents.

Disclosure: No significant relationships.

Keywords: heritability; attention deficit hyperactivity disorder; Adult

EPV0022

Self-compassion mediates the relationship between self-esteem and social anxiety symptoms in socially anxious individuals

P. Holas^{1,2*}, M. Kowalczyk³, I. Krejtz³, K. Wisiecka³ and T. Jankowski⁴

¹Faculty Of Psychology, University of Warsaw, Warsaw, Poland;

²Department Of Psychology, University of Warsaw, Warsaw, Poland;

³Psychology, SWPS University of Social Sciences and Humanities,

Warsaw, Poland and ⁴Psychology, John Paul II Catholic University of Lublin, Lublin, Poland

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1639

Introduction: Fear of evaluation and a negative view of the self are the key characteristics of social anxiety, which is one of the most prevalent anxiety problem. Self-esteem refers to views of oneself, including individual’s personal feelings towards self, whereas self-compassion refers to caring attitude toward oneself. Both constructs are two distinct positive views of the self, and were found to be related to each other, well-being and good mental health. To date, however, little is known, how they interplay in people with predominantly negative view of themselves, that is in socially anxious individuals.