

Original Research

# Effect of a Novel Training Program in Patients With Chronic Shoulder Pain Based on Implicit Motor Learning: Pilot and Feasibility Study.

Annelies Maenhout<sup>a</sup>, Wieger Heijnen, Peter Glashouwer, Lore Quatacker, Luna Praet, Dorien Borms

Keywords: motor control, chronic shoulder pain, rehabilitation, exercise, telerehabilitation

<https://doi.org/10.26603/001c.90284>

---

## International Journal of Sports Physical Therapy

Vol. 19, Issue 1, 2024

---

### Background

Implicit motor learning has been shown to be effective for learning sports-related motor skills. It facilitates automaticity of movements and thereby improves performance in multitasking and high-pressure environments. Motor learning to develop motor skills and neuroplastic capacities is not sufficiently incorporated in musculoskeletal rehabilitation. Especially in patients with chronic pain conditions like shoulder pain this approach might benefit over traditional exercise programs.

### Purpose/hypothesis

The aim of this study was to investigate the feasibility and clinical outcome of a new implicit motor learning exercise program in a group of patients with chronic shoulder pain.

### Study design

Pilot and feasibility cohort study

### Methods

Twenty-six patients with chronic shoulder pain performed a 6-week home exercise program with weekly remote follow up by a physiotherapist. The program comprised five exercises designed to challenge overall body balance, simultaneously engaging the upper limbs in a range of reaching tasks. The tasks included reaching above the head, at and below waist level, in various directions. No instructions on correct performance were provided to foster external focus. Feasibility was assessed by (1) recruitment rate, (2) follow up rate, (3) subjective experience, (4) self-reported adverse events and (5) self-reported adherence of subjects. Clinical effects of the program were assessed with (1) the Shoulder Pain and Disability Index (SPADI), (2) the Auto-Constant score, (3) the numeric rating scale (NRS) at rest and at night, (4) the patient specific functional scale (PSFS), (5) the avoidance endurance questionnaire (AEQ), (6) patient acceptable symptom state (PASS) and (7) a global rating of change (GROC).

### Results

The study protocol was feasible in terms of follow up rate (16w for 28 patients), exercise adherence (77.1%± 29.41), and adverse events (no serious, 5 light adverse events). Statistically significant improvements were observed for SPADI ( $p < 0.001$ ), NRS at rest

---

<sup>a</sup> Corresponding author:  
Annelies Maenhout, PhD, PT  
Department of Rehabilitation Sciences  
Ghent University  
De Pintelaan 185, 9000 Ghent, Belgium  
Phone number: 0032 3320268  
Email: [annelies.maenhout@ugent.be](mailto:annelies.maenhout@ugent.be)

( $p=0.033$ ), at night ( $p=0.29$ ), PSFS ( $p<0.001$ ) and PASS ( $p<0.001$ ) after only six weeks training.

## Conclusion

This study reveals promising results of another way of looking at exercise for patients with chronic shoulder pain. Both feasibility and clinical effects of the program on pain and function was acceptable. Future studies should incorporate a control group, provide longer follow up and include objective measurements.

## Level of evidence

2b

## INTRODUCTION

Shoulder disorders are the third most common musculoskeletal disorder after low back and neck pain.<sup>1,2</sup> When shoulder pain becomes chronic, it may seriously impair work and leisure participation, disturb sleep, and decrease health-related quality of life.<sup>3,4</sup> A conservative approach including exercise is the mainstay in the treatment of chronic shoulder complaints.<sup>5,6</sup> Although exercise therapy has proven its effectiveness in chronic musculoskeletal pain there is still a lot unknown about which approach is best for conditions of the shoulder.<sup>7</sup> Exercise programs mainly focus on scapular posture, scapular/rotator cuff strengthening, and anterior/posterior shoulder stretching.

Besides improving posture and increasing strength and range of motion, it might be important to prescribe exercises to optimize movement behavior in patients with chronic shoulder pain. Hodges and Tucker<sup>8</sup> reported changes at multiple levels of the motor system in patients with chronic pain resulting in altered movement behavior with the objective to “protect” the tissues from further pain or injury, or threatened pain or injury. These long-lasting neuromuscular adaptations in response to pain may offer short-term benefit, but have potential long-term consequences due to factors such as increased load, decreased movement, and decreased variability and are advised to address in patients with chronic musculoskeletal pain.<sup>8-10</sup> In the majority of clinical practice, patients are provided with explicit guidance for proper movement when retraining motor control. This guidance commonly pertains to coordinating the movements of the patient’s body, including instructions on scapular posture and muscle recruitment. A small alteration in the wording of instructions can result in a substantial influence on both performance and motor learning outcomes. When patients are directed to focus their attention on how their movements affect the environment—an external focus of attention—it leads to more effective and efficient movements.<sup>11,12</sup> A movement pattern is considered more efficient or economical if the same movement outcome is achieved with less energy expended. In contrast, directing attention internally towards one’s own movements results in a more conscious form of control that limits the motor system and disrupts automatic control processes.<sup>13</sup> The aim of implicit motor learning methods is to minimize the amount of explicit knowledge about movement execution and/or muscle recruitment during learning

(internal focus), reduce the reliance on the working memory, and promote a more automatic control process.<sup>14</sup>

Implicit learning is an approach in which self-organization is key. The principle of self-organization emphasizes that coordinated movements emerge naturally from the interactions among the components of the motor system. Instead of being rigidly controlled from outside, the body adapts and organizes itself to achieve task goals while exploring different movement patterns. This concept highlights the importance of allowing learners to explore and discover their own movement solutions, promoting adaptability and skill development. In sports science, there is increasing evidence that implicit motor learning is superior to explicit learning when it comes to learning new skills and improving performance.<sup>15,16</sup> There are a number of disciplines in which implicit learning has been successfully applied, including pediatric physiotherapy, gait training for stroke patients, and ACL rehabilitation, but implicit learning is rarely used in the context of shoulder rehabilitation.<sup>14,17,18</sup>

Considering the expected duration of rehabilitation within chronic shoulder pain patients, telerehabilitation seems a suitable medium for this population.<sup>19</sup> During the COVID-19 pandemic physiotherapists widely used telerehabilitation to coach patients in their home exercise programs. In systematic reviews, both Seron et al. and Gava et al. concluded that telerehabilitation could be comparable to in-person rehabilitation to reduce pain and improve physical function in musculoskeletal conditions generally and shoulder pain specifically, but more high quality research is needed.<sup>20,21</sup> The main advantage of telerehabilitation is that both the healthcare professional and the patient can carry out the treatment in their own environment without travel time or necessary transport.

The promising results of implicit motor learning in other conditions and the possibility of implicit learning to change motor control deficits evidenced in chronic pain conditions gave rise to this study. The aim of this study was to investigate the feasibility and clinical outcome of a new implicit motor learning exercise program in a group of patients with chronic shoulder pain.

## MATERIALS AND METHODS

### STUDY DESIGN

The feasibility study was a single group pre-post intervention design. The Institutional Review Board of Ghent

University Hospital (B.U.N.: B6702020000611) approved the study.

## PARTICIPANTS

Adults with chronic shoulder pain were recruited on social media and via flyers in public buildings and physiotherapy practices. Potential candidates were first screened through an online questionnaire to check whether they met inclusion criteria (questions on duration of shoulder pain, the experience of an unstable feeling in the shoulder, dislocations/subluxations, mobility restrictions of the shoulder, results of medical imaging, medical diagnosis). To be included patients had to have had shoulder pain for at least six months for which they consulted a medical doctor (MD). Participants were excluded if they were currently in therapy or had undergone shoulder surgery or sustained a shoulder fracture. Moreover, patients that were expected not to benefit the intervention were excluded namely patients with an unstable shoulder (shoulder dislocation/instability, labral lesion), capsular stiffness (frozen shoulder, severe osteoarthritis) or with inflammatory nociceptive pain mechanisms (bursitis, calcific tendinopathy, rheumatic disorders), as diagnosed by the MD, were excluded. All participants signed an informed consent prior to participation.

## OUTCOMES

Feasibility outcomes consisted of recruitment rate, follow up rate (% of participants who successfully completed the program and were reevaluated after six weeks), subjective reports regarding experience with the content of the program (custom-crafted statements on the content of this program scored with Likert scale eg. "I understood what was expected", "I thought the exercises were easy to do"), self-reported adverse events, and self-reported adherence. Adherence was scored weekly as 0 (did not meet minimal exercise adherence of 15', 3x/week) or 1 (meets minimal exercise adherence of 15', 3x/week). A score of six out of six (100%) was defined as perfect adherence. Based on this outcome the feasibility of a future RCT and the need for protocol modification were assessed.


At baseline and after the six-week home exercise program patients were requested to complete an online questionnaire package that consisted of demographic variables and several patient reported outcome measures (PROMs). Pain and function were evaluated with the Shoulder Pain and Disability Index (SPADI)<sup>22</sup> and Auto-Constant score.<sup>23</sup> The SPADI was chosen due to excellent reported reliability (ICC $\geq$ 0.90) in most studies, with reported minimal detectable change (MDC) of 18 points and minimal clinically important difference (MCID) between 8 and 13 points.<sup>24, 25</sup> The Auto-Constant score (patient self-report based) and the Constant score (clinician-based) have been shown to correlate well but no information on test-retest reliability of the Auto-Constant score is available.<sup>23,26</sup> This alternative to the Constant score allows remote assessment which was necessary at the time of the study due to COVID precautions. Self-perceived active mobility and strength was

measured with the Auto-Constant score. In this questionnaire patients have to select the pictures of a person lifting a weight or performing a specific movement that correspond with the level of lifting or moving they estimated themselves to be capable of.<sup>23</sup> To assess patterns of fear-avoidance and endurance-related responses to pain the Avoidance Endurance Questionnaire (AEQ) was used.<sup>27</sup> According to Hasenbring et al.,<sup>27</sup> inactive, avoidant behavior would drive disuse-related muscle weakness, deconditioning, and metabolic changes in the musculoskeletal structures and the central nervous system, potentially leading to peripheral and central sensitization and increased pain perception. In contrast, in overactive behavior, inappropriate muscle forces might expose mainly passive structures (vertebral joints, ligaments, connective tissue) to increased stress and repetitive strain, causing microdamage, laxity, and inflammation. The AEQ was developed to assess emotional, cognitive, and behavioral fear-avoidance and endurance responses to pain, and has been validated and repeatedly applied in different chronic pain populations.<sup>27-29</sup> This self report questionnaire consists of three parts (pain affective, cognitive and behavioural responses). More detailed information on the subscales of the AEQ can be found in Appendix 2.

Each item was scored on a 7 point-Likert scale (0: never, 1: almost never, 2: seldom, 3: sometimes, 4: often, 5: most of the time, and 6: always). Higher scores represent more thoughts that correspond with avoidance or endurance of pain. All subscales Chronbach's alpha were in range from 0.76 to 0.91, which showed acceptable internal consistency.<sup>27</sup> The participants were asked to fill out the questionnaire considering the pain they experienced in the past 14 days.

Overall satisfaction with the shoulder condition was evaluated before and after the 6-week program with the Patient Acceptable Symptom State (PASS).<sup>30</sup> After the program patients were asked to score the rate of change between -5 and 5 (Global rating of change (GROC)).<sup>31,32</sup> A score of -5 corresponds with shoulder complaints being a lot worse than before the program, 0 with no change, and 5 with full recovery of shoulder complaints. A systematic review and meta-analysis on measurement properties concluded that GROC shows excellent test-retest reliability (ICC = 0.84) but as recall bias might influence validity it is advised to combine with other outcomes to track changes in symptoms.<sup>33</sup>

During weekly follow up, patients were asked to rate their pain at rest and at night on a numeric rating scale (NRS) as well as their ability to perform three important functional activities, using the Patient Specific Function Scale (PSFS).<sup>34,35</sup> For NRS MCID was found to be 2.17 points in a population with shoulder pain.<sup>36</sup> The PSFS shows moderate to good test-retest reliability (ICC 0.71) and MCID was shown to be 1.2 points.<sup>35</sup> The advantage of the PSFS is that each patient can rate functional ability during daily life tasks that are relevant for them.



**Figure 1. Home exercise program**

#### INTERVENTION

For specific content reporting of the home based exercise program the Consensus on Exercise Reporting Template was followed.<sup>37</sup> A general instruction video (<https://youtu.be/s11NShRi35Y>) and videos with exercise instructions (Appendix 1) were accessible online.

The program consisted of five exercises (Figure 1 and Appendix 1 detailed instructions). The exercise program was based on the principles of implicit motor learning.<sup>38</sup>

#### SELF ORGANIZATION

To elicit unconscious automatic bottom-up organization of motor control of the upper extremities, overall balance was challenged during the exercises.<sup>38</sup> Each exercise had ten levels of difficulty to maintain total body balance (Appendix 1). To determine the level at the start, the 70-30% rule was used. Participants were instructed to select the level at which they could execute the exercise approximately seven out of ten times without losing balance.

To encourage participants to expand their range of motor control possibilities and engage their shoulder muscles beyond their comfort zone, they were consistently prompted to reach their maximum extent during all exercises. While performing the exercise, individuals should exert additional effort to reach the target and fully utilize their shoulder mobility. To achieve this, subjects were directed to position themselves in a manner where the target was placed just beyond their normal reach.

#### EXTERNAL FOCUS INSTRUCTIONS

During all exercises the focus was on the target (sliding objects over the floor, reaching towards the wall, touching toes) and no instructions were provided on how to perform the exercise. To correct performance in case of discomfort or low back pain for example analogies were used to elicit the desired movement behavior implicitly throughout the

program such as for example: “imagine your legs both in a cast as you perform the exercises, now imagine that your legs are out of the casts” or “imagine your legs stuck in concrete, now the concrete is broken and you are free again”.

#### SELF-CONTROLLED LEARNING AND ENHANCED EXPECTATIONS

In the majority of rehabilitation scenarios, clinicians typically dictate the specifics of the training session. This includes decisions about the sequencing of tasks, the duration of practice, and when or whether instructions or demonstrations are provided. Consequently, patients often play a relatively passive role, with clinicians maintaining general control over most practice aspects. However, self-controlled learning, such as allowing patients to have some influence, requesting feedback, or selecting exercises, emerges as a potent instrument in the realm of motor learning. From a motivational perspective, patients were encouraged by the therapists (two last year master physiotherapy students that were specifically trained and under the supervision of two experienced senior shoulder physiotherapists) to take maximal responsibility in their training program and received the autonomy to make choices.<sup>13, 39</sup> During the six weeks participants were asked to practice with a minimum of three times a week. The therapists suggested that there may be superior outcome if the patients practiced more than three times per week. Duration of one exercise session was instructed to be 15 to 20 minutes but participants were allowed to perform shorter sessions multiple times a day. The sequence and duration of each exercise was not set. Participants could choose when and how much a specific exercise was done as long as a total of 15-20 minutes was reached. To control the pain (during and immediately after exercise, day to day, and week to week), participants were instructed to follow the pain monitoring model (Appendix 1).<sup>40,41</sup> When pain increased or fatigue began, participants were instructed to switch to another exercise.

Follow-up with the participants was provided by the therapists during a weekly video or phone call according to individual preference. During these calls pain, PSFS, the level of difficulty achieved for each exercise (and whether this was challenging enough), adverse events, and the frequency and duration of training sessions were discussed.

## STATISTICAL ANALYSIS

Data were analyzed with SPSS statistical software (IBM SPSS Statistics, Version 27.0). Descriptive statistics were used to assess the feasibility outcomes. A Shapiro-Wilk test was combined with a histogram to check normal distribution. Paired sample *t*-tests and Wilcoxon signed-rank tests were used to compare SPADI, AEQ, Auto-Constant scores and NRS at rest, at night, and during patient specific activities (PSFS) pre to post. Differences in the PASS were analyzed with a McNemar test. Statistically significant differences were considered at  $p < 0.05$ . Post hoc power analysis was conducted based on the SPADI.

## RESULTS

A group of 28 participants (Mean age= 34.9 years  $\pm 13.59$ ; male/female= 10/17) was recruited. Baseline characteristics are summarized in Table 2. Medical diagnoses of their shoulder pain differed among the group most of which were related to the rotator cuff (shoulder impingement, subacromial impingement, rotator cuff related shoulder pain, subacromial pain syndrome, rotator cuff tendinopathy, rotator cuff muscle “sprain”) however, some patients did not have clear medical diagnoses (such as “functional shoulder pain”). Twenty-six patients completed the study and two were lost to follow-up (one spontaneous improvement of pain before start of the study and one decided to consult an orthopedic surgeon for second opinion).

## FEASIBILITY OUTCOME

- **Recruitment rate:** A total of 16 weeks was required to screen 96 patients of which 28 patients met the inclusion criteria and were eligible for participation in this study.
- **Follow up rate:** 92.9% of participants successfully completed follow up assessment after 6 weeks.
- **Subjective experience with the content of the program:** The outcome of the questions regarding the personal experience of the exercise program is visualized in [Figure 2](#). Majority of participants (66.6%) indicated their expectations regarding the exercise program were fulfilled. Therapist guidance, follow up and content of the exercise program received positive feedback from all participants. Moreover, 62.9% of participants felt the exercise program had positive effects on their shoulder pain.
- **Self-reported adverse events:** Five patients reported adverse events during performance of exercises: low back pain (n=3), groin pain (n=1) and knee pain (n=1). All could be resolved by providing information on

correct performance of the exercises or adapting the exercises. No patients reported adverse events after completion of the study.

- **Self-reported adherence:** The mean adherence to the exercise program was  $77.1 \pm 29.41\%$ . It can be observed in [Figure 3](#) that the adherence was higher during the first two weeks, decreased after that, and increased again towards the end of the program.

## EFFECT OF THE PROGRAM ON PROMS

An overview of the pre- and post-intervention scores for the PROMs and the corresponding p-values are presented in [Table 1](#). Post hoc power analysis based on the SPADI resulted in a power of 0.94.

SPADI pain scores were significantly lower after six weeks of training (median difference [95%CI]= -14.0 [-22.0; -7.0];  $p < 0.001$ ). SPADI function scores were significantly improved after the exercise program (median difference [95%CI]= -10.0 [-15.0; -5.6];  $p < 0.001$ ). Total SPADI scores improved significantly (mean difference [95%CI]= -13.6 [-19.8; -7.4];  $p < 0.001$ ).

In the Auto-Constant questionnaire, significant improvement was attained for pain, self-perceived strength, and ROM. The pain scores decreased significantly (median difference [95%CI]= -3.0 [-4.5; -1.5];  $p < 0.001$ ), strength scores increased significantly (median difference [95%CI]= 2.5 [0.0; 5.0];  $p = 0.008$ ) and ROM scores improved significantly (median difference [95%CI]= 2.0 [1.0; 3.5];  $p = 0.001$ ). Activity scores did not change significantly (mean difference [95%CI]= 0.5 [-0.2; 1.2];  $p = 0.168$ ) and the total Auto-Constant score was not significantly different after intervention (mean difference [95%CI]= 3.0 [-0.1; 6.1];  $p = 0.060$ ).

Before the program, 23.1% of the participants was satisfied with their shoulder condition, whereas after the study 69.3% were satisfied as measured with the PASS. This proportional change was significant ( $p < 0.001$ ).

The mean NRS pain score at rest improved significantly through the research (median difference [95%CI]= -0.5 [-1.5; 0.0];  $p = 0.033$ ) as did the pain scores at night (median difference [95%CI]= -0.8 [-1.5; 0.0];  $p = 0.029$ ). For the PSFS the participants reported a significantly lower score reflecting improved ability to perform functional activities after the program (mean difference [95%CI]= -2.4 [-3.4; -1.5];  $p < 0.001$ ).

Regarding the AEQ, the Pain Emotional Reactions score improved significantly after the intervention. The Anxiety Depression Scale (ADS) showed a significant mean difference [95%CI] of -0.6 [-0.8; -0.3] ( $p < 0.001$ ). Additionally, the Positive Mood Scale (PMS) significantly improved after training with a median difference [95%CI] of 0.5 [0.2; 1.0] ( $p = 0.007$ ). For the subscale Pain Cognitive Reactions, only the Help Hopelessness Scale (HHS) reached a significant difference (mean difference [95%CI]= -0.7 [-1.0; -0.3];  $p = 0.002$ ).

The self-perceived change in the condition of the shoulder after six weeks of exercising as measured with the GROC showed a mean change of 1.9 points ( $\pm 1.89$ ). The distribution of the scores are presented in [Figure 4](#).





Figure 2. Personal experience regarding the exercise program (expressed as the percentage of participants in each answer category)

DISCUSSION

The aim of this research was to determine whether a novel home exercise program conducted via telerehabilitation based on implicit motor learning was effective for patients with chronic shoulder pain and whether the study protocol was feasible. The results showed that the study protocol was feasible in terms of follow up rate, exercise adherence, and adverse events. Statistically significant improvements were observed for the SPADI, NRS at rest and at night, and PASS after only six weeks training.

Twenty-six participants completed the six-week exercise program at home with weekly remote follow-up. The results

showed that the study protocol was feasible with respect to follow-up rate and acceptable in terms of personal experience with the program, exercise adherence and adverse events. Majority of participants (66.6%) indicated their expectations regarding the exercise program to be fulfilled. Therapist guidance, follow up, and content of the exercise program received positive feedback from all participants. Moreover, 62.9% of participants felt the exercise program had positive effects on their shoulder pain. This is an important finding, since perception of participants about the exercise program is crucial for adherence to the treatment and thus successful rehabilitation.<sup>42,43</sup> Although patient and therapist never met in person due to COVID-19 mea-



**Figure 3. Mean adherence (percentage) over 6 weeks (100% equals exercising as prescribed at least 3 days a week, at least 15 minutes)**

tures, participants indicated that they understood the instructions on the phone or video-calls, when combined with the YouTube videos.

Frequency of follow-up also appeared important in view of adverse events. Providing instructions on how to adapt the exercises was key in preventing pain in other joints such as low back and knee to persist. Hence, the communication used in this study appeared suitable for telerehabilitation.

The clinical outcome results are very promising after only six weeks of training. There was a 13.6 point decrease in the primary outcome (SPADI score). This result is higher than the MCID of 8-13 points but lower than the MDC of the SPADI of 18 points.<sup>24</sup> The magnitude of this difference is comparable to other studies investigating the effect of more traditional exercise programs with strength and mobility exercises.<sup>44,45</sup> Littlewood et al. for example showed improvement of SPADI scores after three months with 12.4 points in the self-managed group and 16.7 points in the usual physiotherapy group in patients with subacromial pain syndrome.<sup>45</sup> Especially considering the study population (>6 months shoulder pain) a longer duration is recommended given that earlier research indicated longer lasting complaints may require a prolonged rehabilitation.<sup>46</sup> The improvement in pain was significant for pain at rest and at night. The amount of change was smaller than the MCID of 2.17 points but comparable to changes seen in other intervention studies in this population.<sup>47,48</sup> It is important to note that the small improvement might also be related to the rather low pain scores at baseline (1.8-1.9) which is most likely the result of exclusion of serious shoulder pathology such as frozen shoulder in the current study. A mean change in shoulder complaints of 1.9 points on

the GROCC represents that participants consider the change valuable regarding their health status.<sup>31</sup>

After the program it was shown that pain emotional reactions, as measured with the AEQ, significantly improved, meaning that patients had fewer depressive thoughts and anxiety and were in a better mood. This may relate to a happier lifestyle and thus a higher quality of life. In contrast, no significant changes were seen in participant's coping reactions. Pain cognitive reactions also did not change significantly except for the HHS. The HHS represents the lack of hope and the feeling of impossibility to become pain-free.<sup>49</sup> In other words, participants believed in the advantages of exercising and had a more optimistic point of view on the future after study completion. Combining the exercise program with behavioral change techniques could be an interesting direction for future research.

The current study attained a high percentage of compliance of 77%. Compared to other studies this adherence rate with home-based training was rather high. For example, Burns et al.<sup>45</sup> reported an adherence of 41% with a home exercise program for patients with rotator cuff pathology. Possible reasons for this strong adherence might be the weekly follow-up calls, the dynamic nature of the exercises and the freedom of planning practice. Freedom of planning their own practice was perceived as an advantage by 96% of participants. Rehabilitation outcome strongly depends on treatment adherence. Interestingly, in this study it was observed that adherence decreased to the lowest value at week four and then started improving again towards the end of the program. A potential cause for the motivational dip could be that the novelty of the exercises was gone, and the end of the training period still seemed a long way off. NRS scores at rest and at night increased between week four

**Table 1. Patient reported outcome measurements before and after the intervention.**

		Pre (mean ± SD)	Post (mean ± SD)	p-value
SPADI	Pain	36.0 ± 21.50	20.8 ± 19.21	<0.001*
	Function	26.6 ± 20.81	14.6 ± 16.81	<0.001*
	<b>Total</b>	<b>31.3 ± 20.25</b>	<b>17.7 ± 17.75</b>	<b>&lt;0.001*</b>
Auto-Constant	Pain	6.7 ± 3.07	3.8 ± 2.98	<0.001*
	Strength	14.4 ± 4.97	17.1 ± 5.32	0.008*
	ROM	16.5 ± 4.81	19.3 ± 3.46	0.001*
	Activity	12.8 ± 2.06	13.3 ± 1.76	0.168
	<b>Total</b>	<b>50.5 ± 8.52</b>	<b>53.4 ± 5.43</b>	<b>0.060</b>
PASS		0.2 ± 0.43	0.7 ± 0.47	<0.001*
AEQ: pain emotional reactions	ADS	1.5 ± 1.02	0.9 ± 0.82	<0.001*
	PMS	4.2 ± 1.52	4.7 ± 0.82	0.007*
AEQ: pain cognitive reactions	HHS	2.1 ± 1.31	1.4 ± 1.12	0.002*
	CTS	0.5 ± 0.89	0.4 ± 0.69	0.471
	TSS	2.7 ± 1.37	2.8 ± 1.44	0.737
AEQ: pain coping reactions (light pain)	APAS	1.3 ± 0.98	1.3 ± 0.87	0.951
	ASAS	1.5 ± 0.55	1.4 ± 0.42	0.169
	PPS	2.5 ± 0.75	2.6 ± 0.74	0.340
	HDS	2.9 ± 0.98	2.8 ± 0.75	0.700
	BES (PPS + HDS)	5.4 ± 1.55	5.4 ± 1.32	0.859
AEQ: pain coping reactions (severe pain)	APAS	1.7 ± 0.57	1.4 ± 0.81	0.089
	ASAS	2.1 ± 0.48	2.0 ± 0.46	0.146
	PPS	3.0 ± 0.90	2.9 ± 1.02	0.615
	HDS	3.0 ± 1.26	3.2 ± 1.25	0.475
	BES (PPS + HDS)	6.1 ± 1.93	6.1 ± 2.00	0.895
NRS	Pain @ rest	1.8 ± 1.68	1.2 ± 1.91	0.033*
	Pain @ night	1.9 ± 2.05	1.1 ± 1.94	0.029*
	PSFS	4.6 ± 1.73	2.3 ± 2.00	<0.001*

\* indicates a statistically significant change at  $p < 0.05$ ; SD = Standard Deviation; SPADI= Shoulder Pain and Disability Index; ROM= Range of Motion; PASS= Patient Acceptable Symptom State; AEQ= Avoidance-Endurance Questionnaire; ADS = Anxiety Depression Scale; PMS= Positive Mood Scale; HHS= Help Hopelessness Scale; CTS= Catastrophizing Thoughts Scale; TSS= Thought Suppression Scale; APAS= Avoidance of Physical Activities Scale; ASA= Avoidance of Social Activities Scale; PPS= Pain Persistence Scale; HDS= Humor Distraction Scale; BES= Behavioral Endurance Scale; NRS= Numeric Rating Scale; PSFS= Patient Specific Function Scale.

and five probably causing the participants to see the benefit of the exercise program which in turn increased the adherence. After six weeks of training 70.4% of participants indicated they planned to continue to perform the exercises after study completion. The setup of this study with a home-based program and maximal responsibility for patients might be advantageous for patients' self-efficacy, autonomy, and active coping behavior.


The training program only consisted of exercises based on implicit motor learning with external focus, no manual therapy or strength training was included. The underlying mechanism that could explain the clinical improvements might be related to motor learning resulting in more efficient and less rigid movement patterns. Studies investigating EMG muscle recruitment during a variety of exercises comparing external focus of attention to internal focus of attention showed lower muscle recruitment in the first.<sup>11, 12,50-55</sup> In these studies, where the executed movement remains constant and only the focus of attention is shifted, a

decrease in muscle recruitment could indicate enhanced efficiency in performing the exercise. This could be beneficial during functional movements in daily activities. During follow up, some participants spontaneously reported that they noticed starting to use their affected shoulder more during daily activities and a significant improvement was found in their ability to perform individually reported activities as measured with the PSFS. A future study might include objective measurements of arm use in daily life. The placebo effect due to being included in a study and receiving attention from the therapist cannot be out ruled as there was no control group.

#### LIMITATIONS

A first limitation of this study is the lack of a control group. In addition to ruling out the influence of natural recovery, incorporating a control group would provide clarity on how the outcomes of this novel exercise program compare to a





**Figure 4. Global rating of change**

A score of -5: shoulder complaints a lot worse than before the program, 0: no change, 5: full recovery of shoulder complaints

conventional function-based program. The latter typically includes a range of isolated shoulder strengthening and mobility exercises.

The second limitation is that no objective data on strength and mobility could be collected due to COVID measures at the time of the study. To evaluate exercise adherence objective data would also be desirable. Since the follow-up calls were not anonymous, socially desirable responses might unconsciously have been encouraged.

Finally, study duration can be considered a limitation. A future trial should have a longer training program of at least 12 weeks and include long term follow-up after six months up to one year.

**CONCLUSION**

The results of the current study provide promising results of a novel way designing shoulder exercises for patients with chronic shoulder pain. The home-based, six-week exercise program based on implicit motor learning produced

significant improvements in pain and function in patients with chronic shoulder pain. The study protocol was feasible in terms of follow up rate, exercise adherence and adverse events. Participants especially appreciated the freedom to plan their exercise sessions and instructions and follow up provided by the therapists. Recruitment rate was rather low. Future studies should incorporate a control group, provide longer follow up, and include objective measurements to analyze changes in strength, range of motion, and use of the affected upper limb during daily life.

.....  
**CONFLICT OF INTEREST**

The authors declare that they have no conflict of interest.

Submitted: February 02, 2023 CST, Accepted: October 30, 2023 CST

© The Author(s)



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

## REFERENCES

1. Djade CD, Porgo TV, Zomahoun HTV, Perrault-Sullivan G, Dionne CE. Incidence of shoulder pain in 40 years old and over and associated factors: A systematic review. *Eur J Pain*. 2020;24(1):39-50. doi:10.1002/ejp.1482
2. Kuijpers T, van Tulder MW, van der Heijden GJ, Bouter LM, van der Windt DA. Costs of shoulder pain in primary care consultants: a prospective cohort study in The Netherlands. *BMC Musculoskelet Disord*. 2006;7(1):83. doi:10.1186/1471-2474-7-83
3. Serrano-Aguilar P, Kovacs FM, Cabrera-Hernández JM, Ramos-Goñi JM, García-Pérez L. Avoidable costs of physical treatments for chronic back, neck and shoulder pain within the Spanish National Health Service: a cross-sectional study. *BMC Musculoskelet Disord*. 2011;12(1):287. doi:10.1186/1471-2474-12-287
4. Ackerman IN, Fotis K, Pearson L, et al. Impaired health-related quality of life, psychological distress, and productivity loss in younger people with persistent shoulder pain: a cross-sectional analysis. *Disabil Rehabil*. 2022;44(15):3785-3794. doi:10.1080/09638288.2021.1887376
5. Kroll HR. Exercise therapy for chronic pain. *Phys Med Rehabil Clin N Am*. 2015;26(2):263-281. doi:10.1016/j.pmr.2014.12.007
6. Naugle KM, Fillingim RB, Riley JL III. A meta-analytic review of the hypoalgesic effects of exercise. *J Pain*. 2012;13(12):1139-1150. doi:10.1016/j.jpain.2012.09.006
7. Dueñas L, Aguilar-Rodríguez M, Voogt L, et al. Specific versus non-specific exercises for chronic neck or shoulder pain: A systematic review. *J Clin Med*. 2021;10(24):5946. doi:10.3390/jcm10245946
8. Hodges PW, Tucker K. Moving differently in pain: a new theory to explain the adaptation to pain. *Pain*. 2011;152(3 Suppl):S90-S98. doi:10.1016/j.pain.2010.10.020
9. Hodges PW. Pain and motor control: From the laboratory to rehabilitation. *J Electromyogr Kinesiol*. 2011;21(2):220-228. doi:10.1016/j.jelekin.2011.01.002
10. Schabrun SM, Jones E, Kloster J, Hodges PW. Temporal association between changes in primary sensory cortex and corticomotor output during muscle pain. *Neuroscience*. 2013;235:159-164. doi:10.1016/j.neuroscience.2012.12.072
11. Wulf G, Dufek JS, Lozano L, Pettigrew C. Increased jump height and reduced EMG activity with an external focus. *Hum Mov Sci*. 2010;29(3):440-448. doi:10.1016/j.humov.2009.11.008
12. Chua LK, Jimenez-Diaz J, Lewthwaite R, Kim T, Wulf G. Superiority of external attentional focus for motor performance and learning: Systematic reviews and meta-analyses. *Psychol Bull*. 2021;147(6):618-645. doi:10.1037/bul0000335
13. Wulf G, Lewthwaite R. Optimizing performance through intrinsic motivation and attention for learning: The OPTIMAL theory of motor learning. *Psychon Bull Rev*. 2016;23(5):1382-1414. doi:10.3758/s13423-015-0999-9
14. Gokeler A, Neuhaus D, Benjaminse A, Grooms DR, Baumeister J. Principles of Motor Learning to Support Neuroplasticity After ACL Injury: Implications for Optimizing Performance and Reducing Risk of Second ACL Injury. *Sports Med*. 2019;49(6):853-865. doi:10.1007/s40279-019-01058-0
15. Kal E, Prosée R, Winters M, van der Kamp J. Does implicit motor learning lead to greater automatization of motor skills compared to explicit motor learning? A systematic review. *PLoS One*. 2018;13(9):e0203591. doi:10.1371/journal.pone.0203591
16. Navarro M, van der Kamp J, Schor P, Savelsbergh GJP. Implicit learning increases shot accuracy of football players when making strategic decisions during penalty kicking. *Hum Mov Sci*. 2018;61:72-80. doi:10.1016/j.humov.2018.07.004
17. Kal E, Winters M, van der Kamp J, et al. Is Implicit Motor Learning Preserved after Stroke? A Systematic Review with Meta-Analysis. *PLoS One*. 2016;11(12):e0166376. doi:10.1371/journal.pone.0166376
18. Jarus T, Ghanouni P, Abel RL, et al. Effect of internal versus external focus of attention on implicit motor learning in children with developmental coordination disorder. *Res Dev Disabil*. 2015;37:119-126. doi:10.1016/j.ridd.2014.11.009
19. Janela D, Costa F, Molinos M, et al. Asynchronous and Tailored Digital Rehabilitation of Chronic Shoulder Pain: A Prospective Longitudinal Cohort Study. *J Pain Res*. 2022;15:53-66. doi:10.2147/jpr.s343308

20. Seron P, Oliveros MJ, Gutierrez-Arias R, et al. Effectiveness of Telerehabilitation in Physical Therapy: A Rapid Overview. *Phys Ther.* 2021;101(6). doi:10.1093/ptj/pzab053
21. Gava V, Ribeiro LP, Barreto RPG, Camargo PR. Effectiveness of physical therapy given by telerehabilitation on pain and disability of individuals with shoulder pain: A systematic review. *Clin Rehabil.* 2022;36(6):715-725. doi:10.1177/02692155221083496
22. Kc S, Sharma S, Ginn KA, Reed D. Measurement properties of translated versions of the Shoulder Pain and Disability Index: A systematic review. *Clin Rehabil.* 2021;35(3):410-422. doi:10.1177/0269215520963199
23. Chelli M, Levy Y, Lavoué V, Clowez G, Gonzalez JF, Boileau P. The "Auto-Constant": Can we estimate the Constant-Murley score with a self-administered questionnaire? A pilot study. *Orthop Traumatol Surg Res.* 2019;105(2):251-256. doi:10.1016/j.otsr.2018.11.023
24. Roy JS, MacDermid JC, Woodhouse LJ. Measuring shoulder function: a systematic review of four questionnaires. *Arthritis Rheum.* 2009;61(5):623-632. doi:10.1002/art.24396
25. Paul A, Lewis M, Shadforth MF, Croft PR, Van Der Windt DA, Hay EM. A comparison of four shoulder-specific questionnaires in primary care. *Ann Rheum Dis.* 2004;63(10):1293-1299. doi:10.1136/ard.2003.012088
26. Levy O, Haddo O, Massoud S, Mullett H, Atoun E. A patient-derived Constant-Murley score is comparable to a clinician-derived score. *Clin Orthop Relat Res.* 2014;472(1):294-303. doi:10.1007/s11999-013-3249-3
27. Hasenbring MI, Hallner D, Rusu AC. Fear-avoidance- and endurance-related responses to pain: development and validation of the Avoidance-Endurance Questionnaire (AEQ). *Eur J Pain.* 2009;13(6):620-628. doi:10.1016/j.ejpain.2008.11.001
28. Gajjar H, Titze C, Levenig C, et al. Psychological pain responses in athletes and non-athletes with low back pain: Avoidance and endurance matter. *Eur J Pain.* 2019;23(9):1649-1662. doi:10.1002/ejp.1442
29. Karimi Ghasem Abad S, Akhbari B, Salavati M, Saeedi A, Seydi M, Shakoorianfard MA. Translation, reliability, and validity of the avoidance endurance questionnaire in Iranian subjects with chronic non-specific neck pain. *J Family Med Prim Care.* 2020;9(7):3565-3573. doi:10.4103/jfmpc.jfmpc\_194\_20
30. Kvien TK, Heiberg T, Hagen KB. Minimal clinically important improvement/difference (MCII/MCID) and patient acceptable symptom state (PASS): what do these concepts mean? *Ann Rheum Dis.* 2007;66(Suppl 3):iii40-iii41. doi:10.1136/ard.2007.079798
31. Kamper SJ, Maher CG, Mackay G. Global rating of change scales: a review of strengths and weaknesses and considerations for design. *J Man Manip Ther.* 2009;17(3):163-170. doi:10.1179/jmt.2009.17.3.163
32. Kamper S. Global Rating of Change scales. *Aust J Physiother.* 2009;55(4):289. doi:10.1016/s0004-9514(09)70015-7
33. Bobos P, Ziebart C, Furtado R, Lu Z, MacDermid JC. Psychometric properties of the global rating of change scales in patients with low back pain, upper and lower extremity disorders. A systematic review with meta-analysis. *J Orthop.* 2020;21:40-48. doi:10.1016/j.jor.2020.01.047
34. Koehorst ML, van Trijffel E, Lindeboom R. Evaluative measurement properties of the patient-specific functional scale for primary shoulder complaints in physical therapy practice. *J Orthop Sports Phys Ther.* 2014;44(8):595-603. doi:10.2519/jospt.2014.5133
35. Hefford C, Abbott JH, Arnold R, Baxter GD. The patient-specific functional scale: validity, reliability, and responsiveness in patients with upper extremity musculoskeletal problems. *J Orthop Sports Phys Ther.* 2012;42(2):56-65. doi:10.2519/jospt.2012.3953
36. Michener LA, Snyder AR, Leggin BG. Responsiveness of the numeric pain rating scale in patients with shoulder pain and the effect of surgical status. *J Sport Rehabil.* 2011;20(1):115-128. doi:10.1123/jsr.20.1.115
37. Slade SC, Dionne CE, Underwood M, Buchbinder R. Consensus on Exercise Reporting Template (CERT): Explanation and Elaboration Statement. *Br J Sports Med.* 2016;50(23):1428-1437. doi:10.1136/bjssports-2016-096651
38. Profeta VLS, Turvey MT. Bernstein's levels of movement construction: A contemporary perspective. *Hum Mov Sci.* 2018;57:111-133. doi:10.1016/j.humov.2017.11.013
39. O'Neil J, Pelletier L, Bilodeau M, Egan M, Marshall S, Sveistrup H. A physiotherapist's perception of their own behavior compared to the perception of their behavior by persons with TBI within the context of telerehabilitation: A self-determination theory perspective. *Physiother Theory Pract.* 2022;39(8):1650-1661. doi:10.1080/09593985.2022.2046219

40. Silbernagel KG, Brorsson A, Lundberg M. The majority of patients with Achilles tendinopathy recover fully when treated with exercise alone: a 5-year follow-up. *Am J Sports Med.* 2011;39(3):607-613. doi:10.1177/0363546510384789
41. Thomeé R. A comprehensive treatment approach for patellofemoral pain syndrome in young women. *Phys Ther.* 1997;77(12):1690-1703. doi:10.1093/ptj/77.12.1690
42. Essery R, Geraghty AW, Kirby S, Yardley L. Predictors of adherence to home-based physical therapies: a systematic review. *Disabil Rehabil.* 2017;39(6):519-534. doi:10.3109/09638288.2016.1153160
43. Burns D, Boyer P, Razmjou H, Richards R, Whyne C. Adherence Patterns and Dose Response of Physiotherapy for Rotator Cuff Pathology: Longitudinal Cohort Study. *JMIR Rehabil Assist Technol.* 2021;8(1):e21374. doi:10.2196/21374
44. Heron SR, Woby SR, Thompson DP. Comparison of three types of exercise in the treatment of rotator cuff tendinopathy/shoulder impingement syndrome: A randomized controlled trial. *Physiotherapy.* 2017;103(2):167-173. doi:10.1016/j.physio.2016.09.001
45. Littlewood C, Bateman M, Brown K, et al. A self-managed single exercise programme versus usual physiotherapy treatment for rotator cuff tendinopathy: a randomised controlled trial (the SELF study). *Clin Rehabil.* 2016;30(7):686-696. doi:10.1177/0269215515593784
46. Chester R, Shepstone L, Daniell H, Sweeting D, Lewis J, Jerosch-Herold C. Predicting response to physiotherapy treatment for musculoskeletal shoulder pain: a systematic review. *BMC Musculoskelet Disord.* 2013;14(1):203. doi:10.1186/1471-2474-14-203
47. Pekyavas NO, Ergun N. Comparison of virtual reality exergaming and home exercise programs in patients with subacromial impingement syndrome and scapular dyskinesis: Short term effect. *Acta Orthop Traumatol Turc.* 2017;51(3):238-242. doi:10.1016/j.aott.2017.03.008
48. Struyf F, Nijs J, Mollekens S, et al. Scapular-focused treatment in patients with shoulder impingement syndrome: a randomized clinical trial. *Clin Rheumatol.* 2013;32(1):73-85. doi:10.1007/s10067-012-2093-2
49. Erlenwein J, Przemeczek M, Degenhart A, et al. The Influence of Chronic Pain on Postoperative Pain and Function After Hip Surgery: A Prospective Observational Cohort Study. *J Pain.* 2016;17(2):236-247. doi:10.1016/j.jpain.2015.10.013
50. Calatayud J, Vinstrup J, Jakobsen MD, Sundstrup E, Colado JC, Andersen LL. Influence of different attentional focus on EMG amplitude and contraction duration during the bench press at different speeds. *J Sports Sci.* 2018;36(10):1162-1166. doi:10.1080/02640414.2017.1363403
51. Coratella G, Tornatore G, Longo S, et al. The Effects of Verbal Instructions on Lower Limb Muscles' Excitation in Back-Squat. *Res Q Exerc Sport.* 2022;93(2):429-435. doi:10.1080/02701367.2020.1840496
52. Greig M, Marchant D. Speed dependant influence of attentional focusing instructions on force production and muscular activity during isokinetic elbow flexions. *Hum Mov Sci.* 2014;33:135-148. doi:10.1016/j.humov.2013.08.008
53. Kristiansen M, Samani A, Vuillerme N, Madeleine P, Hansen EA. External and Internal Focus of Attention Increases Muscular Activation During Bench Press in Resistance-Trained Participants. *J Strength Cond Res.* 2018;32(9):2442-2451. doi:10.1519/jsc.0000000000002613
54. Lohse KR, Sherwood DE. Thinking about muscles: the neuromuscular effects of attentional focus on accuracy and fatigue. *Acta Psychol.* 2012;140(3):236-245. doi:10.1016/j.actpsy.2012.05.009
55. Marchant DC. Attentional focusing instructions and force production. *Front Psychol.* 2010;1:210.

## SUPPLEMENTARY MATERIALS

### **Appendix 1**

Download: <https://ijspt.scholasticahq.com/article/90284-effect-of-a-novel-training-program-in-patients-with-chronic-shoulder-pain-based-on-implicit-motor-learning-pilot-and-feasibility-study/attachment/187251.pdf>

---

### **Appendix 2**

Download: <https://ijspt.scholasticahq.com/article/90284-effect-of-a-novel-training-program-in-patients-with-chronic-shoulder-pain-based-on-implicit-motor-learning-pilot-and-feasibility-study/attachment/187252.docx>

---