

Technology-Based Interventions for Nursing Home Residents: Implications for Nursing Home Practice Amid and Beyond the Influence of COVID-19: A Systematic Review Protocol

Zhaohui Su (✉ szh@utexas.edu)

University of Texas Health Science Center at San Antonio <https://orcid.org/0000-0003-2005-9504>

Kylie Meyer

University of Texas Health Science Center at San Antonio

Yue Li

University of Rochester Medical Center

Dean McDonnell

Institute of Technology Carlow

Nitha Mathew Joseph

University of Texas Health Science Center at Houston

Xiaoshan Li

Beijing Normal University-Hong Kong Baptist University United International College

Yan Du

University of Texas Health Science Center at San Antonio

Shailesh Advani

National Institute of Health

Ali Cheshmehzangi

University of Nottingham

Junaid Ahmad

Peshawar Medical College

Claudimar Pereira da Veiga

Universidade Federal do Parana

Roger Yat-Nork Chung

University of Hong Kong

Jing Wang

University of Texas Health Science Center at San Antonio

Keywords: COVID-19, novel coronavirus, SARS-CoV2, nursing home residents, older adults, underserved populations, technology-based interventions, socio-ecological model

DOI: <https://doi.org/10.21203/rs.3.rs-56102/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: A growing number of technology-based interventions are used to support the health and quality of life of nursing home residents. The onset of COVID-19 and recommended social distancing policies that followed led to an increased interest in technology-based solutions to provide healthcare and promote health. Yet, there are no comprehensive resources on technology-based healthcare solutions that describe their efficacy for nursing home residents. This systematic review will identify technology-based interventions designed for nursing home residents and describe the characteristics and effects of these interventions concerning the distinctive traits of nursing home residents and nursing facilities. Additionally, this paper will present practical insights into the varying intervention approaches that can assist in the delivery of broad digital health solutions for nursing home residents amid and beyond the impact of COVID-19.

Methods: Databases including PubMed, PsycINFO, CINAHL, and Scopus will be used to identify articles related to technology-based interventions for nursing home residents published between January 1st, 2020 to July 7th, 2020. Titles, abstracts, and full-texts papers will be reviewed against the eligibility criteria. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses procedures will be followed for the reporting process, and implications for existing interventions and research evaluated by a multidisciplinary research team.

Results: NA—protocol study

Conclusions: Our study will fill critical gaps in the literature by providing a review of technology-based interventions tested in the nursing home setting. As the older adult population grows, there is an urgent need to identify effective technology-based interventions that can address the distinctive characteristics and preferences of nursing home residents. Clear and comprehensive understanding of how available technology-based health solutions facilitate healthcare for nursing home residents will shed light on the approaches open to residents to fend off the negative health consequences amid and beyond the influence of COVID-19.

Systematic Review Registrations: PROSPERO [CRD 42020191880](https://www.crd.york.ac.uk/PROSPERO/record/CRD42020191880)

Background

Nursing homes have been described as a "ground zero" throughout the coronavirus outbreak [1-4]. While the final impact of COVID19 (coronavirus) on the short and longer-term health outcomes is still unclear as the pandemic continues to unfold [5], what is clear is that nursing home residents have suffered some of the gravest consequences of this pandemic so far [6]. Contributing to over 44% of COVID-19 deaths within the United States, residents and workers in nursing homes account for approximately 56,143 of coronavirus deaths (as of July 9th, 2020) [7]. However, a more harrowing realization is that the nursing home residents who have died from COVID-19 died without the care or company of their family members [8-10]. These staggering numbers underscore the urgent need for healthcare researchers to understand

factors that make nursing home residents more vulnerable to COVID-19, and to identify practical solutions that can address these factors in a timely fashion.

Nursing homes and nursing facilities provide long-term service and support for individuals living with chronic or disabling conditions who are unable to live at home independently [11-13]. Often living with multiple morbidities, in the United States approximately 85.1% of nursing home residents are 65 years and older, with 75.8% of these individuals experiencing from hypertension, 58.9% living with Alzheimer's Disease, and 53.0% of residents living with depression [13]. Three sets of factors likely contribute to the alarming COVID-19 death rates seen in nursing homes: (1) characteristics of nursing home residents [14-18], (2) distinctive attributes of nursing home facilities [1, 6, 9, 10, 19], and (3) the micro and macro-level supports available to nursing home residents [2, 20-24]. On a micro level, research indicates that nursing home residents are more susceptible to infection and fatal outcomes from COVID-19 because they are often older adults living with medical conditions that compromise the immune systems and lower their ability to combat the virus [13, 19, 25, 26]. Additionally, they often lack specific medical skills, such as utilizing telemedicine tools, or they may have a physical or cognitive impairment that impedes their ability to take care of their health and wellbeing [27-31]. The macro perspective focus on the unique characteristics of nursing home facilities, as they are typically operating on a close and shared-living environment - conditions that are ideal for the spread of the virus [10, 19, 32]. Further, nursing homes often lack adequate healthcare resources or infrastructure needed to curb the impact of COVID-19. For example, numerous studies have indicated a lack of investment in training programs for nursing home staff, in addition to high turnover rates [33, 34], that management teams are often ineffective [35, 36], and that the nursing home infrastructure is often too outdated [32, 37-39].

The third set of factors that contribute to nursing home residents' vulnerability to COVID-19 centers on social supports available to these adults. Nursing home residents often have limited access to micro-level social support, including support from family (e.g., infrequent visitation) [40], local community [41], and organizations (e.g., inexperienced or inadequately trained staff) [42]. Furthermore, nursing home residents often have limited macro-level social support. This is evidenced by harmful social norms (e.g., age-related discrimination)[43-45] and inadequate policy support that facilitates healthy aging and quality of life (e.g., insufficient regulatory oversight to ensure quality care in nursing homes) [4, 20, 46, 47].

The above areas of inquiry resonate with the core principles of the bioecological model [48-51], which highlights the way individuals are influenced by a series of synergistic interactions between intrapersonal and interpersonal factors (e.g., residents, resident families), organizational characteristics (e.g., nursing homes), policy (e.g., legislative response) and the social/community (e.g., ageism) context, and how these processes can change over time (See Figure 1.) To successfully and effectively protect nursing home residents from global health crises like COVID-19, stakeholders such as policymakers, healthcare professionals, informal caregivers, and older adults themselves all need to contribute to the change-making process [2, 46, 52, 53]. While some effective changes are resource-intensive, time-consuming, and need concerted efforts from multilevel stakeholders to achieve, there are cost-effective, efficient, and

accessible health solutions available to nursing home residents, such as technology-based interventions [54, 55].

Technology-based interventions can be considered as the use of technology (e.g., digital devices like tablets and wearable devices, communication platforms) to manage or support health promotion strategies that aim to produce accessible and affordable health solutions to a target audience. Compared to traditional health solutions, such as face-to-face consultations, technology-based interventions have the potential to deliver healthcare more effectively and can mitigate geographic and access-related limitations that, as studies show, can play a significant role within nursing homes [56-62]. The evidence further suggests that technology-based interventions can help free healthcare professionals from repetitive work and allow them to make more meaningful contributions in delivering healthcare solutions to the care recipients [63-65].

Telemedicine and other technology-based solutions are particularly crucial given circumstances rendered by the COVID-19 pandemic, such as the limited ability for some healthcare providers to enter residences or for residents to visit their healthcare team for primary care visits. Limiting the exposure to infection through the use of telemedicine may assist in situations where a resident is required to attend a hospital appointment and return to a residence, thus alleviating the potential risk for a virus to spread to others [66]. Further, technologies that support residents' ability to remain in contact with families and friends outside of skilled care settings may reduce the adverse effects of loneliness and social isolation that is more common among nursing home residents compared to community-dwelling older adults [17, 67].

While technology-based solutions have potential in delivering health solutions to nursing home residents [68, 69], there is limited awareness of the benefits and delivery options for state-of-art technology-based interventions specifically designed for nursing home residents. By factoring for the distinctive characteristics of nursing home residents and nursing home facilities, the main focus of this systematic review is to identify and evaluate technology-based interventions tailored explicitly for nursing home residents. Additionally, this research will present practical insights into the varying intervention approaches that can assist in the delivery of broad digital health solutions for nursing home residents amid and beyond the impact of COVID-19.

Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) procedures will be adhered to in the reporting process [70]. This systematic review is registered with the International Prospective Register of Systematic Reviews (PROSPERO) system (CRD 42020191880); these measures are to avoid unnecessary study duplication [71, 72], increase research rigor [73, 74], improve study comparability and replicability [75], and ultimately, promote quality and transparency in research [76].

Inclusion and exclusion criteria

Based on the research aim, inclusion criteria were set a priori (Table 1). In this study, nursing homes are defined as housing arrangements for older adults who are 65 years and older [77, 78] and no longer able to live independently at home. Technology-based interventions are defined as “the use of technology to manage or support health promotion strategies that aim to produce accessible and affordable health solutions to a target audience” [79]. Articles will be excluded if (1) the study sample did not include a majority nursing home residents (i.e., nursing home residents \leq 50% of the total research population), (2) the study did not include and discuss technology-based health solutions designed for nursing home residents, (3) the study did not adopt a randomized controlled trial design (e.g., studies with quasi-experimental design were excluded), and (4) the authors did not report original empirical findings on intervention outcomes (e.g., research protocols and systematic review studies were excluded from the review).

Search strategy

Databases including PubMed, PsycINFO, CINAHL, and Scopus, will be searched for eligible articles. A search strategy was developed in consultation with a librarian experienced in systematic review methods. Search terms used to locate articles will center on three concepts: nursing home residents, technology-based interventions, and randomized controlled trials. An example PubMed search string is illustrated in Table 2.

Study selection

Following the search, all citations will be collated and uploaded to Mendeley, and duplicate studies will be removed. Titles and then abstracts will be screened by two principle reviewers (ZS and XL) independently. The same screening procedure will be adopted in the full-text article review process on selected article abstracts. Reasons for exclusion will be recorded and detailed in the PRISMA flowchart. Discrepancies between reviewers will be resolved via group discussions using videoconferencing and email correspondence to reach a consensus.

Study quality assessment

The Cochrane Collaboration evaluation framework will be adopted to examine risk of bias of the included study [80]. The framework has seven domains: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and any other source of bias. Scores from the first five items will be used to determine whether the included study has a low or a high risk of bias [80]. A study is considered as having a low risk of bias if it rated as "low" on three or more of these five items [80, 81]. The risk of bias will be evaluated independently by two reviewers (ZS and XL). Any discrepancy regarding the risk of bias will be resolved by consensus via group discussions.

Data extraction and synthesis

Data on study design, sample characteristics (i.e., sample size and sample details), intervention characteristics (i.e., technology type, intervention application, intervention exposure, and intervention materials), outcome variables assessed, and research findings will be extracted by mainly eight reviewers (ZS, XL, DMD, AC, JA, SA, KM, and YD). Findings from the included studies will be narratively synthesized to examine the characteristics and effects of the interventions. Gaining a more structured understanding of the interventions, the multidisciplinary study team will organize insights on intervention application and outcomes in tandem with the distinctive traits of nursing home residents and the overall nursing home environment. Due to the heterogeneity found within the articles included, meta-analyses are not considered.

Results

NA: As this is a protocol study

Discussion

There is a growing body of technology-based interventions designed to support the health and quality of life of nursing home residents [56–62]. The onset of COVID-19 and recommended social distancing policy led to an increased interest in reliance on technology-based solutions [82, 83]. However, while some useful insights are available, research has yet to provide comparative insight into the state of development of these interventions and how current evidence apply in the context of the COVID-19 pandemic. The use of the socio-ecological model, combined with multidisciplinary expertise, provides a framework to present practical insights on how these interventions can be utilized to deliver health solutions to nursing home residents amid and beyond the impact of COVID-19.

This research fills a critical gap in the literature by consolidating, in one place, the evidence for technology-based interventions empirically tested with nursing home populations. As the older adult population grows, there is an urgent need to identify effective technology-based interventions that can address the distinctive characteristics and preferences of nursing home residents [84, 85]. Improving person-centered care and the delivery of effective care solutions to nursing home residents, especially as the pandemic continues, is of critical importance. Comprehensive understanding of how available technology-based health solutions facilitate healthcare for nursing home residents can help shed light on approaches that are available to these residents to fend off the negative health consequences amid and beyond the influence of COVID-19. While the COVID-19 pandemic has revealed troubling vulnerabilities in the long-term care system across the globe, it also shows how telemedicine can support nursing home residents and their families. Technology can also assist clinicians in connecting with patients when in-person medical visits are difficult or dangerous (e.g., in rural settings, following natural disasters). Telemedicine and other technology-based interventions have the potential to provide a comprehensive range of benefits. This research also serves as a platform for care institutions and policymakers to inform government policies and further justify the role technology can play in strengthening the service provision across nursing homes and facilities.

Conclusions

NA: As this is a protocol study

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

No

Competing interests

None.

Funding

None.

Authors' contributions

ZS developed the research idea and wrote and manuscript. KM, YL, DMD, NMJ, XL, YD, SA, AC, JA, CPV, YNC, and JW reviewed and revised the manuscript.

Acknowledgements

The authors wish to express their gratitude to Emme Lopez, the academic librarian who helped with finalizing the search strategy. Gratitude also goes to the editors and reviewers for their constructive input.

Abbreviations

COVID-19: covronavirus disease 2019

References

1. Barnett ML, Grabowski DC. Nursing homes are ground zero for COVID-19 pandemic. JAMA Health Forum. 2020;1(3):e200369–9.

2. Pillemer K, Subramanian L, Hupert N. *The importance of long-term care populations in models of COVID-19*. JAMA, 2020.
3. Benson JJ, et al. Online social support groups for informal caregivers of hospice patients with cancer. *European Journal of Oncology Nursing*. 2020;44:101698.
4. Li Y, et al., *COVID-19 infections and deaths among Connecticut nursing home residents: Facility correlates*. *Journal of the American Geriatrics Society*, 2020. n/a(n/a).
5. Cutler D. How will COVID-19 affect the health care economy? *JAMA Health Forum*. 2020;1(4):e200419–9.
6. Fallon A, et al., *COVID-19 in nursing homes*. *QJM*, 2020: p. hcaa136.
7. Kaiser Family Foundation, *State data and policy actions to address coronavirus*. 2020, Kaiser Family Foundation.
8. Davidson PM, Szanton SL, *Nursing homes and COVID-19: We can and should do better*. *Journal of Clinical Nursing*, 2020. n/a(n/a).
9. Comas-Herrera A, et al., *Mortality associated with COVID-19 outbreaks in care homes: Early international evidence*. International Long Term Care Policy Network, 2020.
10. McMichael TM, et al. Epidemiology of Covid-19 in a long-term care facility in King County, Washington. *N Engl J Med*. 2020;382(21):2005–11.
11. Backman A, et al. Characteristics of highly rated leadership in nursing homes using item response theory. *J Adv Nurs*. 2017;73(12):2903–13.
12. National Institute for Aging. *Residential facilities, assisted living, and nursing homes*. 2017 [cited 2020 June 14]; Available from: <https://www.nia.nih.gov/health/residential-facilities-assisted-living-and-nursing-homes>.
13. U.S. Department of Health And Human Services, *Long-term care providers and services users in the United States, 2015–2016*, in *Vital and Health Statistics*. 2019, Centers for Disease Control and Prevention.
14. Dowd JB, et al., *Demographic science aids in understanding the spread and fatality rates of COVID-19*. *Proceedings of the National Academy of Sciences*, 2020. 117(18): p. 9696.
15. CDC COVID-19 Response Team. Severe outcomes among patients with coronavirus disease 2019 (COVID-19)—United States, February 12–March 16, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(12):343–6.
16. Wherton J, et al. Designing assisted living technologies 'in the wild': Preliminary experiences with cultural probe methodology. *BMC Med Res Methodol*. 2012;12:188.
17. Hado E, Friss Feinberg L. *Amid the COVID-19 pandemic, meaningful communication between family caregivers and residents of long-term care facilities is imperative*. *Journal of Aging & Social Policy*, 2020: p. 1–6.
18. Liu Y-B, et al. Social support, health literacy, and health care utilization among older adults. *Asia Pacific Journal of Public Health*. 2019;31(4):359–66.

19. Etard J-F, et al., *Potential lethal outbreak of coronavirus disease (COVID-19) among the elderly in retirement homes and long-term facilities, France, March 2020*. Euro surveillance: bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin, 2020. 25(15): p. 2000448.
20. Gurwitz JH. COVID-19, post-acute care preparedness and nursing homes: Flawed policy in the fog of war. *J Am Geriatr Soc*. 2020. 10.1111/jgs.16499. p.
21. Laditka SB, et al. Providing shelter to nursing home evacuees in disasters: Lessons from Hurricane Katrina. *Am J Public Health*. 2008;98(7):1288–93.
22. Schott N, Tietjens M. Exploring the mediating role of social support and fall efficacy on the association between falls and physical activity: A cross-sectional study in an assisted-living population. *J Aging Phys Act*. 2019;27(1):53–60.
23. Grabowski DC, Joynt KE, Maddox. Postacute care preparedness for COVID-19: Thinking ahead. *JAMA*. 2020;323(20):2007–8.
24. White EM, Aiken LH, McHugh MD. Registered nurse burnout, job dissatisfaction, and missed care in nursing homes. *J Am Geriatr Soc*. 2019;67(10):2065–71.
25. Applegate WB, Ouslander JG. COVID-19 presents high risk to older persons. *J Am Geriatr Soc*. 2020;68(4):681–1.
26. Centers for Disease Control and Prevention. *Coronavirus disease 2019 (COVID-19): Older adults*. 2020; Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/older-adults.html>.
27. Scott Kruse C, et al. Evaluating barriers to adopting telemedicine worldwide: A systematic review. *Journal of Telemedicine Telecare*. 2016;24(1):4–12.
28. Nouri S, et al., *Addressing equity in telemedicine for chronic disease management during the Covid-19 pandemic*. NEJM Catalyst Innovations in Care Delivery, 2020. 1(3).
29. Berkowsky RW, Czaja SJ. Challenges associated with online health information seeking among older adults, *Aging, Technology and Health*, R. Pak and A.C. McLaughlin, Editors. 2018, Academic Press: San Diego. pp. 31–48.
30. Piper AM, Brewer R, Cornejo R. Technology learning and use among older adults with late-life vision impairments. *Univ Access Inf Soc*. 2017;16(3):699–711.
31. Cutilli CC, et al., *Health literacy, health disparities, and sources of health information in U.S. older adults*. *Orthopaedic Nursing*, 2018. 37(1).
32. Benson NM, Öngür D, Hsu J. COVID-19 testing and patients in mental health facilities. *The Lancet Psychiatry*. 2020;7(6):476–7.
33. Fosse A, et al. Nursing home physicians' assessments of barriers and strategies for end-of-life care in Norway and the Netherlands. *J Am Med Dir Assoc*. 2017;18(8):713–8.
34. Quigley DD, et al. COVID-19 preparedness in nursing homes in the midst of the pandemic. *J Am Geriatr Soc*. 2020. 10.1111/jgs.16520. p.

35. Glette MK, et al. Nursing home leaders' and nurses' experiences of resources, staffing and competence levels and the relation to hospital readmissions - a case study. *BMC Health Serv Res.* 2018;18(1):955.
36. Vaughn VM, et al., *Characteristics of healthcare organisations struggling to improve quality: Results from a systematic review of qualitative studies.* *BMJ Quality & Safety*, 2019. 28(1): p. 74.
37. Solis J, et al., *Structural vulnerability in the United States revealed in three waves of novel coronavirus disease (COVID-19).* *The American Journal of Tropical Medicine and Hygiene*, 2020: p. tpmd200391.
38. Zimmerman S, et al. The need to include assisted living in responding to the COVID-19 pandemic. *J Am Med Dir Assoc.* 2020;21(5):572–5.
39. Khan F, Driessen J. Bridging the telemedicine infrastructure gap: Implications for long-term care in rural America. *Public Policy Aging Report.* 2018;28(3):80–4.
40. Miller VJ. Investigating barriers to family visitation of nursing home residents: A systematic review. *Journal of Gerontological Social Work.* 2019;62(3):261–78.
41. Stone PW, et al. The expansion of National Healthcare Safety Network enrollment and reporting in nursing homes: Lessons learned from a national qualitative study. *Am J Infect Control.* 2019;47(6):615–22.
42. Hales C, et al., *The care of older adults with extreme obesity in nursing homes: A collective case study.* medRxiv, 2019: p. 19013326.
43. Jimenez-Sotomayor MR, Gomez-Moreno C, and E. Soto-Perez-de-Celis, *Coronavirus, ageism, and Twitter: An evaluation of tweets about older adults and COVID-19.* *Journal of the American Geriatrics Society*, 2020. n/an/a.
44. Han SD, Mosqueda L. Elder abuse in the COVID-19 era. *J Am Geriatr Soc.* 2020. 10.1111/jgs.16496. p.
45. Fraser S, et al., *Ageism and COVID-19: What does our society's response say about us?* *Age and ageing*, 2020: p. afaa097.
46. Comas-Herrera A, et al., *COVID-19: Implications for the support of people with social care needs in England.* *Journal of Aging & Social Policy*, 2020: p. 1–8.
47. Ko M, Wagner L, Spetz J. *Nursing home implementation of health information technology: Review of the literature finds inadequate investment in preparation, infrastructure, and training.* *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*, 2018. 55: p. 0046958018778902.
48. Bronfenbrenner U. Toward an experimental ecology of human development. *Am Psychol.* 1977;32(7):513–31.
49. Bronfenbrenner U. *The ecology of human development: Experiments by nature and by design.* Cambridge: Harvard University Press; 1979.
50. McLeroy KR, et al. An ecological perspective on health promotion programs. *Health Educ Q.* 1988;15(4):351–77.

51. Bronfenbrenner U, Morris P. A., *The bioecological model of human development*, in *Volume 1: Theoretical Models of Human Development*. 2007: *Handbook of child psychology*.
52. Cooper E, et al. Priorities for the professional development of registered nurses in nursing homes: A Delphi study. *Age Ageing*. 2016;46(1):39–45.
53. Cutter CM, Nelson C, Abir M, *Accountability to population health in the COVID-19 pandemic: Designing health care delivery within a social responsibility framework*. Population Health Management, 2020.
54. Chiu C-J, Wu CH. Information and communications technology as a health promotion method for older adults in assisted-living facilities: Three-arm group-randomized trial. *JMIR Aging*. 2019;2(1):e12633.
55. Banbury A, et al. Adding value to remote monitoring: Co-design of a health literacy intervention for older people with chronic disease delivered by telehealth - The telehealth literacy project. *Patient Educ Couns*. 2020;103(3):597–606.
56. Ohannessian R, Duong TA, Odone A. Global telemedicine implementation and integration within health systems to fight the COVID-19 pandemic: A call to action. *JMIR Public Health Surveill*. 2020;6(2):e18810.
57. Smith AC, et al., *Telehealth for global emergencies: Implications for coronavirus disease 2019 (COVID-19)*. *Journal of Telemedicine and Telecare*, 2020: p. 1357633 × 20916567.
58. Wang J, et al. Connecting smartphone and wearable fitness tracker data with a nationally used electronic health record system for diabetes education to facilitate behavioral goal monitoring in diabetes care: Protocol for a pragmatic multi-site randomized trial. *JMIR Res Protoc*. 2018;7(4):e10009.
59. Zhou X, et al. The role of telehealth in reducing the mental health burden from COVID-19. *Telemedicine e-Health*. 2020;26(4):377–9.
60. Mette LA, et al. Reaching high-risk underserved individuals for cancer genetic counseling by video-teleconferencing. *J Community Support Oncol*. 2016;14(4):162–8.
61. Hollander JE, Carr BG. Virtually perfect? Telemedicine for Covid-19. *N Engl J Med*. 2020;382(18):1679–81.
62. Wang J, et al., *A behavioral lifestyle intervention enhanced with multiple-behavior self-monitoring using mobile and connected tools for underserved individuals with type 2 diabetes and comorbid overweight or obesity: Pilot comparative effectiveness trial*. *JMIR mHealth uHealth*, 2018. 6(4): p. e92.
63. Keyworth C, et al. What maximizes the effectiveness and implementation of technology-based interventions to support healthcare professional practice? A systematic literature review. *BMC Med Inform Decis Mak*. 2018;18(1):93.
64. Ehn M, Johansson A-C, Revenäs Å. *Technology-based motivation support for seniors' physical activity—A qualitative study on seniors' and health care professionals' views*. *International Journal of Environmental Research and Public Health*, 2019. 16(13).

65. Bagot KL, et al. Sustainable implementation of innovative, technology-based health care practices: A qualitative case study from stroke telemedicine. *Journal of Telemedicine Telecare*. 2018;26(1–2):79–91.
66. Naylor MD, Hirschman KB, McCauley K. Meeting the transitional care needs of older adults with COVID-19. *Journal of Aging Social Policy*. 2020;32(4–5):387–95.
67. Prieto-Flores ME, et al. Factors associated with loneliness of noninstitutionalized and institutionalized older adults. *J Aging Health*. 2011;23(1):177–94.
68. Mahmood S, et al. Global preparedness against COVID-19: We must leverage the power of digital health. *JMIR Public Health Surveill*. 2020;6(2):e18980.
69. Banskota S, Healy M, Goldberg EM. 15 smartphone apps for older adults to use while in isolation during the COVID-19 pandemic. *West J Emerg Med*. 2020;21(3):514–25.
70. Moher D, et al. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA statement. *PLoS Medicine*. 2009;6(7):e1000097.
71. Stewart L, Moher D, Shekelle P. Why prospective registration of systematic reviews makes sense. *Syst Rev*. 2012;1:7.
72. Chang SM, Slutsky J. Debunking myths of protocol registration. *Syst Rev*. 2012;1:4.
73. Xu C, et al. Protocol registration or development may benefit the design, conduct and reporting of dose-response meta-analysis: Empirical evidence from a literature survey. *BMC Med Res Methodol*. 2019;19(1):78.
74. dos Santos MBF, et al. Protocol registration improves reporting quality of systematic reviews in dentistry. *BMC Med Res Methodol*. 2020;20(1):57.
75. Shokraneh F. *Reproducibility and replicability of systematic reviews*. *World Journal of Meta-Analysis*, 2019. 7(3).
76. Ramstrand N, et al. Promoting quality and transparency in clinical research. *Prosthet Orthot Int*. 2019;43(5):474–7.
77. Lemoyne SE, et al. Appropriateness of transferring nursing home residents to emergency departments: A systematic review. *BMC Geriatr*. 2019;19(1):17.
78. van den Berg MEL, et al. Understanding the barriers and enablers to using outdoor spaces in nursing homes: A systematic review. *Gerontologist*. 2020;60(4):e254–69.
79. Su Z, et al., *Understanding technology-based interventions for caregivers of cancer patients: A systematic review-based concept analysis*. *Journal of Medical Internet Research*, 2020.
80. Higgins JP, et al., *Cochrane handbook for systematic reviews of interventions*. 2019: John Wiley & Sons.
81. van Tulder M, et al. Updated method guidelines for systematic reviews in the cochrane collaboration back review group. *Spine (Phila Pa 1976)*. 2003;28(12):1290–9.
82. Goodman-Casanova JM, et al. Telehealth home support during COVID-19 confinement for community-dwelling older adults with mild cognitive impairment or mild dementia: Survey study. *J*

Med Internet Res. 2020;22(5):e19434.

83. Gao Z, et al., *Virtual reality exercise as a coping strategy for health and wellness promotion in older adults during the COVID-19 pandemic*. Journal of Clinical Medicine, 2020. 9(6).
84. Schneider JE, et al. Economic evaluation of passive monitoring technology for seniors. Aging Clin Exp Res. 2020;32(7):1375–82.
85. Sapci AH, Sapci HA. Innovative assisted living tools, remote monitoring technologies, artificial intelligence-driven solutions, and robotic systems for aging societies: Systematic review. JMIR Aging. 2019;2(2):e15429.

Tables

Table 1. Study inclusion criteria

Category	Criteria
Study population	Adults (≥ 65 years old) living in nursing homes
Intervention/health solution	Technology-based interventions (e.g., digital tools such as smartphones and tablets, sensor devices, internet-based programs)
Key variable	Detailed descriptions of the technology-based interventions (i.e., purpose of the intervention, use of technology, application of the interventions, intervention exposure, outcome variables assessed/measured, and whether the design of the intervention material is tailored to nursing home residents)
Study type	Original research (i.e., research that reports original and empirical research findings)
Study design	Randomized controlled trials
Study outcome	Empirical reporting of the effect of the intervention (i.e., qualitative designs excluded)

Table 2. Example PubMed search string

Concept	Search string
Nursing homes	<p>“nursing home*”[MeSH] OR “nursing home*”[TIAB] OR “nursing homes”[MeSH] OR “nursing homes”[TIAB] OR “residential home*”[MeSH] OR “residential home*”[TIAB] OR “caring home*”[TIAB] OR “home for the aged” [MeSH] “home for the aged” [TIAB] OR “long term care”[MeSH] OR “long term care”[TIAB] OR “senior housing”[TIAB] OR “assisted living facilities”[MeSH] OR “assisted living facilities”[TIAB] OR (senior[TIAB] OR geriatric[TIAB] OR elderly[TIAB] OR aged[TIAB] OR elder[TIAB] “older adults”[TIAB]) AND (housing[TIAB] OR living[TIAB])</p>
Technology-based Interventions	<p>“technology”[MeSH] OR “technology”[TIAB] OR “eHealth”[TIAB] OR “telemedicine”[MeSH] OR “telemedicine”[TIAB] OR “tele-medicine”[MeSH] OR “tele-medicine”[TIAB] OR “telehealth”[TIAB] OR “tele-health”[TIAB] OR “connected health”[TIAB] OR “digital health”[TIAB] OR “mHealth”[TIAB] OR “mobile health”[TIAB]</p>
Randomized controlled trials	<p>randomized controlled trial[PT] OR randomized controlled trials as topic[MH] OR random allocation [MH] OR double-blind method[MH] OR single-blind method[MH] OR random*[tw] OR "Placebos"[MeSH] OR placebo[TIAB] OR ((singl*[tw] OR doubl*[tw] OR trebl*[TW] OR tripl*[TW]) AND (mask*[TW] OR blind*[TW] OR dumm*[TW]))</p>

Figures

Figure 1: An ecological model of factors increasing nursing home resident vulnerability to COVID-19




Figure 1

An ecological model of factors increasing nursing home resident vulnerability to COVID-19

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [NursingHomePRISMA2009checklist.doc](#)