



ORAL PRESENTATION

Open Access

# An altered maturation and adhesion phenotype of dendritic cells in diseased individuals compared to asymptomatic carriers of human T-cell leukemia virus type 1

Sharrón L Manuel<sup>1</sup>, Mohit Sehgal<sup>1</sup>, Zafar K Khan<sup>1</sup>, James J Goedert<sup>2</sup>, Michael R Betts<sup>3</sup>, Pooja Jain<sup>1\*</sup>

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses  
Montreal, Canada. 26-30 June 2013

As critical effectors of antiviral immune response, dendritic cells (DCs) are implicated to play an important role in determining the outcome of HTLV-1 infection. However, a complete understanding of their role in any disease pathogenesis requires extensive assessment of the phenotypic and functional state of DCs. To enable this, we developed a polychromatic antibody cocktail comprising key phenotypic and functional markers of DCs and applied it in a patient cohort from the HTLV-1 endemic region, Jamaica, consisted of seronegative controls, asymptomatic carriers (ACs), ATL, and HAM/TSP patients. This ex vivo analyses included two major subsets of blood DCs, myeloid and plasmacytoid (mDCs and pDCs, respectively). The comparative analyses of results demonstrated a decreased pDC frequency in both ATL and HAM/TSP patients as compared to ACs and seronegative controls. Similarly, CD86 expression on both mDCs and pDCs was significantly higher in HAM/TSP (but not ATL) patients compared to ACs. Interestingly, HLA-DR expression was significantly lower on pDCs of patients as compared to carriers, whereas for mDCs, only the HAM/TSP group had significantly lower expression of HLA-DR. Unlike HAM/TSP individuals, ATL individuals had significantly higher HLA-ABC expression on mDCs compared to ACs. Finally, both mDCs and pDCs of HAM/TSP patients had significantly higher PD-L1 expression compared to ACs. Overall, this study suggests that DCs are differentially regulated between patients (ATL and HAM/TSP)

and carriers of HTLV-1 and could provide an important tool to understand HTLV-1 immunopathogenesis during infection and disease.

**Authors' details**

<sup>1</sup>Drexel Institute for Biotechnology and Virology Research, and the Department of Microbiology and Immunology, Drexel University College of Medicine, Doylestown, PA, USA. <sup>2</sup>National Cancer Institute, Division of Cancer Epidemiology and Genetics, Bethesda, MD, USA. <sup>3</sup>Department of Microbiology and Immunology, University of Pennsylvania School of Medicine, Philadelphia, PA, USA.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-O67

Cite this article as: Manuel et al.: An altered maturation and adhesion phenotype of dendritic cells in diseased individuals compared to asymptomatic carriers of human T-cell leukemia virus type 1. *Retrovirology* 2014 **11**(Suppl 1):O67.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Drexel Institute for Biotechnology and Virology Research, and the Department of Microbiology and Immunology, Drexel University College of Medicine, Doylestown, PA, USA

Full list of author information is available at the end of the article