



ORIGINAL ARTICLE

Evaluation of the complementary feeding practices, dietary intake, and nutritional status of infants on a cow's milk protein elimination diet



Juliana Frizzo ^a, Vanessa C.C. Rodrigues ^a, Patrícia G.L. Speridião ^{b,c},
Mauro B. Morais ^{b,d,*}

^a Universidade Federal de São Paulo, Departamento de Pediatria, Divisão de Gastroenterologia Pediátrica, Programa de Pós-Graduação em Nutrição, São Paulo, SP, Brazil

^b Universidade Federal de São Paulo, Departamento de Pediatria, Divisão de Gastroenterologia Pediátrica, São Paulo, SP, Brazil

^c Universidade Federal de São Paulo, Curso de Nutrição, Departamento de Saúde, Educação e Sociedade, Santos, SP, Brazil

^d Escola Paulista de Medicina, Gastroenterologia Pediátrica, São Paulo, SP, Brazil

Received 22 March 2021; accepted 1 June 2021

Available online 22 July 2021

KEYWORDS

Complementary feeding;
Infant nutrition;
Milk hypersensitivity (Milk allergy);
Nutritional status

Abstract

Objective: To evaluate the complementary feeding practices, food intake, and nutritional status of infants on a cow's milk protein elimination diet.

Methods: A cross-sectional and observational study was conducted to compare infants aged 4–18 months who were on a cow's milk protein elimination diet with a control group of healthy infants without any dietary restrictions. General information on the child's health, demographic data, and food consumption were collected.

Results: The study included 96 infants in the elimination diet group and 99 in the control group. In the elimination diet group, the median age (in months) of introduction of solid foods (5.0×4.0 ; $p < 0.001$) and water (5.5×4.0 ; $p < 0.05$) was later, consumption of soft drinks and industrialized cookies was less frequent ($p < 0.05$), and a lower index of complementary feeding inadequacies (2.75×3.50 ; $p < 0.001$) was observed. The elimination diet group presented lower individual values of Z scores for weight/age, weight/height, and body mass index/age, although they were fed with higher amounts of energy (117.4×81.3 kcal/kg of weight; $p < 0.001$) and macro-and micronutrients, except for vitamin A. In the elimination diet group, breast milk and its substitutes contributed to more than 67% of energy intake. Although calcium consumption was a deficit in 31.5% of the infants, none received supplementation.

Conclusion: Infants on an elimination diet presented more adequate complementary feeding practices and higher nutritional intake, despite lower body weight values.

* Corresponding author.

E-mail: maurobmorais@gmail.com (M.B. Morais).

Introduction

An elimination diet for cow's milk and dairy products is recommended for infants with a suspected or confirmed diagnosis of cow's milk protein allergy (CMPA).^{1,2} However, the elimination diet is followed by a greater number of infants in the initial phase, before diagnostic confirmation, and also by patients who for various reasons do not undergo a diagnostic confirmation test. An elimination diet should always be adequate to ensure full growth and development.

Until 6 months of age, exclusive breastfeeding should be maintained, and the breastfeeding mother should eliminate cow milk from her diet. Infants who are not breastfed should be fed special formulas without cow's milk protein.^{1–3} Complementary feeding should be started after 6 months of age. Complementary feeding is defined as all foods offered in addition to breast milk during the period that child will not be exclusively breastfed. The same definition may be used for infants receiving artificial feeding.^{3,4} According to national and international guidelines, complementary feeding for infants with CMPA should follow the recommendations for healthy children. Hence, no restriction or delay in the introduction of foods, including potentially allergenic foods, should be done^{1,3,4} as often occurs in healthcare practice.^{5,6} Complementary feeding is important to define lifelong eating habits and behaviors.^{4,7}

Despite the relevance of the quality of diet for infants on an elimination diet, there are only a few studies on this subject. There are pieces of evidence of dietary nutrient deficiencies, growth deficits, and low food intake.^{8–12} To the best of the authors's knowledge, no publication on complementary feeding practices of infants on an elimination diet for cow's milk and dairy products is available.

Thus, the objective of this study was to evaluate the complementary feeding practices, dietary intake, and nutritional status classification of infants on a cow's milk protein elimination diet compared to a control group.

Methods

A cross-sectional and observational study approved by the Research Ethics Committee of the Federal University of São Paulo (UNIFESP) was conducted. Infants aged 4–18 months who were on a diet free of cow's milk and dairy products for a minimum period of 1 month owing to suspected or confirmed diagnosis of CMPA were considered for this study. These infants were included regardless of the type of clinical presentation or criteria used to establish the diagnosis. Confirmation of the correct completion of the cow's milk elimination diet was carried out during the individual interview. It was based on a specific question of whether the infant was receiving or not cow's milk. The elimination of cow's milk was also evaluated in the food survey. Infants on the elimination diet for other reasons besides cow's milk allergy, such as vegetarianism, veganism, or galactosemia, were not

admitted to the study. A control group consisting of healthy infants without any dietary restrictions was included for comparison purposes.

The parents/caregivers of children in both the groups were aged ≥ 18 years and they provided written informed consent. In case the volunteers had more than one eligible child, only the youngest child was included. Infants previously diagnosed with other severe diseases or those requiring major dietary changes, healthy infants living with children with CMPA, and infants who weren't correctly complying with the exemption diet were excluded.

The cow's milk protein elimination diet group was recruited in the metropolitan region of São Paulo, at the Food Allergy Clinic of the Pediatric Gastroenterology Department of UNIFESP and at Vila Mariana High-Cost Pharmacy of the Health Department of the State of São Paulo, Brazil, similar to that reported in a previous study.¹³ The latter is a pharmacy from the Health Public System where special formulas are supplied free of charge to infants on an elimination diet living in São Paulo, which complies with the terms established in legislation (Resolution SS-336, of November 27, 2007). The control group was recruited at Vila dos Remédios Basic Public Health Unit and at a church located in the Butantã neighborhood.

The sample size estimative was based on a previous Brazilian study that revealed that children on a cow's milk protein elimination diet had insufficient intake of energy and other nutrients such as calcium. There were differences of more than 40% in the proportions of children with inadequate intake in the study and control groups.⁸ Thus, in the present study, the sample size was calculated to identify a smaller difference between the groups (15%) with expectations of intake below the recommendation in 20% of the infants on an elimination diet and 5% of those on a diet without exclusions.

The data were collected between January 2018 and April 2019, through individual interviews using a standardized protocol that contained the parents/caregivers socioeconomic and educational information, the principal clinical presentation that motivated the suspicion or diagnosis of CMPA; and information on feeding, supplementation, and professionals who provided guidance on complementary feeding. The socio-economic classification was performed using the "ABEP - Brazilian Association of Research Companies" questionnaire (2015).¹⁴ It was expressed in decreasing order of strata, being class A the highest socioeconomic level, followed by classes B1, B2, C1, C2, and DE (the lowest socioeconomic level).

To evaluate the complementary feeding practices, information regarding the age at which food groups were introduced, forms of preparation, and supply of ultra-processed foods were collected. The index of complementary feeding inadequacies developed in Brazil was calculated.¹⁵ This index is specified in Supplementary Material 1. A formula based on a multiple linear regression model was used to estimate the volume of breast milk ingested.^{16,17} Food intake

was evaluated based on the usual daily food intake method.⁸ Data on food intake and complementary feeding practices were compared to the recommendations of the Brazilian Society of Pediatrics,⁴ Food Guide for Children under 2 years of age,⁷ and Institute of Medicine.¹⁸ The DietWin PC¹⁹ version 2.5.4.26856 software was used to calculate macro- and micronutrient intake.

The World Health Organization's Anthro software version 3.2.2 was used to evaluate and classify nutritional status based on weight and height data.²⁰

The infant weight and height measurements were performed on a Welmy[®] pediatric electronic scale, with a division scale of 5 g, a minimum load of 100 g, and capacity of 15 kg, after consultation with a pediatrician and under the supervision of the main researcher. Height was measured using a horizontal stadiometer, with a precision of 1 mm.²¹ When weight and height could not be checked, values measured and registered by the infants' pediatricians were used, since the measurements had been performed in the previous 30 days.

The Z scores for weight/age, weight/height, body mass index (BMI)/age, and height/age were calculated according to the World Health Organization standards.²⁰ The Z scores of infants who had their weight and height values extracted from the pediatrician's records were calculated considering the date of weight and height measurement.

The statistical analysis was performed using the Epi-Info 7.2.3.1 (Center of Disease Control and Prevention, Atlanta, GA, USA) and SigmaPlot 12.5 (Systat Software, San Jose, CA, USA) software. A significance level of 5% was considered in all analyses.

Results

A total of 230 parents/caregivers were interviewed for this study. Twenty infants with severe diseases or conditions that could lead to nutritional deficiencies or dietary restrictions; six with transgressions in the elimination diet; two with siblings already included in the group; and seven with incomplete data collection were excluded. Thus, the study included 96 infants in the elimination diet group and 99 in the control group.

Table 1 shows the demographic characteristics, breastfeeding habits, family history of atopy, and nutritional status of both groups. The groups were similar in terms of age, sex, frequency of premature birth, weight at birth, and frequency of exclusive breastfeeding for a period of 4 months or more. Infants on an elimination diet had a lower frequency of vaginal delivery and breastfeeding, and a lower age at the first contact with the cow's milk protein. Furthermore, these infants had a higher frequency of family history of atopy and CMPA and previous use of antibiotics. Regarding the socio-economic classification, the elimination diet group presented a higher frequency of families belonging to classes A and B (70.8% vs. 37.5%; $p < 0.001$) and mothers with bachelor's degrees (66.7% vs. 25.0%; $p < 0.001$).

The elimination diet group presented lower individual values of Z scores for weight/age, weight/height, and body mass index/age as presented in Table 1. No statistical difference between the groups regarding height/age z score was observed.

In the elimination diet group, 27.1% (26/96) of the infants were fed breast milk with other foods (21 with special formulas and complementary foods, four with complementary foods only, and one with special formula only). Other 70.8% (68/96) were fed special formulas, of which 61 also received complementary foods. Finally, 2.1% (2/96) of the infants were fed with soy drinks as the only substitute for breast milk, although they already had a medical prescription for soy-based infant formulas and were awaiting supply by the government. Of the 90 infants who were fed special formulas, regardless of whether they received breast milk, 47 (52.2%) received free amino acid formula, 36 (40.0%) received formula based on extensively hydrolyzed milk protein, six (6.7%) were on soy-based infant formula alone, and one (1.1%) was fed a formula based on extensively hydrolyzed rice protein. Only one infant had a soy-based infant formula introduced before 6 months of age.

In the control group, 60.6% (60/99) of the infants were breastfed (eight exclusively, 26 with other dairy sources and complementary foods, and 26 with complementary foods only). Other 23.2% (23/99) received infant formula with complementary foods. Finally, 14.1% (14/99) were fed whole cow's milk, and 2.0% (2/99) received dairy products.

In the elimination diet group, other foods than cow's milk were excluded from the diet of 19 infants. The most commonly reported were eggs ($n = 8$), wheat ($n = 7$), and bananas ($n = 7$).

Table 2 presents the results of complementary feeding introduction. Water, leafy vegetables, legumes, wheat, and eggs were introduced later in the elimination diet group. The introduction of tea, tubers, roots, and cereals was also later in this group, although the difference was not statistically significant. Delay was observed in the introduction of potentially allergenic foods, such as egg and wheat. The proportion of children who had already consumed soft drinks and industrialized cookies was higher in the control group.

The index of complementary feeding inadequacies¹⁵ (Supplementary Material 1) was lower in the elimination diet group. Therefore, liquified or sieved soups, which are inadequate inconsistency, had already been consumed by 47.9% and 52.1% of the elimination diet and control groups, respectively.

As shown in Supplementary Material 2A, the elimination diet group had a higher energy and macronutrient intake than the control group. Breast milk and its substitutes provided a higher percentage of energy and macronutrient intake in the elimination diet group than in the control group (Supplementary Material 2B). The median energy and protein intakes per kilogram of body weight were higher in the elimination diet group than in the control group. A higher daily volume of formula was consumed by the infants in the elimination diet group than by those in the control group.

Table 3 shows the consumption of micronutrients provided by breast milk and its substitutes, complementary foods, and supplements. The elimination diet group intake was highest for calcium, zinc, and vitamin C. However, the consumption of vitamin A was lower in the elimination diet group and no difference regarding iron and vitamin D intake was observed between the groups.

Table 4 shows the number and percentage of infants who had a food intake lower than the Recommended Dietary Allowances¹⁸ for calcium, iron, zinc, and vitamin A, C, and

Table 1 Sociodemographic characteristics and anthropometric data of infants fed a cow’s milk protein elimination diet and an unrestricted diet (control group).

Variables	Cow’s milk protein elimination diet group (n = 96)	Control group (n = 99)	p
Age ^a	9,5 (5,6 – 14,6)	8,1 (5,3 – 13,0)	0,205
4 - 5,9 months ^b	27 (28,1%)	30 (30,3%)	0,749
6 - 11,9 months ^b	35 (36,5%)	39 (39,4%)	
12 - 18 months ^b	34 (35,4%)	30 (30,3%)	
Male ^b	54 (56,3%)	45 (45,5%)	0,172
Vaginal birth ^b	32 (33,3%)	51 (51,5%)	0,015
Preterm birth ^b	9 (9,6%) ^d	4 (4,0%)	0,213
Birth weight (g) ^c	3188 ± 649	3219 ± 552	0,924
Exclusive breastfeeding for 4 months or more ^b	41 (42,7%)	48 (48,5%)	0,427
Current breastfeeding ^b	26 (27,1%)	60 (60,6%)	< 0,001
Age of first contact with cows’ milk protein (months) ^a	1,0 (0,0 – 3,0)	4,0 (1,0 – 6,0)	< 0,001
Family history of CMPA ^b	5 (10,0%)	0 (0%)	0,023
High risk for atopy ^b	30 (31,3%)	8 (8,1%)	< 0,001
Antibiotic therapy in the first year of life ^b	49 (47,9%)	25 (30,0%) ^e	0,005
Anthropometric indicators (z-scores)			
Weight-for-age ^a	-0,60 (-1,05; +0,09) ^f	+0,24 (-0,43; +0,88) ^g	< 0,001
Height-for-age ^a	-0,12 (-0,75; +0,59) ^f	-0,22 (-0,86; +0,37) ^h	0,680
BMI-for-age ^a	-0,79 (-1,29; -0,20) ^f	+0,34 (-0,20; +1,12) ⁱ	< 0,001
Weight-for-height ^a	-0,85 (-1,18; -0,13) ^f	+0,32 (-0,04; +1,20) ⁱ	< 0,001

Family history of CMPA, presence of one or more first-degree relatives with a past or current history of cows’ milk protein allergy; High risk for atopy, presence of two or more first-degree relatives with atopy; BMI, body mass index.

Available data: n = 94^d, n = 82^e, n = 43^f, n = 84^g, n = 83^h, n = 82ⁱ.

^a Values are expressed as the median and 25th and 75th percentiles in parentheses, Mann-Whitney test.

^b Values are expressed as the absolute number and percentages in parentheses, Pearson chi-squared test or Fisher Exact test,

^c Values are expressed as the mean and standard deviation, Student’s t-test.

D, including oral supplementations. The control group presented a higher percentage of infants with dietary intake below the Recommended Dietary Allowances for vitamin C.

Specific guidance on complementary feeding by a nutritionist or physician was more frequent in the elimination diet group (80.2%; 77/96) than in the control group (53.8%; 49/9; p < 0.001).

Discussion

In the present study, complementary feeding practices of infants on an elimination diet were more closer to the recommendations, although inadequacies were observed in both groups. Infants on an elimination diet had lower weight values according to score Z of weight-for-age, weight-for-height and BMI-for-age but the height was similar to the control group. Breast milk and infant formulas were contributors to more than 50% of the daily energy intake, as well as most nutrients.

In the present study, complementary feeding practices of infants on an elimination diet were more appropriate, although inadequacies were observed in both groups. Infants on an elimination diet had lower weight values, although they were more frequently guided by pediatricians and nutritionists. Breast milk and infant formulas were contributors to more than 50% of the daily energy intake, as well as most nutrients.

As for complementary feeding, liquids, such as water and tea were introduced earlier in the control group, probably

owning to the belief that the child feels thirsty and that tea has soothing properties. However, this assumption is unfounded.^{4,7} The elimination diet group presented a later introduction of leafy vegetables, legumes, eggs, and wheat.

Regarding ultra-processed foods, cookies and soft drinks were introduced less frequently in the elimination diet group, possibly because of the more frequent nutritional guidance of health care professionals and the presence of allergenic fractions of cow’s milk in some of these products. The control group also consumed thickening agents (starches) more frequently. Ultra-processed foods are characterized by high sugar, fructose, sodium, and saturated and trans-fat content, besides the presence of color additives and preservatives. They are highly contraindicated, especially for children under 18 months of age.^{3,4,7}

This study also identified a higher complementary feeding inadequacies index in the control group.¹⁵ When analyzing individually each item that makes up this index, the elimination diet group showed a greater inadequacy only in the item “late introduction of solid foods (at 7 months of age or more)”. This difference was also observed in a study held in the United Kingdom.²² This practice may be associated with the fear of introducing new foods into the diet of infants with CMPA. There are no studies to support the later introduction of foods, since this conduct may even increase the risk of developing food allergies.^{1,4,11,23,24} It is important to emphasize that even those children who had adequate practices up to the date of the assessment may still have inadequate practices before reaching the cut-off age. Moreover, it is alarming to observe that half of the infants

Table 2 Age of introduction of complementary foods and frequency of introduction of foods unsuitable for the age of infants fed a cow's milk protein elimination diet and an unrestricted diet (control group).

Foods	n ^a	Cow's milk protein elimination diet group (n = 96)	n ^a	Control group (n = 99)	p
Age of introduction of complementary foods (months)^b	88	5,0 (4,0-6,0)	91	4,0 (2,0-6,0)	0,001
Water	85	5,5 (3,8-6,0)	88	4,0 (3,0-6,0)	0,021
Tea	45	5,5 (2,8-7,3)	54	3,3 (2,0-6,0)	0,052
Fruit juice	56	6,0 (4,6-6,0)	65	5,5 (4,0-6,0)	0,477
Whole fruit	82	6,0 (5,0-6,0)	76	6,0 (4,6-6,0)	0,410
Vegetables	72	6,0 (6,0-6,5)	71	6,0 (5,0-6,1)	0,111
Tubers and roots	73	6,0 (6,0-6,5)	73	6,0 (5,0-6,0)	0,083
Leafy vegetables	64	6,4 (6,0-8,5)	36	6,0 (6,0-7,0)	0,014
Cereals	63	7,0 (6,0-8,0)	68	6,0 (5,0-8,0)	0,081
Wheat	52	7,0 (6,0-8,0)	62	6,0 (5,0-7,0)	0,022
Legumes	61	7,0 (6,0-8,3)	64	6,0 (6,0-8,0)	0,013
Meat (beef, chicken and/or pork)	69	6,0 (6,0-7,0)	64	6,0 (6,0-7,0)	0,238
Fish	38	8,3 (7,0-11,0)	34	8,0 (6,0-11,3)	0,415
Eggs	52	8,0 (7,0-10,0)	46	7,0 (6,0-8,0)	0,019
Oilseeds	6	12,0 (6,8-13,0)	13	12,0 (8,0-12,0)	0,889
Frequency of introduction of foods unsuitable for their age^c					
Processed meats		8 (8,3%)		12 (12,1%)	0,383
Sugar or sweets		34 (35,4%)		42 (42,4%)	0,316
Soft drink		6 (6,3%)		17 (17,2%)	0,018
Artificial juice		11 (11,5%)		19 (19,2%)	0,135
Industrialized cookies		24 (25,0%)		40 (40,4%)	0,022

^a Number of infants who had introduced the foods at the time of the study.

^b Values are expressed as the median and 25th and 75th percentiles in parentheses, Mann-Whitney test.

^c Values are expressed as the absolute number and percentages in parentheses, Pearson chi-squared test, or Fisher Exact test.

were fed liquefied or sieved soups, a practice proscribed decades ago.

As for nutritional intake, the results differ from those reported in the literature. Past studies showed lower energy and nutrients intake in infants fed a cow's milk elimination diet.^{8,12} Possibly, the wide access to special CMPA formulas, associated with the lower occurrence of complementary feeding inadequacies, contributed to the higher dietary intake. Moreover, the nutritional contribution of special CMPA formulas was greater than that of the milk sources in the control group, possibly owing to the higher volume of formula ingested, reinforcing their importance in the absence of breastfeeding.^{4,7}

Therefore, the mandatory provision of special CMPA formulas by the government is extremely important for promoting nutritional safety, especially for those on an elimination diet, as they are quite vulnerable to nutritional deficiencies. The soy-based infant formulas, which are less expensive and could be consumed by infants over 6 months of age without any gastrointestinal symptoms, were consumed only by 6.7% of the infants. Moreover, the amino acid formula was the most consumed formula in the elimination diet group (52.2%), although it is not recommended as the first-line treatment for infants with CMPA.^{2,3,25} This result is consistent with that of studies conducted in the United Kingdom (45.5%).²² and the Netherlands (51%),²⁶ even though amino

acid formulas are more expensive and generally indicated for infants with more complex or severe symptoms, failure to thrive, or when a reaction to extensively hydrolyzed formulas¹ is presented.

Although both the groups had met most of their nutritional demands, it is important to note that most infants in the elimination diet group were children of well-educated mothers, who were guided by physicians and nutritionists. The mothers of the infants in the elimination diet group were probably better prepared to perform the complementary feeding correctly than the mothers of the infants in the control group.

None of the groups reported the intake of calcium supplementation, a nutrient often documented as deficient in infants fed an elimination diet. Calcium intake was below the recommended daily value in approximately one-quarter of the cases, indicating that its intake may be underestimated in routine infant healthcare assistance.

Despite the higher nutritional intake, the lower weight values observed in infants on an elimination diet were similar to that reported in previous studies.^{8,27-29} It is still unclear whether the lower weight values are related to poor adherence, multiple allergies, or to inflammation status, which can lead to decreased nutrients bioavailability, inappetence, increased energy requirement, or nutrients loss.³⁰

Table 3 Daily intake of calcium, iron, zinc, vitamin A, C, and D of infants fed a cow’s milk protein elimination diet and an unrestricted diet (control group).

Nutrients	Cow’s milk protein elimination diet group (n = 92)	Control group (n = 99)	p
Calcium (mg)			
Total	559,2 (458,3-766,8)	471,5 (259,8-768,5)	0,027
Complementary foods	59,7 (0,2-101,6)	97,7 (0,9-162,3)	0,016
Breast milk / infant formulas / dairy	509,6 (390,9-694,2)	377,2 (217,7-706,6)	0,004
Supplements	-	-	-
Iron (mg)			
Total	12,0 (8,5-17,5)	13,0 (5,9-21,6)	0,906
Complementary foods	2,2 (0,1-3,7)	2,5 (0,1-4,9)	0,285
Breast milk / infant formulas / dairy	8,6 (4,7-6,3)	7,9 (0,3-1,6)	< 0,001
Supplements	11,0 (6,0-10,0)	10,0 (8,0-10,0)	0,618
Zinc (mg)			
Total	8,1 (5,5-10,2)	6,3 (3,5-9,8)	0,003
Complementary foods	2,9 (0,1-4,2)	2,1 (0,0-5,0)	0,766
Breast milk / infant formulas / dairy	5,0 (3,6-6,2)	2,7 (1,9-5,4)	< 0,001
Supplements	4,0 (2,5-4,8)	2,0 (2,0-2,0)	0,500
Vitamin D (mcg)			
Total	22,2 (18,9-28,1)	23,6 (18,3-27,9)	0,831
Complementary foods	0,0 (0,0-0,0)	0,0 (0,0-0,6)	< 0,001
Breast milk / infant formulas / dairy	11,8 (8,6-14,8)	9,7 (6,2-14,7)	0,006
Supplements	12,6 (12,6-12,6)	12,6 (12,6-12,6)	0,256
Vitamin C (mg)			
Total	114,6 (86,1-155,8)	91,9 (33,9-163,8)	0,002
Complementary foods	41,3 (2,9-95,8)	41,8 (1,8-96,7)	0,664
Breast milk / infant formulas / dairy	63,5 (47,6-85,1)	30,2 (17,1-65,4)	< 0,001
Supplements	39,6 (25,5-39,6)	46,0 (12,0-80,0)	0,889
Vitamin A (mg)			
Total	1265,3 (900,3-1616,0)	1659,2 (1309,9-1978,8)	< 0,001
Complementary foods	85,7 (0,1-167,7)	90,9 (0,0-200,1)	0,497
Breast milk / infant formulas / dairy	522,5 (392,2-754,3)	721,9 (433,6-976,6)	0,010
Supplements	750,0 (750,0-750,0)	750,0 (750,0-750,0)	0,902

Values are expressed as the median and 25th and 75th percentiles in parentheses, Mann-Whitney test.

The height of infants on an elimination diet showed a slight deviation to the left (median Z score = -0.12). However, the lower height values observed in previous studies were not evidenced^{8,11}. The better socioeconomic status and access to adequate nutrition, provided by the special CMPA formulas supported by the government, may have contributed to this result. The control group also showed a deviation to the left with respect to height (median score Z = -0.22), reinforcing the absence of differences between the groups.

The limitations of this study were the socioeconomic and demographic differences observed between the groups, and anthropometric measurements were not performed on all the infants by the main researcher, although only anthropometric data collected by fully qualified healthcare professionals were considered. The lower frequency of breastfeeding in the elimination diet group may have been influenced by the recruitment of volunteers at the pharmacy where patients get access to the special formulas. The results of inadequate food intake should be interpreted considering the sample size and the possibility that future studies with greater power would identify differences not evidenced in

the present study. On the other hand, the present study’s results give a suggestion on how to calculate the sample size in future projects.

The strengths of this study were as follows: this was a pioneering study with a considerable sample size. The age range limited to 18 months may have contributed to minimizing memory bias about complementary feeding practices. Food consumption was performed during the four seasons of the year and data collection and coding were performed by a single researcher, decreasing the risk of bias. Another strength of this study was the fact that children from both groups were not being followed up in a single service, which brings the results closer to real life.

This study reinforces the importance of actions to encourage the maintenance of breastfeeding, the provision of infant formulas by the government, and increased educational actions on complementary feeding to prevent nutritional deficiencies in early childhood.

In conclusion, infants on a cow’s milk elimination diet presented more adequate complementary feeding practices and higher nutritional intake, despite lower body weight values.

Table 4 Infants with food intake below the Recommended Daily Intakes (RDA/AI)¹⁷ for micronutrients on a cow's milk protein elimination diet and control groups.

Nutrients		Cow's milk protein elimination diet group (n = 92)	Control group (n = 99)	p
Feed only (including milk sources)				
Calcium		29 (31,5%)	22 (22,2%)	0,198
Iron		24 (26,1%)	35 (35,4%)	0,219
Zinc		2 (2,2%)	7 (7,1%)	0,171
Vitamin A		5 (5,4%)	9 (9,1%)	0,333
Vitamin C		3 (3,1%)	25 (25,3%)	< 0,001
Vitamin D		41 (44,6%)	42 (42,4%)	0,879
Feed (including milk sources)± supplementation				
	n ^a		n ^a	
Calcium	0	29 (31,5%)	0	0,198
Iron	45	13 (14,1%)	59	0,996
Zinc	3	2 (2,2%)	1	0,171
Vitamin A	70	1 (1,1%)	87	1,000
Vitamin C	8	3 (3,1%)	2	< 0,001
Vitamin D	78	9 (9,8%)	87	0,941

Values are expressed as the absolute number and percentages in parentheses, Pearson chi-squared test, or Fisher Exact test.

RDA/AI (Recommended Dietary Allowances and Adequate Intakes)¹⁷: values corresponding to the recommendations for 0-6 months/ 7-12 months/ 1-3 years: Calcium 200/260/700 mg/day; Iron 0,27/11/7 mg/ day; Zinc 2/3/3 mg/ day; Vitamin A 400/500/300 mcg/ day; Vitamin C 40/50/15 mg/ day; Vitamin D 10/10/15 mcg/day.

^a Number of infants receiving mineral and vitamin supplementation.

Funding

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

Conflicts of Interest

The authors declare no conflicts of interest.

Supplementary materials

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.jpmed.2021.06.005>.

References

- Koletzko S, Niggemann B, Arato A, Dias JA, Heuschkel R, Husby S, et al. Diagnostic approach and management of cow's-milk protein allergy in infants and children: ESPGHAN GI Committee Practical Guidelines. *J Pediatr Gastroenterol Nutr.* 2012;55:221–9.
- Fiocchi A, Brozek J, Schünemann H, Bahna SL, Berg AV, Beyer K, et al. World allergy organization (WAO) diagnosis and rationale for action against cow's milk allergy (DRACMA) guidelines. *Pediatr Allergy Immunol.* 2010;21:1–125.
- Fewtrell M, Bronsky J, Campoy C, Domellof M, Embleton N, Mis NF, et al. Complementary feeding: a position paper by the European society for paediatric gastroenterology, hepatology, and nutrition (ESPGHAN) committee on nutrition. *J Pediatr Gastroenterol Nutr.* 2017;64:119–32.
- Sociedade Brasileira de Pediatria- Departamento de Nutrologia. Manual de alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar. Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. 4 ed. São Paulo (SP).
- Ribeiro CC, Leite G, Morais MB. Knowledge and practice of physicians and nutritionists regarding the prevention of food allergy. *Clin Nutr.* 2013;32:624–9.
- Vandenplas Y, AlFrayh AS, AlMutairi B, Elhalik MS, Green RJ, Haddad J, et al. Physician practice in food allergy prevention in the Middle East and North Africa. *BMC Pediatr.* 2017;17:118.
- Ministério da Saúde. Brasil. Secretaria de Atenção Primária à Saúde. Departamento de Promoção da Saúde. Guia alimentar para crianças menores de 2 anos/ministério da saúde, secretaria de atenção primária à saúde. Departamento de Promoção da Saúde – Brasília: Ministério da Saúde; 2019, 265 p.: il.
- Medeiros LC, Speridião PG, Sdepanian VL, Neto UF, Morais MB. Nutrient intake and nutritional status of children following a diet free from cow's milk and cow's milk by-products. *J Pediatr (Rio J).* 2004;80:363–70.
- Vieira MC, Morais MB, Spolidoro JV, Toporovski MS, Cardoso AL, Araujo GT, et al. A survey on clinical presentation and nutritional status of infants with suspected cow' milk allergy. *BMC Pediatr.* 2010;10:25.
- Foong RX, Meyer R, Dziubak R, Lozinsky AC, Godwin H, Reeve K, et al. Establishing the prevalence of low vitamin D in non-immunoglobulin-E mediated gastrointestinal food allergic children in a tertiary centre. *World Allergy Organ J.* 2017;10:4.
- Boaventura RM, Mendonça RB, Fonseca FA, Mallozi M, Souza FS, Sarni RO. Nutritional status and food intake of children with

- cow's milk allergy. *Allergol Immunopathol (Madr)*. 2019;47:544–50.
12. Meyer R. Nutritional disorders resulting from food allergy in children. *Pediatr Allergy Immunol*. 2018;29:689–704.
 13. Mendonça RB, Solé D, DunnGalvin A, Len CA, Sarni RO. Evaluation of the measurement properties of the Brazilian version of two quality-of-life questionnaires in food allergy – for children and their parents. *J Pediatr (Rio J)*. 2020;96:600–6.
 14. ABEP - Associação Brasileira das Empresas de Pesquisa. Brazilian Criteria 2015 and social class distribution update for 2016 [Cited 2021 May]. Available from: <http://www.abep.org/Servicos/Download.aspx?id=13>
 15. Mais LA, Domene SM, Barbosa MB, Taddei JA. Diagnosis of complementary feeding practices for creating a matrix model for action in primary health care. *Cien Saude Colet*. 2014;19:93–104.
 16. Drewett RF, Woolridge MW, Jackson DA, Imong SM, Mangklabruks A, Wongsawadit L, et al. Relationships between nursing patterns, supplementary food intake and breast-milk intake in a rural Thai population. *Early Hum Dev*. 1989;20:13–23.
 17. Nejar FF, Segall-Corrêa AM, Rea MF, Vianna RP, Panigassi G. Breastfeeding patterns and energy adequacy. *Cad Saude Publica*. 2004;20:64–71.
 18. Institute of medicine. Dietary reference intakes (DRIs): Recommended dietary allowances and adequate intakes, vitamins life-stage vitamin A vitamin C vitamin D vitamin E vitamin K thiamin riboflavin niacin vitamin B 6 folate vitamin B 12. [Cited Mar 2021]. Available from: https://www.nal.usda.gov/sites/default/files/fnic_uploads/recommended_intakes_individuals.pdf.
 19. Karoline P, Cordeiro DA, Castro JL, De Lima SH, Alves CX, Leite LD. Comparação de três programas computacionais utilizados na avaliação de recordatórios alimentares 24 horas. *J Heal Informatics*. 2011;3:13–8.
 20. World Health Organization. WHO Anthro for personal computers, version 3.2.2, 2011: Software for assessing growth and development of the world's children.
 21. Ministério da Saúde. Orientações para a coleta e análise de dados antropométricos em serviços de saúde: norma técnica do Sistema de Vigilância Alimentar e Nutricional. Brasília. 2011.
 22. Maslin K, Dean T, Arshad SH, Venter C. Fussy eating and feeding difficulties in infants and toddlers consuming a cows' milk exclusion diet. *Pediatr Allergy Immunol*. 2015;26:503–8.
 23. Carletti C, Pani P, Monasta L, Knowles A, Cattaneo A. Introduction of complementary foods in a cohort of infants in northeast Italy: do parents comply with WHO recommendations? *Nutrients*. 2017;9:1–11.
 24. Dupont C, Chouraqui JP, Linglart A, Bocquet A, Darmaun D, Feillet F, et al. Nutritional management of cow's milk allergy in children: An update. *Arch Pediatr*. 2018;25:236–43.
 25. Solé D, Silva LR, Cocco RR, Ferreira CT, Sarni RO, Oliveira LC, et al. Brazilian Consensus on Food Allergy: 2018 - Part 2 - diagnosis, treatment and prevention. Joint position paper of the Brazilian society of pediatrics and the Brazilian association of allergy and immunology imunologia. *Arq Asma Alerg Imunol*. 2018;2:39–82.
 26. Petrus NC, Schoemaker AA, Hoek MW, Sprikkelman AB. Remaining symptoms in half the children treated for milk allergy. *Eur J Pediatr*. 2015;174:759–65.
 27. Meyer R, De Koker C, Dziubak R, Venter C, Dominguez-Ortega G, Cutts R, et al. Malnutrition in children with food allergies in the UK. *J Hum Nutr Diet*. 2014;27:227–35.
 28. Meyer R, De Koker C, Dziubak R, Godwin H, Dominguez-Ortega G, Lozinsky AC, et al. The impact of the elimination diet on growth and nutrient intake in children with food protein induced gastrointestinal allergies. *Clin Transl Allergy*. 2016;6:25.
 29. Meyer R, Wright K, Vieira MC, Chong KW, Chatchatee P, Vlieg-Boerstra BJ, et al. International survey on growth indices and impacting factors in children with food allergies. *J Hum Nutr Diet*. 2019;32:175–84.
 30. Venter C, Brown T, Meyer R, Walsh J, Shah N, Nowak-Wegrzyn A, et al. Better recognition, diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: IMAP - an international interpretation of the MAP (Milk Allergy in Primary Care) guideline. *Clin Transl Allergy*. 2017;7:1–9.