


# BMJ Open Associations between initial American Board of Internal Medicine certification and maintenance of certification status of attending physicians and in-hospital mortality of patients with acute myocardial infarction or congestive heart failure: a retrospective cohort study of hospitalisations in Pennsylvania, USA

John J Norcini ,<sup>1</sup> Weifeng Weng,<sup>2</sup> John Boulet,<sup>1</sup> Furman McDonald,<sup>2</sup> Rebecca S Lipner<sup>2</sup>

**To cite:** Norcini JJ, Weng W, Boulet J, *et al.* Associations between initial American Board of Internal Medicine certification and maintenance of certification status of attending physicians and in-hospital mortality of patients with acute myocardial infarction or congestive heart failure: a retrospective cohort study of hospitalisations in Pennsylvania, USA. *BMJ Open* 2022;**12**:e055558. doi:10.1136/bmjopen-2021-055558

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055558>).

Received 17 July 2021  
Accepted 23 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr John J Norcini;  
[john.norcini@gmail.com](mailto:john.norcini@gmail.com)

## ABSTRACT

**Objective** To determine whether internists' initial specialty certification and the maintenance of that certification (MOC) is associated with lower in-hospital mortality for their patients with acute myocardial infarction (AMI) or congestive heart failure (CHF).

**Design** Retrospective cohort study of hospitalisations in Pennsylvania, USA, from 2012 to 2017.

**Setting** All hospitals in Pennsylvania.

**Participants** All 184 115 hospitalisations for primary diagnoses of AMI or CHF where the attending physician was a self-designated internist.

**Primary outcome measure** In-hospital mortality.

**Results** Of the 2575 physicians, 2238 had initial certification and 820 were eligible for MOC. After controlling for patient demographics and clinical characteristics, hospital-level factors and physicians' demographic and medical school characteristics, both initial certification and MOC were associated with lower mortality. The adjusted OR for initial certification was 0.835 (95% CI 0.756 to 0.922;  $p < 0.001$ ). Patients cared for by physicians with initial certification had a 15.87% decrease in mortality compared with those cared for by non-certified physicians (mortality rate difference of 5.09 per 1000 patients; 95% CI 2.12 to 8.05;  $p < 0.001$ ). The adjusted OR for MOC was 0.804 (95% CI 0.697 to 0.926;  $p = 0.003$ ). Patients cared for by physicians who completed MOC had an 18.91% decrease in mortality compared with those cared for by MOC lapsed physicians (mortality rate difference of 6.22 per 1000 patients; 95% CI 2.0 to 10.4;  $p = 0.004$ ).

**Conclusions** Initial certification was associated with lower mortality for AMI or CHF. Moreover, for patients whose physicians had initial certification, an additional advantage was associated with its maintenance.

## Strengths and limitations of this study

- The study is based on a sizeable number of patients and physicians and extensive sensitivity analyses were conducted.
- This is a retrospective observational study and cannot establish causality.
- Unmeasured institutional effects may have influenced the results, but a hospital fixed effect analysis significantly reduced this possibility.
- Despite our inclusion of a standard set of covariates for both patients and physicians, there might be other unobserved factors that were potentially confounding.
- Our work is based on only two primary diagnoses in one state in the USA so broader sampling is needed to increase the generalisability of the results.

## INTRODUCTION

Initial certification in internal medicine has been offered by the American Board of Internal Medicine (ABIM) since its establishment in 1936. These certificates were of unlimited duration until 1990, when ABIM restricted their validity to 10 years and introduced a recertification or maintenance of certification (MOC) programme.<sup>1</sup> Board certification in internal medicine seeks to recognise physicians who have demonstrated that they have the knowledge, skills and attitudes to provide excellent patient care.<sup>2</sup> Given this mission, it is reasonable to expect

an association between initial board certification, MOC and patient outcomes. There is some evidence of this kind for initial certification but less for MOC. This retrospective observational study seeks to test whether such relationships exist.

The requirements for initial certification in internal medicine have changed over time, but they have been broadly similar since the oral examination was discontinued in 1970. Candidates must have graduated from a US medical school or been certified by the Educational Commission for Foreign Medical Graduates (ECFMG), satisfactorily completed accredited training, have an unrestricted licence to practise medicine and perform successfully on the certifying examination. There has been more variability in the MOC programme over time, but the programme has basically required initial certification, an unrestricted licence to practise medicine, some form of educational experience and successful performance on an assessment.

There is considerable evidence that initial board certification and educational markers of performance are associated.<sup>3–5</sup> While this form of evidence is important for building an argument for the validity of the certificate, it would fall short of the appropriate expectations of patients and physicians if certification were not also associated with the quality of care provided in practice. In this regard, there is reasonable evidence of a relationship with clinical care. For instance, there are several studies showing that certification is positively associated with the process of care provided to patients and their outcomes, while being negatively associated with disciplinary actions by state licensing boards.<sup>6–10</sup> One aim of this study is to gather additional validity evidence based on the association between initial certification and patient outcomes.

Although it is a newer process, there is evidence of the relationship between MOC and educational markers of performance as well.<sup>11</sup> Nevertheless MOC is a controversial programme, and evidence of a relationship with clinical care is essential.<sup>12–14</sup> On this front, studies have demonstrated a positive association of MOC with process of care measures<sup>15 16</sup> and negative associations with diagnostic errors,<sup>17</sup> healthcare cost<sup>18</sup> and state licensing board disciplinary actions.<sup>19–23</sup> However, there is little evidence speaking to the relationship between MOC and patient outcomes. Consequently, the second aim of this study is to determine whether such an association exists for those physicians who already have initial certification and are choosing to maintain it by taking ongoing assessments over the course of their careers.

To achieve our two aims, we analysed all the hospitalisations for acute myocardial infarction (AMI) or congestive heart failure (CHF) in the state of Pennsylvania for a 6-year period starting in 2012. We limited our focus to the patients of self-identified internists. Included in our study was the initial certification status of the attending physicians, whether they were eligible for MOC, and whether they maintained this status. The outcome measure was in-hospital mortality and we controlled for characteristics

of patients and hospitals as well as other physician characteristics.

## METHODS

The Strengthening the Reporting of Observational Studies in Epidemiology reporting guidelines were followed for the study.<sup>24</sup>

### Sources of the data

Data for this study were drawn from the records of the Pennsylvania Health Care Cost Containment Council (PHC4), the American Medical Association (AMA) Physician Masterfile, the ABIM and the ECFMG.

In Pennsylvania, all hospitals (both public and private) are required to send data to PHC4 each time a patient is discharged. Included are patient demographics, primary and secondary diagnoses, discharge status, hospital and the attending physician. The attending physician is identified by the hospital as the individual who has overall responsibility for the medical care and treatment of the patient. Validation and editing processes are applied to the data by PHC4 and hospitals can correct the information. In this study, hospitalisations with a principal diagnosis of AMI or CHF from 1 January 2012 to 31 December 2017 were available for analysis. AMI and CHF were chosen because they occur in sufficient volume and are often used to judge quality of care.<sup>25–27</sup>


The AMA Physician Masterfile contains information on practising physicians in the USA. For this study, we limited the analysis to self-identified general internists who were the attending physicians for patients with AMI and/or CHF in the PHC4 database (we explored the inclusion of self-designated hospitalists as well but not all are internists, and the group was too small (n=60) for analysis). The initial certification status of the self-designated internists, their eligibility for MOC, their initial subspecialty certifications, as well as whether they retained their certification, if eligible, were available from the files of the ABIM. In addition, an indication of whether the physician was an international medical school graduate (IMG) was available from the ECFMG.

### Data elements

From our data sources, we extracted patient age, sex, race, ethnicity, principal and secondary diagnoses, year of admission, discharge status and hospital. For the attending physicians, we knew whether they were US medical graduates (USMGs) or international medical school graduates (IMGs), whether they had initial certification in internal medicine, their sex, age, and year of initial certification, the year they were due for MOC and whether they maintained their certification.

From the data for patients, we calculated a modified version of the Charlson Comorbidity Index.<sup>28</sup> This offered a measure of the severity of illness of the patients in the study. We also combined data for race and ethnicity to a single variable, where white, non-Hispanic patients were

Figure 1. Flow Diagram of Identifying Physician Sample



**Figure 1** Flow diagram of the physicians in the study. AMI, acute myocardial infarction; CHF, congestive heart failure; MOC, maintenance of certification.

in one group and all others were in another. We did this because some of the subgroups were too small for reasonable analysis. Finally, we ascertained whether the patient died while in the hospital based on their discharge status.

For the hospitals in the study, we developed two measures. The first was an indicator of whether the hospital was in a rural county by reference to a list maintained by the Pennsylvania Office of Rural Health.<sup>29</sup> The second, an indicator of institutional experience, was a count of the number of patients with AMI or CHF treated by study physicians in the hospital.

### Physicians

Figure 1 presents a flow diagram that identifies the physicians included in the study. From the group of all self-identified internists who were attending physicians to patients in the PHC4 data, we eliminated those who were certified in internal medicine but had a subspecialty certificate as well (n=431) and those who did not

care for patients with AMI or CHF (n=274). To determine if initial certification was associated with better patient outcomes, we compared the in-hospital mortality of the patients whose attending physician had initial certification (n=2238) with those who did not (n=428).

To address the question of whether MOC was associated with better patient outcomes, we eliminated from the 2238 physicians with initial certification those who were certified prior to 1990 or were within 10 years of initial certification at the end of the study (ie, by 31 December 2017). The remaining 820 physicians were MOC eligible for some or all the time between 1 January 2012 and 31 December 2017.

During the period of study, a physician's status may have changed from year to year for a variety of reasons. For example, some physicians may have allowed their certification to lapse but restored it at a later point in time, while others may have been within 10 years of

**Table 1** Characteristics of the patients and hospitals by attending physician certification status

	Physicians		All hospitalisations	
	Never certified	Initial certification		
Patient age: n (%)				
≤49	1200 (5)	8391 (5.2)	9591 (5.2)	
50–64	4368 (18.3)	30299 (18.9)	34667 (18.8)	
65–74	4912 (20.6)	33209 (20.7)	38121 (20.7)	
75–84	6499 (27.2)	43184 (27)	49683 (27.0)	
≥85	6900 (28.9)	45153 (28.2)	52053 (28.3)	
Total	23879 (13)	160236 (87)	184115 (100)	P<0.05
Patient sex: n (%)				
Female	11998 (50.2)	78861 (49.2)	90859 (49.4)	Missing=3
Male	11881 (49.8)	81372 (50.8)	93253 (50.6)	P<0.005
Patient race/ethnicity: n (%)				
White, non-Hispanic	4083 (17.1)	27913 (17.4)	31996 (17.4)	Missing=3
Other	19796 (82.9)	132323 (82.6)	152119 (82.6)	P=0.22
Patient condition: n (%)				
Acute myocardial infarction	6575 (27.5)	41669 (26)	48224 (26.2)	
Congestive heart failure	17304 (72.5)	118567 (74)	135871 (73.8)	P<0.0001
Patient comorbid conditions: mean (SD)	1.012 (1.112)	0.953 (1.117)	0.960 (1.117)	P<0.0001
Patient deaths: n (% hospitalisations for that condition)				
Acute myocardial infarction	315 (4.8)	1553 (3.7)	1868 (3.9)	P<0.0001
Congestive heart failure	494 (2.9)	2664 (2.3)	3158 (2.3)	P<0.0001
Total	809 (3.4)	4217 (2.6)	5026 (2.7)	P<0.0001
Facility location: n (%)				
Non-rural	17170 (71.9)	122850 (76.7)	140020 (76)	
Rural	6709 (28.1)	37386 (23.3)	37386 (24)	P<0.0001
Facility volume/1000: mean (SD)	4.41 (3.08)	5.89 (3.55)	5.70 (3.52)	P<0.0001

initial certification for part of the study (eg, they acquired initial certification in 2005 and were not eligible for MOC until 2015). Consequently, we determined the status of the physician in the year they treated the patient, and in figure 1 we present physician-years (ie, the number of years physicians were eligible for MOC) as well as the number of physicians.

### Analysis

Descriptive statistics were calculated for the hospitalisations and institutions, both for the total patient population and for the subpopulation treated by the MOC group. To determine the adjusted relationship between initial certification and patient mortality, we applied a multivariable logistic regression model which was adjusted for (1) the comorbidity index, (2) the condition (with CHF as the reference), (3) patient age, (4) patient sex (with female as the reference), (5) patient race/ethnicity (with white/non-Hispanic as the reference), (6) whether the physician was an IMG (with USMG as the reference), (7) physician sex (with male as the reference), (8) physician age, (9) hospital location (with non-rural as the reference),

and (10) the number of hospitalisations in the institution for AMI or CHF. We applied a similar model to determine the adjusted relationship between MOC and patient mortality. Generalised estimating equations (Stata V.14, StataCorp) were applied to adjust for the clustering of patients within physicians and physicians within hospitals. We used the margins function to simulate the percentage point change in mortality if patients who were cared for by never-certified internists had instead been cared for by ever-certified internists. We did the same for patients who were cared for by internists who did not maintain their certification versus those who had been cared for by internists who maintained their certification.

### Sensitivity analyses

Although we included hospital volume and location in our analyses, it is possible that additional unmeasured institutional characteristics, such as on-site percutaneous coronary intervention or coronary artery bypass graft surgery, might influence the findings. To rule out these and other unknown potential institutional confounds,

**Table 2** Estimated adjusted ORs with 95% CI for in-hospital mortality for patients of all physicians

Parameter	OR (95% CI)	P value
<b>Patient characteristics</b>		
Acute myocardial infarction	2.071 (1.937 to 2.214)	<0.001
Comorbidity index	1.119 (1.091 to 1.149)	<0.001
Sex (female)	1.113 (1.051 to 1.179)	<0.001
Age	1.046 (1.039 to 1.054)	<0.001
White, non-Hispanic	1.028 (0.573 to 1.844)	0.93
Race/ethnicity × age	1.004 (0.996 to 1.011)	0.34
<b>Physician characteristics</b>		
Ever certified	0.835 (0.756 to 0.922)	<0.001
Sex		
Female	0.923 (0.847 to 1.006)	0.07
Missing	0.644 (0.411 to 1.009)	0.06
Age	1.005 (1.002 to 1.009)	<0.01
USMG	0.974 (0.906 to 1.048)	0.48
<b>Facility</b>		
Volume/1000	0.997 (0.987 to 1.008)	0.64
Rural location	1.228 (1.128 to 1.337)	<0.001

USMG, US medical graduate.

we also ran a hospital fixed effect regression model to control the effect of all hospital-level factors.

Likewise, in our models we controlled for condition (AMI vs CHF) but it is possible that the findings could be driven primarily by one or the other. Consequently, we conducted separate analyses for AMI and CHF.

In preliminary analyses we found that there was a significant difference in age between the white non-Hispanic patients (mean=76.3, SD=13.0) and other patients (mean=65.8, SD=14.6). Consequently, we included the interaction of age and race/ethnicity in our multivariate models. Other interactions were tested but not included since they did not alter the interpretation of the results. Likewise, we tested whether a non-linear version of age might be more appropriate, but again it did not make an appreciable difference.

### Patient and public involvement

No patients were involved.

## RESULTS

### Initial certification

There were 184155 hospitalisations for AMI or CHF in the study with an average of 71.5 per physician. The attending physicians for 160236 hospitalisations had initial certification. The mean age of the physicians for these hospitalisations was 44.5 (SD=10.0) and 27.0% of

them were managed by women (sex was missing for 771 (0.5%) hospitalisations). The attending physicians for the remaining 23879 hospitalisations did not have initial certification, the mean age of the physicians for these hospitalisations was 52.8 (SD=12.3) and 21.1% of them were managed by women (sex was missing for 235 (1%) hospitalisations).

Table 1 presents the characteristics of the hospitalisations and facilities broken down by whether the attending physician achieved initial certification or was never certified. The patients of the doctors who achieved initial certification were slightly younger and slightly less likely to be female (49.2% vs 50.2%) than the patients of the doctors who were never certified. The difference between groups in terms of race/ethnicity was not statistically significant. As a group, a slightly lower proportion of the patients of the certified doctors had a primary diagnosis of AMI (26% vs 27.5%) and they had slightly fewer comorbid conditions (1.012 vs 0.953). The raw mortalities of the patients of certified doctors were lower (2.6% vs 3.4%).

Most hospitalisations were in facilities located in non-rural settings (76%). However, a smaller proportion of the hospitalisations of the initially certified group were in rural counties (23.3% vs 28.1%). In addition, the hospitalisations of the initially certified group were in hospitals with higher volume (5.89 vs 4.41).

Table 2 shows the results of the multivariable generalized estimating equations (GEE) analysis which incorporates characteristics of the patients, physicians and facilities. After adjustment, initial certification was associated with lower mortality. The adjusted OR was 0.835 (95% CI 0.756 to 0.922;  $p<0.001$ ). Patients cared for by physicians with initial certification had a 15.87% decrease in mortality compared with those cared for by non-certified physicians (mortality rate difference of 5.09 per 1000 patients; 95% CI 2.12 to 8.05;  $p<0.001$ ).

In the hospital fixed effect sensitivity analysis which incorporates the same patient and physician characteristics and controls for all hospital-level factors, the adjusted OR for initial certification was 0.826 (95% CI 0.754 to 0.905;  $p<0.001$ ) (online supplemental table 1). In the sensitivity analysis which models patients with AMI and CHF separately, the adjusted OR for initial certification was 0.827 (95% CI 0.715 to 0.957;  $p=0.01$ ) for the AMI model and 0.842 (95% CI 0.747 to 0.95;  $p<0.01$ ) for CHF model (online supplemental table 2). Whether using GEE or hospital fixed effect or fitting patients with AMI and CHF in one combined model or two separate models, all analyses indicate that after adjustment, initial certification was associated with lower mortality and the adjusted ORs were similar across different models.

### Maintenance of certification

The attending physicians for 40468 hospitalisations maintained their certification. The mean age of the physicians for these hospitalisations was 49.8 (SD=6.2), and 22.5% of them were managed by women (sex was missing for 284 hospitalisations). The attending physicians for the

**Table 3** Characteristics of the patients by attending physician certification status for those MOC eligible

	Physicians		All hospitalisations	
	Lapsed certification	Certified		
Patient age: n (%)				
≤49	506 (3.6)	1810 (4.5)	2316 (4.3)	
50–64	2116 (15.1)	7081 (17.5)	9197 (16.9)	
65–74	2800 (20.0)	8214 (20.3)	11 014 (20.2)	
75–84	4040 (28.8)	11 105 (27.6)	15 145 (27.8)	
≥85	4534 (32.4)	12 258 (30.3)	16 792 (30.8)	
Total	13 996 (25.7)	40 468 (74.3)	54 464 (100%)	P<0.0001
Patient sex: n (%)				
Female	7159 (51.2)	20 113 (49.7)	27 272 (50.1)	Missing=1
Male	6837 (48.9)	20 354 (50.3)	27 191 (49.9)	P=0.003
Patient race/ethnicity: n (%)				
White, non-Hispanic	12 125 (86.6)	33 964 (83.9)	46 089 (84.6)	Missing=1
Other	1871 (13.4)	6404 (16.1)	8375 (15.4)	P<0.0001
Patient condition: n (%)				
Acute myocardial infarction	3455 (24.7)	10 112 (25.0)	13 567 (24.9)	
Congestive heart failure	10 541 (75.3)	30 356 (75.0)	40 897 (75.1)	P=0.48
Comorbid conditions: mean (SD)	0.9303 (1.0986)	0.9658 (1.1177)	0.9566 (1.1122)	P<0.0001
In-hospital deaths: n (% hospitalisations for that condition)				
Acute myocardial infarction	171 (5.0)	380 (3.8)	551 (4.1)	P<0.003
Congestive heart failure	299 (2.8)	666 (2.2)	965 (2.4)	P<0.0003
Total	470 (3.4)	1046 (2.6)	1516 (2.8)	P<0.0001
Facility location: n (%)				
Non-rural	10 710 (76.5)	31 244 (77.2)	41 954 (77.0)	
Rural	3286 (23.5)	9224 (22.8)	12 510 (23.0)	P=0.10
Facility volume/1000: mean (SD)	5.11 (3.34)	5.82 (3.58)	5.64 (3.54)	P<0.0001

MOC, maintenance of certification.

remaining 13 996 hospitalisations were eligible for MOC but did not maintain it, the mean age of the physicians for these hospitalisations was 52.6 (SD=6.3) and 23.0% of them were managed by women (no missing data).

Table 3 presents the characteristics of the hospitalisations and facilities for attending physicians who had achieved initial certification and were eligible for MOC, broken into two groups depending on whether the physician had maintained his/her certification in the year the patient was treated.

The patients of the doctors who maintained their certification were slightly younger and more often male (50.3% vs 48.9%) than the patients of the doctors who did not. Moreover, a greater proportion of non-white/Hispanic patients were cared for by physicians who maintained their certification (16.1% vs 13.4%). There was not a statistically significant difference between groups of patients in terms of their primary diagnosis and the patients of doctors who maintained their certification had slightly more comorbid conditions. The raw mortalities of

the patients of doctors who maintained their certification were lower (2.6% vs 3.4%).

Again, most hospitalisations were in facilities located in non-rural settings (77%). There was not a statistically significant difference in hospital location between the patients of the doctors who maintained their certification and those who did not. The hospitalisations of doctors who maintained certification were in facilities with higher volume (5.82 vs 5.11).

Table 4 shows the results of the multivariable GEE analysis which incorporates characteristics of the patients, physicians and facilities. After adjustment, MOC was associated with lower mortality. The adjusted OR was 0.804 (95% CI 0.697 to 0.926; p=0.003). Patients cared for by physicians who completed MOC had an 18.91% decrease in mortality compared with those cared for by MOC lapsed physicians (mortality rate difference of 6.22 per 1000 patients; 95% CI 2.0 to 10.4; p=0.004).

In the hospital fixed effect sensitivity analysis which incorporates the same patient and physician characteristics

**Table 4** Estimated adjusted ORs with 95% CI for in-hospital mortality for patients of physicians eligible for MOC

Parameter	OR (95% CI)	P value
<b>Patient characteristics</b>		
Acute myocardial infarction	2.077 (1.839 to 2.347)	<0.001
Comorbidity index	1.159 (1.107 to 1.215)	<0.001
Sex (female)	1.07 (0.959 to 1.194)	0.23
Age	1.054 (1.04 to 1.068)	<0.001
White, non-Hispanic	2.61 (0.901 to 7.561)	0.08
Race/ethnicity × age	0.991 (0.977 to 1.004)	0.18
<b>Physician characteristics</b>		
MOC	0.804 (0.697 to 0.926)	<0.01
Sex		
Female	0.919 (0.775 to 1.089)	0.33
Missing	0.718 (0.26 to 1.983)	0.52
Age	0.998 (0.987 to 1.009)	0.72
USMG	0.978 (0.853 to 1.123)	0.76
<b>Facility</b>		
Volume/1000	0.997 (0.978 to 1.016)	0.77
Rural location	1.228 (1.044 to 1.444)	0.01

MOC, maintenance of certification; USMG, US medical graduate.

and controls for all hospital-level factors, the adjusted OR for MOC was 0.794 (95% CI 0.695 to 0.908;  $p < 0.01$ ) (online supplemental table 3). In the sensitivity analysis which models patients with AMI and CHF separately, the adjusted OR for MOC was 0.853 (95% CI 0.678 to 1.074;  $p = 0.18$ ) for the AMI model and 0.777 (95% CI 0.66 to 0.915;  $p < 0.01$ ) for the CHF model (online supplemental table 4). Both the main GEE analysis and sensitivity analyses indicate that after adjustment, MOC was associated with lower mortality and the adjusted ORs were similar across different models, except for AMI model. Due to the small sample size, the adjusted OR, even though it was of similar magnitude, was not statistically significant.

## DISCUSSION

The first aim of this study was to ascertain whether there was an association between the initial certification status of attending physicians and the in-hospital mortality of their patients with AMI or CHF. After adjustment for physician, patient and hospital characteristics, initial certification was associated with a 15.87% decrease in in-hospital mortality. These findings are consistent with others reported in the literature.<sup>30 31</sup>

For our analysis of MOC, we focused only on physicians who had initial certification and eliminated those who were certified prior to 1990 or within 10 years of completing training at the end of the study. After

adjustment, the MOC status of these physicians was associated with an 18.91% decrease in in-hospital mortality. This adds patient outcomes to the literature on the association of MOC with disciplinary actions and process of care measures.<sup>15–23</sup>

Initial certification assesses the medical knowledge and judgement needed for independent practice and MOC assesses whether physicians remain current throughout their careers. Thus, we propose that the mechanism for the associations we found is that physicians with greater medical knowledge and better medical judgement are, as a group, providing better care which results in improved outcomes, including decreased mortality. Our findings are concordant with other studies associating increased medical knowledge with better care as demonstrated by process measures, costs, state medical board disciplinary actions and specifically diagnostic knowledge as measured by MOC with emergency department visits, hospitalisations and death.<sup>8–10 15–18 21 32</sup>

The magnitude of the associations we found is clinically significant. In fact, the relative decrease in mortality associated with certification and MOC is similar in magnitude to those observed for the use of aspirin in AMI (23%) and for the decrease in all-cause mortality when aspirin is used for the secondary prevention of cardiovascular and cerebrovascular events (18%).<sup>33 34</sup> Ours is an observational study and we are not claiming causality based on our results alone; however, given our hypothesised mechanism, it is plausible that physicians who are staying current in medical knowledge and practice may be more likely to practise in accordance with the best available evidence thus mediating better outcomes.

We found that increased physician age was associated with increased patient mortality. Lending credence to our model is that our finding is concordant with many studies that have demonstrated negative associations of age with educational and patient-relevant outcomes.<sup>3 35 36</sup> Based on our proposed mechanism, it is plausible that increased age, and in particular time from completion of medical training, may make staying current in knowledge and practice more challenging.<sup>37 38</sup> Regrettably, age is not modifiable, but medical knowledge is. Our study showed that, controlling for age, those physicians who stayed current as evidenced by MOC had better patient outcomes.<sup>39</sup>

Of course, our work has the limitations associated with retrospective observational studies. By definition, such studies alone are unable to establish causality. However, for prediction or prognosis investigations, cohort studies are appropriate, and the hospital fixed effect analysis significantly reduced the possibility of unmeasured institutional differences. Despite our inclusion of a standard set of covariates for both patients (age, race, sex and primary and comorbid conditions) and physicians (age, sex, IMG status), there might be other unobserved factors that were potentially confounding. For instance, we did not have access to clinical data for the studied conditions (eg, left ventricular ejection fraction or B-type natriuretic

peptide concentrations). However, we have no reason to believe these would vary consistently by physicians' certification status and the measure we used incorporates a broad range of potential morbidities specific to AMI and CHF.

Finally, our work is based on two primary diagnoses in one state in the USA and we could not include cardiologists because there were too few for appropriate analysis. However, Pennsylvania is the fifth most populous state in the USA with a population exceeding 13 million spread across a large, geographically diverse area with both urban and rural communities. Notably, our findings are concordant with studies associating lower mortality in hospitals with more board-certified cardiologists in Japan where board certification also requires a periodic test of medical knowledge consistent with the mechanism we have proposed.<sup>40</sup> Nonetheless, broader sampling across US states and other countries with other disciplines and other diagnoses would increase confidence in the generalisability of our results.

Our study raises questions which could serve as the focus of future work as additional data become available. For instance, we had to exclude cardiologists and physicians certified prior to 1990 from our MOC analyses; yet it is important to know if initial certification without continued maintenance has similar associations with care outcomes among these physicians. Our study is based on hospital care and there are many conditions that are primarily managed in the outpatient setting now that may lend themselves to similar analyses if data sets are available to assess them. Finally, there are several studies demonstrating an association with patient outcomes not just based on whether a physician passed the initial certification or MOC examination, but on how well they performed on the examination (ie, score).<sup>17 21 32 41 42</sup> If a similar association (ie, a 'dose effect') existed with decreased mortality in a population like ours, it would lend further evidence to the underlying medical knowledge of the physician as a factor contributing to these improved outcomes.

Despite the limitations of our study and the opportunities for future investigations, we demonstrated a clear relationship between initial certification and mortality. Moreover, for patients whose physicians had initial certification, there was an additional associated advantage if they maintained it. At this point, it is unknown whether this advantage is related to the educational aspects of the programme, the assessment component or both. Future work is needed to address this issue. Regardless of mechanism, the successful engagement of general internists with MOC is associated with lower in-hospital mortality in their patients with AMI or CHF.

#### Author affiliations

<sup>1</sup>FAIMER, Philadelphia, Pennsylvania, USA

<sup>2</sup>American Board of Internal Medicine, Philadelphia, Pennsylvania, USA

**Contributors** JJN, WW, JB, FM, and RSL take full responsibility for the work, had access to the data, and controlled the decision to publish. JJN, WW, JB, FM and RSL made substantial contributions to the conception and design of the work. JJN, WW, JB, FM and RSL made substantial contributions to the acquisition, analysis or interpretation of data for the work. JJN drafted the paper. JJN, WW, JB, FM and RSL revised the paper critically for important intellectual content. JJN, WW, JB, FM and RSL gave final approval of the version to be published. JJN, WW, JB, FM and RSL agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** WW, FM and RSL are paid employees of the American Board of Internal Medicine.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Ethics approval** The Institutional Review Board at Advarra approved this study as exempt (Pro00031598).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available. The data are from the files of the Pennsylvania Health Care Cost Containment Council, the American Board of Internal Medicine, the Educational Commission for Foreign Medical Graduates and the American Medical Association.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

John J Norcini <http://orcid.org/0000-0002-8464-4115>

#### REFERENCES

- Levinson W, Holmboe E. Maintenance of certification: 20 years later. *Am J Med* 2011;124:180–5.
- American Board of Internal Medicine. Becoming certified. Available: <https://www.abim.org/certification/default.aspx> [Accessed 07 Jan 2021].
- McDonald FS, Jurich D, Duhigg LM, *et al*. Correlations between the USMLE step examinations, American College of physicians in-training examination, and ABIM internal medicine certification examination. *Acad Med* 2020;95:1388–95.
- Hauer KE, Vandergrift J, Hess B, *et al*. Correlations between ratings on the resident annual evaluation summary and the internal medicine milestones and association with ABIM certification examination scores among US internal medicine residents, 2013–2014. *JAMA* 2016;316:2253–62.
- Jeffe DB, Andriole DA. Factors associated with American Board of medical specialties member board certification among US medical school graduates. *JAMA* 2011;306:961–70.
- Norcini JJ, Lipner RS, Kimball HR. Certifying examination performance and patient outcomes following acute myocardial infarction. *Med Educ* 2002;36:853–9.
- Chen J, Rathore SS, Wang Y, *et al*. Physician board certification and the care and outcomes of elderly patients with acute myocardial infarction. *J Gen Intern Med* 2006;21:238–44.
- Lipner RS, Young A, Chaudhry HJ, *et al*. Specialty certification status, performance ratings, and disciplinary actions of internal medicine residents. *Acad Med* 2016;91:376–81.



- 9 Kopp JP, Ibáñez B, Jones AT, *et al.* Association between American Board of surgery initial certification and risk of receiving severe disciplinary actions against medical licenses. *JAMA Surg* 2020;155:e200093.
- 10 Reid RO, Friedberg MW, Adams JL, *et al.* Associations between physician characteristics and quality of care. *Arch Intern Med* 2010;170:1442–9.
- 11 Vandergrift JL, Gray BM, Weng W. Do state continuing medical education requirements for physicians improve clinical knowledge? *Health Serv Res* 2018;53:1682–701.
- 12 Levinson W, Holmboe E. Maintenance of certification in internal medicine: facts and misconceptions. *Arch Intern Med* 2011;171:174–6.
- 13 Teirstein PS, Topol EJ. The role of maintenance of certification programs in governance and professionalism. *JAMA* 2015;313:1809–10.
- 14 Lipner RS, Hess BJ, Phillips RL. Specialty board certification in the United States: issues and evidence. *J Contin Educ Health Prof* 2013;33 Suppl 1:S20–35.
- 15 Gray B, Vandergrift J, Landon B, *et al.* Associations between American Board of internal medicine maintenance of certification status and performance on a set of healthcare effectiveness data and information set (HEDIS) process measures. *Ann Intern Med* 2018;169:97–105.
- 16 Gray BM, Vandergrift JL, Lipner RS. Association between the American Board of Internal Medicine's General Internist's Maintenance of Certification Requirement and Mammography Screening for Medicare Beneficiaries. *Women's Health Issues* 2018;28:35–41.
- 17 Gray BM, Vandergrift JL, McCoy RG, *et al.* Association between primary care physician diagnostic knowledge and death, hospitalisation and emergency department visits following an outpatient visit at risk for diagnostic error: a retrospective cohort study using Medicare claims. *BMJ Open* 2021;11:e041817.
- 18 Gray BM, Vandergrift JL, Johnston MM, *et al.* Association between imposition of a maintenance of certification requirement and ambulatory care-sensitive hospitalizations and health care costs. *JAMA* 2014;312:2348–57.
- 19 Nelson LS, Duhigg LM, Arnold GK, *et al.* The association between maintaining American Board of emergency medicine certification and state Medical board disciplinary actions. *J Emerg Med* 2019;57:772–9.
- 20 Kinney CL, Raddatz MM, Sliwa JA, *et al.* Association of participation in the American Board of physical medicine and rehabilitation maintenance of certification program and physician disciplinary actions. *Am J Phys Med Rehabil* 2020;99:325–9.
- 21 McDonald FS, Duhigg LM, Arnold GK, *et al.* The American Board of internal medicine maintenance of certification examination and state Medical board disciplinary actions: a population cohort study. *J Gen Intern Med* 2018;33:1292–8.
- 22 Zhou Y, Sun H, Macario A, *et al.* Association between performance in a maintenance of certification program and disciplinary actions against the medical licenses of Anesthesiologists. *Anesthesiology* 2018;129:812–20.
- 23 Jones AT, Kopp JP, Malangoni MA. Association between maintaining certification in general surgery and loss-of-license actions. *JAMA* 2018;320:1195–6.
- 24 Network E. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. Available: <https://www.equator-network.org/reporting-guidelines/strobe/> [Accessed February 9, 2020].
- 25 Krumholz HM, Chen J, Wang Y, *et al.* Comparing AMI mortality among hospitals in patients 65 years of age and older: evaluating methods of risk adjustment. *Circulation* 1999;99:2986–92.
- 26 Krumholz HM, Lin Z, Keenan PS, *et al.* Relationship between hospital readmission and mortality rates for patients hospitalized with acute myocardial infarction, heart failure, or pneumonia. *JAMA* 2013;309:587–93.
- 27 Schmaltz SP, Williams SC, Chassin MR, *et al.* Hospital performance trends on national quality measures and the association with joint Commission accreditation. *J Hosp Med* 2011;6:454–61.
- 28 Quan H, Sundararajan V, Halfon P, *et al.* Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005;43:1130–9.
- 29 The center for rural Pennsylvania. Available: <https://www.rural.palegislatore.us/ruralcounties.html> [Accessed 18 Jan 2021].
- 30 Prystowsky JB, Bordage G, Feinglass JM. Patient outcomes for segmental colon resection according to surgeon's training, certification, and experience. *Surgery* 2002;132:663–72.
- 31 Norcini JJ, Kimball HR, Lipner RS. Certification and specialization: do they matter in the outcome of acute myocardial infarction? *Acad Med* 2000;75:1193–8.
- 32 Holmboe ES, Wang Y, Meehan TP, *et al.* Association between maintenance of certification examination scores and quality of care for medicare beneficiaries. *Arch Intern Med* 2008;168:1396–403.
- 33 Second International Study of Infarct Survival Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. ISIS-2. *Lancet* 1988.
- 34 Weisman SM, Graham DY. Evaluation of the benefits and risks of low-dose aspirin in the secondary prevention of cardiovascular and cerebrovascular events. *Arch Intern Med* 2002;162:2197–202.
- 35 McDonald FS, Zeger SL, Kolars JC. Factors associated with medical knowledge acquisition during internal medicine residency. *J Gen Intern Med* 2007;22:962–8.
- 36 Marco CA, Wahl RP, House HR, *et al.* Physician age and performance on the American Board of emergency medicine concert examination. *Acad Emerg Med* 2018;25:891–900.
- 37 Dellinger EP, Pellegrini CA, Gallagher TH. The aging physician and the medical profession: a review. *JAMA Surg* 2017;152:967–71.
- 38 Norcini JJ, Boulet JR, Opalek A, *et al.* Patients of doctors further from medical school graduation have poorer outcomes. *Med Educ* 2017;51:480–9.
- 39 McDonald FS, Zeger SL, Kolars JC. Factors associated with medical knowledge acquisition during internal medicine residency. *J Gen Intern Med* 2007;22:962–8.
- 40 Yoneyama K, Kanaoka K, Okayama S, *et al.* Association between the number of board-certified cardiologists and the risk of in-hospital mortality: a nationwide study involving the Japanese registry of all cardiac and vascular diseases. *BMJ Open* 2019;9:e024657.
- 41 Gray BM, Vandergrift JL, Weng W, *et al.* Clinical knowledge and trends in physicians' prescribing of opioids for new onset back pain, 2009–2017. *JAMA Netw Open* 2021;4:e2115328.
- 42 Vandergrift JL, Weng W, Gray BM. The association between physician knowledge and inappropriate medications for older populations. *J Am Geriatr Soc* 2021;69:3584–94.