

Research

Open Access

Oral vaccination with a recombinant *Salmonella* vaccine vector provokes systemic HIV-1 subtype C Gag-specific CD4⁺ Th1 and Th2 cell immune responses in mice

Nyasha Chin'ombe¹, William R Bourn^{1,4}, Anna-Lise Williamson^{1,2} and Enid G Shephard*^{1,3}

Address: ¹Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Observatory 7925, Cape Town, South Africa, ²National Health Laboratory Services, Groote Schuur Hospital, Cape Town, South Africa, ³Department of Medicine, Faculty of Health Sciences, University of Cape Town, Observatory 7925, Cape Town, South Africa and ⁴Kapa Biosystems (Pty) Ltd, Observatory 7925, Cape Town, South Africa

Email: Nyasha Chin'ombe - Nyasha.Chinombe@uct.ac.za; William R Bourn - william.bourn@kapabiosystems.com; Anna-Lise Williamson - Anna-Lise.Williamson@uct.ac.za; Enid G Shephard* - Enid.Shephard@uct.ac.za

* Corresponding author

Published: 25 June 2009

Received: 30 April 2009

Virology Journal 2009, 6:87 doi:10.1186/1743-422X-6-87

Accepted: 25 June 2009

This article is available from: <http://www.virologyj.com/content/6/1/87>

© 2009 Chin'ombe et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Recombinant *Salmonella* vaccine vectors may potentially be used to induce specific CD4⁺ T cell responses against foreign viral antigens. Such immune responses are required features of vaccines against pathogens such as human immunodeficiency virus type 1 (HIV-1). The aim of this study was to investigate the induction of systemic HIV-1-specific CD4⁺ T helper (Th) responses in mice after oral immunization with a live attenuated *Salmonella* vaccine vector that expressed HIV-1 subtype C Gag. Groups of BALB/c mice were vaccinated orally three times (4 weeks apart) with this recombinant *Salmonella*. At sacrifice, 28 days after the last immunization, systemic CD4⁺ Th1 and Th2 cytokine responses were evaluated by enzyme-linked immunospot assay and cytometric bead array. HIV-1 Gag-specific IgG1 and IgG2a humoral responses in the serum were determined by enzyme-linked immunosorbent assay.

Results: Mice vaccinated with the recombinant *Salmonella* elicited both HIV-1-specific Th1 (interferon-gamma (IFN- γ) and tumour necrosis factor-alpha (TNF- α)) and Th2 (interleukin-4 (IL-4) and interleukin-5 (IL-5)) cytokine responses. The vaccine induced 70 (IFN- γ) spot-forming units (SFUs)/10e6 splenocytes and 238 IL-4 SFUs/10e6 splenocytes. Splenocytes from vaccinated mice also produced high levels of Th1 and Th2 cytokines upon stimulation with a Gag CD4 peptide. The levels of IFN- γ , TNF- α , IL-4 and IL-5 were 7.5-, 29.1-, 26.2- and 89.3-fold above the background, respectively. Both HIV-1 Gag-specific IgG1 and IgG2a antibodies were detected in the sera of vaccinated mice.

Conclusion: The study highlights the potential of orally-delivered attenuated *Salmonella* as mucosal vaccine vectors for HIV-1 Subtype C Gag to induce Gag-specific CD4⁺ Th1 and Th2 cellular immune responses and antibodies which may be important characteristics required for protection against HIV-1 infection.

Background

Attenuated *Salmonella* bacterial vaccines may be exploited for use as vectors for the oral delivery of HIV-1 antigens to both the mucosal and systemic compartments of the immune system. The bacteria provoke potent mucosal and systemic immune responses when administered by the oral route [1-4]. After oral administration, the bacteria are taken-up by professional phagocytes in the gut; they can then spread throughout the intestinal lymphatic tissues and reach the systemic compartments such as the liver and the spleen. In the phagocytes, the bacteria are found in *Salmonella*-containing vacuoles, or phagosomes, and the antigens are predominantly targeted to the MHC Class II presentation pathway, thereby provoking mainly the CD4+ Th1 and Th2 responses [5]. The induction of antigen-specific CD4 Th1 and Th2 responses is important for protection against infection by various types of pathogens. CD4+ Th1 cells produce cytokines such as IFN- γ , IL-2 and TNF- α , while CD4+ Th2 cells produce cytokines such as IL-4 and IL-5 [6-8]. In the case of viral infection, CD4+ Th1 and Th2 cells play a critical role in maintaining CD8+ T cell and antibody responses, respectively [9,10]. These cells are, therefore also indirectly important in their control of viral replication and viremia [11]. In the present study, we investigated the induction of systemic antigen-specific CD4+ Th1 and Th2 cell responses in mice that had been orally vaccinated with a recombinant *Salmonella enterica* serovar Typhimurium *aroC* vaccine vector that expressed codon-optimized HIV-1 subtype C Gag antigen.

Methods

Bacterial strains and culture conditions

Escherichia coli SCS110 cells (Stratagene, USA) were used for genetic manipulations and cloning. The *Salmonella enterica* serovar Typhimurium Δ *aroC* mutant vaccine strain (TML-MD58) was supplied by Microscience Pty Ltd (United Kingdom). This mutant has a deletion in the *aroC* gene, which encodes chorismate synthase, an enzyme that is crucial for the biosynthesis of tryptophan, tyrosine, phenylalanine, para-aminobenzoic acid and 2,3-dihydroxybenzoate [12]. 2YT media (supplemented, where necessary, with ampicillin and the aromatic amino acids) was used for culture of the recombinant *Salmonella*.

Construction of a Gag expression cassette

To construct the *Salmonella* Gag-expression plasmid, a codon-optimized HIV-1 *gag* gene, synthesized for us by Genart (USA), was cloned by standard recombinant DNA protocols [13] into pGEM+GFP, a plasmid designed to express green fluorescent protein that we previously constructed [14]. The *gfp* gene in pGEM+GFP was replaced with the *gag* gene and the plasmid designated pGEM+Gag was generated. The expression of Gag was under the control of the *E. coli lac* (lactose) promoter. Competent Δ *aroC*

Salmonella enterica serovar Typhimurium mutant was transformed with the expression plasmid (pGEM+Gag) and this resulted in the generation of a recombinant *Salmonella* vaccine clone, designated, *aroC*+Gag. The parent cloning vector, pGEM[®]-T Easy (Promega), was used as a negative control plasmid to generate a vaccine designated *aroC*+pGEM. The expression of HIV-1 Gag in the recombinant *Salmonella* bacteria, *aroC*+Gag, was assessed by SDS-PAGE according to standard protocols. The expression of the Gag protein was further confirmed using the Roche Elecsys[®] HIV p24 Ag assay (Roche) according to manufacturer's recommendations.

Salmonella vaccine stocks

Stocks of recombinant *Salmonella* bacterial vaccines were prepared from culture colonies of *aroC*+Gag (test vaccine) or *aroC*+pGEM (negative control vaccine). A single colony was inoculated into 200 ml of 2YT liquid media supplemented with ampicillin (100 μ g/ml) and aromatic amino acids and grown at 37°C with vigorous aeration. The bacterial cells were harvested when they reached logarithmic phase (OD₆₀₀ of 0.8) by centrifugation and washed once with PBS (pH 7.4). The vaccine was suspended in PBS with 15% glycerol and stored at -80°C prior to use for immunization.

Immunization of mice and preparation of splenocytes

Animal work was approved by the University of Cape Town Animal Ethics Committee. Female H-2^d BALB/c mice (8–10 weeks old) were purchased from South African Vaccine Producers (Johannesburg, South Africa) and were housed (5 per group) at the University of Cape Town Animal Unit. The mice were allowed to adapt to the new environment for at least 10 days before immunization. They were bled on day 0 prior to being inoculated by intragastric gavage with 1 \times 10⁸ colony forming units (CFUs)/mouse of either the *Salmonella* test vaccine (*aroC*+Gag) or the negative control (*aroC*+pGEM) and again on days 28 and 56 prior to the booster *Salmonella* vaccine being given. Blood was also taken on day 84 just prior to sacrifice of the mice for harvest of spleens. Serum was collected and used in ELISA assays to measure Gag-specific antibody responses.

Spleens were harvested from the mice at sacrifice on day 84. A pool of 5 spleens per group was made, then passed through a 70- μ m cell strainer to obtain a single cell suspension of splenocytes. Splenocytes were suspended in R10 medium (RPMI-1640 with 10% heat inactivated fetal calf serum, 15 mM β -mercaptoethanol, 100 U penicillin and 100 μ g streptomycin per ml). Red cells were lysed using erythrocyte lysing solution (0.15 M NH₄Cl, 10 mM KHCO₃, 0.1 mM Na₂EDTA). The splenocytes were suspended at a concentration of 5 \times 10⁶ cells per ml for use in enzyme-linked immunospot (ELISPOT) assays or at a

concentration of 15×10^6 cells per ml for stimulation with Gag peptides to quantify Gag-specific cytokines released into the supernatant.

IFN- γ and IL-4 ELISPOT assays

Gag-specific CD4⁺ T cells secreting IFN- γ or IL-4 in the spleen were enumerated using IFN- γ and IL-4 ELISPOT assay kits (BD Pharmingen) as previously described [15]. Briefly, splenocytes were cultured in triplicate wells at 0.5×10^6 cells/well in a final volume of 200 μ l with either R10 medium only (background responses) or R10 medium containing the Gag CD4 peptide (NPPIPVGDYKRWIILGLNK, an MHC class II-restricted, CD4 binding peptide) at 4 μ g/ml. After incubation (37°C, 5% CO₂) for 24 hours (IFN- γ ELISPOT assay) or 48 hours (IL-4 ELISPOT assay), the cells secreting IFN- γ or IL-4 were detected using Nova Red substrate (Vector Labs) according to the manufacturer's instructions. Spots were counted using CTL Analyzer and ImmunoSpot Version 3.2 software (Cellular Technology, USA). The mean number of spots \pm SD in triplicate wells was calculated and expressed as spot-forming units (SFU) per 10^6 splenocytes. Differences in immune responses between vaccine groups were analyzed by a two-sample student's t-test. A $p < 0.05$ was considered statistically significant. The differences in response between stimulated cells and unstimulated cells within the same vaccine group were also analyzed by a student's t-test and $p < 0.05$ was considered statistically significant.

Quantification of Gag-specific CD4⁺ Th1 and Th2 cytokine production

Splenocytes (1.5×10^6) were cultured (37°C, 5% CO₂) in a final volume of 200 μ l R10 medium only (background responses) or R10 medium containing the Gag CD4 peptide (as used in the ELISPOT assays) at 4 μ g/ml [15]. Culture supernatants were harvested at 48 hours and the content of Gag-specific Th1 cytokines (IFN- γ and TNF- α) and Th2 cytokines (IL-4 and IL-5) were quantified using a mouse Th1/Th2 Cytokine Bead Array (CBA) assay (BD Pharmingen) according to manufacturer's instructions. Results were expressed as pg cytokine per 10^6 splenocytes. Differences in immune responses between vaccine groups were analyzed by a two-sample student's t-test and $p < 0.05$ was considered statistically significant. The differences in response between stimulated cells and unstimulated cells within the same group were also analyzed by a student's t-test and $p < 0.05$ was considered statistically significant.

HIV-1 Gag antibody assay

Antibodies (total IgG) to HIV Gag were detected in mouse serum (pooled from 5 mice per group) collected on day 0, 28, 56 and 84 and Gag-specific IgG1 and IgG2a detected at day 84 using HIV-1 Gag (P55) enzyme linked immunosorbent assay (ELISA). Briefly, flat-bottom, 96-well Max-

iSorp ELISA plates (AEC-Amersham) were coated with 50 μ g of HIV-1 P55 (Quality Biological Inc. USA) in 50 μ l 0.1 M bicarbonate buffer (pH 9.5) overnight at 4°C. Plates were washed with PBS-Tween (PBS containing 0.3% Tween 20, Merck) and blocked with blocking solution (PBS containing 0.3% Tween 20, 1% goat serum, 3% milk powder) overnight at 4°C. Mouse serum samples were diluted (1/100), and 100 μ l added to duplicate wells and the reaction incubated overnight at 4°C. After washing with PBS-Tween containing 1% goat serum and 1% milk powder, total Gag (P55)-specific IgG, was detected by the addition of 100 μ l/well of biotinylated goat anti-mouse IgG (Southern Biotechnology, Birmingham, AL). After incubation at 37°C for 2 hours, plates were washed with PBS-Tween and streptavidin-alkaline phosphatase (50 μ l/well) was added and the plates incubated a further 1 hour at 37°C. To detect HIV-1 Gag (P55)-specific IgG1 and IgG2a isotypes, alkaline phosphatase-conjugated antibodies specific for IgG1 or IgG2a (Serotec) were used. Alkaline phosphatase was reacted with 50 μ l/well para-nitrophenyl phosphate (PNP) diluted to a final concentration of 1 mg/ml. The optical density (OD) was read at 405 nm and means of duplicates was calculated. For total IgG, the ratios of vaccinated to prebleed were calculated and represented graphically. The OD readings were represented graphically for IgG1 and IgG2a. Differences in immune responses between vaccine groups were analyzed by a two-sample student's t-test and a p value < 0.05 was considered statistically significant.

Results

Construction of a codon-optimized HIV-1 Gag-expressing Salmonella vaccine vector

A prokaryotic expression plasmid that contained a wild-type HIV-1 Subtype C gag gene was initially constructed. As Gag expression was poor, probably because of the presence of bacterial rare codons (data not shown), we redesigned the recombinant vaccine. For this the wild-type HIV-1 gag gene was replaced with a synthesised gene that reflected codon usage in *Salmonella* and the expression plasmid, pGEM+Gag, containing the codon-optimized gag was successfully constructed (Figure 1A). The plasmid contained the HIV-1 gag gene fused to the β -galactosidase α -fragment and the expression was constitutively driven by the *E. coli lac* promoter and other lac operon transcription and translation domains found in pGEM-Teasy plasmid. The recombinant plasmid-carrying *Salmonella* expressed very high levels of HIV Gag (Figure 1B). The Gag protein band was visible in Coomassie blue-stained SDS-PAGE gels (Figure 1B, Lane 2). The high-level of Gag protein expressed by the *Salmonella* was further confirmed by Roche Elecsys® HIV p24 Ag Capture assay (Figure 1C). The expression of codon-optimized Gag was higher than previously found with wild-type Gag (data not shown).




Figure 1
The HIV-1 subtype C Gag expression plasmid (pGEM+Gag) and Gag expression by Salmonella. (A) The HIV Gag expression cassette contained the *gag* gene fused in-frame with the β -galactosidase α -gene and expression was under the *E. coli lac* (*lactose*) promoter. The plasmid contained an *E. coli* origin of replication (*ori*) and ampicillin resistance gene (*AmpR*). (B) The relative expression of the Gag by the recombinant *Salmonella* vaccine (*aroC+Gag*) was determined by SDS-PAGE and (C) the Roche Elecsys[®] HIV p24 Ag assay. In the Roche Elecsys[®] HIV p24 Ag assay, total bacterial protein lysate was diluted 1/1000 in water and the cut-off index was calculated by the Elecsys[®] 2010 analyzer using readings from the negative and positive calibrators.

Oral vaccination of mice with a recombinant Salmonella induces Th1 and Th2 cytokine producing CD4+ T cells

CD4+ Th1/Th2 T cells induced in the spleens of mice in response to oral vaccination with the recombinant codon-optimized HIV-1 Gag-expressing *Salmonella* vaccine vector, *aroC+Gag*, were evaluated using ELISPOT assays. Mice vaccinated with *aroC+Gag* developed HIV-1 Gag-specific CD4+ Th1 (IFN- γ) and Th2 (IL-4) cells in the spleen. The frequency of Gag-specific IFN- γ spot-forming units from *aroC+Gag* was above the background ($p < 0.05$) (Figure 2A). Cells from the *aroC+Gag* group responded to Gag CD4 peptide stimulation. The number of IL-4-producing cells from *aroC+Gag* was also significantly higher than the number of cells producing IL-4 from the negative vaccine control ($p < 0.05$) (Figure 2B). It was, therefore, evident that both Gag-specific IFN- γ and IL-4 cytokines were elicited by the recombinant *Salmonella* vaccine vector.

A CBA assay was used to quantify Gag-specific IFN- γ , TNF- α , IL-4 and IL-5 cytokines secreted by the splenocytes upon antigenic stimulation. The amounts of CD4+ Th1 (IFN- γ and TNF- α) and Th2 (IL-4 and IL-5) cytokines secreted by the splenocytes into the supernatant were quantified. The levels of Gag-specific Th1 and Th2 cytokines were all significantly elevated above the background ($p < 0.05$) (Figure 3) in mice vaccinated with *aroC+Gag*. The levels of Th1 cytokines, IFN- γ and TNF- α were 7.5-fold and 29.1-fold above the background, respectively (Figure 3A and 3B). The Th2 cytokine levels, IL-4 and IL-5 were 26.2- and 89.3-fold above the background, respectively (Figure 3C and 3D). The CBA results, therefore, confirmed that vaccination of mice orally with the *Salmonella* vaccine vector induced systemic Gag-specific Th1 and Th2 cytokine responses.

Oral vaccination of mice with recombinant Salmonella induces Gag-specific antibodies

HIV-1 Gag-specific humoral immune responses in mice vaccinated with the HIV Gag-expressing *Salmonella* vaccine were evaluated on days 28, 56 and 84. Anti-Gag total IgG and IgG subtypes IgG2a and IgG1 in the serum of vaccinated mice were determined. A very low serum HIV-1 Gag specific IgG response (1.52-fold OD₄₀₅ reading above prebleed) was detected on day 28 in serum (1/100 dilution) in mice vaccinated with *aroC+Gag* and this was not significantly above the control ($p > 0.05$) (Figure 4A). On day 56, the antibody response was boosted significantly (5-fold OD₄₀₅ reading above prebleed) ($p < 0.05$). The Gag-specific antibody response was further boosted by day 84 (22-fold OD₄₀₅ reading above prebleed) ($p < 0.05$) (Figure 4A). HIV-1 Gag-specific IgG responses were confirmed using the New LAV Blot I HIV-1 Western blotting kit (Biorad). The serum from the *aroC+Gag*-vaccinated mice reacted specifically with Gag bands (P55, P40, P24/25 and P17/18) on the blot (results not shown). Gag-spe-




Figure 2
HIV-1 subtype C Gag-specific CD4+ Th1 and Th2 cell responses as evaluated by IFN-γ and IL-4 ELISPOT assays. Groups of mice (5 mice per group) were vaccinated three times (day 0, 28 and 56) with live recombinant *Salmonella* vaccine that expressed HIV-1 Subtype C Gag (aroC+Gag) or an antigen-negative *Salmonella* control vaccine (aroC+pGEM). Mice were sacrificed on day 84, and splenocytes were prepared from isolated spleens pooled from 5 mice per group then used in IFN-γ (A) and IL-4 (B) ELISPOT assays with R10 medium only (negative assay control) or with the Gag CD4 peptide. Bars are the mean number of spots from triplicate wells and are expressed as SFU/10e6 splenocytes. Differences in immune responses between vaccine groups were analyzed by a two-sample student's t-test. A p < 0.05 was considered statistically significant. The differences in response between stimulated cells and unstimulated cells within the same vaccine group were also analyzed by a student's t-test and a p < 0.05 was considered statistically significant. A p > 0.05 was considered not statistically significant (NS).

cific serum IgG subtypes measured on day 84 indicated the presence of significant (p < 0.01) levels of IgG1 and IgG2a (p < 0.05) (Figure 4B). The Gag-specific Ig2a response was slightly greater than the IgG1 response. In summary, both Th2-related IgG1 and Th1-related IgG2a antibody responses were induced in the mice vaccinated with the HIV Gag-expressing *Salmonella* vaccine.




Figure 3
HIV-1 subtype C Gag-specific CD4+ Th1 and Th2 cell responses as evaluated by the amount of cytokines secreted by stimulated cells. Groups of mice (5 per group) were vaccinated with live recombinant *Salmonella* vaccine that expressed HIV-1 Subtype C Gag (aroC+Gag) or antigen-negative *Salmonella* control vaccine (aroC+pGEM) as indicated in Figure 2. Splenocytes isolated and pooled from 5 mice per group on day 84 were incubated in R10 medium only (negative assay control), or stimulated with a Gag CD4 peptide for 48 hrs. Cytokines released into the supernatant were quantified using a mouse Th1/Th2 cytokine bead array (CBA) assay. (A) IFN-γ, (B) TNF-α, (C) IL-4 and (D) IL-5. Each bar in the graphs represents the average picogram amount of cytokine produced per 10e6 splenocytes in 48 hours of stimulation with media or Gag CD4 peptide, for triplicate responses. A p < 0.05 was considered statistically significant. A p > 0.05 was considered not statistically significant (NS).




Figure 4
HIV-1 subtype C Gag-specific serum IgG responses in mice vaccinated with recombinant *Salmonella* vaccine vector. Groups of mice (5 per group) were vaccinated with live recombinant *Salmonella* vaccine that expressed HIV-1 Subtype C Gag (aroC+Gag) or an antigen-negative *Salmonella* control vaccine (aroC+pGEM) as indicated in Figure 2. Serum (pooled from 5 mice per group) was isolated from blood taken before each vaccination on day 0, 28 and 56 and just before sacrifice on day 84. (A) The HIV-1 Gag-specific IgG for each group of mice with a 1/100 serum dilution. The data are the ratio of the OD_{405 nm} for vaccinated mice and the OD_{405 nm} for the day 0 serum (pre-bleed). (B) The HIV-1 Gag-specific IgG1 and IgG2a were measured in serum of each group of mice on Day 84 with a 1/100 serum dilution. Each bar represents the mean OD_{405 nm} value. Differences in antibody responses between vaccine groups at different time points were analyzed by a two-sample student's t-test and a $p < 0.05$ was considered statistically significant. A $p > 0.05$ was considered not statistically significant (NS).

Discussion

Delivery of heterologous antigens through the mucosal surface by recombinant *Salmonella* vectors is a powerful strategy for inducing both mucosal and systemic immune responses. After oral vaccination, the *Salmonella* bacteria invade the mucosal surfaces and spread through the

mesenteric lymph nodes to distal sites, such as spleen and liver [16,17]. This normally results in the induction of both mucosal and systemic cellular and humoral immune responses [18-20]. In the current study, we investigated the immunogenicity of a recombinant HIV-1 Subtype C Gag-expressing attenuated live *Salmonella enterica* serovar Typhimurium vaccine vector in mice after oral immunization. The HIV-1 gag gene was codon-optimized to reflect codons commonly used by *Salmonella* bacteria. The presence of rare codons in foreign genes may affect mRNA and plasmid stability and, in some cases, protein synthesis and bacterial growth [21-24]. In this study, we found that the expression of the HIV-1 Gag in *Salmonella* was improved when the gene was codon-optimized confirming published data [25]. However, other studies have shown that the level of antigen expression was decreased in *Salmonella* when the gene was codon-optimized [26]. Codon-optimization of genes for expression in recombinant *Salmonella* vaccine vectors has been reported to have an impact on the nature, breadth and magnitude of the immune responses induced after vaccination. It has been shown that antigen-specific immune responses against a *Salmonella*-based vaccine that expressed human papillomavirus type 16 L1 improved after codon-optimization [26]. The expression of measles virus (MV) epitopes in a *Salmonella* vaccine vector has been shown to be enhanced by codon-optimization [27]. Oral vaccination of MV-susceptible mice with the recombinant *Salmonella* vector induced MV-specific serum antibodies and CD4+ T cell response [26]. Codon-optimization of HIV-1 gag for expression in *Salmonella* resulted in enhanced mucosal immunity in vaccinated mice [25].

Our recombinant HIV-1 Gag-expressing *Salmonella* vaccine induced specific Th1 and Th2 cytokine responses in the spleen. This indicates that the recombinant bacteria successfully delivered the heterologous HIV-1 antigen to the systemic immune system after oral vaccination. Live *Salmonella* can be taken up by antigen-presenting cells by the process of phagocytosis and the bacteria are able to reside and replicate in the phagosomes. The *Salmonella* antigens are, therefore, presented to MHC class II-restricted CD4+ T cells [28-30]. This elicits antigen-specific Th1 and/or Th2 cytokine responses [31-34]. The two types of responses (CD4+ Th1 and Th2) induced by the recombinant *Salmonella* vector in this study are crucial for vaccines that are required to induce both cell-mediated and antibody responses for protection against infection by a number of pathogens. CD4+ Th1 cytokines such as IFN- γ and TNF- α provide protective immunity against intracellular pathogens such as viruses. They promote CD8+ T cell responses and B cell class-switching to IgG2a [35,36]. In contrast, CD4+ Th2 cytokines such as IL-4 and IL-5 promote B cell class switching to neutralizing antibodies such as IgG1 and they further regulate the magni-

tude of Th1 cytokine responses [36-38]. In this study, we also tested for the induction of Gag-specific CD8+ T cells in vaccinated mice. No specific CD8+ T cells were detectable by ELISPOT and CBA assays (results not shown). This was not unexpected; the secretion of antigens from the *Salmonella* bacteria has been shown to result in induction of antigen-specific CD8+ T cell responses in vaccinated mice [39-41]. Our HIV-1 Gag antigen was expressed inside the bacterial cytoplasm as inclusion bodies and was therefore unlikely to induce potent antigen-specific CD8+ T cell responses.

The nature of immune responses found in this study has relevance to the field of HIV-1 vaccinology. In HIV-1 infection, CD4+ Th1 and Th2 cells play regulatory roles in controlling infection and replication [42,43]. HIV-infected long-term non-progressors have been found to have strong CD4+ T cell responses to HIV-1 antigens such as Gag [44]. Recent studies have also suggested that HIV-specific CD4+ Th1 cell that produce INF- γ and IL-2 are important in long-term reduction of HIV viremia [45,46]. Other studies have shown that the loss of CD8+ T cell responses in HIV-1 infection could be reversed by vaccine-induced CD4+ Th cell responses [47]. Although CD4+ Th cells provide immunological help to CD8+ T cells and B cells, they can also play a more direct role in antiviral activity [48]. CD4+ T cells, like CD8+ T cells, have cytolytic activities against HIV-infected cells and can provide protective immune responses [49-52]. The cytokines, such as IFN- γ and TNF- α , secreted by activated CD4+ T cells have direct antiviral activities [53]. HIV-1 vaccines should, therefore, provoke both specific CD8+ and CD4+ T cell responses, so as to maximize the chance of preventing or controlling infection.

The development of an HIV-1 vaccine that elicits protective humoral immune responses is still a challenge to the scientific community. Such antibody responses should be able to neutralize many strains of the virus if they are to be useful. Although HIV-1 Gag is not a target for neutralizing antibodies, antibodies to Gag may play a role in other responses such as antibody-mediated cellular cytotoxicity and complement-mediated lysis of HIV-1 virions [54]. In the current study, we investigated whether *Salmonella* that express codon-optimized Gag could induce HIV-1 specific antibodies. Gag-specific IgG responses were induced in vaccinated mice. Further characterization of the anti-HIV-1 Gag antibody responses induced after secondary vaccinations with aroC+Gag showed the presence of both IgG1 and IgG2a subclasses. This result was supported by the finding that aroC+Gag induced Gag-specific CD4+ Th1 and Th2 cytokines. These cytokines produced by the Th1 and Th2 cells were most probably responsible for the induction of heavy-chain isotype switching to both IgG2a and IgG1 respectively [36]. In summary, our results

highlight the potential of using recombinant *Salmonella* as a vector for HIV-1 antigens. A *Salmonella*-delivered HIV-1 vaccine would be convenient for mass-vaccinations and inexpensive to produce. This would be advantageous for developing countries, where the HIV/AIDS is most prevalent, and the pandemic urgently needs to be brought under control.

Conclusion

In conclusion, the study showed that oral vaccination of mice with *Salmonella Typhimurium* vector expressing codon-optimized HIV-1 Gag could result in systemic HIV-1-specific CD4+ Th1 and Th2 cell immune responses, together with IgG1 and IgG2a humoral responses. Vaccines that provoke this type of immune response may be important in the prevention or control of HIV-1 infection.

Abbreviations

CBA: cytometric bead array; ELISA: enzyme-linked immunosorbent assay; ELISPOT: enzyme-linked immunospot; Gag: HIV group antigen; HIV-1: human immunodeficiency virus type 1; IFN- γ : interferon-gamma; IgG: immunoglobulin G; IL-4: interleukin 4; IL-5: interleukin 5; NS: not significant; OD: optical density; SDS-PAGE: sodium dodecyl sulphate-polyacrylamide gel electrophoresis; SFUs: spot-forming units; Th: T-helper; TNF- α : Tumour necrosis factor alpha; 2YT: 2 \times Yeast Tryptone

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NC, WRB, AW and EGS planned and designed the experiment. NC performed all the experiments. NC, WRB, AW and EGS all participated in the drafting of the manuscript. All the authors read and approved the manuscript.

Acknowledgements

This work was supported financially by a grant from the South African Aids Vaccine Initiative (SAAVI) of the Medical Research Council of South Africa. The *Salmonella* mutant was supplied by Microscience Pty Ltd (UK). We are grateful to members of the University of Cape Town Animal Unit and Sharon Makhubela, Shireen Galant, Desiree Bowers and Anke Binder for assistance with the immunology assays.

References

1. Fouts TR, Tuskan RG, Chada S, Hone DM, Lewis GK: **Construction and immunogenicity of *Salmonella typhimurium* vaccine vectors that express HIV-1 gp120.** *Vaccine* 1995, **13**:1697-1705.
2. Dietrich G, Griot-Wenk M, Metcalfe IC, Lang AB, Viret JF: **Experience with registered mucosal vaccines.** *Vaccine* 2003, **21**:678-683.
3. Capozzo AV, Cuberos L, Levine MM, Pasetti MF: **Mucosally delivered *Salmonella* live vector vaccines elicit potent immune responses against a foreign antigen in neonatal mice born to naive and immune mothers.** *Infect Immun* 2004, **72**:4637-4646.
4. Spreng S, Dietrich G, Weidinger G: **Rational design of *Salmonella*-based vaccination strategies.** *Methods* 2006, **38**:133-143.

5. Hess J, Schaible U, Raupach B, Kaufmann SH: **Exploiting the immune system: toward new vaccines against intracellular bacteria.** *Adv Immunol* 2000, **75**:1-88.
6. Mosmann TR, Sad S: **The expanding universe of T-cell subsets: Th1, Th2 and more.** *Immunol Today*. 1986, **17**(3):138-146.
7. Openshaw P, Murphy EE, Hosken NA, Maino V, Davis K, Murphy K, O'Garra A: **Heterogeneity of intracellular cytokine synthesis at the single-cell level in polarized T helper 1 and T helper 2 populations.** *J Exp Med* 1995, **182**:1357-1367.
8. Sornasse T, Larenas PV, Davis KA, de Vries JE, Yssel H: **Differentiation and stability of T helper 1 and 2 cells derived from naïve human neonatal CD4+ T cells, analyzed at the single-cell level.** *J Exp Med* 1996, **184**:473-483.
9. Hasenkruug KJ, Brooks DM, Dittmer U: **Critical role for CD4(+) T cells in controlling retrovirus replication and spread in persistently infected mice.** *J Virol* 1998, **72**:6559-6564.
10. Janssen EM, Lemmens EE, Wolfe T, Christen U, von Herrath MG, Schoenberger SP: **CD4+ T cells are required for secondary expansion and memory in CD8+ T lymphocytes.** *Nature* 2003, **421**:852-856.
11. Rosenberg ES, Billingsley JM, Caliendo AM, Boswell SL, Sax PE, Kalams SA, Walker BD: **Vigorous HIV-1-specific CD4+ T cell responses associated with control of viremia.** *Science* 1997, **278**:1447-1450.
12. Khan SA, Stratford R, Wu T, Mckelvie N, Bellaby T, Hindle Z, Sinha KA, Eltze S, Mastroeni P, Pickard D, Dougan G, Chatfield SN, Brennan FR: **Salmonella typhi and S typhimurium derivatives harbouring deletions in aromatic biosynthesis and Salmonella Pathogenicity Island-2 (SPI-2) genes as vaccines and vectors.** *Vaccine* 2003, **21**:538-548.
13. Sambrook J, Maniatis T, Fritsch EF: *Molecular cloning: a laboratory manual* 2nd edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; 1989.
14. Chin'ombe N, Bourn RW, Williamson A, Shepard EG: **An oral recombinant Salmonella enteric serovar Typhimurium mutant elicits systemic antigen-specific CD8+ T cell cytokine responses in mice.** *Gut Pathogens* 2009, **1**:9.
15. Shephard E, Burgers WA, Van Harmelen JH, Monroe JE, Greenhalgh T, Williamson C, Williamson AL: **A multigene HIV type 1 subtype C modified vaccinia Ankara (MVA) vaccine efficiently boosts immune responses to a DNA vaccine in mice.** *AIDS Res Hum Retroviruses* 2008, **24**:207-217.
16. Bradley DJ, Ghori N, Falkov S: **Salmonella Typhimurium initiates murine infection by penetrating and destroying the specialized epithelial M cells of the Peyer's patches.** *J Exp Med* 1994, **180**:15-23.
17. Everest P, Wain J, Roberts M, Rook G, Dougan G: **The molecular mechanisms of severe typhoid fever.** *Trends Microbiol* 2001, **9**:316-320.
18. Huang Y, Hajishengallis G, Michalek SM: **Induction of protective immunity against Streptococcus mutans colonization after mucosal immunization with attenuated Salmonella enterica serovar typhimurium expressing an S. mutans adhesin under the control of in vivo-inducible nirB promoter.** *Infect Immun* 2001, **69**:2154-2161.
19. Chen H, Schifferli DM: **Construction, characterization, and immunogenicity of an attenuated Salmonella enterica serovar typhimurium pgtE vaccine expressing fimbriae with integrated viral epitopes from the spiC promoter.** *Infect Immun* 2003, **71**:4664-4673.
20. Salam MA, Katz J, Zhang P, Hajishengallis G, Michalek SM: **Immunogenicity of Salmonella vector vaccines expressing SBR of Streptococcus mutans under the control of a T7-nirB (dual) promoter system.** *Vaccine* 2006, **24**:5003-5015.
21. Zahn K: **Overexpression of an mRNA dependent on rare codons inhibits protein synthesis and cell growth.** *J Bacteriol* 1996, **178**:2926-2933.
22. Ejdebäck M, Young S, Samuelsson A, Karlsson BG: **Effects of codon usage and vector-host combinations on the expression of spinach plastocyanin in Escherichia coli.** *Protein Expr Purif* 1997, **11**:17-25.
23. Baneyx F: **Recombinant protein expression in Escherichia coli.** *Curr Opin Biotechnol* 1999, **10**:411-421.
24. Kim S, Lee SB: **Rare codon clusters at 5'-end influence heterologous expression of archaeal gene in Escherichia coli.** *Protein Expr Purif* 2006, **50**:49-57.
25. Tsunetsugu-Yokota Y, Ishige M, Murakami M: **Oral attenuated Salmonella enterica serovar Typhimurium vaccine expressing codon-optimized HIV type 1 Gag enhanced intestinal immunity in mice.** *AIDS Res Hum Retroviruses* 2007, **23**:278-286.
26. Baud D, Ponci F, Bobst M, De Grandi P, Nardelli-Haeffliger D: **Improved efficiency of a Salmonella-based vaccine against human papillomavirus type 16 virus-like particles achieved by using a codon-optimized version of L1.** *J Virol* 2004, **78**:12901-12909.
27. Spreng S, Gentschev I, Goebel W, Weidinger G, Meulen V, Niewiesk S: **Salmonella vaccines secreting measles virus epitopes induce protective immune responses against measles virus encephalitis.** *Microbes Infect* 2000, **2**:1687-1692.
28. Svensson M, Stockinger B, Wick MJ: **Bone marrow-derived dendritic cells can process bacteria for MHC-I and MHC-II presentation to T cells.** *J Immunol* 1997, **158**:4229-4236.
29. Yrlid U, Wick MJ: **Salmonella-induced apoptosis of infected macrophages results in presentation of a bacteria-encoded antigen after uptake by bystander dendritic cells.** *J Exp Med* 2000, **191**:613-624.
30. Kalupahana RS, Mastroeni P, Maskell D, Blacklaws BA: **Activation of murine dendritic cells and macrophages induced by Salmonella enterica serovar Typhimurium.** *Immunology* 2005, **115**:462-472.
31. Galdiero M, De Martino L, Marcatili A, Nuzzo I, Vitiello M, Cipollaro de l'Ero G: **Th1 and Th2 cell involvement in immune response to Salmonella typhimurium porins.** *Immunology* 1998, **94**:5-13.
32. Pascual DW, Hone DM, Hall S, van Ginkel FW, Yamamoto M, Walters N, Fujihashi K, Powell RJ, Wu S, Vancott JL, Kiyono H, McGhee JR: **Expression of recombinant enterotoxigenic Escherichia coli colonization factor antigen I by Salmonella typhimurium elicits a biphasic T helper cell response.** *Infect Immun* 1999, **67**:6249-6256.
33. Kang HY, Curtiss R 3rd: **Immune responses dependent on antigen location in recombinant attenuated Salmonella typhimurium vaccines following oral immunization.** *FEMS Immunol Med Microbiol* 2003, **37**:99-104.
34. Jun S, Gilmore W, Callis G, Rynda A, Haddad A, Pascual DW: **A live diarrheal vaccine imprints a Th2 cell bias and acts as an anti-inflammatory vaccine.** *J Immunol* 2005, **175**:6733-6740.
35. Finkelman FD, Holmes J, Katona IM, Urban JF, Beckmann JP, Park LS, Schooley KA, Coffman RL, Mosmann TR, Paul WE: **Lymphokine control of in vivo immunoglobulin isotype selection.** *Ann Rev Immunol* 1990, **8**:303-333.
36. Spellberg B, Edwards JE Jr: **Type 1/Type 2 immunity in infectious diseases.** *Clin Infect Dis* 2001, **32**:76-102.
37. Stavnezer J: **Antibody class switching.** *Adv Immunol* 1996, **61**:79-146.
38. O'Garra A, Arai N: **The molecular basis of T helper 1 and T helper 2 cell differentiation.** *Trends Cell Biol* 2000, **10**:542-450.
39. Rüssmann H, Shams H, Poblete F, Fu Y, Galán JE, Donis RO: **Delivery of epitopes by the Salmonella type III secretion system for vaccine development.** *Science* 1998, **281**:565-568.
40. Chen LM, Briones G, Donis RO, Galán JE: **Optimization of the delivery of heterologous proteins by the Salmonella enterica serovar Typhimurium type III secretion system for vaccine development.** *Infect Immun* 2006, **74**:5826-5833.
41. Panthel K, Meinel KM, Sevil Domènech VE, Trülzsch K, Rüssmann H: **Salmonella type III-mediated heterologous antigen delivery: a versatile oral vaccination strategy to induce cellular immunity against infectious agents and tumors.** *Int J Med Microbiol* 2008, **298**:99-103.
42. Norris PJ, Moffett HF, Yang OO, Kaufmann DE, Clark MJ, Addo MM, Rosenberg ES: **Beyond help: direct effector functions of human immunodeficiency virus type 1-specific CD4(+) T cells.** *J Virol* 2004, **78**:8844-8851.
43. Jansen CA, van Baarle D, Miedema F: **HIV-specific CD4+ T cells and viremia: who's in control?** *Trends Immunol* 2006, **27**:119-124.
44. Kalams SA, Buchbinder SP, Rosenberg ES, Billingsley JM, Colbert DS, Jones NG, Shea AK, Trocha AK, Walker BD: **Association between virus-specific cytotoxic T-lymphocyte and helper responses in human immunodeficiency virus type 1 infection.** *J Virol* 1999, **73**:6715-6720.
45. Emu B, Sinclair E, Favre D, Moretto WJ, Hsue P, Hoh R, Martin JN, Nixon DF, McCune JM, Deeks SG: **Phenotypic, functional, and kinetic parameters associated with apparent T-cell control**

- of human immunodeficiency virus replication in individuals with and without antiretroviral treatment. *J Virol* 2005, **79**:14169-14178.
46. Martinez V, Costagliola D, Bonduelle O, N'go N, Schnuriger A, Theodorou I, Clauvel JP, Sicard D, Agut H, Debre P, Rouzioux C, Autran B: **Combination of HIV-1-specific CD4 Th1 cell responses and IgG2 antibodies is the best predictor for persistence of long-term nonprogression.** *J Infect Dis* 2005, **191**:2053-2063.
 47. Lichterfeld M, Kaufmann DE, Yu XG, Mui SK, Addo MM, Johnston MN, Cohen D, Robbins GK, Pae E, Alter G, Wurcel A, Stone D, Rosenberg ES, Walker BD, Altfeld M: **Loss of HIV-1-specific CD8+ T cell proliferation after acute HIV-1 infection and restoration by vaccine-induced HIV-1-specific CD4+ T cells.** *J Exp Med* 2004, **200**:701-712.
 48. Hasenkrug KJ, Brooks DM, Robertson MN, Srinivas RV, Chesebro B: **Immunoprotective determinants in friend murine leukemia virus envelope protein.** *Virology* 1998, **248**:66-73.
 49. Orentas RJ, Hildreth JE, Obah E, Polydefkis M, Smith GE, Clements ML, Siliciano RF: **Induction of CD4+ human cytolytic T cells specific for HIV-infected cells by a gp160 subunit vaccine.** *Science* 1990, **248**:1234-1237.
 50. Manickan E, Francotte M, Kuklin N, Dewerchin M, Molitor C, Gheysen D, Slaoui M, Rouse BT: **Vaccination with recombinant vaccinia viruses expressing ICP27 induces protective immunity against herpes simplex virus through CD4+ Th1+ T cells.** *J Virol* 1995, **69**:4711-4716.
 51. Hasenkrug KJ, Dittmer U: **The role of CD4 and CD8 T cells in recovery and protection from retroviral infection: lessons from the Friend virus model.** *Virology* 2000, **272**:244-249.
 52. Zaunders JJ, Dyer WB, Wang B, Munier ML, Miranda-Saksena M, Newton R, Moore J, Mackay CR, Cooper DA, Saksena NK, Kelleher AD: **Identification of circulating antigen-specific CD4+ T lymphocytes with a CCR5+, cytotoxic phenotype in an HIV-1 long-term nonprogressor and in CMV infection.** *Blood* 2004, **103**:2238-2247.
 53. Norris PJ, Sumaroka M, Brander C, Moffett HF, Boswell SL, Nguyen T, Sykulev Y, Walker BD, Rosenberg ES: **Multiple effector functions mediated by human immunodeficiency virus-specific CD4(+) T-cell clones.** *J Virol* 2001, **75**:9771-9779.
 54. Huber M, Trkola A: **Humoral immunity to HIV-1: neutralization and beyond.** *J Intern Med* 2007, **262**:5-25.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

