

# Cytotoxicity of some edible mushrooms extracts over liver hepatocellular carcinoma cells in conjunction with their antioxidant and antibacterial properties

Gökhan Sadi, Buğrahan Emsen, Abdullah Kaya, Aytaç Kocabaş, Seval Çınar, Deniz İrem Kartal<sup>1</sup>

Department of Biology, Kamil Özdağ Faculty of Science, Karamanoğlu Mehmetbey University, Karaman, <sup>1</sup>Department of Biology, Biochemistry Graduate Programme, Middle East Technical University, Ankara, Turkey

Submitted: 30-09-2014

Revised: 01-12-2014

Published: 27-05-2015

## ABSTRACT

**Background:** Mushrooms have been valued for their nutritive content and as traditional medicines; several important medicinal properties of mushrooms have been recognized worldwide. **Objective:** The purpose of this study was to elucidate the cell growth inhibitory potential of four edible mushrooms; *Coprinus comatus* (O.F. Mull.) Pers. (Agaricaceae), *Tricholoma fracticum* (Britzelm.) Kreisel (Tricholomataceae), *Rhizopogon luteolus* Fr. and Nordholm (Rhizopogonaceae), *Lentinus tigrinus* (Bull.) Fr. (Polyporaceae) on hepatocellular carcinoma (HepG2) cells in conjunction with their antioxidant and antibacterial capacities. **Materials and Methods:** Five different extracts of edible mushrooms were obtained using water, methanol, acetone, *n*-hexane and chloroform as solvent systems for cytotoxic, antioxidant and antibacterial properties. **Results:** *C. comatus* showed substantial *in vitro* cytotoxic activity against HepG2 cell lines with all extracts especially with chloroform 50% inhibition (IC<sub>50</sub> value of 0.086 mg/ml) and acetone (IC<sub>50</sub> value of 0.420 mg/ml). Chloroform extract of *C. comatus* had maximum amount of β-carotene (25.94 μg/mg), total phenolic content (76.32 μg/mg) and lycopene (12.00 μg/mg), and *n*-hexane extract of *L. tigrinus* had maximum amount of flavonoid (3.67 μg/mg). While chloroform extract of *C. comatus* showed the highest 2,2-diphenyl-1-picrylhydrazyl (DPPH) capturing activity (1.579 mg/ml), the best result for metal chelating activity was obtained from methanolic extract (0.842 mg/ml). Moreover, all tested mushrooms demonstrated antibacterial activity and *n*-hexane extract of *L. tigrinus* and acetone extracts of *T. fracticum* were the most active against tested microorganism. **Conclusion:** These results indicate that different extracts of investigated mushroom have considerable cytotoxic, antioxidant and antibacterial properties and may be utilized as a promising source of therapeutics.

**Key words:** Cancer cells, cell toxicity, edible macrofungi, radical scavenging

## INTRODUCTION

Wild edible mushrooms have been consumed in fresh or dried forms by human for a long period due to its organoleptic properties such as flavor and texture.<sup>[1-3]</sup> Although thousands of mushrooms are present in nature, few of them are accepted as edible and commercial.<sup>[4,5]</sup> Studies on nutritional values of mushrooms showed richness in protein, dietary fibers, vitamins and minerals along with low fat and energy<sup>[2,3,6,7]</sup> enables to use as edible. Local consumption of wild mushrooms are also increasing because

of their medicinal properties<sup>[4,6]</sup> due to the presence of secondary metabolites having pharmaceutical importance.<sup>[1,6]</sup>

Functional foods are new emerging area of food science and defined as food, which exerts health improving effects other than its nutritional value. Whereas, nutraceuticals are defined as concentrated phytochemicals derived from food and applied as a supplement.<sup>[8]</sup> Mushrooms are rich in nutraceuticals having antioxidant, antitumor and antimicrobial features.<sup>[1,4,9,10]</sup> There has been also a continuous search for new antimicrobial substances for decades and mushrooms are of interest for several researchers. Albeit nearly 60 antimicrobial compounds have been isolated from mushrooms; only the compounds from microscopic fungi have been present in the market as antibiotics until now.<sup>[11]</sup>

### Address for correspondence:

Dr. Gökhan Sadi, Department of Biology, Kamil Özdağ Faculty of Science, Karamanoğlu Mehmetbey University, 70100, Karaman, Turkey.  
E-mail: sadi@kmu.edu.tr

### Access this article online

#### Website:

www.phcog.com

#### DOI:

10.4103/0973-1296.157665

#### Quick Response Code:



Antioxidants protect the living systems from free radicals, which results in oxidative damage of protein and nucleic acids. Even though humans are well protected against free radicals, many diseases such as cancer may result from the uncontrolled production of reactive oxygen species.<sup>[12]</sup> Therefore, antioxidant supplementation either as foods or drugs helps people to reduce the risk resulted from oxidative stress. Drugs of herbal origin have been used in traditional systems of medicine since ancient times, and many researchers have benefited from medicinal plants to prevent oxidative stress.<sup>[13-17]</sup>

Although, antioxidant capacities of mushrooms are well-established, anticancerogenic properties of wild edible mushrooms in clinical studies<sup>[8]</sup> is of immense important due to very few reports. Therefore, this study focused on analysis of bioactive feature of some wild edible mushrooms. In this concern, we elucidated the cell growth inhibitory potential of four edible mushrooms; *Coprinus comatus* (O.F. Mull.) Pers. (Agaricaceae), *Tricholoma fracticum* (Britzelm.) Kreisel (Tricholomataceae), *Rhizopogon luteolus* Fr. and Nordholm (Rhizopogonaceae), *Lentinus tigrinus* (Bull.) Fr. (Polyporaceae) on liver hepatocellular carcinoma (HepG2) cells in conjunction with their antioxidant and antibacterial capacities.

## MATERIALS AND METHODS

### Collection and identification of mushroom samples

Samples were collected in 2012 from different localities within Karaman province of Central Anatolian Region of Turkey. Necessary morphological and ecological characteristics of the samples were recorded, and they were photographed in their natural habitats. Then they were brought to the fungarium and microstructural properties were investigated. Comparing the obtained macroscopic and microscopic data with literature,<sup>[18]</sup> they were identified as *T. fracticum* (Kaya 7482), *L. tigrinus* (Kaya 7454), *C. comatus* (Kaya 7483) and *R. luteolus* (Kaya 7455). The samples are kept at Karamanoğlu Mehmetbey University, Science Faculty, Department of Biology.

### Extraction of bioactive ingredients

After drying under mild heat evaporator, entire mushroom samples (cap and stripe) were powdered with liquid nitrogen, mortar and paste. Then bioactive ingredients were extracted by 250 ml of different solvent systems such as, water, methanol, chloroform, acetone and *n*-hexane using Soxhlet extraction apparatus throughout 2 days. After extraction, solvents were evaporated with rotary evaporator (IKA, Staufen Germany) under vacuum to dryness and lyophilized to get ultra dry powders that were solubilized with minimum amount of sterile distilled water.

They were diluted to different concentrations according to experiments before the experimental setup.

### Determination of cell growth inhibitory potential of mushrooms on hepatocellular carcinoma cells

Cell viability was quantified using 2,3-bis-(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide (XTT) *in vitro* cellular toxicity assay.<sup>[19]</sup> Accordingly, HepG2 cells were cultured in RPMI 1640 medium with L-glutamine and 25 mM HEPES (Lonza, USA) supplemented with 10% fetal bovine serum and 0.2% gentamycin sulfate. They were grown in a humidified incubator at 37°C with 5% CO<sub>2</sub>. After growing to expected confluency (90%), they were harvested by 2 ml 0.05% trypsin/ethylenediaminetetraacetic acid (EDTA), complete detachment was obtained by incubation at 37°C for 5 min. Later, trypsin activity was brought to end by the addition of 5 ml growth medium. Cell suspension was counted with trypan blue dye and cell concentration was adjusted to 100,000 cells/ml. In sterile microtiter culture disc, 50 µl HepG2 cell in fresh complete medium and 50 µl extracts in different concentration (0.5–10 mg/ml) were mixed and incubated for 48 h or 72 h at 37°C in 5% CO<sub>2</sub>.

Cell growth inhibitory potential of the mushrooms on HepG2 cells was studied using commercial XTT cell proliferation kit (Biological Industries, Israel). 100 µl of activation solution (PMS-N methyl dibenzopyrazine methyl sulfate) was added to 5 ml XTT reagent (supplied with kit) and mixed. Then, 50 µl of this solution was added to each well containing cells and extracts and incubated for 10 h with activated XTT under 5% CO<sub>2</sub> at 37°C in a humidified incubator. Finally, the intensity of the formazan was measured at 415 nm with Multiscan Go microplate reader (Thermo Scientific, USA), and 50% inhibition (IC<sub>50</sub>) values were calculated.

### Determination of antioxidant potentials

Possible antioxidant potential of different extracts of tested mushrooms were evaluated by measuring their total flavanoids, β-carotene and lycopene amount and phenolic contents as well as reducing power, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging and metal chelating activity. The methods used to evaluate these indicators of antioxidative capacity are given in the following parts.

### Determination of total phenolic contents

Concentration of total phenolic compounds in different extracts of *T. fracticum*, *L. tigrinus*, *C. comatus* and *R. luteolus* was determined spectrophotometrically according to the method described previously by<sup>[20]</sup> with slight modifications. In this method, gallic acid with various concentrations (0.01–1.0 mM) was used as a standard

phenolic compound. This method was adapted to microtiter plate measurements in triplicate. A volume of 20  $\mu\text{l}$  of standards or different extracts (10 mg/ml) were mixed with the same amount of Folin and Ciocalteu's phenol reagent (2N) and kept in dark for 3 min. Afterward, 20  $\mu\text{l}$  of 35% sodium carbonate (w/v) and 140  $\mu\text{l}$   $\text{dH}_2\text{O}$  were added to start incubation period for 10 min. After that, the absorbance was measured at 725 nm and the results were calculated as mean  $\pm$  standard error of mean (SEM) from gallic acid calibration curve and expressed as mg of gallic acid equivalents/mg of extracts.

#### Determination of total flavonoid contents

Total flavonoids of water, methanol, chloroform, acetone and *n*-hexane extracts obtained from edible mushrooms were determined using protocols reported by<sup>[21]</sup> with slight modifications. A volume of 50  $\mu\text{l}$  of extracts (10 mg/ml) were mixed with 215  $\mu\text{l}$  of ethyl alcohol (80% v/v), 5  $\mu\text{l}$  of aluminum nitrate (10% w/v) and 5  $\mu\text{l}$  potassium acetate (1 M) in microtiter plates and incubated for 40 min at room temperature. After reading at 415 nm, total flavonoid contents were calculated according to following equation:

$$\text{Total flavonoid contents } (\mu\text{g}/\text{mg extract}) = (A_{415} + 0.01089)/0.002108$$

#### Determination of $\beta$ -carotene and lycopene contents

Different extracts with water, methanol, *n*-hexane, acetone and chloroform that were obtained from four edible mushroom was re-extracted with 10 ml of acetone: hexane (4:6) mixture and filtered through Whatman No. 4 filter paper in order to determine  $\beta$ -carotene and lycopene contents. After filtration, absorbance of the filtrates was measured at 453, 505 and 663 nm.  $\beta$ -carotene and lycopene contents were determined according to following equations.<sup>[21]</sup>

$$\beta\text{-carotene content (mg}/100 \text{ mg)} = 0.216 A_{663} - 0.304 A_{505} + 0.452 A_{453}$$

$$\text{Lycopene content (mg}/100 \text{ mg)} = -0.0458 A_{663} + 0.372 A_{505} - 0.0806 A_{453}$$

#### DPPH radical scavenging activity

Free radical scavenging activities (RSA) of extracts obtained from *T. fracticum*, *L. tigrinus*, *C. comatus* and *R. luteolus* were determined by monitoring DPPH reduction. Gallic acid (0.01–0.5 mM) was used as standard antioxidant molecule and to determine RSA, 20  $\mu\text{l}$  of various concentrations of standard or extracts were mixed with 180  $\mu\text{l}$  of DPPH solution (0.06 mM in methanol) and incubated for 1 h at dark in microtiter plates. Blank measurements without standard or extracts were also performed. After reading absorbance at 517 nm,

reduction of the DPPH radical was determined as percent discoloration of DPPH which was calculated according to following formula;

$$\text{Radical Scavenging Activity (RSA) (\%)} = 100$$

$$\times \frac{\text{DPPH absorbance} - \text{DPPH and extract absorbance}}{\text{DPPH absorbance}}$$

The extract concentrations providing the  $\text{IC}_{50}$  was calculated from the graph of RSA versus extract amount that was used for the comparison of different extracts of tested mushrooms.<sup>[22]</sup>

#### Determination of metal chelating activity

Chelating abilities of various extracts from edible mushrooms were determined using EDTA as a standard chelating agent.<sup>[23]</sup> Different concentrations (50  $\mu\text{l}$ ) of extracts (2, 4, 6, 8, 10 mg/ml) and standard (0.1–5 mM) were added to microplate wells and mixed with 10  $\mu\text{l}$  ferrozine (5 mM), 5  $\mu\text{l}$  iron (II) chloride (2 mM) and 185  $\mu\text{l}$  absolute methanol. After incubation for 10 min at room temperature, absorbances were read at 562 nm, and  $\text{IC}_{50}$  values were calculated.

#### Determination of reducing power

Reducing powers of different extracts were determined according to the method<sup>[24]</sup> with an adaptation to microplate measurement. Gallic acid (0.01–0.1 mM) was used as a standard antioxidant. In this method, various concentrations of 50  $\mu\text{l}$  mushroom extracts (2, 4, 6, 8, 10 mg/ml) were mixed with 75  $\mu\text{l}$  phosphate buffer (0.2 M pH: 6.6) and 75  $\mu\text{l}$  potassium ferricyanide (1% w/v) in a total volume of 200  $\mu\text{l}$  and incubated at 50°C for 20 min. After adding 75  $\mu\text{l}$  trichloroacetic acid (10% w/v), samples were centrifuged for 10 min at 1000  $\times g$ . Supernatant (75  $\mu\text{l}$ ) were transferred to another microtiter plate and mixed with 75  $\mu\text{l}$  distilled water and 15  $\mu\text{l}$  iron (III) chloride (0.1% w/v). After reading the absorbance at 700 nm, effective concentrations ( $\text{EC}_{50}$ ) at which the absorbance was 0.5 for reducing power was calculated.

#### Determination of antibacterial activities

The antibacterial activities of different extracts of four edible mushrooms (40  $\mu\text{g}/\mu\text{l}$ ) were determined against *Bacillus subtilis* (Ehrenberg) Cohn, *Bacillus licheniformis*, *Staphylococcus aureus* Rosenbach, *Escherichia coli* T. Escherich and *Agrobacterium tumefaciens* Smith and Townsend by the disc diffusion susceptibility method.<sup>[25]</sup> Penicillin, gentamicin and tetracycline were used as the standard antibiotic discs. Microorganisms were grown in Müller Hilton broth at 35°C and 28°C for 18 h and then cell density were adjusted according to 0.5 McFarland standards. After that, 100  $\mu\text{l}$  of adjusted microorganism suspension was inoculated onto Müller Hilton Agar by spread plate



technique. A volume of 40  $\mu\text{l}$  of extract (40  $\mu\text{g}/\mu\text{L}$ ) loaded discs and standard antibiotic discs were placed on inoculated Petri dishes. Negative controls were prepared with water alone. Inhibition zones (IZ) around the discs after overnight incubation were measured to determine antibacterial activities of the extracts. The study was performed in triplicates of each sample.

### Statistical analyses

All the assays were carried out at least in triplicate measurements. The results are expressed as mean values and SEM. Antioxidant, antibacterial and cytotoxicity activities of mushrooms were compared with Analysis of Variance (ANOVA) test followed by the appropriate *post-hoc* test (Duncan Test) and values with  $P < 0.05$  were considered as significantly different.  $\text{IC}_{50}$  and  $\text{EC}_{50}$  values were calculated with Probit Regression Analysis and associated 95% confidence limits for each treatment. Relations among the variables were tested by Bivariate Correlation Analysis. These calculations were carried out using Statistical Package for Social Sciences (SPSS®, version 21.0, IBM Corporation, Armonk, NY, USA).

## RESULTS AND DISCUSSION

Naturally occurring substances present in edible mushrooms have beneficial effects on health. Several species of mushroom show free RSA and high growth inhibitory potential against cancer cells and microorganisms<sup>[23,26-33]</sup> that's why, mushrooms have recently become attractive as nutritionally beneficial foods and as a source material for the development of some drugs. This study presents the comparison of four different edible mushrooms; *T. fracticum*, *L. tigrinus*, *C. comatus* and *R. luteolus* which are consumed locally in different regions of Turkey in terms of their important medicinal properties such as cytotoxic, antioxidant and antibacterial effects.

### Cell growth inhibitory potential of mushrooms on liver hepatocellular carcinoma cells

Liver tissue is involved in the metabolism of various nutrients including edible mushrooms and their digestive products in the form of glucose, amino acids and fatty acids. Furthermore, it contains thousands of enzymes essential to perform vital metabolic functions to metabolize both beneficial and harmful substances and store some of the compounds having antioxidant properties.<sup>[26,34,35]</sup>

Cancer diseases are one of the main causes of death worldwide and hepatocellular carcinoma (also called malignant hepatoma) is the most common type of liver cancer.<sup>[26]</sup> Liver HepG2 cells are the model cell lines to study liver functions *in vitro* since they maintain several

liver properties. Our study reports on the investigation into the cytotoxic activity of different extracts of four edible mushrooms on HepG2 cells that have not been reported previously.

To study the growth inhibitory activity of the four edible mushrooms *in vitro*, HepG2 cell lines were incubated with various concentrations of four edible mushrooms obtained from different solvent extraction. The potentially cytotoxic effect of the mushroom extracts was assessed using *in vitro* XTT assay. Findings from our study clearly demonstrated that elevated concentrations of mushroom extracts are cytotoxic to human liver carcinoma (HepG2) cells. In particular, *C. comatus* displayed the strongest growth inhibitory potential on HepG2 cells. However, all extracts of *C. comatus* are more effective anticancer agent than other mushrooms tested with very low  $\text{IC}_{50}$  values (0.086 mg/ml for chloroform and 0.42 mg/ml for acetone extracts). In addition to *C. comatus*, *L. tigrinus* also possess high degree of growth inhibitory potential over HepG2 cells for all extracts ( $\text{IC}_{50}$  value of 0.376 mg/ml for acetone, 0.434 mg/ml for *n*-hexane and 2.854 mg/ml for chloroform). In contrast, *T. fracticum* was the least effective cytotoxic material among the all tested mushroom since nearly tenfold higher amount of extract was needed to obtain  $\text{IC}_{50}$  values as compared with the most effective *C. comatus*.  $\text{IC}_{50}$  value of acetone extract was 3.851 mg/ml, 5.164 mg/ml for chloroform and 51.982 mg/ml for water extract of *T. fracticum* that was significantly higher than those of other mushrooms. Cytotoxic activities of all tested extracts were summarized in Table 1.

### Antioxidant components

Substances that can inhibit or retard oxidation of substances are called antioxidants and recently there has been increasing interest in discovering natural antioxidants, especially those of macrofungal origin. Various macrofungi species, plants, lichens and algae have antioxidant constituents which are important materials due to the fact that these components have anti-inflammatory, antiallergic, antiviral, anti-aging, anticarcinogenic properties and behave like biological response modifiers.<sup>[36-39]</sup> Therefore, they are used in medicinal and pharmaceutical fields by many researchers. The present study was undertaken with the aim to carry out a systematic comparative antioxidant study on different extracts gained from four edible mushrooms using different *in vitro* methods, and to find any correlation between the antioxidant activity and bioactive ingredients such as total phenolic, total flavonoid, lycopene and  $\beta$ -carotene contents of the extracts. To evaluate the total antioxidant properties of four edible mushrooms, three different assays were carried out namely; scavenging activity on DPPH radicals, reducing power and metal chelating activity.

**Table 1: 48 and 72 h IC<sub>50</sub> values (mg/ml) of different extracts obtained from four mushrooms for cytotoxicity test**

Treatment	Species	48 h IC <sub>50</sub> (limits)	72 h IC <sub>50</sub> (limits)
Acetone <sup>a</sup>	<i>T. fracticum</i>	3.851 (3.529-4.194) K <sup>b</sup>	3.079 (2.827-3.331) M <sup>b</sup>
	<i>L. tigrinus</i>	0.376 (0.093-0.650) B	0.362 (0.100-0.610) D
	<i>C. comatus</i>	0.420 (0.143-0.692) C	0.307 (0.082-0.544) B
	<i>R. luteolus</i>	1.539 (1.289-1.763) F	1.410 (1.177-1.618) G
Chloroform <sup>a</sup>	<i>T. fracticum</i>	5.164 (4.617-5.827) O	3.675 (3.324-4.052) O
	<i>L. tigrinus</i>	2.854 (1.398-60.414) I	2.576 (1.329-34.778) L
	<i>C. comatus</i>	0.086 (0.015-0.149) A	0.078 (0.009-0.141) A
	<i>R. luteolus</i>	2.167 (1.843-2.491) H	1.824 (1.518-2.106) H
<i>n</i> -Hexane <sup>a</sup>	<i>T. fracticum</i>	16.548 (4.347-4656.340) S	11.962 (3.729-570.714) S
	<i>L. tigrinus</i>	0.434 (0.400-0.471) D	0.322 (0.301-0.341) C
	<i>C. comatus</i>	0.864 (0.721-1.188) E	0.523 (0.481-0.576) E
	<i>R. luteolus</i>	1.604 (1.437-1.762) G	1.327 (1.166-1.474) F
Methanol <sup>a</sup>	<i>T. fracticum</i>	11.096 (9.743-13.133) Q	8.786 (7.516-10.629) Q
	<i>L. tigrinus</i>	7.446 (6.607-8.607) P	5.046 (4.613-5.558) P
	<i>C. comatus</i>	3.899 (3.590-4.238) L	3.217 (2.921-3.541) N
	<i>R. luteolus</i>	4.319 (3.992-4.684) N	2.445 (2.222-2.677) K
Water <sup>a</sup>	<i>T. fracticum</i>	51.982 (22.611-548.396) T	44.063 (20.480-330.053) T
	<i>L. tigrinus</i>	3.902 (3.356-4.584) M	2.201 (1.947-2.469) I
	<i>C. comatus</i>	3.404 (2.960-3.969) J	2.339 (2.041-2.669) J
	<i>R. luteolus</i>	12.641 (10.116-17.344) R	11.519 (9.058-16.161) R

<sup>a</sup>Extract; <sup>b</sup>Values followed by different capital letters in the same column differ significantly at  $P < 0.05$ . *T. fracticum*: *Tricholoma fracticum*, *L. tigrinus*: *Lentinus tigrinus*, *C. comatus*: *Coprinus comatus*, *R. luteolus*: *Rhizopogon luteolus*, IC<sub>50</sub>: 50% inhibition

### Scavenging ability on DPPH radicals

The model of scavenging DPPH radical is a widely used method to evaluate the free RSA of antioxidants. DPPH is a stable free radical, and when antioxidants react with DPPH, unpaired electron is paired off and the DPPH solution is decolorized. Therefore, decolorization degree demonstrates the radical scavenging activity of an antioxidant containing materials.

Previous studies showed that extracts or chemical constituents obtained from various mushrooms possess DPPH scavenging activities due to their antioxidant compounds. Studies carried out with *Laetiporus sulphureus*;<sup>[40]</sup> *Pleurotus ostreatus*;<sup>[31,41]</sup> *Lentinus edodes*, *Sparassis crispa* and *Mycoleptodonoides aitchisonii*;<sup>[42,43]</sup> *Stereum hirsutum*;<sup>[44]</sup> *Russula virescens*;<sup>[45]</sup> *Pleurotus eryngii*;<sup>[46]</sup> *Macrolepiota procera*;<sup>[27,47]</sup> *Pleurotus flabellatus*;<sup>[38]</sup> *Russula laurocerasi*;<sup>[48]</sup> *Suillus bovinus*;<sup>[49]</sup> *Phellinus fastuosus*, *Phellinus grenadensis*, *Phellinus merrillii* and *Phellinus badius*;<sup>[50]</sup> *Lenzites betulina*;<sup>[51]</sup> *Lactarius piperatus*;<sup>[52]</sup> *Cordyceps militaris*;<sup>[53]</sup> *Tricholoma portentosum* and *Lactarius deliciosus*;<sup>[54]</sup> *Pholiota nameko*<sup>[55]</sup> revealed the high DPPH scavenging activities of these fungi correlated with their antioxidant compounds.

In the present study, all the mushroom species showed DPPH free RSA that were found to rise with increasing concentration. Maximum concentrations (10 mg/ml) of methanol extracts from *R. luteolus* and *T. fracticum* and chloroform extract from *C. comatus* showed DPPH

scavenging activities over 70% (71.38, 71.19 and 70.28%, respectively). Among all the mushroom extracts, highest activity was obtained from the lowest concentration (2 mg/ml) of chloroform extract from *C. comatus* (53.56%). Comparing activity rates of the extracts of four mushrooms, chloroform extracts gave best results and all activity rates were over 60%. The scavenging effects of four mushrooms (in chloroform extract) on DPPH radical were in the descending order of *C. comatus* > *T. fracticum* > *L. tigrinus* > *R. luteolus*. The lowest activity values belonged to acetone extract. In acetone extract, DPPH scavenging activity rates of four mushrooms ranged from 13.67% to 40.08% for maximum concentration (10 mg/ml) and was in the order of *L. tigrinus* < *T. fracticum* < *C. comatus* < *R. luteolus* [Figure 1]. The chloroform extract from *C. comatus* showed strong DPPH radical scavenger that was reflected by its low IC<sub>50</sub> value (1.579 mg/ml). Other extracts that had low IC<sub>50</sub> values (under 3 mg/ml) are methanol and chloroform extract from *R. luteolus* (2.398 and 2.518 mg/ml) and methanol extract from *C. comatus* (2.566 mg/ml). IC<sub>50</sub> value of standard (gallic acid) was found as 0.037 mg/ml and there was statistically ( $P < 0.05$ ) significant difference between this value and the values belonging to all mushroom extracts [Table 2].


### Chelating abilities of metal ions

The presence of metal ions; especially Fe<sup>2+</sup> lead the production of reactive species such as hydroxyl radicals and metal-oxygen complex in biological systems and cellular

**Table 2: IC<sub>50</sub> values (mg/ml) of different extracts from four mushrooms and standard for scavenging activity on DPPH radicals**

Treatment	Species	IC <sub>50</sub> (limits)	Slope±SEM
Acetone <sup>a</sup>	<i>T. fracticum</i>	44.122 (27.256-106.624) Q <sup>c</sup>	1.214±0.187
	<i>L. tigrinus</i>	68.591 (36.524-249.140) R	1.359±0.242
	<i>C. comatus</i>	26.226 (18.622-46.629) P	1.195±0.163
	<i>R. luteolus</i>	17.089 (13.072-26.471) N	1.087±0.147
Chloroform <sup>a</sup>	<i>T. fracticum</i>	4.731 (4.094-5.406) I	1.114±0.134
	<i>L. tigrinus</i>	4.472 (3.693-5.267) H	0.902±0.133
	<i>C. comatus</i>	1.579 (0.568-2.415) B	0.581±0.133
	<i>R. luteolus</i>	2.518 (1.203-3.490) D	0.549±0.132
<i>n</i> -Hexane <sup>a</sup>	<i>T. fracticum</i>	128.817 (37.953-16039.858) U	0.471±0.145
	<i>L. tigrinus</i>	85.547 (31.499-2261.451) T	0.524±0.144
	<i>C. comatus</i>	76.708 (29.311-1778.957) S	0.521±0.143
	<i>R. luteolus</i>	15.857 (12.214-24.349) M	1.045±0.145
Methanol <sup>a</sup>	<i>T. fracticum</i>	3.257 (2.642-3.800) G	1.095±0.134
	<i>L. tigrinus</i>	7.451 (5.669-12.070) K	0.542±0.132
	<i>C. comatus</i>	2.566 (1.734-3.242) E	0.815±0.133
	<i>R. luteolus</i>	2.398 (1.571-3.070) C	0.810±0.133
Water <sup>a</sup>	<i>T. fracticum</i>	10.161 (8.152-14.595) L	0.846±0.136
	<i>L. tigrinus</i>	22.638 (13.523-87.929) O	0.566±0.137
	<i>C. comatus</i>	6.072 (5.121-7.409) J	0.866±0.134
	<i>R. luteolus</i>	3.193 (0.817-4.783) F	0.366±0.131
Gallic acid <sup>b</sup>	-	0.037 (0.026-0.048) A	0.601±0.053

<sup>a</sup>Extract, <sup>b</sup>Standard solution, <sup>c</sup>Values followed by different capital letters in the same column differ significantly at  $P < 0.05$ . *T. fracticum*: *Tricholoma fracticum*, *L. tigrinus*: *Lentinus tigrinus*, *C. comatus*: *Coprinus comatus*, *R. luteolus*: *Rhizopogon luteolus*, SEM: Standard error of mean, DPPH: 2,2-diphenyl-1-picrylhydrazyl, IC<sub>50</sub>: 50% inhibition



**Figure 1:** Scavenging abilities of extracts in different concentrations from four mushrooms on 2,2-diphenyl-1-picrylhydrazyl (DPPH) radicals

damage eventually occurs with metal catalyzed reactions.<sup>[56]</sup> To prevent oxidative stress caused by metal ions, edible mushrooms that have metal chelating abilities are very

important in foods. In the determination of metal chelating activities, extracts of the mushroom species was used to suppress the formation of ferrous and ferrozine complex,

indicating their chelating activity and capture ferrous ion before the production of ferrozine in the system.

Many researchers have previously examined chelating activities of different extracts of some mushroom species and chemical constituents of different fungi in their studies and achieved positive results.<sup>[38,40,45,46,57-62]</sup>

The present study has also been focused on metal chelating activities of different extracts from *T. fracticum*, *L. tigrinus*, *C. comatus* and *R. luteolus*. Correlations between chelating capacities of various extracts of four mushrooms and their concentrations were observed. Methanol extracts at concentrations of 8 and 10 mg/ml and chloroform extract at concentration of 10 mg/ml of of *C. comatus* displayed over 70% metal chelating activity (70.17, 78.01 and 70.52%, respectively). Methanol extract gave higher results compared to other extracts. For the chelating effect of ferrous ion, four mushrooms in methanol extract at concentration of 10 mg/ml were effective in the following order: *C. comatus* > *R. luteolus* > *T. fracticum* > *L. tigrinus* [Figure 2].

As shown in Table 3, for chelating on ferrous ions, highly potent inhibitions were found by methanol extract from *C. comatus* (IC<sub>50</sub> = 0.842 mg/ml), methanol extract from *R. luteolus*, (IC<sub>50</sub> = 2.397 mg/ml) and chloroform extract from *C. comatus* (IC<sub>50</sub> = 5.650 mg/ml) when compared with standard (EDTA) inhibitor (IC<sub>50</sub> = 0.158 mg/ml) [Table 3].

### Determination of reducing power

The reducing capacity of a compound may serve as a significant indicator of its potential antioxidant activity.<sup>[63]</sup> In the used protocol, an increase in the color of the test solution depends on the reducing power of each compound. The presence of reducers (antioxidants) causes the conversion of the Fe<sup>3+</sup>/ferricyanide complex used in this method to the ferrous form. Higher absorbance indicated higher reducing power of the selected extracts.

Previously, researchers reported that many mushroom species have reducing power.<sup>[52,54,60]</sup> They also reported that methanolic extracts of some mushroom species had high reducing power activity.

In the present study, reducing power of the various extracts of edible mushrooms increased with concentration. Based on the reducing power, among the mushroom extracts predominant one was acetone extract from *R. luteolus* (EC<sub>50</sub> = 1.754 mg/ml). While the highest reducing powers in chloroform and methanol extracts were belonged to *C. comatus* (EC<sub>50</sub> = 4.045 and 2.541 mg/ml, respectively), in *n*-hexane and water extracts *R. luteolus* showed the highest reducing capacities (EC<sub>50</sub> = 4.869 and 2.468 mg/ml, respectively) [Table 4].

However, as shown in Figure 3, *L. tigrinus* demonstrably had the lowest reducing activity in acetone and chloroform

**Table 3: IC<sub>50</sub> values (mg/ml) of different extracts from four mushrooms and standard for chelating on ferrous ions**

Extract	Species	IC <sub>50</sub> (Limits)	Slope±SEM
Acetone <sup>a</sup>	<i>T. fracticum</i>	10.488 (9.109-12.697) J <sup>c</sup>	1.476±0.148
	<i>L. tigrinus</i>	10.996 (9.848-12.669) K	2.086±0.171
	<i>C. comatus</i>	11.632 (9.944-14.480) M	1.448±0.150
	<i>R. luteolus</i>	24.468 (17.846-41.007) T	1.261±0.165
Chloroform <sup>a</sup>	<i>T. fracticum</i>	8.946 (7.616-11.224) F	1.096±0.138
	<i>L. tigrinus</i>	9.564 (8.525-11.079) H	1.706±0.151
	<i>C. comatus</i>	5.650 (5.157-6.208) D	1.654±0.140
	<i>R. luteolus</i>	66.157 (30.383-457.340) U	0.687±0.152
<i>n</i> -Hexane <sup>a</sup>	<i>T. fracticum</i>	13.339 (10.739-18.677) P	1.117±0.144
	<i>L. tigrinus</i>	15.088 (11.953-21.729) S	1.154±0.147
	<i>C. comatus</i>	12.740 (11.099-15.336) O	1.891±0.169
	<i>R. luteolus</i>	14.664 (12.115-19.359) R	1.458±0.157
Methanol <sup>a</sup>	<i>T. fracticum</i>	7.467 (6.597-8.712) E	1.287±0.139
	<i>L. tigrinus</i>	14.119 (11.809-18.229) Q	1.523±0.158
	<i>C. comatus</i>	0.842 (0.207-1.504) B	0.618±0.136
	<i>R. luteolus</i>	2.397 (1.460-3.141) C	0.725±0.133
Water <sup>a</sup>	<i>T. fracticum</i>	12.337 (10.619-15.147) N	1.629±0.156
	<i>L. tigrinus</i>	9.513 (8.136-11.841) G	1.193±0.140
	<i>C. comatus</i>	11.534 (10.052-13.875) L	1.681±0.156
	<i>R. luteolus</i>	9.596 (8.712-10.818) I	2.082±0.163
EDTA <sup>b</sup>	–	0.158 (0.135-0.183) A	1.207±0.064

<sup>a</sup>Extract, <sup>b</sup>Standard solution, <sup>c</sup>Values followed by different capital letters in the same column differ significantly at *P*<0.05. EDTA: Ethylenediaminetetraacetic acid, *T. fracticum*: *Tricholoma fracticum*, *L. tigrinus*: *Lentinus tigrinus*, *C. comatus*: *Coprinus comatus*, *R. luteolus*: *Rhizopogon luteolus*, SEM: Standard error of mean, IC<sub>50</sub>: 50% inhibition



**Table 4: EC<sub>50</sub> values (mg/ml) of different extracts from four mushrooms and standard for reducing power**


Treatment	Species	EC <sub>50</sub>
Acetone <sup>a</sup>	<i>T. fracticum</i>	5.757 K <sup>c</sup>
	<i>L. tigrinus</i>	15.185 T
	<i>C. comatus</i>	4.429 G
	<i>R. luteolus</i>	1.754 B
Chloroform <sup>a</sup>	<i>T. fracticum</i>	6.899 N
	<i>L. tigrinus</i>	11.368 R
	<i>C. comatus</i>	4.045 E
	<i>R. luteolus</i>	6.667 M
<i>n</i> -Hexane <sup>a</sup>	<i>T. fracticum</i>	8.493 O
	<i>L. tigrinus</i>	5.664 J
	<i>C. comatus</i>	8.722 P
	<i>R. luteolus</i>	4.869 H
Methanol <sup>a</sup>	<i>T. fracticum</i>	6.624 L
	<i>L. tigrinus</i>	5.498 I
	<i>C. comatus</i>	2.541 D
	<i>R. luteolus</i>	4.104 F
Water <sup>a</sup>	<i>T. fracticum</i>	13.011 S
	<i>L. tigrinus</i>	9.850 Q
	<i>C. comatus</i>	17.139 U
	<i>R. luteolus</i>	2.468 C
Gallic acid <sup>b</sup>	-	0.365 A

<sup>a</sup>Extract, <sup>b</sup>Standard solution, <sup>c</sup>Values followed by different capital letters in the same column differ significantly at  $P < 0.05$ . *T. fracticum*: *Tricholoma fracticum*, *L. tigrinus*: *Lentinus tigrinus*, *C. comatus*: *Coprinus comatus*, *R. luteolus*: *Rhizopogon luteolus*, EC<sub>50</sub>: Effective concentrations

extracts. Whereas in *n*-hexane and water extracts, *T. fracticum* had minimum reducing power. Similarly, in methanol extract, lower reducing power were determined in *L. tigrinus* and *T. fracticum*.


#### Amount of bioactive components working as antioxidants

Significant differences between the chemical composition of wild and commercial edible mushrooms have been reported.<sup>[6]</sup> Natural antioxidants derived from macrofungi are of considerable interest since these compounds have important constituents with free radical scavenging ability and hence may contribute directly to the antioxidative action. Among them, phenolic compounds, flavonoids,  $\beta$ -carotene and lycopene contents correlates well with the antioxidant potential of the edible mushrooms. Numerous studies carried out by scientists with some mushrooms demonstrated that these materials had antioxidant components such as  $\beta$ -carotene, flavonoid, lycopene and phenol.<sup>[31,38,41,52,54,60-62,64,65]</sup> Previous studies also revealed that many other edible mushrooms such as *Agaricus bisporus*,<sup>[60,62]</sup> *Amanita vaginata*,<sup>[66]</sup> *Grifola frondosa*,<sup>[61]</sup> *P. eryngii*,<sup>[46]</sup> *P. ostreatus*,<sup>[31,41]</sup> *Russula delica*<sup>[60]</sup> and *S. bovinus*<sup>[49]</sup> had high rate of phenol, flavonoids, ascorbic acid,  $\beta$ -carotene and lycopene. For that reason, various extracts from *T. fracticum*, *L. tigrinus*, *C. comatus* and *R. luteolus* were analyzed



**Figure 2:** Chelating abilities of extracts in different concentrations from four mushrooms on ferrous ions





**Figure 3:** Reducing powers of various extracts from four mushrooms

for their antioxidant components including  $\beta$ -carotene, flavonoids, lycopene and total phenols.

As for the total phenolic contents, highest quantities were found in chloroform and acetone extract of *C. comatus* (76.32 and 61.09  $\mu\text{g}/\text{mg}$ , respectively). At the same time, considerable phenolic content was determined in water extract of *L. tigrinus* (46.76  $\mu\text{g}/\text{mg}$ ). The total phenolic content in other mushroom extracts was lower than 37.00  $\mu\text{g}/\text{mg}$ .

Considering flavonoid content, *n*-hexane and water extract of *L. tigrinus* showed the maximum amounts (3.67 and 3.22  $\mu\text{g}/\text{mg}$  respectively). The flavonoid content in other mushroom extracts was lower than 3.00  $\mu\text{g}/\text{mg}$ .

Among the mushroom extracts, chloroform extract of *C. comatus* had the highest  $\beta$ -carotene content (25.94  $\mu\text{g}/\text{mg}$ ). Furthermore acetone and methanolic extracts of this mushroom had higher  $\beta$ -carotene level than the other edible mushrooms (23.32 and 6.55  $\mu\text{g}/\text{mg}$ , respectively), *n*-hexane and water extracts of *L. tigrinus* (17.28 and 13.13  $\mu\text{g}/\text{mg}$ , respectively) had also considerable amount.

Chloroform and acetone extract of *C. comatus* revealed higher lycopene content (12.00 and 9.16  $\mu\text{g}/\text{mg}$ , respectively) as compared with other mushroom extracts. However, lycopene contents in other mushroom extracts were found in lower amounts (0.96-6.25  $\mu\text{g}/\text{mg}$ ) [Table 5].

In this study, it was also determined that there is a positive correlation between antioxidant activity and bioactive constituents of tested edible mushrooms [Table 6]. Our results suggested that phenolic acids and flavonoids may be the second major contributors for the antioxidant activity as the  $\text{IC}_{50}$  values of radical scavenging activity of various soluble fractions of edible mushrooms and also the contents of phenolics or flavonoids exhibited significant correlation. Present study also revealed that DPPH radical scavenging activity was in significant correlation with the amount of phenolics, lycopene, flavonoids and  $\beta$ -carotene. Furthermore, reducing capacity of edible mushrooms was in positive correlation with the amount of flavonoids. Therefore, many foods that have a high rate of phenolics or flavonoids can be used for the purpose of medical therapies.

#### Antibacterial activities of different extracts

Mushrooms are rich sources of natural antibiotics, and many of the secondary metabolites may combat with many bacteria and viruses.<sup>[67]</sup> With an increasing number of bacteria developing resistance to commercial antibiotics, extracts and derivatives from mushrooms hold great promise for novel medicines. Therefore, antibacterial activity of tested edible mushrooms were analyzed by disc diffusion method on two Gram-negative (*E. coli* and *A. tumefaciens*) and three Gram-positive (*B. subtilis*, *B. licheniformis*, *S. aureus*) bacteria. All four tested mushrooms showed antibacterial activity at least on one

**Table 5: Antioxidant contents of different extracts from four edible mushrooms**

Content (µg/mg)	Species	Extract <sup>a</sup>				
		Acetone	Chloroform	n-Hexane	Methanol	Water
β-Carotene	<i>T. fracticum</i>	18.70±0.78	18.40±1.18	14.70±0.25	7.03±0.19	12.11±0.18
	<i>L. tigrinus</i>	2.58±1.36	9.41±2.58	17.28±1.10	2.71±0.08	13.13±0.15
	<i>C. comatus</i>	23.32±0.28	25.94±0.63	4.80±0.85	6.55±0.30	12.87±0.94
	<i>R. luteolus</i>	8.02±0.19	6.11±0.10	3.42±0.81	4.85±0.20	5.41±0.19
Flavonoid	<i>T. fracticum</i>	1.51±0.05	1.36±0.17	2.17±0.10	1.60±0.24	1.93±0.01
	<i>L. tigrinus</i>	0.41±0.07	0.94±0.02	3.67±0.05	1.16±0.13	3.22±0.01
	<i>C. comatus</i>	0.88±0.04	2.88±0.09	0.27±0.05	2.33±0.14	0.72±0.01
	<i>R. luteolus</i>	2.28±0.20	0.63±0.08	1.23±0.05	2.99±0.07	1.62±0.26
Total phenols	<i>T. fracticum</i>	32.72±0.12	30.36±0.34	5.93±0.04	2.52±0.38	36.81±0.09
	<i>L. tigrinus</i>	18.66±0.21	24.74±0.07	19.36±0.29	3.81±0.13	46.76±1.19
	<i>C. comatus</i>	61.09±0.32	76.32±1.11	5.02±0.01	8.40±0.28	32.37±16.19
	<i>R. luteolus</i>	12.16±0.72	7.70±0.60	8.03±0.19	4.13±0.23	6.69±0.18
Lycopene	<i>T. fracticum</i>	6.12±0.37	6.25±0.50	5.94±0.02	5.69±0.27	3.37±0.18
	<i>L. tigrinus</i>	1.06±0.21	3.52±1.68	3.43±2.14	1.51±0.05	4.21±0.08
	<i>C. comatus</i>	9.16±0.32	12.00±0.48	1.99±0.11	3.87±0.01	5.41±0.38
	<i>R. luteolus</i>	4.44±0.11	1.08±0.03	1.77±1.33	3.11±0.06	0.96±0.03

<sup>a</sup>Each value is expressed as mean±SEM (n=3). *T. fracticum*: *Tricholoma fracticum*; *L. tigrinus*: *Lentinus tigrinus*; *C. comatus*: *Coprinus comatus*; *R. luteolus*: *Rhizopogon luteolus*; SEM: Standard error of mean

**Table 6: Correlations between amounts of bioactive components and antioxidant activities**

Correlation	Pearson correlation coefficient				
	Acetone <sup>b</sup>	Chloroform <sup>b</sup>	n-Hexane <sup>b</sup>	Methanol <sup>b</sup>	Water <sup>b</sup>
DPPH activity	0.287	0.984 <sup>a</sup>	-0.687	0.955 <sup>a</sup>	-0.522
β-Carotene					
DPPH activity	0.751	0.979 <sup>a</sup>	-0.306	0.574	-0.934
Flavonoid					
DPPH activity	0.087	0.954 <sup>a</sup>	-0.083	0.372	-0.767
Total phenols					
DPPH activity	0.514	0.983 <sup>a</sup>	-0.680	0.827	-0.218
Lycopene					
Metal chelating activity	0.335	0.956 <sup>a</sup>	0.760	0.731	-0.533
β-Carotene					
Metal chelating activity	-0.693	0.952 <sup>a</sup>	0.604	0.806	0.714
Flavonoid					
Metal chelating activity	0.446	0.970 <sup>a</sup>	0.113	0.649	-0.247
Total phenols					
Metal chelating activity	0.087	0.982 <sup>a</sup>	0.965 <sup>a</sup>	0.484	-0.692
Lycopene					
Reducing power activity	0.062	0.858	-0.330	0.335	-0.980 <sup>a</sup>
β-Carotene					
Reducing power activity	0.961 <sup>a</sup>	0.956 <sup>a</sup>	0.126	0.586	-0.075
Flavonoid					
Reducing power activity	-0.285	0.924	0.439	0.968 <sup>a</sup>	-0.898
Total phenols					
Reducing power activity	0.246	0.902	-0.677	-0.021	-0.897
Lycopene					

<sup>a</sup>Correlation is significant at the 0.05 level; <sup>b</sup>Extract. DPPH: 2,2-diphenyl-1-picrylhydrazyl

microorganism. On the other hand, 10 out of 20 extracts (five different extracts of each four mushrooms, total 20 extracts) had stronger antimicrobial potential. Table 7 summarizes the antibacterial effects of edible mushroom extracts. Accordingly, the largest IZ was measured as 13 mm with chloroform extract of *R. luteolus* on *B. subtilis*. Besides, wide range of activity was observed with

*n*-hexane extract of *L. tigrinus*. It inhibited the growth of entire microorganism used in this survey, including *E. coli*, because, none of the other extracts did inhibit the growth of *E. coli*. Strong antibacterial activity of *L. tigrinus* was followed by *T. fracticum* in such that its acetone extract showed antimicrobial activity against all microorganisms other than *E. coli*. The results revealed that for all tested

**Table 7: Antibacterial activities of different extracts obtained from four mushrooms**

Species	Extract	<i>Escherichia coli</i>	<i>Bacillus subtilis</i>	<i>Bacillus licheniformis</i>	<i>Staphylococcus aureus</i>	<i>Agrobacterium tumefaciens</i>
<i>Tricholoma fracticum</i> (IZ; mm)	Water	-	-	6	-	-
	<i>n</i> -Hexane	-	6	8	-	8
	Chloroform	-	6	6	-	-
	Acetone	-	6	6	9	6
	Methanol	-	-	-	-	-
<i>Lentinus tigrinus</i> (IZ; mm)	Water	-	8	6	-	-
	<i>n</i> -Hexane	6	6	8	6	8
	Chloroform	-	-	-	-	-
	Acetone	-	-	-	-	-
	Methanol	-	-	-	-	-
<i>Coprinus comatus</i> (IZ; mm)	Water	-	-	6	-	-
	<i>n</i> -Hexane	-	-	-	-	-
	Chloroform	-	-	-	-	-
	Acetone	-	-	-	-	-
	Methanol	-	-	-	-	-
<i>Rhizopogon luteolus</i> (IZ; mm)	Water	-	-	-	-	-
	<i>n</i> -Hexane	-	-	6	-	6
	Chloroform	-	13	6	-	6
	Acetone	-	-	8	6	7
	Methanol	-	-	-	-	-

IZ: Inhibition zones

extracts *Bacillus* sp. were more susceptible, since at least one extract from this mushroom showed antibacterial activity. Moreover, water extract of *C. comatus* has antibacterial activity against *B. licheniformis*. Indeed it is well known that Gram-negative microorganism more resistant to antibacterial than Gram-positives.

This study provided consistent result with earlier reports that mushrooms produce wide variety of antimicrobial compounds.<sup>[68,69]</sup> It was previously reported that *C. comatus* had antimicrobial activity against *S. aureus* and *B. subtilis*<sup>[70]</sup> whereas in this study it showed its activity against *B. licheniformis*. This discrepancy could result from cultivation from the different environment because climate, environment and the soil type can effect chemical composition of the mushrooms. On the other hand, similar result was obtained with *L. tigrinus* that it showed a wide range of antimicrobial activity.

## CONCLUSION

Edible macrofungi have gained increasing reputation in recent decades, not only due to their nutritive value, but also for their possible health benefits and therapeutic uses. Therefore, wild beneficial sources bring new natural products into the pharmaceutical industry. Our results on four edible mushrooms elucidates that, *C. comatus* showed substantial *in vitro* cytotoxic activity against HepG2 cell lines that was in parallel with its high  $\beta$ -carotene, total phenolic and lycopene contents. High antioxidant potential of *C. comatus*

obtained from highest DPPH capturing, and metal chelating activity was in significant positive correlation with the abovementioned bioactive ingredients. As for antibacterial activity, *L. tigrinus* and *T. fracticum* were the most active against tested microorganism. These results indicate that different extracts of investigated edible mushrooms have considerable cytotoxic, antioxidant and antibacterial properties and may be utilized as a promising source of therapeutics. Therefore, edible mushrooms might provide an appropriate source of antioxidant and cytotoxic natural compounds and could also be searched as a potent antibacterial drug against infectious diseases. *C. comatus* deserves further studies due to its incredible cytotoxic and antioxidant potential, and it can be a promising mushroom that would be benefitted in the field of pharmacology with safer and better effectiveness.

## ACKNOWLEDGMENT

The authors are grateful to Karamanoğlu Mehmetbey University Scientific Research Fund for financial support (37-M-12).

## REFERENCES

- Cheung LM, Cheung PC, Ooi VE. Antioxidant activity and total phenolics of edible mushroom extracts. *Food Chem* 2003;81:249-55.
- Cheung PC. The nutritional and health benefits of mushrooms. *Nutr Bull* 2010;35:292-9.
- Beluhan S, Ranogajec A. Chemical composition and non-volatile components of croatian wild edible mushrooms. *Food Chem* 2011;124:1076-82.

4. Barros L, Baptista P, Correia D, Casal S, Oliveira B, Ferreira I. Fatty acid and sugar compositions, and nutritional value of five wild edible mushrooms from Northeast Portugal. *Food Chem* 2007;105:140-5.
5. Reis FS, Barros L, Martins A, Ferreira IC. Chemical composition and nutritional value of the most widely appreciated cultivated mushrooms: An inter-species comparative study. *Food Chem Toxicol* 2012;50:191-7.
6. Barros L, Cruz T, Baptista P, Estevinho LM, Ferreira IC. Wild and commercial mushrooms as source of nutrients and nutraceuticals. *Food Chem Toxicol* 2008;46:2742-7.
7. Kalač P. Chemical composition and nutritional value of European species of wild growing mushrooms: A review. *Food Chem* 2009;113:9-16.
8. Espín JC, García-Conesa MT, Tomás-Barberán FA. Nutraceuticals: Facts and fiction. *Phytochemistry* 2007;68:2986-3008.
9. Fan L, Pan H, Soccol AT, Pandey A, Soccol CR. Advances in mushroom research in the last decade. *Food Technol Biotechnol* 2006;44:303-11.
10. Palacios I, Lozano M, Moro C, D'Arrigo M, Rostagno MA, Martínez JA, *et al.* Antioxidant properties of phenolic compounds occurring in edible mushrooms. *Food Chem* 2011;128:674-8.
11. Ramesh Ch, Pattar MG. Antimicrobial properties, antioxidant activity and bioactive compounds from six wild edible mushrooms of western ghats of Karnataka, India. *Pharmacognosy Res* 2010;2:107-12.
12. Wong JY, Chye FY. Antioxidant properties of selected tropical wild edible mushrooms. *J Food Compos Anal* 2009;22:269-77.
13. Kotan E, Alpsoy L, Anar M, Aslan A, Agar G. Protective role of methanol extract of *Cetraria islandica* (L.) against oxidative stress and genotoxic effects of AFB1 in human lymphocytes *in vitro*. *Toxicol Ind Health* 2011;27:599-605.
14. Ceker S, Agar G, Nardemir G, Anar M, Kizil HE, Alpsoy L. Investigation of anti-oxidative and anti-genotoxic effects of *Origanum vulgare* L. essential oil on human lymphocytes *in vitro*. *J Essent Oil Bear Plants* 2012;15:997-1005.
15. Türkez H, Togar B. Olive (*Olea europaea* L.) leaf extract counteracts genotoxicity and oxidative stress of permethrin in human lymphocytes. *J Toxicol Sci* 2011;36:531-7.
16. Turkez H, Togar B, Polat E. Olive leaf extract modulates permethrin induced genetic and oxidative damage in rats. *Cytotechnology* 2012;64:459-64.
17. Türkez H, Togar B. Aluminium phosphide-induced genetic and oxidative damages *in vitro*: Attenuation by *Laurus nobilis* L. leaf extract. *Indian J Pharmacol* 2013;45:71-5.
18. Breitenbach J, Kränzlin F. Fungi of Switzerland. Vol. 1-5. Lucerne: Verlag Mykologia;1984-2000.
19. Abe K, Matsuki N. Measurement of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction activity and lactate dehydrogenase release using MTT. *Neurosci Res* 2000;38:325-9.
20. Taga MS, Miller EE, Pratt DE. Chia seeds as a source of natural lipid antioxidants. *J Am Oil Chem Soc* 1984;61:928-31.
21. Pal J, Ganguly S, Tahsin KS, Acharya K. *In vitro* free radical scavenging activity of wild edible mushroom, *Pleurotus squarrosulus* (Mont.) Singer. *Indian J Exp Biol* 2010;48:1210-8.
22. Turkoglu A, Duru ME, Mercan N, Kivrak I, Gezer K. Antioxidant and antimicrobial activities of *Laetiporus sulphureus* (Bull.) Murrill. *Food Chem* 2007;101:267-73.
23. Oyetayo VO, Dong CH, Yao YJ. Antioxidant and antimicrobial properties of aqueous extract from *Dictyophora indusiata*. *Open Mycol J* 2009;3:20-6.
24. Karamac M, Amarowicz R, Weidner S, Abe S, Shahidi F. Antioxidant activity of rye caryopses and embryos extracts. *Czech J Food Sci* 2000;20:209-14.
25. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 1966;45:493-6.
26. Costantini S, Colonna G, Castello G. A holistic approach to study the effects of natural antioxidants on inflammation and liver cancer. *Cancer Treat Res* 2014;159:311-23.
27. Fernandes Â, Barreira JC, Antonio AL, Oliveira MB, Martins A, Ferreira IC. Effects of gamma irradiation on chemical composition and antioxidant potential of processed samples of the wild mushroom *Macrolepiota procera*. *Food Chem* 2014;149:91-8.
28. Ito T, Urushima H, Sakaue M, Yukawa S, Honda H, Hirai K, *et al.* Reduction of adverse effects by a mushroom product, active hexose correlated compound (AHCC) in patients with advanced cancer during chemotherapy – The significance of the levels of HHV-6 DNA in saliva as a surrogate biomarker during chemotherapy. *Nutr Cancer* 2014;66:377-82.
29. Kim SP, Nam SH, Friedman M. *Hericium erinaceus* (Lion's Mane) mushroom extracts inhibit metastasis of cancer cells to the lung in CT-26 colon cancer-transplanted mice. *J Agric Food Chem* 2013;61:4898-904.
30. Wang S, Zhu F, Meckling KA, Marcone MF. Antioxidant capacity of food mixtures is not correlated with their antiproliferative activity against MCF-7 breast cancer cells. *J Med Food* 2013;16:1138-45.
31. Fontes Vieira PA, Gontijo DC, Vieira BC, Fontes EA, de Assunção LS, Leite JP, *et al.* Antioxidant activities, total phenolics and metal contents in *Pleurotus ostreatus* mushrooms enriched with iron, zinc or lithium. *LWT Food Sci Technol* 2013;54:421-5.
32. Alves MJ, Froufe HJ, Costa AF, Santos AF, Oliveira LG, Osório SR, *et al.* Docking studies in target proteins involved in antibacterial action mechanisms: Extending the knowledge on standard antibiotics to antimicrobial mushroom compounds. *Molecules* 2014;19:1672-84.
33. Petrovic J, Glamoclija J, Stojkovic DS, Ciric A, Nikolic M, Bukvicki D, *et al.* *Laetiporus sulphureus*, edible mushroom from Serbia: Investigation on volatile compounds, *in vitro* antimicrobial activity and *in situ* control of *Aspergillus flavus* in tomato paste. *Food Chem Toxicol* 2013;59:297-302.
34. Vázquez-Velasco M, González-Torres L, López-Gasco P, Bastida S, Benedí J, Sánchez-Reus MI, *et al.* Liver oxidation and inflammation in Fa/Fa rats fed glucomannan/spirulina-surimi. *Food Chem* 2014;159:215-21.
35. Tavakol HS, Farzad K, Fariba M, Abdolkarim C, Hassan G, Seyed-Mostafa HZ, *et al.* Hepatoprotective effect of *Matricaria chamomilla* L. in paraquat induced rat liver injury. *Drug Res (Stuttg)* 2015;65:61-4.
36. Nichols JA, Katiyar SK. Skin photoprotection by natural polyphenols: Anti-inflammatory, antioxidant and DNA repair mechanisms. *Arch Dermatol Res* 2010;302:71-83.
37. Christudas IV, Kumar PP, Sunil C, Vajravijayan S, Sundaram RL, Siril SJ, *et al.* *In vitro* studies on  $\alpha$ -glucosidase inhibition, antioxidant and free radical scavenging activities of *Hedyotis biflora* L. *Food Chem* 2013;138:1689-95.
38. Dasgupta A, Rai M, Acharya K. Chemical composition and antioxidant activity of a wild edible mushroom *Pleurotus flabellatus*. *Int J PharmTech Res* 2013;5:1655-63.
39. Dulay RM, Arenas MC, Kalaw SP, Reyes RG, Cabrera EC. Proximate composition and functionality of the culinary-medicinal tiger sawgill mushroom, *Lentinus tigrinus* (higher basidiomycetes), from the Philippines. *Int J Med Mushrooms* 2014;16:85-94.



40. Lung MY, Huang WZ. Antioxidant potential and antioxidant compounds of extracts from the medicinal sulphur polypore, *Laetiporus sulphureus* (Higher basidiomycetes) in submerged cultures. *Int J Med Mushrooms* 2013;15:569-82.
41. Ejelonu OC, Akinmoladun AC, Elekofehinti OO, Olaleye MT. Antioxidant profile of four selected wild edible mushrooms in Nigeria. *J Chem Pharm Res* 2013;5:286-95.
42. Lee MR, Hou JG, Begum S, Xue JJ, Wang YB, Sung CK. Comparison of constituents, antioxidant potency, and acetylcholinesterase inhibition in *Lentinus edodes*, *Sparassis crispa*, and *Mycoleptodonoides aitchisonii*. *Food Sci Biotechnol* 2013;22:1747-51.
43. Kim MJ, Chu WM, Park EJ. Antioxidant and antigenotoxic effects of shiitake mushrooms affected by different drying methods. *J Korean Soc Food Sci Nutr* 2012;41:1041-8.
44. Lee J, Hong JH, Kim JD, Ahn BJ, Kim BS, Kim GH, *et al.* The antioxidant properties of solid-culture extracts of basidiomycetous fungi. *J Gen Appl Microbiol* 2013;59:279-85.
45. Hasnat MA, Pervin M, Debnath T, Lim BO. DNA protection, Total phenolics and antioxidant potential of the mushroom *Russula virescens*. *J Food Biochem* 2014;38:6-17.
46. Yildirim NC, Turkoglu S, Yildirim N, Ince OK. Antioxidant properties of wild edible mushroom *Pleurotus eryngii* collected from Tunceli province of Turkey. *Dig J Nanomater Biostruct* 2012;7:1647-54.
47. Fernandes Â, Barros L, Barreira JC, Antonio AL, Oliveira MB, Martins A, *et al.* Effects of different processing technologies on chemical and antioxidant parameters of *Macrolepiota procera* wild mushroom. *LWT Food Sci Technol* 2013;54:493-9.
48. Khatua S, Roy T, Acharya K. Antioxidant and free radical scavenging capacity of phenolic extract from *Russula laurocerasi*. *Asian J Pharm Clin Res* 2013;6:156-60.
49. Pereira E, Oliveira I, Baptista P. Guttation droplets of the edible mushroom *Suillus bovinus* as a new source of natural antioxidants. *Sci Hortic (Amsterdam)* 2012;148:89-92.
50. Ayala-Zavala JF, Perez-Carlon JJ, Esqueda M, Gonzalez-Aguilar GA, Leyva JM, Cruz-Valenzuela MR, *et al.* Polar fractionation affects the antioxidant properties of methanolic extracts from species of genus *Phellinus* quel.(higher Basidiomycetes). *Int J Med Mushrooms* 2012;14:563-73.
51. Liu K, Wang JL, Gong WZ, Xiao X, Wang Q. Antioxidant activities *in vitro* of ethanol extract and fractions from mushroom, *Lenzites betulina*. *J Food Biochem* 2013;37:687-93.
52. Barros L, Baptista P, Ferreira IC. Effect of *Lactarius piperatus* fruiting body maturity stage on antioxidant activity measured by several biochemical assays. *Food Chem Toxicol* 2007;45:1731-7.
53. Reis FS, Barros L, Calhela RC, Ciric A, van Griensven LJ, Sokovic M, *et al.* The methanolic extract of *Cordyceps militaris* (L.) Link fruiting body shows antioxidant, antibacterial, antifungal and antihuman tumor cell lines properties. *Food Chem Toxicol* 2013;62:91-8.
54. Ferreira IC, Baptista P, Vilas-boas M, Barros L. Free-radical scavenging capacity and reducing power of wild edible mushrooms from northeast Portugal: Individual cap and stipe activity. *Food Chem* 2007;100:1511-6.
55. Zhang Y, Liu Z, Ng TB, Chen Z, Qiao W, Liu F. Purification and characterization of a novel antitumor protein with antioxidant and deoxyribonuclease activity from edible mushroom *Pholiota nameko*. *Biochimie* 2014;99:28-37.
56. Soltaninejad K, Kebriaeezadeh A, Minaiee B, Ostad SN, Hosseini R, Azizi E, *et al.* Biochemical and ultrastructural evidences for toxicity of lead through free radicals in rat brain. *Hum Exp Toxicol* 2003;22:417-23.
57. Wong SP, Leong LP, Hoe J, Koh W. Antioxidant activities of aqueous extracts of selected plants. *Food Chem* 2006;99:775-83.
58. Liu Y, Du YQ, Wang JH, Zha XQ, Zhang JB. Structural analysis and antioxidant activities of polysaccharide isolated from Jinqian mushroom. *Int J Biol Macromol* 2014;64:63-8.
59. Nandi AK, Samanta S, Maity S, Sen IK, Khatua S, Devi KS, *et al.* Antioxidant and immunostimulant  $\beta$ -glucan from edible mushroom *Russula albonigra* (Krombh.) Fr. *Carbohydr Polym* 2014;99:774-82.
60. Elmastas M, Isildak O, Turkekul I, Temur N. Determination of antioxidant activity and antioxidant compounds in wild edible mushrooms. *J Food Compos Anal* 2014;20:337-45.
61. Yeh JY, Hsieh LH, Wu KT, Tsai CF. Antioxidant properties and antioxidant compounds of various extracts from the edible basidiomycete *Grifola frondosa* (Maitake). *Molecules* 2011;16:3197-211.
62. Ji H, Du A, Zhang L, Li S, Yang M, Li B. Effects of drying methods on antioxidant properties and phenolic content in white button mushroom. *Int J Food Eng* 2012;8.
63. Meir S, Kanner J, Akiri B, Philosoph-Hadas S. Determination and involvement of aqueous reducing compounds in oxidative defense systems of various senescing leaves. *J Agric Food Chem* 1995;43:1813-9.
64. Barros L, Falcão S, Baptista P, Freire C, Vilas-boas M, Ferreira IC. Antioxidant activity of *Agaricus* sp. mushrooms by chemical, biochemical and electrochemical assays. *Food Chem* 2008;111:61-6.
65. Sulkowska-Ziaja K, Muszynska B, Motyl P, Pasko P, Ekiert H. Phenolic compounds and antioxidant activity in some species of polyporoid mushrooms from Poland. *Int J Med Mushrooms* 2012;14:385-93.
66. Paloi S, Acharya K. Antioxidant activities and bioactive compounds of polyphenol rich extract from *Amanita vaginata* (Bull.) Lam. *Int J PharmTech Res* 2013;5:1645-54.
67. Barros L, Calhela RC, Vaz JA, Ferreira IC, Baptista P, Estevinho LM. Antimicrobial activity and bioactive compounds of Portuguese wild edible mushrooms methanolic extracts. *Eur Food Res Technol* 2006;225:151-6.
68. Yamaç M, Bilgili F. Antimicrobial activities of fruit bodies and/or mycelial cultures of some mushroom isolates. *Pharm Biol* 2006;44:660-7.
69. Ranadive KR, Belsare MH, Deokule SS, Jagtap NV, Jadhav HK, Vaidya JG. Glimpses of antimicrobial activity of fungi from World. *J New Biol Rep* 2013;2:142-62.
70. Dyakov MY, Kamzolkina OV, Shtaer OV, Bisko NA, Poedinok NL, Mikhailova OB, *et al.* Morphological characteristics of natural strains of certain species of basidiomycetes and biological analysis of antimicrobial activity under submerged cultural conditions. *Microbiology* 2011;80:274-85.

**Cite this article as:** Sadi G, Emsen B, Kaya A, Kocabas A, Çınar S, Kartal DI. Cytotoxicity of some edible mushrooms extracts over liver hepatocellular carcinoma cells in conjunction with their antioxidant and antibacterial properties. *Phcog Mag* 2015;11:6-18.

**Source of Support:** The authors are grateful to Karamanoğlu Mehmetbey University Scientific Research Fund for financial support (37-M-12), **Conflict of Interest:** None declared.