

MEETING ABSTRACT

Open Access

Sentinel node biopsy and radical lymph node dissection for advanced melanoma in the elderly

V Desiato*, S Perrotta, GL Benassai, G Quarto, G Benassai, G Limite

From XXIII Annual Meeting of the Italian Society of Geriatric Surgery
Lecce, Italy. 2-4 December 2010

Background

The majority of indications for surgery in melanoma are for the treatment of primary tumor and lymph node metastases. During the last decade, the Sentinel Node Biopsy (SNB), from a research procedure, has become standard of care in most institutions. SNB is normally considered for patients with melanoma > 1 mm and generally about 20% are positive; however, the risk of a positive SNB in a melanoma < 1 mm is still 5%. Usually when SNB is positive a complete lymphadenectomy is performed.

Materials and methods

In the period 2004-2009, 18 elderly patients (median age 68 years) affected by cutaneous melanoma (mean Breslow's thickness = 3.77 mm), after SNB histologically confirmed regional lymph node involvement, underwent complete lymph node dissection (CLND). We treated 11 of them with groin dissection, in 3 cases bilateral; 4 patients underwent axillar dissection, in one case bilateral; 2 patients underwent neck dissection and another patient underwent groin-axillar dissection. We treated bilateral groin involvement with laparoscopic access for dissection of lumbar-aortic, iliac and obturator lymph nodes.

Results

Disagreeing with literature, 12/18 (67%) of these patients had positive lymph nodes, a high percentage if compared with younger patients' data. Currently the average follow-up is 25 months. In our sample CLND has a crucial prognostic role (16% vs 41% of deceased in CLND - and CLND + patients respectively).

* Correspondence: vin_des@hotmail.it

Dipartimento Universitario di Chirurgia Generale, Geriatrica, Oncologica e Tecnologie Avanzate Facoltà di Medicina e Chirurgia, Università degli Studi di Napoli Federico II, Naples, Italy

Conclusions

Elderly melanoma patients are characterized by a higher tumor stage and, in patients with nodal metastases, the prognosis is independently affected by older age. In case of positive SNB the CLND plays a notable prognostic role and a presumable therapeutic role.

Published: 24 August 2011

References

1. Marsden JR, Newton-Bishop JA, et al: Revised U.K. guidelines for the management of cutaneous melanoma 2010. *Br J Dermatol* 2010, **163**(2):238-56.
2. Testori A, Stanganelli I, et al: Diagnosis of melanoma in the elderly and surgical implications. *Surg Oncol* 2004, **13**(4):211-21.
3. Kruijff S, Bastiaannet E, et al: Detection of Melanoma Nodal Metastases; Differences in Detection Between Elderly and Younger Patients Do Not Affect Survival. *Ann Surg Oncol* 2010, **17**:3008-3014.

doi:10.1186/1471-2318-11-S1-A9

Cite this article as: Desiato et al: Sentinel node biopsy and radical lymph node dissection for advanced melanoma in the elderly. *BMC Geriatrics* 2011 **11**(Suppl 1):A9.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

