


Perception and Experience of Dupilumab in Atopic Dermatitis: A Real-Life Study

Léa Antoine¹, Eve Puzenat¹, Dragos Popescu², Romain Charollais³, Flora Dresco³, Anne-Sophie Dupond², Dominique Salard⁴, Marie-Christine Drobacheff-Thiebaut¹, Anaïs Zanella¹, Anne Ducournau², Irène Gallais-Serezal¹, François Aubin^{1,5} 

¹Department of Dermatology, University Hospital, Besançon, France; ²Department of Dermatology, Hôpital Nord Franche Comté, Trevenans, France; ³Department of Dermatology, Centre Hospitalier, Vesoul, France; ⁴Department of Dermatology, Centre Hospitalier, Pontarlier, France; ⁵Inserm 1098 RIGHT, University of Franche Comté, Besançon, France

Correspondence: François Aubin, Centre hospitalier Universitaire, Service de Dermatologie, 3 Bd Fleming, Besançon, 25000, France, Email francois.aubin@univ-fcomte.fr

Purpose: There are few data on the practical use of dupilumab by the patients and on the patients' experience with this treatment.

Objective: The objective of our study was to describe the experience and perception of dupilumab treatment in patients with atopic dermatitis (AD).

Patients and Methods: We conducted a multicenter retrospective observational study including adult patients with moderate to severe AD treated with dupilumab between January 2017 and December 2021. Clinical characteristics were collected and a questionnaire was sent to all patients. It consisted of different parts including the injection method and different numeric rating scales (NRS) evaluating the patient's satisfaction and the constraints related to the treatment.

Results: Eighty-two patients were included and the information was available for 77 patients who responded to the questionnaire. Injection of dupilumab was performed by a nurse in 47% (n=36) of patients and 43% (n=33) were autonomous. Injections were performed by a family member for 7 patients or by the general practitioner (1 patient). A wearing-off of the beneficial effect of dupilumab was reported by 47% of patients leading to shorten the dosing interval. In contrast, dose spacing was reported by 9 patients (11%). After a mean follow-up time of 29.7 ± 10.7 months (median: 27 months), drug survival was 72%. From the patients' perspective, the mean patient's satisfaction NRS score was 7.5 ± 1.8, and the constraints related to the treatment were scored at 3.1 ± 2.1 on NRS.

Conclusion: Although AD treatments may contribute to the burden of the disease, dupilumab was associated with a lower burden score, likely reflecting both treatment efficacy and easy of use and patient satisfaction.

Keywords: atopic dermatitis, dupilumab, real life, experience

Introduction

Real-world data on the long-term data of dupilumab are increasing and are reassuring in terms of both efficacy and safety.¹⁻⁴

However, few data exist on the practical use of dupilumab by the patients and on the patient experience with this treatment.⁵ The objective of our study was to describe the experience and perception of dupilumab treatment in patients with AD.

Patients and Methods

Study Design and Patients

We conducted a multicenter retrospective observational study in the 4 dermatology units of the Franche Comté, a well-defined administrative area located in eastern France. After informed consent (Comité de Protection des Personnes, CHU Besançon, 2022-015), all adult patients with moderate to severe AD treated with dupilumab between January 2017 and December 2021 were included.

Outcomes

Data were collected on computerized records and included: demographics, comorbidities, treatment history, side effects, associated treatments, and clinical severity scores: the scoring AD (SCORAD), the Dermatology Life Quality Index (DLQI), and the pruritus and sleep 11-points numeric rating scales (NRS). In addition, a questionnaire ([Supplementary File](#)) was sent to all patients. It consisted of different parts including the injection method, the management of adverse events, and different NRS evaluating the patient's satisfaction, the constraints related to the treatment, the quality of medical information, the patients' apprehension to dupilumab and the reasons for a possible discontinuation of the treatment.

Analysis

Statistical analysis was performed with Excel software (Microsoft Office, France). Continuous variables were expressed as mean \pm standard deviation (SD) and categorical variables as number (percentage).

Results

Demographic Data

Eighty-two patients were included. The mean age was 39.4 ± 13 years with a M/F sex ratio of 1.5. AD had started in childhood in 83% of the patients. Atopic background was present in 70%. Prior to the introduction of dupilumab, 17% of patients were only treated with topical compounds and 83% of patients had previously received an average of 1.15 systemic treatments, including cyclosporine (48%), methotrexate (42%), phototherapy (34%), oral corticosteroids (18%), or another systemic treatment (7%, azathioprine or mycophenolate mofetil). The mean follow-up time was 29.2 months \pm 10.7 (11–57).

Use of Dupilumab

The information was available for 77 patients who responded to the questionnaire. Injection of dupilumab was performed by a nurse in 47% of patients (n=36), 43% (n=33) of patients were autonomous. Injections were performed by a family member for 7 patients or by the general practitioner (1 patient).

Dupilumab was combined with topical steroids in 80% (n=63) of patients, most often occasionally (77%, n=48). After a mean duration of treatment of 13.5 months \pm 1.7, a wearing-off of the beneficial effect of dupilumab or an end-of-dose effect was reported by 47% (n=36) of patients. In this situation, the dosing interval was shortened to 10 days by 3% of patients (n=2). In contrast, dose spacing was reported by 9 patients (11%) along with a maintained efficacy. The spacing pattern was one injection every 3 to 4 weeks (n=5) or no injections in the summer (n=4). The reasons for dose spacing were remission in summer, forgetfulness, good clinical response, or to reduce ocular adverse events (2 patients).

Tolerance

During treatment, 58% of patients (n=45) reported no problems while pain was experienced by 32% (n=25) of patients, 10% (n=8) reported an injection site reaction, 2.5% used an EMLA patch (n=2), and 11.6% (n=9) reported other effects such as post-injection inflammation (n=5). Seventy-six percent (n=60) of patients reported at least one systemic adverse event (AE). Altogether, 91 AE were reported by patients including ocular AE (n=54, 59%), increased fatigue (n=8), increased herpes recurrence (n=6), headaches (n=5), weight gain (n=5), tendency to rhinoparyngitis (n=5), allergy to dupilumab (n=2) but without allergologic confirmation, mucosal dryness (n=2). Twenty-seven percent (n=21) of the patients had contracted COVID-19 while on dupilumab, but no severe forms were reported.

Discontinuation of Dupilumab

After a mean follow-up time of 29.7 ± 10.7 months (median: 27 months), 23 patients (28%) had permanently discontinued dupilumab (72% drug survival). Seven patients (9%) had discontinued treatment for a mean of 10 ± 6 months before restarting dupilumab. The reasons for permanent discontinuation were loss of efficacy (26%, n=6), ineffectiveness and ocular adverse events (22%, n=5), ocular adverse events (17%, n=4), other causes (13%, n=3) including COVID 19 infection (n=1), rash, treatment constraints, ongoing cold sore, pregnancy (13%, n=3) and clear improvement (9%, n=2).

The therapeutic strategies at the end of the treatment were as follows: a new systemic treatment was introduced in 56% of patients (n=13). For 3 of them, it was a conventional treatment (ciclosporin, phototherapy), for 10 patients, it was a new generation treatment (baricitinib, nemolizumab, tralokinumab, upadacitinib). Ten patients were continuing with topical treatment (dermocorticoids or topical tacrolimus) as monotherapy. Four percent (n=3) of patients in this study were lost to follow-up.

Overall Patient Experience of Dupilumab

Five patients did not respond to the questionnaire. Thirty-five percent of patients had no apprehension before starting treatment and 53% of patients rated the quality of medical information prior to treatment initiation as greater than 8/10. From the patients' perspective, the mean patient's satisfaction score was 7.5 ± 1.8 , and the constraints related to the treatment were scored at 3.1 ± 2.1 .

Discussion

Several real-life studies have confirmed the efficacy and safety of dupilumab in AD.²⁻⁴ However, few studies have looked at how patients use dupilumab in practice and their perception of the treatment.

In our series, the administration of dupilumab was mainly performed by a nurse and 43% of patients reported self-injection. It has been demonstrated that self-injection was associated with an increase in therapeutic compliance, notably due to greater flexibility in the time and place of administration.⁶ In addition, improved adherence could be associated with better disease control, increased patient quality of life and economic benefits to the health care system.^{7,8}

Although AD treatments may contribute to the burden of the disease, dupilumab was associated with a lower burden score,⁹ likely reflecting treatment efficacy and easy of use and patient satisfaction.

Altogether, in our study, patients were satisfied with the dupilumab treatment as shown by the satisfaction NRS score of 7.5 and the low constraint NRS score (3.1).

Although the majority of patients (65%) expressed apprehension before starting the treatment, they assessed the drawbacks of side effects against the benefits, and the vast majority of patients believed that dupilumab was worth using. A recent study [5] underlined the importance of education and good patient–healthcare professionals communication which enable patients to manage their disease and treatment expectations.

Furthermore, the satisfaction of patients may explain the dupilumab persistence. In our study, after a mean follow-up time of 29.7 ± 10.7 months (median: 27 months), drug survival was 72%. A drug survival rate of 89% after 26.3 months of treatment with dupilumab in a cohort of 112 AD patients had previously been reported.¹⁰ Similar high persistence was confirmed with overall drug survival rates in 402 patients of 91% and 88% after 1 and 2 years, respectively.¹¹ In a retrospective study that included 1963 patients with AD,¹² 77% of patients remained in treatment for 12 months. In an observational study including 288 AD patients treated with systemic therapy, the median duration of dupilumab treatment was 14.9 months.¹³

However, discontinuation of dupilumab treatment may occur. In a prospective multicenter registry including 402 AD patients, 9% discontinued dupilumab treatment mainly for AE.¹¹ In a recent retrospective multicentric study,¹⁴ 150 AD patients (15.5%) out of 968 patients discontinued treatment after a median treatment duration of 5 months. The main reasons for discontinuation were AE (41%), lack of efficacy (15%), lack of efficacy and AE (15%), planned pregnancy (8%), disease remission (4%) or various other reasons (17%). In contrast, in our study, the main reason for permanent discontinuation (28%) were loss of efficacy (26%) followed by ineffectiveness and ocular adverse events (22%), ocular adverse events (17%), other cause (13%), pregnancy (13%), and disease remission (9%). We did not observe any impact of the COVID-19 pandemic on the dupilumab treatment.

This difference could be explained by the appearance of new therapies in AD (baricitinib, upadacitinib and tralokinumab) that may favour the switch of treatments.

Although a longer dosing interval is feasible in some patients presenting AE, it is still not recommended that patients with a good clinical response should reduce the total number of injections and burden of treatment. In our study, dose spacing was rarely reported (11%) but in all cases with a maintained efficacy.

In a recent French multicentre retrospective study¹⁵ including 1017 AD patients for whom a dupilumab dosing interval (> every two weeks) had been implemented, dose spacing was introduced for response, AE and for both AE and response in 55 (63%), 18 (20%) and 15 (17%) patients, respectively. The authors concluded that a longer dupilumab dosing interval in patients achieving a clinically relevant improvement is linked to maintained efficacy in 2 out of 3 patients. However, in the “LIBERTY AD SOLO-CONTINUE” study, high-responding patients taking dupilumab every 4 weeks maintained an EASI-75 response in only 58% of cases,¹⁶ and continued response over time was most consistently maintained with dupilumab administered weekly or every 2 weeks. Longer dosage intervals resulted in a diminution of response. Conversely, we observed an end-of-dose effect in 47% of patients leading to the shortening of the dosing interval of dupilumab injection. To our knowledge the wearing-off of the beneficial effect of dupilumab has never been reported.

The limitations of our study are the retrospective nature and the small number of participants, the absence of a control group, the recall bias, and the declarative aspect of the responses.

We do not have a clear explanation for the male to female ratio of 1.5 observed in our study which is different from previous studies.¹⁷ Even though females had a higher AD prevalence in most countries, relatively little is known regarding differences between females and males concerning the course and characteristics of AD. However, sex did not seem to influence the self-rated severity of AD,¹⁸ but the impact of sex on the dupilumab prescription is unknown.

Conclusions

Our study shows that treatment with dupilumab is associated with a high level of satisfaction among treated patients. The hospital follow-up and the possibility to easily personalize the treatment according to clinical responses or side effects help explain the excellent perception of the treatment by the patients. Education and good patient-dermatologist communication allow patients to better understand their AD and their expectations of treatment.

Data Sharing Statement

The data that support the findings of this study are available upon reasonable request from the corresponding author, [FA]. The data are not publicly available due to their containing information that could compromise the privacy of research participants.

Statement of Ethics

All patients provided written informed consent.

Acknowledgments

The authors wish to thank all patients who took part in this study and Elisabeth Homassel for her technical assistance.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, the study design, execution, acquisition of data, analysis and interpretation, or in all of these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wollenberg A, Barbarot S, Bieber T, et al. Consensus-based European guidelines for treatment of atopic eczema (atopic dermatitis) in adults and children: part II. *J Eur Acad Dermatol Venereol*. 2018;32:850–878. doi:10.1111/jdv.14888

2. Faiz S, Giovannelli J, Podevin C, et al. Effectiveness and safety of dupilumab for the treatment of atopic dermatitis in a real-life French multicenter adult cohort. *J Am Acad Dermatol*. 2019;81:143-51. doi:10.1016/j.jaad.2019.02.053
3. Beck LA, Deleuran M, Bissonnette R, et al. Dupilumab provides acceptable safety and sustained efficacy for up to 4 years in an open-label study of adults with moderate-to-severe atopic dermatitis. *Am J Clin Dermatol*. 2022;23:393–408. doi:10.1007/s40257-022-00685-0
4. Spekhorst LS, de Graaf M, Zuihthoff NPA, et al. Dupilumab drug survival and associated predictors in patients with moderate to severe atopic dermatitis: long-term results from the daily practice bioday registry. *JAMA Dermatol*. 2022;158:1048–1056. doi:10.1001/jamadermatol.2022.3014
5. Ameen M, Meller S, Pinter A, Shear NH, Soria A; The BADEL Study Group. Perception and experience of biologic therapy in atopic dermatitis: a qualitative focus group study of physicians and patients in Europe and Canada. *Dermatol Ther*. 2021;11:2159–2177. doi:10.1007/s13555-021-00631-8
6. Ito M, Kamata M, Uchida H, et al. Introduction of self-injection increases compliance with dupilumab administration in adult patients with atopic dermatitis: a single-center retrospective study. *J Am Acad Dermatol Int*. 2020;1:91-2.
7. Chilton F, Collett RA. Treatment choices, preferences and decision-making by patients with rheumatoid arthritis: treatment choices in patients with RA. *Musculoskeletal Care*. 2008;6:1-14. doi:10.1002/msc.110
8. Kivitz A, Cohen S, Dowd JE, et al. Clinical assessment of pain, tolerability, and preference of an autoinjection pen versus a prefilled syringe for patient self-administration of the fully human, monoclonal antibody adalimumab: the TOUCH trial. *Clin Ther*. 2006;28:1619–1629. doi:10.1016/j.clinthera.2006.10.006
9. Nic Dhonncha E, O'Connor C, Cosgrave N, Murphy M. Burden of treatment in adult patients with atopic dermatitis. *J Eur Acad Dermatol Venereol*. 2022;36:e467–e468. doi:10.1111/jdv.17954
10. Khosravi H, Zhang S, Anderson AM, Ferris LK, Choudhary S, Patton T. Dupilumab drug survival, treatment failures, and insurance approval at a tertiary care center in the United States. *J Am Acad Dermatol*. 2019;82:1023–1024. doi:10.1016/j.jaad.2019.12.034
11. Spekhorst LS, Ariëns LFM, van der Schaft J, et al. Two-year drug survival of dupilumab in a large cohort of difficult-to-treat adult atopic dermatitis patients compared to cyclosporine A and methotrexate: results from the BioDay registry. *Allergy*. 2020;75:2376–2379. doi:10.1111/all.14324
12. Silverberg JI, Guttman-Yassky E, Gadkari A, et al. Real-world persistence with dupilumab among adults with atopic dermatitis. *Ann Allergy Asthma Immunol*. 2021;126:40–45. doi:10.1016/j.anai.2020.07.026
13. de Bruin-Weller M, Pink AE, Ferrucci SM, et al. Use of systemic therapies in adults with atopic dermatitis: 12-month results from the European prospective observational study in patients eligible for systemic therapy for atopic dermatitis (EUROSTAD). *J Dermatological Treat*. 2022;33:2565–2570. doi:10.1080/09546634.2022.2038361
14. Marniquet ME, Seneschal J, Darrigade AS, et al. Reasons for discontinuation of dupilumab in adult atopic dermatitis in clinical practice. *Br J Dermatol*. 2022;186:733–735. doi:10.1111/bjd.20883
15. Jendoubi F, Shourik J, Seneschal J, et al. Longer dupilumab dosing intervals in adult patients with atopic dermatitis: experience from a French multicentre retrospective cohort study. *Br J Dermatol*. 2022;187:602–603. doi:10.1111/bjd.21628
16. Worm M, Simpson EL, Thaçi D, et al. Efficacy and safety of multiple dupilumab dose regimens after initial successful treatment in patients with atopic dermatitis: a randomized clinical trial. *JAMA Dermatol*. 2020;156:131–143. doi:10.1001/jamadermatol.2019.3617
17. Barbarot S, Auziere S, Gadkari A, et al. Epidemiology of atopic dermatitis in adults: results from an international survey. *Allergy*. 2018;73:1284–1293. doi:10.1111/all.13401
18. Johansson EK, Bergström A, Kull I, et al. Prevalence and characteristics of atopic dermatitis among young adult females and males-report from the Swedish population-based study BAMSE. *J Eur Acad Dermatol Venereol*. 2022;36:698–704. doi:10.1111/jdv.17929

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>