



ORAL PRESENTATION

Open Access

Role of Ras isoforms in $\gamma\delta$ T-cell development

Ana V Marín-Marín¹, Eva Jiménez Pérez², Eugenio Santos³, Edgar Fernández-Malavé^{1*}

From 7th European Workshop on Immune-Mediated Inflammatory Diseases
Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Introduction

The small guanine nucleotide binding proteins of the Ras family (H-, K-, and N-ras isoforms) couple surface receptors, including T-cell antigen receptors, to a variety of cellular responses. Ras proteins are highly homologous and expressed ubiquitously, raising questions as to their functional specificity. H-ras and N-ras have been shown to be dispensable for development in the thymus but critical for peripheral Th1 differentiation of $\alpha\beta$ T cells [1]. In the $\gamma\delta$ T-cell lineage, in contrast, the specific function of Ras isoforms had not been addressed.

Aim

We aimed at identifying the specific roles of H-ras and N-ras in $\gamma\delta$ T-cell development, particularly, in the generation of $\gamma\delta$ T-cell subsets defined by expression of CD27.

Mice and methods

Mice deficient for H-ras or N-ras were analyzed by multi-parametric flow cytometry. In addition, N-ras-deficient mice with impaired $\alpha\beta$ T-cell development were generated by breeding with CD3 δ KO mice and analyzed.

Results

Ras isoform-deficient mice exhibited normal frequencies and numbers of $\gamma\delta$ T cells in the thymus, which expressed normal levels of surface $\gamma\delta$ TCR, but a consistent decrease of CD27 $^+$ $\gamma\delta$ T cells in peripheral lymphoid organs, compared to controls. Conversely, in mice lacking mature $\alpha\beta$ T cells, N-ras deficiency resulted in reduced numbers of CD27 $^+$ $\gamma\delta$ T cells, with concomitant increase of the CD27 $^-$ population, in the thymus but not in the periphery. This suggests that signals from $\alpha\beta$ T-cell thymocytes for developing CD27 $^+$ $\gamma\delta$ T cells are partly N-ras-dependent.

Conclusions

As previously shown for $\alpha\beta$ T cells, H-ras and N-ras are dispensable for intrathymic development of $\gamma\delta$ T cells, but they could be involved in finely regulating survival and/or expansion of $\gamma\delta$ T cells in the periphery, in agreement with recent findings in mice lacking RasGRP1 [2], a guanine nucleotide exchange factor for Ras.

Acknowledgements

The study was supported by grant FIS PI11/02198 from Instituto de Salud Carlos III to EF-M.

Author details

¹Dept. of Immunology, Faculty of Medicine, Complutense University, Madrid, Spain. ²Cellular Biology Dept., Faculty of Medicine, Complutense University, Madrid, Spain. ³Center for Cancer Research, Salamanca, Spain.

Published: 28 November 2012

References

- Iborra S, Soto M, Stark-Aroeira L, Castellano E, Alarcón B, Alonso C, Santos E, Fernández-Malavé E: H-ras and N-ras are dispensable for T-cell development and activation but critical for protective Th1 immunity. *Blood* 2011, 117:5102-5111.
- Chen Y, Ci X, Gorentla B, Sullivan SA, Stone JC, Zhang W, Pereira P, Lu J, Zhong XP: Differential requirement of RasGRP1 for $\gamma\delta$ T cell development and activation. *J Immunol* 2012, 189:61-71.

doi:10.1186/1479-5876-10-S3-O6

Cite this article as: Marín-Marín et al.: Role of Ras isoforms in $\gamma\delta$ T-cell development. *Journal of Translational Medicine* 2012 10(Suppl 3):O6.

¹Dept. of Immunology, Faculty of Medicine, Complutense University, Madrid, Spain

Full list of author information is available at the end of the article