

ARTICLE

Received 14 Jun 2011 | Accepted 28 Jul 2011 | Published 30 Aug 2011

DOI: 10.1038/ncomms1461

A segmental genomic duplication generates a functional intron

Uffe Hellsten^{1,*}, Julie L. Aspden^{2,3,*}, Donald C. Rio^{2,3} & Daniel S. Rokhsar^{1,2}

An intron is an extended genomic feature whose function requires multiple constrained positions—donor and acceptor splice sites, a branch point, a polypyrimidine tract and suitable splicing enhancers—that may be distributed over hundreds or thousands of nucleotides. New introns are therefore unlikely to emerge by incremental accumulation of functional sub-elements. Here we demonstrate that a functional intron can be created *de novo* in a single step by a segmental genomic duplication. This experiment recapitulates *in vivo* the birth of an intron that arose in the ancestral jawed vertebrate lineage nearly half-a-billion years ago.

¹ DOE Joint Genome Institute, Walnut Creek, California 94598, USA. ² Center for Integrative Genomics, University of California Berkeley, California 94707, USA. ³ Department of Molecular and Cell Biology, University of California Berkeley, California 94707, USA. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to U.H. (email: UHellsten@lbl.gov).

The appearance of a new intron that precisely splits an exon without disrupting the corresponding peptide sequence is a very rare event in vertebrate genomes. No such intron gains have been documented in the human, mouse, rat, or dog lineages since their common mammalian ancestor¹, nor in a comparison between the pufferfish *Fugu* and *Tetraodon*². Nevertheless, a few credible cases of vertebrate intron gain have been documented in teleost fish³. In these examples, the novel intron sequences showed recognizable similarity to the surrounding coding exons, and appeared at AG|GT sites embedded within coding sequence. This observation suggested that recent tandem duplication of an AGGT motif-containing coding sequence could have led to the formation of the intron, an idea originally put forward by Rogers⁴. If the 5'-GT and 3'-AG in the duplicated region were recognized by the spliceosome as donor (5'-splice site) and acceptor (3'-splice site) signals, the redundant duplicated region would be excised from the primary transcript, leaving the translated peptide unaltered by the segmental genomic duplication (Fig. 1).

Here we apply a bioinformatic approach to look for early vertebrate-specific intron gains. In addition to the requirement that the intron be absent in invertebrate orthologues, we require that paralogues from the whole-genome duplications at the base of vertebrate evolution contain examples of genes with and without the intron. We find only one example of such an intron gain, namely within the *ATP2A* family, where an intron is present in the human *ATP2A1* gene, but not in *ATP2A2*. We test the segmental duplication scenario by creating artificially duplicated constructs of the intronless gene and demonstrate in live human cells that the redundant region can be spliced out, in essence reenacting a plausible creation mechanism for the intron in the *ATP2A1* gene.

Results

Discovery of the vertebrate-specific *ATP2A1* intron gain. We conducted a genome-wide search for pairs of human paralogues indicated by conserved synteny to have originated in one of the early vertebrate-specific whole-genome duplications. Such pairs




Figure 1 | Proposed mechanism for intron birth. Extant lancelet and lamprey *ATP2A* genes, and human and zebrafish *ATP2A2* genes, are intronless in the region shown, reflecting the ancestral chordate condition, but human and zebrafish *ATP2A1* genes are interrupted by an intron between the first and second nucleotides of codon G310 (coordinate with respect to human amino acid sequence of *ATP2A1* isoform a). The peptide sequence is fully conserved, so only synonymous amino acid codon substitutions are seen in the nucleotide sequence. A segmental tandem duplication encompassing this region would produce a potential intron with consensus donor and acceptor splice sites, including a polypyrimidine tract. The sequence of the intronless human *ATP2A2* gene is used in this example.

were grouped by sequence similarity with putative orthologues from both vertebrates and invertebrates, and intron splice sites within conserved areas of the coding sequence were identified. Of 252 splice sites that meet strict criteria (Methods) only one, found in *ATP2A1*, had the signature of a vertebrate-specific gain.

The *ATP2A* gene family encodes sarco/endoplasmic reticulum calcium ATPases (SERCAs) whose dysfunction has been associated with several human diseases⁵. *ATP2A* genes found outside of the jawed vertebrates are intronless near the motif AAIPEGLPAV, reflecting the ancestral condition (Fig. 1; Table 1). Humans and other tetrapods encode three paralogues *ATP2A1*, *ATP2A2*, and *ATP2A3* that encode *SERCA1*, *SERCA2*, and *SERCA3*, respectively. These genes originated from an ancestral chordate gene by two rounds of duplication and subsequent loss of one copy (Table 1; Methods). All vertebrate *ATP2A1* genes include a novel intron between the first and second nucleotides of the G310 codon at an AGGT motif (Fig. 1). In contrast, the *ATP2A2* and *ATP2A3* genes retain the ancestral intronless state at this position. The intron in the *ATP2A1* gene splits a single ancestral exon (exon 8 in human *ATP2A2*) into two exons (exons 8 and 9 in human *ATP2A1*). As the intron is shared by tetrapods and teleost fish, it evidently arose more than 420 million years ago.

We can infer much of the ancestral vertebrate *ATP2A* sequence around G310, because its nucleotide sequence is highly constrained by the perfect conservation of the amino residues, leaving only synonymous coding positions to vary. A segmental duplication containing the AGGT motif contains most of the sequence elements required for recognition by the U2 (major class) spliceosome (Fig. 1). These motifs include several other consensus nucleotides around the donor and acceptor sites⁶ ((A/C)AG|GT(A/G)AGT ... CAG|G) beyond the GT and AG dinucleotides, as well as a polypyrimidine tract (7 of the 8 nucleotides at position -12 to -5 near the acceptor are pyrimidines), and a potential branchpoint A residue with consensus (YTNAV) at position -48 (not shown).

Reenacting the birth of an intron. To test the hypothesis that a functional intron can be produced *de novo* by an appropriate segmental genomic duplication, we designed a mini-gene construct that contains a duplication of *ATP2A2* exon 8. The construct was transiently transfected into HEK 293 and HeLa cells, and the resulting messenger RNA characterized. We propose that the duplicated

Table 1 | Number of *ATP2A* genes with and without an intron at the motif AAIPEGLPAV in 15 species.

Species	# <i>ATP2A</i> with intron	# <i>ATP2A</i> w/o intron
Human	1	2
Mouse	1	2
Opossum	1	2
Platypus	1	2
Chicken	1	2
Frog	1	2
Zebrafish	3	4
Fugu	3	2
Stickleback	2	4
Medaka	2	4
Lamprey	0	1
Sea squirt	0	4 (tandem)
Lancelet	0	2 (tandem)
Sea urchin	0	1
Sea anemone	0	2

Tetrapods (that is, Human through Frog above) all have three copies, of which one has the intron. These copies can be shown to have originated from two rounds of whole-genome duplication (WGD) and subsequent single-copy loss. Teleost fish (Zebrafish-Medaka) have 5-7 copies, consistent with having undergone an extra round of WGD. Because fugu and zebrafish have more than two copies with an intron, the insertion of the intron probably happened between the two rounds of WGD. No genes with this intron are found outside vertebrates (Lamprey and below).




Figure 2 | Reconstruction of duplicated splice sites undergoes splicing at low levels. (a) Schematic diagram of mini-gene constructs transfected into HEK 293 and HeLa cells including *ATP2A2* duplicated exon 8/9 (D), *ATP2A2* Single (C1) and *ATP2A2* Single with 6 bp insert (C2). The sequence corresponding to exon 8 is shaded light grey, exon 9 dark grey, 6 bp insert black and vector sequences are shown by dashed lines. (b) Diagram of the RNase protection probe along with annotations of what sequences each part of the probe will hybridize to. (c) Schematic representation of potential mRNA species from the transfections and the corresponding RNA probe fragments that their presence will lead to in the RNase protection assay. (d) Phosphorimage of RNase protection assay products from HEK 293 cells with DNA size marker sizes (in nt ssDNA) indicated on the right, and what size RNA fragments the protected probe bands correspond to on the left (ssRNA). Transfections were performed in triplicate. (e) Phosphorimage of RNase protection assay products from HeLa cells with DNA size marker sizes (in nt ssDNA) indicated on the right, and what size RNA fragments the protected probe bands correspond to on the right (ssRNA). Transfections were performed in triplicate.

exon 8 in *ATP2A2* becomes exons 8 and 9 in *ATP2A1* through the creation of a new intron (Fig. 2a). The duplicated AGGT motifs span the borders between the exon 8 and 9 regions, and our hypothesis is that the spliceosome will recognize the AG|gt...ag|GT sequences as splice sites and remove the central exon 9 and exon 8 (Fig. 2a). The segmental duplication could be any length, as long as the 5'-GT and 3'-AG are separated by more than the minimal functional intron length, about 60 nucleotides.

The duplicated nature of the construct rendered standard reverse-transcription PCR ineffective at distinguishing between the presence of unspliced and spliced mRNA products from the *ATP2A2* duplicated minigene, so we employed RNase protection assays instead. To clearly differentiate spliced from unspliced mRNA, we cloned an extra 6bp into the 5' copy of exon 8 near its 3' end that distinguishes it from the 3' copy of exon 8 (Fig. 2a). The RNase protection probe was designed to take advantage of this small difference between the two exon 8s (Fig. 2b) and will lead to the production of four possible protected RNA probe fragments (Fig. 2c). Two control mini-gene plasmids, C1 and C2, were also transfected to act as markers for these predicted RNase protection fragments (Fig. 2a). *ATP2A2* Single (C1) will give rise to a fragment at the size matching unspliced mRNA from the duplicated construct, 194 nt (Fig. 2c). *ATP2A2* single with 6 bp insert (C2) will

produce a protected fragment corresponding in length to spliced mRNA (215 nt). RNase protection assays were performed with total RNA from the transfections of these three mini-genes along with a no-transfection control and a probe-alone control in HEK 293 cells (Fig. 2d) and HeLa cells (Fig. 2e). The probe-alone assay leads to the production of low-level non-specific protected fragments, presumably from internal secondary structures in the radiolabelled probe that are RNase resistant (Fig. 2e, lane 5). The 179 nt protected probe fragment in the no-transfection control (Fig. 2d, lane 1 and Fig. 2e, lane 4) confirms the presence of endogenous *ATP2A2* mRNA transcripts.

Our experiments show that segmental duplication can create a functional intron. Using the duplicated *ATP2A2* construct, we clearly observed a 215 nt protected probe fragment whose size corresponds precisely to the spliced mRNA control (Fig. 2d, lanes 2 and 4, 5, 6, and Fig. 2e, lanes 1 and 3). Quantification of the relative abundance of these protected probe RNA fragments from the *ATP2A2* duplication indicates that 15.9% ($\pm 1.7\%$) in HEK 293 cells and 5.9% ($\pm 1.2\%$) in HeLa cells, of the mRNA is spliced. To rule out the possibility that the 215 nt RNase-protection RNA product resulted from rearrangements at the DNA level rather than splicing, plasmid DNA was recovered from HEK 293 cells transfected with the *ATP2A2* duplication construct. Characterization of these




Figure 3 | Plasmids do not undergo DNA rearrangement during transfection. (a) Schematic of DNA mini-gene constructs used in transfections as described in Figure 2a. Restriction sites are shown along with sizes of DNA digestion fragments. (b) Agarose gels showing diagnostic digests of DNA plasmids recovered from HEK 293 cells transfected with *ATP2A2* duplicated 8/9 with insert (D). Control digests from untransfected plasmids are shown; *ATP2A2* duplicated 8/9 with insert (D), *ATP2A2* single (C1) and *ATP2A2* single with 6 bp insert (C2), along with DNA size markers (m). All the recovered DNA plasmids were the same size as the transfected DNA plasmid *ATP2A2* duplicated 8/9 with insert (D).

plasmids with restriction enzyme digestions (Fig. 3a) revealed that no such DNA arrangement has taken place (Fig. 3b). Although clear evidence for splicing was observed, the majority of protected fragments were 194 and 200 nt, indicating that most of the expressed mRNA was unspliced. Despite the modest level of splicing, we find that the spliceosome can recognize the duplicated splice sites, and that these alone are sufficient to allow the new intron to be excised from mRNA.

Discussion

The genes of early eukaryotes likely contained many more introns than found in present day eukaryotic genomes, with subsequent genome evolution dominated by intron loss⁷. This suggests an early epoch of massive intron invasion, the mechanism of which has long since been inactivated. In contrast, relatively recent intron gains are very uncommon, particularly within vertebrates. The intron in *ATP2A1* discussed in this work is the only example of such an intron gain that we found out of 252 candidate introns within coding regions that are highly conserved across the lancelet, sea urchin, and human genomes (Methods). Such recent intron gains are almost certainly caused by a mechanism different than that responsible for the original genomic invasion of introns⁸.

Our results show that a short intragenic tandem duplication can insert a novel U2-type intron into a protein-coding gene, leaving the corresponding peptide sequence unchanged. The novel intron

described here was produced by segmental duplication of an AGGT site within coding sequence. Tandem duplications are common in genomes; on the scale of a single gene, Lynch and Conery⁹ estimate the order of ~100 gene duplications per genome per million years, and smaller-scale duplications are even more prevalent¹⁰. The newly created intron is accurately spliced *in vivo*, albeit at a modest level of ~16% in HEK 293 and ~6% in HeLa cells. The level of spliced mRNA may differ in the fast twitch muscle cells in which *ATP2A1* is normally expressed. The splicing efficiency of the originally duplicated sequence of the ancestral vertebrate gene could also have been modulated by synonymous sequence differences relative to our human-genome-based construct, and/or differences in the length and position of the duplicated region.

Mutations of the *ATP2A1* gene are associated with Brody disease^{11–12}, an autosomal recessive muscle disorder characterized by impaired relaxation of fast-twitch muscles after exercise. A similar recessive disorder associated with an *ATP2A1* mutation has been described in cattle¹³, and the *ATP2A1* zebrafish mutant *accordion* also shows related behavioural defects¹⁴. The recessive nature of these hereditary disorders implies that vertebrates can tolerate reduced levels of *ATP2A1* protein product (SERCA1). Thus, the ancient intragenic tandem duplication that produced the intron-bearing allele in the proto-vertebrate *ATP2A1* could have initially spread nearly neutrally through the ancestral population even without 100% splicing efficiency. For the small population sizes characteristic of vertebrates¹⁵, such an allele could rise to modest frequency and even become fixed, if homozygotes are not at too high of a disadvantage. An allele with 50% splicing efficiency in a homozygous state, for example, would nominally produce the same level of protein product as a heterozygote. Once the intron-bearing allele is fixed, secondary mutations could then emerge to incrementally improve splicing efficiency.

The precise gain of an intron as described here is conceptually different from the exon gains previously reported in primates, in which the insertion of ALU elements into existing exons creates a new alternatively spliced exon and adds sequence to the final peptide product¹⁶. Recruitment of other sequence elements to form or extend exons has also been described¹⁷.

Most other mechanisms for intron gain that have been proposed differ fundamentally from the mechanism documented here, in that they are expected to be accompanied by deletion or insertions within the resulting coding sequence. In contrast, the mechanism we have demonstrated here generates a precisely inserted new intron without any disruption of the surrounding coding sequence. Two examples of probably very recent intron gains have been described in the water flea *Daphnia*, in which novel introns are still polymorphic in the population¹⁸. In contrast to the mechanism described here, the newly born introns in *Daphnia* do not show similarity to flanking (or any known) sequence, and their origin is unknown. This suggests that other intron-creation mechanisms besides the one shown here are also active.

Methods

Genomewide search for vertebrate-specific intron gains. To identify 'ohnologues', that is, pairs of human paralogues, probably originated in a whole-genome duplication, we assigned position IDs to all loci in the genome, numbering them in the order in which they occur. We used the ENSEMBL models version 55, longest transcript at each locus, and aligned the corresponding 23,266 peptides to each other using BLASTp¹⁹ with an *e*-value cutoff of 10^{-20} . We next identified tandem expanded families, here defined as clusters of neighbouring genes with peptide similarity, allowing a maximum of two intervening genes on any strand. Such clusters were reduced by retaining only the gene with the longest transcript. Genes with strong (*e*-value $< 10^{-20}$) similarity to more than twenty other genes (after removing tandem duplicates) were also eliminated to avoid confounding effects of large gene families such as zinc-finger, kinases, or olfactory receptors.

Next, we identified pairwise reciprocal highest scoring hits between the remaining genes, restricting further analysis to pairwise hits with scores of at least 60% of that of the maximum of each of the members' reciprocal best hit scores. This left us with 9,852 loci that were re-numbered in strict consecutive orders.

We implemented an algorithm similar to that described by Blanc *et al.*²⁰ to detect clusters of adjacent genes with sequence similarity to clusters elsewhere in the genome. The mapping of each such gene to its counterpart in the other cluster can be visualized as rungs in a ladder, defining blocks of conserved synteny. To account for the considerable scrambling of gene order by large-scale inversions during half a billion years of evolution, we allowed up to 15 intervening genes between any two rungs. Furthermore, we required each block to contain at least 5 pairs of genes.

This analysis resulted in 153 blocks of intragenomic conserved synteny containing a total of 1,007 duplicated pairs of paralogues. These are all expected to be *bona fide* ohnologues; identical analysis on randomly scrambled gene IDs yielded no false positives.

As we aimed to identify early vertebrate intron gains, we chose two outgroups to the vertebrates, namely the Florida lancelet *Branchiostoma floridae* (a chordate) and the purple sea urchin *Strongylucentrotus purpuratus* (an echinoderm). For the lancelet, we used the gene annotation by JGI²¹, and, for the sea urchin, the NCBI gene build 2. Reciprocal highest-scoring BLASTp hits yielded 8,501 candidate orthologues between the lancelet and the sea urchin. If both genes in such a pair had mutual best hits to the same human gene, or the same above identified duplicated paralogous pair, we defined a BHHS (lancelet—human 1—human 2—sea urchin) cluster of orthologues. In total, 426 such clusters were defined.

Multiple sequence alignments of the BHHS peptide clusters were performed using clustalW²². From these alignments, we extracted gap-free regions flanked by fully conserved amino residues and with no stretch of more than five non-conserved amino residues using custom PERL scripts. The positions and phases of all intron splice sites within such blocks were mapped, and we retained only splice sites flanked by regions with at least six of eight amino residues fully conserved. Finally, we excluded from the analysis sites within 4 amino residues from a non-overlapping splice site in another species, because such cases are mostly caused by gene models with inaccurate intron-exon boundaries.

A total of 252 intron splice sites met these stringent criteria. These were evaluated against the wider set of species shown in Table 1. The signature of a post-duplication vertebrate intron gain would contain a splice site in only one of the two human copies and none of the invertebrate orthologues. In the set examined, we found only a single candidate: the eighth intron in the *ATP2A1* gene is absent in all invertebrate orthologues examined, as well as vertebrate paralogues *ATP2A2* and *ATP2A3*. The distribution of genes with and without this intron in amniotes and teleost fish strongly suggests that this intron gain happened between the two rounds of whole-genome duplication at the base of vertebrate evolution (Table 1).

RNase protection assays. Mini-gene reporter constructs were generated using genomic PCR of the *ATP2A2* gene and cloned into pCDNA3.1 (Invitrogen) between *KpnI* and *XhoI* sites (5'-ggcgggtggtacacacattgctgctgg-3'; 5'-ggcggctcgagcctcagactgacatctgg-3'). Overlapping PCR was used to generate the *ATP2A2* duplication from *ATP2A2* (5'-aacagatgctgctgacgggtacaacattgctgctgg-3'; 5'-cctgacagctgacatctgg-3') and quick-change mutagenesis was employed to insert the extra 6 bp into exon 8 (5'-cctggctgtagcaggtgattcattcctgaagtc-3'; 5'-gacctcaggaatgacactgactgacagcagg-3'). HEK 293 and HeLa cells were grown in standard conditions in DME medium with 5% fetal calf serum. HEK 293 cells (1.5 × 10⁶) were transfected with 4 μg DNA Lipofectamine 2000 (Invitrogen). 0.3 μg plasmid DNA was transfected into 2 × 10⁶ HeLa cells using Effectene (Qiagen). Cells were collected after 48 h and total RNA purified using RNeasy mini kits (Qiagen). 20 pmol of ³²P-labelled RNA probe, transcribed with T7 polymerase from a PCR fragment generated from *ATP2A2* single with insert (5'-cctggctgtagcaggtg-3'; 5'-taatgactactataggatgctcttcgctgacgtcaccctctagactgagcctg-3'), was hybridized to 10 μg total RNA at 45 °C for 16 h. After cooling to 4 °C, the RNA was incubated with RNases A and T1 at room temperature for 60 min. Following proteinase K treatment, phenol/chloroform extraction and ethanol precipitation, protected RNA fragments were resuspended in formamide dyes and run out on 6–8% denaturing polyacrylamide gels. The resulting dried gels were exposed to a phosphorimager screen and bands were quantified with ImageQuant (GE Healthcare).

Detailed explanation of RNase protection assay. The design of the probe and the duplicated polymorphic construct will lead to the production of four possible protected RNA probe fragments (Fig. 2c). Two of these will be protected in the presence of unspliced mRNA at 194 nt (21 + 12 + 167 nt) and 200 nt (12 + 167 + 15 nt) (Fig. 2c). However, if any *ATP2A2* duplicated mRNA is spliced, it will hybridize to the probe in such a way as to protect a larger fragment, at 215 nt (21 + 12 + 167 + 15 nt). The presence of endogenous *ATP2A2* mRNA, with no intron present will lead to production of a 179 nt fragment (12 + 167 nt). At the 5' end of the RNA probe is a section of probe sequence that is not complementary to any *ATP2A2* sequence and, therefore, will be digested by the RNases. Its presence creates a difference in protection fragment length between the input probe (240 nt) and the potential protection fragments, allowing confirmation that RNase treatment is working (Fig. 2d, lane 7 and Fig. 2e, lane 6).

Sequences. Exon 8 sequence is shown in bold; exon 9 sequence in standard typeface; 6 bp insert is in italics; vector sequence is dashed underlined; RNA flap sequence is double underlined and T7 promoter sequence is single underlined.

DNA sequence of *ATP2A2* duplication construct with insert.

5'-GGTACAAACATTGTGCTGGGAAAGCTATGGGAGTGGTGGTAGC
AACTGGAGTTAACACCCGAAATGGCAAGATCCGGGATGAAATGGTGG
CAACAGAACGAGAGAGAACCACCCTTCAGCAAAAAGTATGAAATTT
GGGGAACAGCTTCCAAAGTCATCTCCCTTATTGTCATTGCAGTCTGGAT
CATAAATATGGGCACTTCAATGACCCGGTGTATGGAGGGTCTGGATC
AGAGGTGCTATTACTACTTTAAAATTCAGTGGCCCTGGCTGTAGCAG
GTGATTCCATTCTGAAAGGTGCTGCCTGCAGTCATACCACCTGCCTGG
CTCTTGGAACTCGCAGAAATGGCAAAGAAAAATGCCATTGTTGGAAGCC
TCCCGTCTGTGAAACCCCTTGGTTGACTTCTGTTATCTGCTCAGACAA
GACTGGTACACTTACAACAAACCAGATGTCAGTCTGCAGGGGTACAAA
CATTGCTGCTGGGAAAGCTATGGGAGTGGTGGTAGCAACTGGAGTTAA
CACCGAAATGGCAAGATCCGGGATGAAATGGTGGCAACAGACAGGA
GAGAACCACCCTTCAGCAAAAAGTATGAAATGGGGAACAGCTTTC
CAAAGTCATCTCCCTTATTGTCATTGTCAGTGGATCATAAATATTGGG
CACTTCAATGACCCGGTTCATGGAGGGTCTGGATCAGAGGTGCTATTTA
CTACTTTAAAATTCAGTGGCCCTGGCTGTAGCAGCCATTCTCGAAGGT
CTGCTGCAGTCATACCACCTGCCTGGCTCTGGAAGTCCGAGAATGG
CAAAGAAAAATGGCATTGTTGCAAGCTCCCGTCTGCGAAACCCCTT
GGTTGACTTCTGTTATCTGCTCAGACAAGACTGGTACACTTACAAC
AAACAGATGTCAGTCTGCAGGCTCGAGTCTAGAGGG-3'

DNA sequence of RNase probe PCR product. 5'-CCCTGGCTGTAGCAG
GTGATTCCATTCTGAAAGGTGCTGCCTGCAGTCATACCACCTGCCTG
GCTCTGGAATCTCGCAGAAATGGCAAAGAAAAATGCCATTGTTGGAAGCC
TCCCGTCTGTGAAACCCCTTGGTTGACTTCTGTTATCTGCTCAGACAA
GACTGGTACACTTACAACAAACCAGATGTCAGTCTGCAGGCTCGAGTCTA
GAGGGGTGACGTCGAGCGAAAGGACATCCCTATAGTGTAGTCTGATTA-3'

Recovery of plasmids from transfected cells. Plasmid DNA was extracted from transfected HEK 293 cells and transformed into bacteria²³. From here, they were purified with minipreps (Fermentas) and underwent restriction enzyme digests. The resulting DNA fragments were separated on agarose gels.

References

- Coulombe-Huntington, J. & Majewski, J. Characterization of intron loss events in mammals. *Genome Res.* **17**, 23–32 (2007).
- Loh, Y. H., Brenner, S. & Venkatesh, B. Investigation of loss and gain of introns in the compact genomes of pufferfishes (Fugu and Tetraodon). *Mol. Biol. Evol.* **25**, 526–535 (2008).
- Venkatesh, B., Ning, Y. & Brenner, S. Late changes in spliceosomal introns define clades in vertebrate evolution. *Proc. Natl Acad. Sci. USA* **96**, 10267–10271 (1999).
- Rogers, J. H. How were introns inserted into nuclear genes? *Trends Genet.* **5**, 213–216 (1989).
- Hovnanian, A. SERCA pumps and human diseases. *Subcell. Biochem.* **45**, 337–363 (2007).
- Lim, L. P. & Burge, C. B. A computational analysis of sequence features involved in recognition of short introns. *Proc. Natl Acad. Sci. USA* **98**, 11193–11198 (2001).
- Koonin, E. V. Intron-dominated genomes of early ancestors of eukaryotes. *J. Hered.* **100**, 618–623 (2009).
- Catania, F., Gao, X. & Scofield, D. G. Endogenous mechanisms for the origins of spliceosomal introns. *J. Hered.* **100**, 591–596 (2009).
- Lynch, M. & Conery, J. S. The evolutionary fate and consequences of duplicate genes. *Science* **290**, 1151–1155 (2000).
- Tanay, A. & Siggia, E. D. Sequence context affects the rate of short insertions and deletions in flies and primates. *Genome Biol.* **9**, R37 (2008).
- Odermatt, A. *et al.* Mutations in the gene-encoding SERCA1, the fast-twitch skeletal muscle sarcoplasmic reticulum Ca²⁺ ATPase, are associated with Brody disease. *Nat. Genet.* **14**, 191–194 (1996).
- Odermatt, A. *et al.* The mutation of Pro789 to Leu reduces the activity of the fast-twitch skeletal muscle sarco(endo)plasmic reticulum Ca²⁺ ATPase (SERCA1) and is associated with Brody disease. *Hum. Genet.* **106**, 482–491 (2000).
- Drogemuller, C. *et al.* Identification of a missense mutation in the bovine *ATP2A1* gene in congenital pseudomyotonia of Chianina cattle: an animal model of human Brody disease. *Genomics* **92**, 474–477 (2008).
- Hirata, H. *et al.* Accordion, a zebrafish behavioral mutant, has a muscle relaxation defect due to a mutation in the ATPase Ca²⁺ pump SERCA1. *Development* **131**, 5457–5468 (2004).
- Lynch, M. *The Origins of Genome Architecture* (Sinaur, 2007).
- Lev-Maor, G., Sorek, R., Shomron, N. & Ast, G. The birth of an alternatively spliced exon: 3' splice-site selection in Alu exons. *Science* **300**, 1288–1291 (2003).
- Sorek, R. The birth of new exons: mechanisms and evolutionary consequences. *RNA* **13**, 1603–1608 (2007).
- Omilian, A. R., Scofield, D. G. & Lynch, M. Intron presence-absence polymorphisms in *Daphnia*. *Mol. Biol. Evol.* **25**, 2129–2139 (2008).
- Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

20. Blanc, G., Hokamp, K. & Wolfe, K. H. A recent polyploidy superimposed on older large-scale duplications in the Arabidopsis genome. *Genome Res.* **13**, 137–144 (2003).
21. Putnam, N. H. *et al.* The amphioxus genome and the evolution of the chordate karyotype. *Nature* **453**, 1064–1071 (2008).
22. Thompson, J. D., Gibson, T. J. & Higgins, D. G. Multiple sequence alignment using ClustalW and ClustalX. *Curr. Protoc. Bioinformatics* Chapter 2, Unit 2 3 (2002).
23. Beall, E. L. & Rio, D. C. Drosophila IRBP/Kup70 corresponds to the mutagen-sensitive mus309 gene and is involved in P-element excision *in vivo*. *Genes Dev.* **10**, 921–933 (1996).

Acknowledgements

We thank Karen Clyde for HeLa cells, Malik Francis for MC1061 RecA⁻ cells, Robert Tjian for DME medium and trypsin, ELIM for sequencing facilities, Matt Taliaferro and Melissa Harrison for helpful discussions. The work conducted by the US Department of Energy Joint Genome Institute is supported by the Office of Science of the US Department of Energy under Contract No. DE-AC02-05CH11231. Research in the Rio lab is supported by NIH R01 GM61987. Research in the Rokhsar lab is supported by a grant from R.A. Melmon and the Gordon and Betty Moore Foundation.

Author contributions

U.H. and J.L.A. equally contributed to this work. U.H. conceived the general idea for the experiment and carried out the bioinformatics analysis. J.L.A. performed the experiments and data analysis, and developed the detailed protocols. D.D.R. contributed ideas and supervision for the experiments. D.S.R. contributed ideas and supervision for the bioinformatics. All authors contributed to the writing, comments, and discussions of the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Hellsten, U. *et al.* A segmental genomic duplication generates a functional intron. *Nat. Commun.* **2**:454 doi: 10.1038/ncomms1461 (2011).

License: This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>