

An insight into more factors affecting adherence to medications in the elderly

Syed Maaz Tariq
Mohammad Hasan
Syed Ali Haider

Department of MBBS, Jinnah Sindh
Medical University, Karachi, Pakistan

Dear editor

We read the article “Medication adherence and beliefs about medication in elderly patients living alone with chronic diseases” by Park et al¹ thoroughly. They were right in stating that identification of barriers to patient adherence to a medication holds vital importance, and despite a proper plan devised by the physician, if the patient does not adhere to medication, management of a chronic illness can prove to be challenging.¹

As part of a society where “quacks”, alternative/herbal medicine, and faith “healers” are quite popular, this has an impact on patient approaches to medication.² Therefore, we would like to carry forward the discussion from this perspective with a broader viewpoint adjoining some factors other than patient beliefs that influence their attitude toward medication. A study was conducted in Malaysia that concluded that more than 40% people showed positive attitudes toward complementary and alternative medication (CAM).² Conventional medications prescribed for chronic illnesses often cause unwanted side effects. On the other hand, CAM has substantially fewer unwanted effects, and these outcomes make the elder patients opt for it instead of physician-prescribed drugs.³ In addition to this, dismissive attitudes of health-care providers toward old patients further discourages doctor–patient relationships in the elderly. Generally, seniors are considered a burden on economy, with many other stereotypes.⁴ This leads to poor care of this population, who then seek alternative easy methods for treatment. Concealment of CAM is a general trend within an elderly population, which further makes it difficult for doctors to realize that the patient has switched treatment.²

This is a wakeup call for health-care professionals to improve their communication with elderly people and approach them in such a way that can convince them to stick to their required medication. The doctor’s role is of much importance in assessing whether or not the alternative treatment taken by the patient is suitable or if it is debilitating the condition, let alone improving it. Enhancement of physician–patient communication is an effective method to improve adherence.⁵

It is reported that for chronic diseases, patients withdraw from medication within the initial few months of treatment.⁵ This is where the role of health-care workers is of great importance. There ought to be proper follow-up of each elderly patient, in order to make sure they take medicines in a timely fashion. Despite studies suggesting elderly people will be a large part of the population in the near future, there remains less development in the field of geriatrics. As such, geriatric programs should be introduced in medical colleges to develop the young minds of future doctors to deal with this situation. There is a need for further studies to assess such factors as alternative/herbal

Correspondence: Syed Maaz Tariq
Flat A2, Block H, Mariam Centre
Phase I, North Nazimabad, Karachi,
74600, Pakistan
Email maaztariqsyed@gmail.com

medicine and spiritual “healers”, since this would result in more insightful outcomes in this domain. Living alone could also be a factor for lower adherence, and thus to evaluate this aspect further, Park et al’s¹ results could be compared to those elderly living with their families.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Park HY, Seo SA, Yoo HY, Lee KH. Medication adherence and beliefs about medication in elderly patients living alone with chronic diseases. *Patient Prefer Adherence*. 2018;12:175–181.
2. Islahudin F, Shahdan IA, Mohamad-Samuri S. Association between belief and attitude toward preference of complementary alternative medicine use. *Patient Prefer Adherence*. 2017;11:913–918.
3. Ennis E. Complementary and alternative medicines (CAMs) and adherence to mental health medications. *BMC Complement Alternat Med*. 2014;14:93.
4. Lovell M. Caring for the elderly: changing perceptions and attitudes. *J Vasc Nur*. 2006;24:22–26.
5. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med*. 2005; 353:487–497.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Patient Preference and Adherence ‘letters to the editor’ section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Patient Preference and Adherence editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>