



Contents lists available at ScienceDirect

# Journal of Ginseng Research

journal homepage: <https://www.sciencedirect.com/journal/journal-of-ginseng-research>



## Review Article

# Ginseng, a promising choice for SARS-CoV-2: A mini review



Zubair Ahmed Ratan <sup>a, b</sup>, Fazla Rabbi Mashrur <sup>a</sup>, Nusrat Jahan Runa <sup>c</sup>, Ki Woong Kwon <sup>d</sup>, Hassan Hosseinzadeh <sup>b</sup>, Jae Youl Cho <sup>d, e, f\*</sup>

<sup>a</sup> Department of Biomedical Engineering, Khulna University of Engineering & Technology, Khulna, Bangladesh

<sup>b</sup> School of Health and Society, University of Wollongong, NSW, Australia

<sup>c</sup> Sylhet MAG Osmani Medical College, Sylhet, Bangladesh

<sup>d</sup> Department of Integrative Biotechnology, Sungkyunkwan University, Suwon, Republic of Korea

<sup>e</sup> Department of Biocosmetics, Sungkyunkwan University, Suwon, Republic of Korea

<sup>f</sup> Research Institute of Biomolecule Control and Biomedical Institute for Convergence at SKKU, Suwon, Republic of Korea

## ARTICLE INFO

### Article history:

Received 29 December 2021

Received in revised form

15 January 2022

Accepted 16 January 2022

Available online 22 January 2022

### Keywords:

Korean ginseng

SARS-CoV-2

Anti-viral immunity

Anti-viral activity

## ABSTRACT

The current Covid-19 pandemic has changed the entire world and brought so many unprecedented challenges to the scientific community. More than 5 million people died due to the SARS-CoV-2 outbreak. For many thousands of years, ginseng, the traditional herb has been used for various infectious diseases by traditional healers. Ginseng showed promising antiviral effects by modulating both natural and acquired immunity. Ginseng might be used as a potential therapeutic agent to prevent SARS-CoV-2 infection along with the vaccine. In this current review, we offer an alternative approach for SARS-CoV-2 prevention during this unprecedented pandemic.

© 2022 The Korean Society of Ginseng. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

SARS-CoV-2 caused significant casualties with huge social and economic impacts all over the world, and it imposed unprecedented challenges in the entire health systems of the world [1]. The Covid-19 has affected all the territories of the world, since its first appearance in Wuhan, China [2,3]. SARS-CoV-2 is a single-stranded, positive-sense, RNA-enveloped virus [4]. In the recent past, Ebola, Zika and SARS viruses also brought considerable attention to the scientific community [5]. The “Black death” (plague) from the fourteenth century or the “Spanish Flu of 1918” has had profound effects on society’s human history [6]. Despite many challenges, the discovery and development of the SARS-CoV-2 vaccine are remarkable, however, still, there is lack of standard pharmaceutical cure for this deadly infection [7]. However, with conventional therapies, the use of traditional medicines can play a

pivotal role in Covid treatment. China, Korea, and some of the African countries have had a long standing tradition of using herbal medicine [8,9]. Interestingly, ginseng, “the king of all herbs”, might play a significant role during Covid-19 infection as it can modulate both natural and acquired immunity during viral infection [10].

Traditional use of ginseng could be traced for thousands of years in China and Korea [11]. Ginseng belongs to the Araliaceae family, however, it has more than eight to thirteen species under this genus, and among those the *P. ginseng* is commonly used, widely famous as “Asian or Chinese ginseng”. Interestingly, the word “Panax”, named after the Greek word “panacea,” which means “the cure for all diseases” [12]. Studies suggested active components of ginseng, ginsenosides can act as anti-viral or anti-bacterial agents [10,13]. Different *in vitro* and animal studies revealed the antiviral action of ginseng (Table 1). The influenza A or B viruses are most common cause of seasonal flu [14]. One study revealed that red ginseng could be helpful in preventing this lethal infection of the H5N1 influenza virus [15]. Interestingly another study found that ginsenosides can act as a mucosal adjuvant against the influenza virus [16]. One clinical study suggested that ginseng can induce genetic defects in the nef gene in AIDS (acquired immunodeficiency syndrome) patients [17]. Moreover, the study found that

\* Corresponding author. Department of Integrative Biotechnology, Sungkyunkwan University, Suwon, Republic of Korea.

E-mail addresses: [zubairahmed@bme.kuet.ac.bd](mailto:zubairahmed@bme.kuet.ac.bd), [zar241@uowmail.edu.au](mailto:zar241@uowmail.edu.au) (Z.A. Ratan), [rabbi.mashrur@gmail.com](mailto:rabbi.mashrur@gmail.com) (F. Rabbi Mashrur), [nusratrunga17@gmail.com](mailto:nusratrunga17@gmail.com) (N.J. Runa), [nexus0322@naver.com](mailto:nexus0322@naver.com) (K.W. Kwon), [hassanh@uow.edu.au](mailto:hassanh@uow.edu.au) (H. Hosseinzadeh), [jaecho@skku.edu](mailto:jaecho@skku.edu) (J.Y. Cho).

**Table 1**

Antiviral effect of ginseng.

Viral Infection	Study type	Sample	Result	Ref.
Human immunodeficiency virus (HIV)-1	Clinical study	Red ginseng extract	Combination with zidovudine (antiretroviral medication) showed promising antiviral effects by preserving CD4 <sup>+</sup> T cell counts	[19–21]
Hepatitis B virus	In vitro study	Ginsenoside Rg3	The secretion of HBsAg, HBeAg in HBV-infected HepG2.2.15 cells was inhibited by ginsenoside Rg3	[18]
Hepatitis A virus	In vitro study	ginsenosides Rb1 and Rg1	ginsenosides Rb1 and Rg1 decreased the HAV titer	[22]
Rotavirus	In vitro study	hot water extract of ginseng	Pectic polysaccharides from <i>Panax ginseng</i> inhibited rotavirus infection	[23]
Influenza virus	In vitro study, in vivo study	Korean Red Ginseng (KRG)	Ginseng treatment can improve the viability of human alveolar epithelial A549 cells	[24]

ginsenoside Rg3 can inhibit the JNK/AP-1 signaling in Hepatitis B virus-infected HepG2.2.15 cells [18].

The SARS-CoV-2 virus is continuously mutating as the countries are racing to vaccinate people as fast as possible. Pneumonia is the common complications of SARS-CoV-2 infection. This infection attenuates body's immune system as a result pneumonia occurs. Previous scientific evidence suggests the antiviral properties of ginseng, which could be effective against Covid 19 as well. Ginseng could act as an adjuvant in SARS-CoV-2 by modulating both natural and acquired immunity.

## 2. Impact of ginseng on immunity

Ginseng is well-known for its adaptogenic response to immune functions [25,26]. Innate immunity which is regarded as the body's initial line of protection against numerous infections. This immunity refers to nonspecific defensive mechanism mediated by a varied range of receptors from multiple protein families [27]. This immune system involves both physical and chemical barriers. Macrophages act as an effector cell in the innate immune system that are capable of secreting both antimicrobial and pro-inflammatory mediators [28]. Studies found that ginseng showed the immunomodulatory effect on macrophage, the wild ginseng extract can upregulate the Nitric oxide (NO) production, and the cytokines production including IL-1 $\alpha$ , IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and granulocyte-macrophage colony-stimulating factor [29]. Ginseng can increase the phagocytic activities of the reticuloendothelial system and showed potential antitumor effects against sarcoma in animal model [30]. Ginsan, a fraction of *P. ginseng*, also showed immunopotentiation effect on peritoneal macrophages [31]. Dendritic cells (DCs), play an important role in linking with innate and adaptive immune system, and these cells are responsible for antigen-specific immune responses [32]. Ginseng also showed immunomodulatory effects on DCs, one study found that it can suppress the cytokine production and also the maturation of LPS-treated DCs with downregulating the CD4 (+) T cells [33]. Another study revealed that it can enhance the surface co-stimulatory molecules, including MHC class II, CD40, CD80, and CD86, on bone marrow-derived DCs (BMDC) [34]. Natural killer (NK) cells play a pivotal role in innate immune responses by combating infection [35]. Study found that *P. ginseng* can increase the natural killing activity in animal model [36,37]. Another clinical study revealed that *Panax ginseng* can significantly enhance the NK cells in AIDS patients [38].

Adaptive immunity is a type of defensive mechanism that offers a more specialized store of recollection for both self-antigens and foreign antigens. This response mechanism emerged from immunologic memory and is characterized by tightly controlled communication between antigen-presenting cells and lymphocytes. Lymphocytes' characteristic fighters are B and T cells, which are continually created from progenitor stem cells found in the

bone marrow and thymus [35]. Ginseng also showed immunomodulatory effects on the cell-mediated immunity, study found that the red ginseng extract can immunosuppress the regulation in animal model by promoting IFN- $\gamma$  + CD4 + T cells, IFN- $\gamma$  T cell (reg) production [39]. Different studies also revealed that ginseng can significantly increase the antibody-dependent cellular cytotoxicity and NK cell activity in *in vitro* and *in vivo* studies [40]. One *in vivo* study attributed that the Rb1, fraction of ginseng adjuvanted vaccine can stimulate the titers of antigen specific IgG1, IgG2a and IgG2b and helped in balancing Th1 and Th2 immune response [41].

## 3. Potential of ginseng in treating SARS-CoV-2

Researchers used different computational techniques to investigate a natural chemical ingredient of ginseng as a potential inhibitor of the primary protease of SARS-CoV-2. Fallah et al.(2021) [42] used seven compounds from *Panax ginseng* and performed molecular docking, the glide docking program (Table 2). Flor-alginoside B extracted from *Panax ginseng*, showed promising binding affinity (-8.618 kcal/mol) with the receptor-binding domain of SARS-CoV-2. This could be a potential candidate for combating the SARS-CoV-2 infection. Han et al. [43] investigated Shenmai Injection's (consists of red ginseng + ophiopogon japonicus) molecular interaction network route in the treatment of COVID-19 patients having heart disease and they found that Shenmai Injection can be effective in treating COVID-19 by reducing cytokine storms. Li et al. [44] used Shenfu decoction to treat patients with severe COVID-19 and the study revealed that, it could play an essential role in regulating the anti-viral process, regulating immunity, and reducing inflammation. Moreover, Wang et al. [45] found that the active chemicals in Shengmai injection (consists of red ginseng, ophiopogon japonicus, and schisandra chinensis) can modulate several signaling pathways including CASP3, PTGS2, NOS2, NOS3 which might play a pivotal role in COVID-19 treatment. Furthermore, another study [46] proposed that *Panax ginseng* can reduce chronic fatigue syndrome and Fibromyalgia, the common symptoms of COVID-19. Again, Han et al.(2021) [47] experimented with fermented Ginseng (black) and reported that it not only stops SARS-CoV-2 from replicating in the cell, but it also lowers the quantity of viral RNA (vRNA) copies floating around in the extracellular environment.

## 4. Ginseng as an adjuvant

Adjuvants are compounds or components that can improve vaccine efficacy [49], ginseng and its ingredients ginsenosides could be potential adjuvants to improve Covid infection. Previously, *P. ginseng* has been used to improve immunity, memory, menopausal symptoms, cardiovascular dysfunction, neurological disorder [12,50–52]. The human immune system is a complex network that defends the body from different infections. Poor nutrition,


**Table 2**

Antiviral components in ginseng.

Country	Compounds	Study Type	Findings	Ref.
Iran	Seven compounds from Panax ginseng	Molecular docking, Glide docking program.	Floralginsenoside B extracted from <i>P.ginseng</i> , showed promising effect, it had strong binding affinity with the receptor-binding domain of SARS-CoV-2.	[42]
China	Shenmai Injection (Red ginseng + ophiopogon japonicus)	Network pharmacology	Shenmai injection can treat COVID-19 by reducing cytokine storms, demonstrating a network control mechanism of mutual influence and complicated correlation.	[43]
China	Shenfu decoction (water extract of the dried root of <i>Panax ginseng</i> ) + root of Aconitum carmichaeli Debx	Network pharmacology	To treat patients with severe COVID-19, shenfu decoction played an essential role in regulating the anti-virus process, regulating immunity, and reducing inflammation.	[44]
China	Shengmai injection (Red ginseng + ophiopogon japonicus + schisandra chinensis)	Network pharmacology and Molecular docking	The active chemicals in Shengmai Injection modulate several signaling pathways for anti-inflammatory, immunological modulation, anti-shock, and boosting blood oxygen saturation, all of which may play a role in COVID-19 treatment.	[45]
United States of America Korea	<i>Panax ginseng</i> root extract (HRG80™ Red Ginseng) Fermented <i>Panax ginseng</i> (Black)	Prospective study <i>In vitro</i>	People suffering from chronic fatigue syndrome and Fibromyalgia (symptoms of COVID-19) yield significant improvement Ginseng lower the quantity of viral RNA (vRNA) copies floating around in the extracellular environment.	[46] [48]

certain chronic conditions, burn and drugs can weaken our immune system [53,54]. Natural products are the good source of phytochemicals which are capable of enhancing immune function [10]; interestingly, ginseng has been shown to use as a dietary supplement that can also enhance both types of immunity [55–57].

For example, orally administered ginseng leaf-stem saponins (GSLS) in combination with a live infectious bursal disease vaccination improved both humoral and gut mucosal immune responses in chickens [58]. In other studies, ginsan, a polysaccharide extracted from *Panax ginseng*, showed different immunomodulatory effects



**Fig. 1.** Schematic diagram of anti-viral activity by Korean ginseng. Ginseng is a potent anti-viral agent by enhancing innate and adaptive immunity and adjuvant action.

by inhibiting the nuclear factor-kappa B (NF- $\kappa$ B) pathway and p38 MAP kinase [59]. Moreover, studies narrated the anti-inflammatory effect of ginseng [60–63]. Different adjuvants like toll-like receptor (TLR) agonist, aluminum salt, and emulsions have been used for vaccine formulations in animal models [64]. In this current situation, ginseng should be a choice of adjuvant during Covid infection.

## 5. Conclusions

The Covid-19 pandemic has posed a serious public health threat in the form of a deadly pandemic. To combat this deadly SARS-CoV-2, it is important to identify promising and successful targeted tactics. Apart from chemical and allopathic medications, naturally occurring compounds can play a pivotal role as antiviral agents. The pathogenesis of SARS-CoV-2 is similar to that of other viral infections. Taken together, according to the previous finding, ginseng is a potent anti-viral agent by enhancing innate and adaptive immunity and adjuvant action (Fig. 1), so it could be a game-changer in Covid treatment, however, clinical trials studies are required for further development.

## Declaration of competing interest

The authors declare no conflict of interest.

## Acknowledgements

This research was funded by the Basic Science Research Program through the National Research Foundation of Korea (NRF), the Ministry of Science and ICT, South Korea (2017R1A6A1A03015642) and also supported by the Korean society of ginseng (2020), South Korea.

## References

- [1] Haidere MF, Ratan ZA, Nowroz S, Zaman SB, Jung Y-J, Hosseinzadeh H, Cho JY. COVID-19 vaccine: critical questions with complicated answers. *Biomol Therapeut* 2021;29:1.
- [2] Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, Si H-R, Zhu Y, Li B, Huang C-L, Chen H-D, Chen J, Luo Y, Guo H, Jiang R-D, Liu M-Q, Chen Y, Shen X-R, Wang X, Zheng X-S, Zhao K, Chen Q-J, Deng F, Liu L-L, Yan B, Zhan F-X, Wang Y-Y, Xiao G-F, Shi Z-L. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–3.
- [3] Mashrur FR, Roy AD, Chhoan AP, Sarker S, Saha A, Hasan SMN, Saha S. Impact of demographic, environmental, socioeconomic, and government intervention on the spreading of COVID-19. *Clin Epidemiol Glob Health* 2021;12:100811.
- [4] Pal M, Berhanu G, Desalegn C, Kandi V. Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2): an update. *Cureus* 2020;12. e7423-e7423.
- [5] Zhang L, Zhao W, Sun B, Huang Y, Glänzel W. How scientific research reacts to international public health emergencies: a global analysis of response patterns. *Scientometrics* 2020;1–27.
- [6] Huremović D. Brief history of pandemics (Pandemics throughout history). Psychiatry of pandemics: a mental health response to infection outbreak. 2019. p. 7–35.
- [7] Yimer G, Ekuadzi E, Fasinu P, de Melo AC, Pillai G. Traditional medicines for COVID-19: perspectives from clinical pharmacologists. *Br J Clin Pharmacol* 2021;87:3455–8.
- [8] Anyinam C. Availability, accessibility, acceptability, and adaptability: four attributes of African ethno-medicine. *Soc Sci Med* 1987;25:803–11.
- [9] Chaudhury RR, Rafei UM. Traditional medicine in Asia. Geneva: World Health Organization; 2002.
- [10] Ratan ZA, Youn SH, Kwak Y-S, Han C-K, Haidere MF, Kim JK, Min H, Jung Y-J, Hosseinzadeh H, Hyun SH, Cho JY. Adaptogenic effects of *Panax ginseng* on modulation of immune functions. *J. Ginseng Res.* 2021;45:32–40.
- [11] Nair R, Sellaturay S, Srivatsad S. The history of ginseng in the management of erectile dysfunction in ancient China (3500–2600 BCE). *Indian J Urol* 2012;28:15–20.
- [12] Ratan ZA, Haidere MF, Hong YH, Park SH, Lee J-O, Lee J, Cho JY. Pharmacological potential of ginseng and its major component ginsenosides. *J. Ginseng Res.* 2021;45:199–210.
- [13] Adusei-Mensah F, Agiee RO, Awoniyi LO, David LK, Awoniyi FB, Adeyemi OS, Olawuni A, Adegbite A. Prevention and control of acute respiratory viral infections in adult population: a systematic review and meta-analysis on ginseng-based clinical trials. *medRxiv* 2021. 2021.2007.21260970.
- [14] Taubenberger JK, Morens DM. The pathology of influenza virus infections. *Ann Rev Pathol* 2008;3:499–522.
- [15] Park EH, Yum J, Ku KB, Kim HM, Kang YM, Kim JC, Kim JA, Kang YK, Seo SH. Red Ginseng-containing diet helps to protect mice and ferrets from the lethal infection by highly pathogenic H5N1 influenza virus. *J. Ginseng Res.* 2014;38:40–6.
- [16] Quan FS, Compans RW, Cho YK, Kang SM. Ginseng and Salviae herbs play a role as immune activators and modulate immune responses during influenza virus infection. *Vaccine* 2007;25:272–82.
- [17] Cho Y-K, Kim J-E, Woo J-H. Genetic defects in the nef gene are associated with Korean Red Ginseng intake: monitoring of nef sequence polymorphisms over 20 years. *J. Ginseng Res.* 2017;41:144–50.
- [18] Kang IJ, Choi YJ, Lee SG. Stimulation of TRAF6/TAK1 degradation and inhibition of JNK/AP-1 signalling by ginsenoside Rg3 attenuates hepatitis B virus replication. *Int J Biochem Cell Biol* 2013;45:2612–21.
- [19] Cho YK, Sung H, Lee HJ, Joo CH, Cho GJ. Long-term intake of Korean red ginseng in HIV-1-infected patients: development of resistance mutation to zidovudine is delayed. *Int Immunopharmacol* 2001;1:1295–305.
- [20] Kim BR, Kim JE, Sung H, Cho YK. Long-term follow up of HIV-1-infected Korean haemophiliacs, after infection from a common source of virus. *Hae-mophilia* 2015;21:e1–11.
- [21] Cho Y-K, Kim YK, Lee IC, Choi M, Shin YO. The effect of Korean Red Ginseng (KRG), zidovudine (ZDV), and the combination of KRG and ZDV on HIV-infected individuals. *J Bacteriol Virol* 1996;31:353–60.
- [22] Lee MH, Lee BH, Lee S, Choi C. Reduction of hepatitis A virus on FRhK-4 cells treated with Korean red ginseng extract and ginsenosides. *J Food Sci* 2013;78: M1412–5.
- [23] Baek SH, Lee JG, Park SY, Bae ON, Kim DH, Park JH. Pectic polysaccharides from *Panax ginseng* as the antirotavirus principals in ginseng. *Biomacromolecules* 2010;11:2044–52.
- [24] Lee JS, Hwang HS, Ko E-J, Lee Y-N, Kwon Y-M, Kim M-C, Kang S-M. Immunomodulatory activity of red ginseng against influenza A virus infection. *Nutrients* 2014;6:517–29.
- [25] CHOI K-T. Botanical characteristics, pharmacological effects and medicinal components of Korean *Panax ginseng* CA Meyer. *Acta Pharmacol Sin* 2008;29: 1109–18.
- [26] Kitts DD, Hu C. Efficacy and safety of ginseng. *Publ Health Nutr* 2000;3: 473–85.
- [27] Medzhitov R, Janeway CA. Innate immunity: the virtues of a nonclonal system of recognition. *Cell* 1997;91:295–8.
- [28] Hirayama D, Iida T, Nakase H. The phagocytic function of macrophage-enforcing innate immunity and tissue homeostasis. *Int J Mol Sci* 2017;19.
- [29] Jang HI, Shin HM. Wild *Panax ginseng* (*Panax ginseng* CA Meyer) protects against methotrexate-induced cell regression by enhancing the immune response in RAW 264.7 macrophages. *Am J Chin Med* 2010;38:949–60.
- [30] Youn SH, Lee SM, Han C-K, In G, Park C-K, Hyun SH. Immune activity of polysaccharide fractions isolated from Korean Red Ginseng. *Molecules* 2020;25:3569.
- [31] Shin J-Y, Song J-Y, Yun Y-S, Yang H-O, Rhee D-K, Pyo S. Immunostimulating effects of acidic polysaccharides extract of *Panax ginseng* on macrophage function. *Immunotoxicol Immunopharmacol* 2002;24:469–82.
- [32] Mellman I. Dendritic cells: master regulators of the immune response. *Canc Immunol Res* 2013;1:145–9.
- [33] Lee YJ, Son YM, Gu MJ, Song KD, Park SM, Song HJ, Kang JS, Woo JS, Jung JH, Yang DC, Han SH, Yun CH. Ginsenoside fractions regulate the action of monocytes and their differentiation into dendritic cells. *J. Ginseng Res.* 2015;39:29–37.
- [34] Kang S, Min H. Ginseng, the 'Immunity Boost': the effects of *Panax ginseng* on immune system. *J. Ginseng Res.* 2012;36:354–68.
- [35] Iwasaki A, Medzhitov R. Control of adaptive immunity by the innate immune system. *Nat Immunol* 2015;16:343–53.
- [36] Jie YH, Cammisuli S, Baggiolini M. Immunomodulatory effects of *Panax ginseng* C.A. Meyer in the mouse. *Agents Actions* 1984;15:386–91.
- [37] Kenarova B, Neychev H, Hadjiivanova C, Petkov VD. Immunomodulating activity of ginsenoside Rg1 from *Panax ginseng*. *Jpn J Pharmacol* 1990;54: 447–54.
- [38] See DM, Broumand N, Sahl L, Tilles JG. *In vitro* effects of echinacea and ginseng on natural killer and antibody-dependent cell cytotoxicity in healthy subjects and chronic fatigue syndrome or acquired immunodeficiency syndrome patients. *Immunopharmacology* 1997;35:229–35.
- [39] Heo SB, Lim SW, Jhun JY, Cho MI, Chung BH, Yang CW. Immunological benefits by ginseng through reciprocal regulation of Th17 and Treg cells during cyclosporine-induced immunosuppression. *J. Ginseng Res.* 2016;40:18–27.
- [40] Kim JY, Germolec DR, Luster MI. *Panax ginseng* as a potential immunomodulator: studies in mice. *Immunopharmacol Immunotoxicol* 1990;12:257–76.
- [41] Rivera E, Ekholm Pettersson F, Inganäs M, Paulie S, Grönvik KO. The Rb1 fraction of ginseng elicits a balanced Th1 and Th2 immune response. *Vaccine* 2005;23:5411–9.
- [42] Fallah MS, Bayati M, Najafi A, Behmard E, Javad S. Molecular docking investigation of antiviral herbal compounds as potential inhibitors of SARS-CoV-2 spike receptor. *Biointerf Res Appl Chem* 2021;11:12916–24.
- [43] Han L-W. Network pharmacologic molecular mechanism of Shenmai Injection in treatment of COVID-19 combined with coronary heart disease. *Chin Tradit Herb Drugs* 2020;23:344–44.

- [44] Li X, Lin H, Wang Q, Cui L, Luo H, Luo L. Chemical composition and pharmacological mechanism of shenfu decoction in the treatment of novel coronavirus pneumonia (COVID-19). *Drug Dev Ind Pharm* 2020;46:1947–59.
- [45] Wang L-F, Li H-T, Wang Y, Liu X-L, Chen Q-Y, Xu J, Yang M, Zhang X-F, Wang F. Study on mechanism of Shengmai Injection against novel coronavirus pneumonia based on network pharmacology and molecular docking technology. *Chin Tradit Herb Drugs* 2020;29:77–87.
- [46] Teitelbaum J, Goudie S. An open-label, pilot trial of HRG80™ red ginseng in chronic fatigue syndrome, fibromyalgia, and post-viral fatigue. *Pharmaceutics* 2022;15:43.
- [47] Khanal P, Chikhale R, Dey YN, Pasha I, Chand S, Gurav N, Ayyanar M, Patil BM, Gurav S. Withanolides from *Withania somnifera* as an immunity booster and their therapeutic options against COVID-19. *J Biomol Struct Dyn* 2021;1:1–14.
- [48] Han Y, Yang D-U, Huo Y, Pu J, Lee S-J, Yang D-C, Kang SC. *In vitro* evaluation of anti-lung cancer and anti-COVID-19 effects using fermented black color ginseng extract. *Nat Prod Commun* 2021;16:1934578X211034387.
- [49] Coffman RL, Sher A, Seder RA. Vaccine adjuvants: putting innate immunity to work. *Immunity* 2010;33:492–503.
- [50] Chen F, Eckman EA, Eckman CB. Reductions in levels of the Alzheimer's amyloid beta peptide after oral administration of ginsenosides. *FASEB (Fed Am Soc Exp Biol) J* 2006;20:1269–71.
- [51] Choi S-H, Jung S-W, Lee B-H, Kim H-J, Hwang S-H, Kim H-K, Nah S-Y. Ginseng pharmacology: a new paradigm based on gintonin-lysophosphatidic acid receptor interactions. *Front Pharmacol* 2015;6.
- [52] Leung KW, Wong AS-T. Pharmacology of ginsenosides: a literature review. *Chin Med* 2010;5:20.
- [53] Chapel H, Prevot J, Gaspar HB, Español T, Bonilla FA, Solis L, Drabwell J. Primary immune deficiencies—principles of care. *Front Immunol* 2014;5:627.
- [54] Raje N, Dinakar CJL, Clinics A. Overview Immunodef Disorder 2015;35: 599–623.
- [55] Zhou Y, Yang Z, Gao L, Liu W, Liu R, Zhao J, You J. Changes in element accumulation, phenolic metabolism, and antioxidative enzyme activities in the red-skin roots of *Panax ginseng*. *J. Ginseng Res.* 2017;41:307–15.
- [56] Yun SN, Moon SJ, Ko SK, Im BO, Chung SH. Wild ginseng prevents the onset of high-fat diet induced hyperglycemia and obesity in ICR mice. *Arch Pharm Res (Seoul)* 2004;27:790–6.
- [57] Lee H, Lee S, Jeong D, Kim SJ. Ginsenoside Rh2 epigenetically regulates cell-mediated immune pathway to inhibit proliferation of MCF-7 breast cancer cells. *J. Ginseng Res.* 2018;42:455–62.
- [58] Zhai L, Wang Y, Yu J, Hu S. Enhanced immune responses of chickens to oral vaccination against infectious bursal disease by ginseng stem-leaf saponins. *Poultry Sci* 2014;93:2473–81.
- [59] Friedl R, Moeslinger T, Kopp B, Speckermann PG. Stimulation of nitric oxide synthesis by the aqueous extract of *Panax ginseng* root in RAW 264.7 cells. *Br J Pharmacol* 2001;134:1663–70.
- [60] Kim JH, Yi Y-S, Kim M-Y, Cho JY. Role of ginsenosides, the main active components of *Panax ginseng*, in inflammatory responses and diseases. *J. Ginseng Res.* 2017;41:435–43.
- [61] JKSS Kim JP, Yoo HY, Yu HJ, Yang HKJ. The effect of ginseng on the nutritional status and the immune functions after curative operations on gastric carcinoma patients. *J Korean Surg Soc* 1998;54:854.
- [62] Lee KG, Son SW. Efficacy of Korean red ginseng in the treatment of atopic dermatitis. *J. Ginseng Res.* 2011;35:149.
- [63] Yun TK, Choi SY, Yun HY. Epidemiological study on cancer prevention by ginseng: are all kinds of cancers preventable by ginseng? *J Kor Med Sci* 2001;16:S19.
- [64] Liang Z, Zhu H, Wang X, Jing B, Li Z, Xia X, Sun H, Yang Y, Zhang W, Shi L, Zeng H, Sun B. Adjuvants for coronavirus vaccines. *Front Immunol* 2020;11: 589833.