

RESEARCH ARTICLE

Exploring the insulin secretory properties of the PGD₂-GPR44/DP2 axis *in vitro* and in a randomized phase-1 trial of type 2 diabetes patients

Stanko Skrtic^{1,2*}, Björn Tyrberg¹, Malin Broberg¹, Hans Ericsson¹, Volker Schneck¹, Magnus Kjaer¹, Marcus Hompesch³, Eva-Marie Andersson¹, Erik Ryberg¹, Alexander Aivazidis¹, Charlotte Wennberg Hult¹, Lars Löfgren¹, Linda Morrow³, Joanna Parkinson¹, Tina Rydén-Bergsten¹, Elaine Watkins³, Maria Sörhede Winzell¹

1 Cardiovascular, Renal and Metabolism, IMED Biotech Unit, AstraZeneca Gothenburg, Mölndal, Sweden, **2** Department of Endocrinology, Sahlgrenska University Hospital and Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden, **3** ProSciento Inc, Chula Vista, CA, United States of America

* stanko.skrtic@astrazeneca.com



OPEN ACCESS

Citation: Skrtic S, Tyrberg B, Broberg M, Ericsson H, Schneck V, Kjaer M, et al. (2018) Exploring the insulin secretory properties of the PGD₂-GPR44/DP2 axis *in vitro* and in a randomized phase-1 trial of type 2 diabetes patients. PLoS ONE 13(12): e0208998. <https://doi.org/10.1371/journal.pone.0208998>

Editor: Elena Barengolts, University of Illinois College of Medicine, UNITED STATES

Received: April 9, 2018

Accepted: October 8, 2018

Published: December 17, 2018

Copyright: © 2018 Skrtic et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Clinical Study Data are available from the AstraZeneca AB Institutional Data Access for researchers who meet the criteria for access to confidential data.

Funding: AstraZeneca AB (SE) financed the submitted work. The funder provided support in the form of salaries for authors SS, BT, MB, HE, VS, E-MA, ER, AA, CW-H, LL, JP, TR-B, MSW, but did not have any additional role in the study design, data collection and analysis, decision to publish, or

Abstract

Aims/Hypothesis

GPR44 (*DP2*, *PTGDR2*, *CRTH2*) is the receptor for the pro-inflammatory mediator prostaglandin D₂ (PGD₂) and it is enriched in human islets. In rodent islets, PGD₂ is produced in response to glucose, suggesting that the PGD₂-GPR44/DP2 axis may play a role in human islet function during hyperglycemia. Consequently, the aim of this work was to elucidate the insulinotropic role of GPR44 antagonism *in vitro* in human beta-cells and in type 2 diabetes (T2DM) patients.

Methods

We determined the drive on PGD₂ secretion by glucose and IL-1beta, as well as, the impact on insulin secretion by pharmacological GPR44/DP2 antagonism (AZD1981) in human islets and beta-cells *in vitro*. To test if metabolic control would be improved by antagonizing a hyperglycemia-driven increased PGD₂ tone, we performed a proof-of-mechanism study in 20 T2DM patients (average 54 years, HbA1c 9.4%, BMI 31.6 kg/m²). The randomized, double-blind, placebo-controlled cross-over study consisted of two three-day treatment periods (AZD1981 or placebo) separated by a three-day wash-out period. Mixed meal tolerance test (MMTT) and intravenous graded glucose infusion (GGI) was performed at start and end of each treatment period. Assessment of AZD1981 pharmacokinetics, glucose, insulin, C-peptide, glucagon, GLP-1, and PGD₂ pathway biomarkers were performed.

Results

We found (1) that PGD₂ is produced in human islet in response to high glucose or IL-1beta, but likely by stellate cells rather than endocrine cells; (2) that PGD₂ suppresses

preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing interests: All authors apart from MH, LM and EW are employees or shareholders of AstraZeneca AB. MH, LM, and EW received funding as a contract research organization by AstraZeneca for the clinical study. EW is affiliated with ProSciento Inc. There are no patents, products in development, or marketed products to declare. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials.

both glucose and GLP-1 induced insulin secretion *in vitro*; and (3) that the GPR44/DP2 antagonist (AZD1981) in human beta-cells normalizes insulin secretion. However, AZD1981 had no impact on neither glucose nor incretin dependent insulin secretion in humans (GGI AUC_{C-peptide 1-2h} and MMTT AUC_{Glucose 0-4h} LS mean ratios vs placebo of 0.94 (80% CI of 0.90–0.98, $p = 0.12$) and 0.99 (90% CI of 0.94–1.05, $p = 0.45$), despite reaching the expected antagonist exposure.

Conclusion/Interpretation

Pharmacological inhibition of the PGD₂-GPR44/DP2 axis has no major impact on the modulation of acute insulin secretion in T2DM patients.

Trial registration

ClinicalTrials.gov [NCT02367066](https://clinicaltrials.gov/ct2/show/study/NCT02367066).

Introduction

Type 2 diabetes mellitus (T2DM) is characterized by a relative loss of functional beta-cell mass leading to insufficient insulin secretion during insulin resistance. It is believed that beta-cell insufficiency is caused by a combination of beta-cell death (apoptosis) and loss-of-function. Beta-cell functional loss may to some degree be explained by dedifferentiation to a cell without the capacity to produce or secrete insulin [1]. Therefore, disease modification in T2DM requires functional beta-cell mass restoration [2] through mechanisms preventing apoptosis or restoring loss-of-function. Although the beta-cell insulin secretory machinery is well understood and pharmacological treatments are available that increase insulin secretion, such as sulphonylureas and GLP-1 receptor agonists, there is a lack of drugs on the market that retain efficiency over time and truly modulate disease mechanisms coupled to insulin insufficiency. Prostaglandin D₂ (PGD₂) is a pro-inflammatory mediator where one of its receptors, GPR44 (DP2, *PTGDR2*, *CRT2*) so far has been of particular interest to respiratory medicine [3]. Several selective GPR44/DP2 receptor antagonists were taken forward into clinical development as potential anti-asthmatic therapeutics, including AZD1981 developed by AstraZeneca [4–6]. Interestingly, in the human pancreas GPR44/DP2 is enriched in the islets versus the exocrine tissue [7, 8]. In fact, it is among the more highly expressed islet GPCRs according to a recent GPCRome survey [9]. Nevertheless, the function of this receptor in the endocrine pancreas is completely unknown. GPR44/DP2 is a G_{αi/o}-coupled GPCR with a negative impact on cAMP production [10]. Rodent islets produce prostaglandins (including PGD₂) locally [11] that have a negative impact on beta-cell function, including insulin secretion [11–13]. Consequently, locally produced PGD₂ via GPR44/DP2 may act as a cAMP dependent break on insulin secretion mediated by glucose metabolic stimulation or other GPCRs, such as the GLP-1 receptor. Therefore, antagonizing the GPR44/DP2 pathway in T2DM to restore insulin secretion via a potentially disease driving mechanism may be an opportunity to develop a disease modifying drug in diabetes.

Here we present the first translational study to investigate the potential gluco-metabolic role of the PGD₂-GPR44/DP2 pathway in T2DM, spanning from *in vitro* studies in human islets and beta-cells to a clinical proof-of-mechanism study in T2DM subjects.

Material and methods

Compound and *in vitro* pharmacology

The GPR44/DP2 antagonist, AZD1981, 4-(acetylamino)-3-[(4-chlorophenyl)thio]-2-methyl-1H-indole-1-acetic acid [6], previously a clinical candidate for treatment of asthma [5, 14] was investigated for treatment of T2DM. The biochemical and pharmacological properties of AZD1981 was recently described [14]. Here, the *in vitro* pharmacological properties of AZD1981 was further investigated in the human beta-cell line EndoC-betaH1, which has high expression of GPR44/DP2 (confirmed by qPCR, Fig Ba in [S1 File](#)). Briefly, EndoC-betaH1 cells [15] were plated in fibronectin- and extracellular matrix-coated 96-well plates for the *in vitro* pharmacology analyses and functional assays. Experimental details of the methods are described in the supporting information [S1 File](#).

Insulin and PGD₂ secretion *in vitro* and gene expression analyses

The EC₅₀ for PGD₂ inhibition of glucose stimulated insulin secretion (GSIS) was determined using EndoC-betaH1 cells. The cells were incubated with varying concentrations of the stable PGD₂ analogue 15(R)-15-methyl PGD₂ (Cayman Chemical, USA) at 11.1 mM glucose and insulin secretion was determined. The EC₅₀ value for PGD₂ was calculated from the dose response curve. Since EndoC-betaH1 cells has no expression of L-PGDS (confirmed by qPCR, data not shown) and thus cannot produce PGD₂, the stable PGD₂ analogue was added at EC₈₀ concentration and the effect of the specific GPR44/DP2 antagonist AZD1981 was determined at 11.1 mM glucose or in the presence of a GLP-1 receptor agonist (100 nM exendin-4; Bachem AG, Switzerland).

Human islets (Prodo Laboratories Inc. USA) were used to study PGD₂ secretion, GSIS and for qPCR analysis of the gene expression of enzymes in the PGD₂ synthesis pathway. The acute effect of PGD₂ on GSIS was measured by incubating human islets for 1h in low (2.8mM) or high (11.1mM) glucose, with or without addition of 1 nM 15(R)-15-methyl PGD₂. In other experiments, islets were incubated in high glucose (22.2 mM) with and without addition of 20 ng/ml interleukin 1-beta (IL-1beta) and PGD₂ secretion into the culture media was determined after 24 h (PGD₂ EIA, Cayman Chemical). The expression of genes in the PGD₂ synthesis pathway and GPR44/DP2 were further studied by reanalyzing published human islet single cell sequencing data (MTAB-5061, EBI accession number) [16]. Experimental details are available in the supporting information [S1 File](#).

Clinical study design

This study (<http://clinicaltrials.gov> NCT02367066) was a phase 1, randomised, double-blind, placebo-controlled, multiple-dose, cross-over study conducted at a single centre. The aim was to explore the acute effects of GPR44/DP2 antagonism on both glucose and incretin dependent insulin secretion in T2DM. Hence a cross-over design was chosen with three-day treatment periods upon which steady state of the oral GPR44/DP2 antagonist AZD1981 was to be reached. The study consisted of a run-in, two three-day treatment periods separated by a wash-out period, and follow-up. During each treatment period, the pharmacokinetic (PK) and pharmacodynamic (PD) effects of AZD1981 or placebo were characterized in a mixed meal tolerance test (MMTT) [17] or by a graded glucose infusion (GGI) [18], respectively. A paracetamol test was conducted simultaneously with the MMTT to determine the effects of AZD1981 on gastric emptying.

The primary objectives were defined as change from baseline (day -1 to day 3) for end-points; a) mixed meal tolerance test (MMTT) area under curve (AUC) (0-4h) for plasma glucose, b) MMTT C_{max} and c) graded glucose infusion AUC (1-2h) for plasma C-peptide. The

study was powered to detect therapeutically relevant changes for these primary variables. A number of secondary and exploratory variables all linked to insulin secretion or PK/PD were also assessed.

The study was conducted at ProSciento Inc in Chula Vista, CA, USA and the study protocol was approved by the Institutional Review Board Schulman IRB in Research Triangle Park, NC, USA. All subjects gave their written voluntary informed consent before participation. Patients were enrolled between March 2015 and April 2015. See [Fig 1](#) for study flow chart. More study details are available in the supporting information [S1 File](#).

The enrolled patients were diagnosed with T2DM and had inadequate glycaemic control on a metformin regimen. Patients were adult males or females, had glycosylated haemoglobin (HbA1c) levels between ≥ 58.5 mmol/mol (7.5%) and ≤ 97 mmol/mol (11%) at enrolment, were treated with metformin alone for at least the last three months, and had a fasting glucose level from 3–14 mmol/l (54–252 mg/dl). See [Table 1](#) for subject demographics at screening.

Pharmacokinetic and pharmacodynamics sampling in clinical study

Patients were fasting overnight and all oral medications including metformin were dispensed 75 min prior to the first PD assessment (MMTT). During the MMTT (Ensure Plus drink, 473 ml, 100g carbohydrates), blood sampling was conducted at baseline, 15, 30, 60, 90 and 120 min post the mixed meal intake. During the GGI, by 20% glucose intravenous infusion over 2h with a constant rate GLP-1 infusion the last hour, blood sampling was conducted at 10 min intervals. Blood samples were obtained for assessments of AZD1981 PK on days 2, 3, 8, and 9 concurrent with the MMTT and GGI, and for paracetamol PK on days 3 and 9 during the MMTT. More details are available in the supporting information [S1 File](#).

Analytical techniques in clinical study

Bedside plasma glucose during GGI was measured by an YSI-2300 analyzer (YSI Life Science, USA). For MMTT & GGI plasma insulin, C-peptide, glucagon and total GLP-1 were analyzed by the respective ELISA (Merckodia AB, Sweden) and glucose by a colorimetric method. Plasma paracetamol and AZD1981 concentrations were measured by LC-MS (Covance Bioanalytical services, IN, USA). EDTA plasma and urine was collected on day -1, 3, 6 and 9 for analysis of exploratory PGD₂ biomarkers performed at AstraZeneca. More details are available in the supporting information [S1 File](#).

Sample size calculations & statistical analysis

The clinical study was powered to detect therapeutically relevant changes for the primary variables. The sample size for the MMTT was based on internal studies that estimated the intra-individual standard deviation (SD) for $\ln(\text{MMTT } \text{AUC}_{\text{Glc } 0-4\text{h}})$ to be 0.1. With this estimated SD and an alpha of 0.1, one-sided test and at least 8% difference versus control in $\text{MMTT } \text{AUC}_{\text{Glc } 0-4\text{h}}$, a sample size of 20 evaluable patients was determined to provide at least 80% power. The 8% difference was chosen based on the rationale that it is a clinically relevant magnitude as this is the effect observed by DPP-IV inhibitors [19]. Similarly, the planned sample size for the GGI was determined based on an estimated SD of $\ln(\text{GGI } \text{AUC}_{\text{C-pep } 1-2\text{h}})$ as 0.17. Based on similar power calculations, a sample size of 20 evaluable patients was determined to provide at least 80% power to detect at least a 20% difference versus control for $\text{GGI } \text{AUC}_{\text{C-pep } 1-2\text{h}}$.

Summary statistics are provided for all of the PK and PD variables assessed. The PK evaluable population was used to assess the PK endpoints. PK parameters were derived using actual sampling times and standard non-compartmental methods, and summarized descriptively for both AZD1981 and paracetamol. The three primary PD variables were the $\text{MMTT } \text{AUC}_{\text{Glc } 0-4\text{h}}$,




Fig 1. Study flow chart according to CONSORT 2010 guidelines. The design of a human Proof of Mechanism study to test the Mechanism of Action hypothesis. The phase 1 study was designed as a randomised, double-blind, placebo-controlled, multiple-dose, cross-over study, consisting of a run-in, two treatment periods separated by a wash-out period and a follow-up for each subject. The subjects were T2DM patients on metformin (n = 20).

<https://doi.org/10.1371/journal.pone.0208998.g001>

Table 1. Demographic characteristics. Data are presented as mean ± SD.

	Screening
Gender (female/male)	7/13
Age (y)	54 ± 7.9
Body weight (kg)	87.4 ± 19.1
BMI (kg/m ²)	31.6 ± 4.5
T2DM duration (y)	9.5 ± 4.6
Metformin daily dose (mg)	1830 ± 428
HbA _{1c} (mmol/mol, DCCT %)	79 ± 9.2 (9.4 ± 1.1)
fGlucose (mg/dl)	167.4 ± 39.4
fInsulin (IU/ml)	15.9 ± 7.7
FFA (mEq/l)	0.49 ± 0.2
Total Cholesterol (mg/dl)	182.3 ± 35.2
Triglycerides (mg/dl)	125.7 ± 69
hsCRP (mg/l)	3.2 ± 2.6
eGFR (ml/min/1.73)	104.5 ± 11.7
SBP (mm/Hg)	128.4 ± 10.3
DBP (mm/Hg)	80.9 ± 6
Pulse (beats/min)	69 ± 11
BMR (kcal/d)	1679 ± 332

Abbreviations: BMI = Body Mass Index, BMR = Basal Metabolic Rate (Harris Benedict equation), DBP = Diastolic Blood Pressure, eGFR = estimated Glomerular Filtration Rate, f = fasting, FFA = Free Fatty Acids, HbA_{1c} = Haemoglobin A_{1c}, hsCRP = high sensitive C-Reactive Protein, SBP = Systolic Blood Pressure, T2DM = Type 2 Diabetes Mellitus.

<https://doi.org/10.1371/journal.pone.0208998.t001>

MMTT_{Glc-Cmax} and the GGI AUC_{C-pep 1-2h} (PD evaluable population). The difference in primary variables between AZD1981 and placebo were analyzed using a mixed-effects model that includes treatment (2 levels), treatment sequence (2 levels), and period (2 levels) as fixed effects, baseline values of the dependent variable as a covariate, and subject-within-sequence as a random effect, were used to analyze the difference in MMTT AUC_{Glc 0-4h}, MMTT_{Glc-Cmax} and the GGI AUC_{C-pep 1-2h} between AZD1981 and placebo. We assumed that there was no carryover effect in the crossover study as the 3 day wash-out period should have appropriately removed any impact from carryover between treatment periods. The primary variables were log-transformed prior to analysis and then transformed back to linear scale. Significance was established 1-sided at 10%. Treatment estimates were provided as estimated least square mean ratios with a 80% 2-sided CI. All secondary endpoints of the PD parameters and the pre-hepatic ISR (fasting β-cell responsiveness) was analyzed similarly as those described for the primary PD endpoints. No tests for multiplicity testing were performed for the secondary endpoints. The SAS MIXED procedure was used for the statistical analysis.

Pre-hepatic insulin secretion rate (ISR) over time was calculated using deconvolution of peripheral C-peptide concentrations during MMTT using the method of Hovorka et al [20].

Results

Expression of GPR44/DP2 and PGD₂ synthesis pathway genes

Since the expression of GPR44/DP2 (*PTGDR2*) in the pancreas is restricted to the islets with similar protein expression in both healthy and T2DM [8] we decided to further characterize the expression of GPR44/DP2 and genes related to PGD₂ synthesis in all islet cell types by mining

a single cell sequencing data set [16]. We found that GPR44/DP2 is prominently expressed in beta-cells (Fig 2A), whereas the related receptor DP1 (*PTGDR1*) and the genes in the PGD₂ synthesis pathway mainly are found in pancreatic stellate cells from both healthy and T2DM donors (Fig 2B). GPR44/DP2 mRNA was marginally upregulated in beta-cells from T2DM donors, but several genes in the PGD₂ synthesis pathway were quite substantially upregulated in the stellate cells (Fig 2A and 2B). Interestingly, the total number of immune-activated stellate cells (according to classification by Baron *et al* [21]) was higher in T2DM donors and the expression of the PGD₂-related genes were particularly prominent in this subpopulation of stellate cells (Fig 2C–2E). Other endocrine cells types do not express these genes, except GPR44/DP2 that is expressed also in alpha-, delta-, and PP-cells (Fig A in S1 File). Together, the gene expression pattern suggests that PGD₂ is produced in the islet by activated stellate cells, particularly in T2DM, and may impact beta-cell function through activation of GPR44/DP2.

PGD₂ secretion by human islets

To confirm activation of the GPR44/DP2 pathway during metabolic stress, human islets were treated with high glucose with or without IL-1beta. GPR44/DP2 was highly expressed in intact human islets (Fig 3A, Fig Ba in S1 File), while DP1 was expressed at very low levels (Fig 3B, Fig Bb in S1 File), as expected based on the findings that only a small pool of cells in islets express DP1 (stellate cells, see above). Interestingly, a significant reduction in GPR44/DP2 expression was observed after incubation with high glucose and IL-1beta, while high glucose in itself had no impact (Fig 3A). PGD₂ is produced from arachidonic acid via enzymatic activity of phospholipase A, the cyclooxygenases (*PTGS*) 1 and 2, and lipocalin-prostaglandin D₂ synthase (*PTGDS*). *PTGDS* and *PTGS1* gene expression were not significantly affected by either high glucose or IL-1beta (Fig 3C and 3D), while *PTGS2* was significantly increased by high glucose (35-fold compared to 5.6 mM glucose, $p < 0.01$) and even more prominent by IL-1beta (165-fold compared to control, $p < 0.0001$; Fig 3E).

After incubation of human islet for 24 h, degraded PGD₂ accumulated in the cell culture media at low glucose (5.6 mM) (Fig 3F). A significant increase in PGD₂ production was observed at 22.2 mM glucose with a 2-fold increase ($p < 0.01$), which was further potentiated by IL1-beta with a 10-fold increase in accumulated PGD₂ over 24h ($p < 0.0001$; Fig 3F).

Effect of PGD₂ and the GPR44/DP2 antagonist AZD1981 in human beta cells

Since the endogenous ligand for GPR44/DP2, PGD₂, is unstable, a stable analogue, 15(R)-15-methyl-PGD₂, was used to determine the effects on insulin secretion in human beta-cells (EndoC-betaH1). Activation of GPR44/DP2 by 15(R)-15-methyl-PGD₂ resulted in a dose dependent inhibition of GSIS (Fig 4A). EC₅₀ was calculated using three methods; DMR-dynamic mass distribution [22], cAMP and GSIS (Fig C in S1 File). The potency of AZD1981 on GPR44/DP2 signaling in EndoC cells were established at a constant concentration (150 pM, EC₈₀) of the stable PGD₂ analogue using the same three assays (DMR, cAMP and GSIS). All three assays correlated well and resulted in EC₅₀ values for AZD1981 in the same range (Fig 4B–4D). Additional pharmacological profiling of AZD1981 and GPR44/DP2 in EndoC cells are presented in Figures D and E in S1 File.

Addition of 150 pM 15(R)-15-methyl-PGD₂ to EndoC cells potently inhibited both GSIS as well as exendin-4-potentiated insulin secretion (Fig 3E). PGD₂ inhibition of insulin secretion induced by both glucose and exendin-4, was completely restored by addition of 100 nM AZD1981 (Fig 4E). The inhibitory effect of 15(R)-15-methyl-PGD₂ on GSIS was confirmed in human islets (Fig 4F). The GPR44/DP2 antagonist used in this experiment was however




Fig 2. Tissue expression of GPR44/DP2 and PGD₂ related genes and characterization of pancreatic stellate cells. *PTGDR2* (GPR44/DP2) mRNA expression in human beta-cells (Healthy, n = 197; T2DM, n = 112) (a) and pancreatic stellate cells (Healthy, n = 23; T2DM, n = 31) (b). Heat map of stellate cell signature gene expression sub-classified as quiescent, immune activated (imm) and standard activated (stand) stellate cells according to the description in the main text (c). Quantification of stellate cell subpopulations in healthy and T2D donor human islets (d). Expression of GPR44/DP2 and PGD₂ related genes in subclasses of stellate cells (e). Data are presented as mean ± SEM (a-b), total number of cells (d), and individual cells (e). *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

<https://doi.org/10.1371/journal.pone.0208998.g002>

different from AZD1981 but with similar potency in all three of the in vitro potency assays described for AZD1981 (data not shown).

We demonstrate that GPR44/DP2 is activated in human beta cells by stress induced either by high glucose or high glucose in combination with IL-1beta, resulting in dysregulated glucose and GLP-1 stimulated insulin secretion in vitro, at least partly through production of PGD₂ in stellate cells. Based on that we formulated a hypothesis that stellate cells are activated by glucose and cytokines, producing PGD₂, which in turn acts on GPR44/DP2 on the beta-cells, reducing intracellular cAMP levels leading to inhibition of GSIS (Fig 5). To test the hypothesis that the PGD₂-GPR44/DP2 axis is activated in T2DM and can be a therapeutically relevant target in treatment of T2DM, a mechanistic clinical study was designed and executed using the oral GPR44/DP2 antagonist AZD1981. The study's aim was to explore the acute effects of GPR44/DP2 antagonism on both glucose and incretin dependent insulin secretion in T2DM subjects. Hence, to investigate the insulinotropic action in a physiological context an oral challenge was given through a 4h MMTT [17] and also an intravenous glucose challenge, i.e. a graded glucose (GGI) ranging from a normoglycemic to hyperglycaemic levels [18].

The study outline is described in more detail both the Materials and Method section and illustrated in Fig 1 as well as in the supporting information S1 and S3 Files.

Effects on glucose, insulin and C-peptide after MMTT and GGI in T2DM patients

Treatment with the GPR44/DP2 antagonist AZD1981 at steady state exposure (AZD1981 at 100 mg BID for three days) did not exert any clinically significant insulinotropic effect in T2DM patients on metformin (Fig 6 and Table 2) for any of the co-primary variables in the study. Neither following the insulinotropic challenge in a physiological context during an oral challenge eg a 4h MMTT (MMTT AUC_{Glc 0-4h}, p = 0.12 and MMTT_{Glc-Cmax}, p = 0.06) nor after the intravenous glucose challenge i.e. the graded glucose infusion from a normoglycemic to hyperglycaemic range (GGI AUC_{C-pep 1-2h} p = 0.41) In addition a number of linked secondary variables were also analysed in a similar manner with a mixed effect model to determine any treatment effect of AZD1981 but without any multiplicity test correction. In order to exclude confounding effects by hepatic insulin clearance, C-peptide secretion was evaluated with three parameters, C_{max}, AUC_(0-4h), and deconvoluted prehepatic insulin secretion rate (ISR) according to Hovorka [20]. Notably, the hypothesized insulinotropic effect was not captured by ISR in any subject or time point after neither the MMTT nor the GGI tests. Further none of the secondary variables did show any significant differences from placebo apart from MMTT AUC_{C-peptide 0-4h} and GGI AUC_{Glucose 0-1h} which showed mechanistically contradictory borderline significant effects of p = 0.04 and 0.045 respectively (Fig F in S1 File and Table 2).

Effects on glucagon, GLP1, paracetamol, PK/PD, PGD₂ biomarkers and safety in T2DM patients

To further delineate the mode of action and impact on the other major hormonal islet axes, insulin, glucagon and total GLP-1 were assessed during the MMTT and GGI challenges.

Fig 3. Human islet gene expression and regulation by high glucose with and without IL-1beta. Expression of *PTGDR2* (GPR44/DP2) (a), *PTGDR1* (DP1) (b), *PTGDS* (L-PGDS) (c), *PTGS1* (COX-1) (d) and *PTGS2* (COX-2) (e). *In vitro* human islet PGD₂ secretion (f). Data are presented as mean ± SEM from 5 different human islet donors, where each condition was evaluated in quadruplicates. Glc = glucose. *p<0.05, **p<0.01, ****p<0.0001.

<https://doi.org/10.1371/journal.pone.0208998.g003>

However, as outlined in Table 2, AZD1981 had no significant effect on glucagon secretion or total GLP-1. To fully understand the impact on oral glucose disposal the change in oral paracetamol PK was used to assess the impact on gastric emptying concomitantly with the MMTT. In the paracetamol test, the 90% confidence intervals (CI) of the geometric least squares (LS) mean ratios (AZD1981 to placebo) for C_{max} and AUC_(0-t) were fully contained within the range of 0.80–1.25 (0.90–1.17 for C_{max}, 0.91–1.20 for AUC_(0-t)) and thus the results demonstrated that AZD1981 had no significant effect on gastric emptying in T2DM patients (Fig G in S1 File).

A PK/PD analysis was conducted between AZD1981 exposure (AUC) during either the MMTT or the GGI tests and all of the efficacy endpoints listed in Table 2. However, the PK/PD analysis indicated that no relationship existed between AZD1981 exposure and the MMTT or GGI efficacy variables (Fig H in S1 File).




Fig 4. Effect of PGD₂ on insulin secretion and *in vitro* pharmacology of AZD1981 in the human beta-cell line EndoC- betaH1. Dose response curve for the PGD₂ analogue (15(R)-15-methyl-PGD₂) on insulin secretion, from which the EC₅₀ was determined (a). The *in vitro* potency of the GPR44/DP2 antagonist AZD1981 was determined using DMR (IC₅₀ = 4.9±1nM) (b), cAMP (IC₅₀ = 21±8nM) (c) and by glucose-stimulated insulin secretion (IC₅₀ = 13±2nM) (d). The assays were performed in the human beta cell line EndoC-betaH1 with addition of 150 pM 15(R)-15-methyl-PGD₂. Insulin secretion in EndoC-betaH1 cells was determined with and without addition of 150pM 15(R)-15-methyl-PGD₂ at 11.1 mM glucose and 100nM exendin-4 (GLP-1 receptor agonist). 100nM of AZD1981 was added to restore GSIS. The effect of PGD₂ on human islet insulin secretion was studied in islets from three donors where each incubation condition was run with 5 islets in 6 replicates (f). Data are presented as mean ± SEM. *p<0.05, **p<0.01, ***p<0.001.

<https://doi.org/10.1371/journal.pone.0208998.g004>




Fig 5. Hypothesis for the endogenous PGD₂-GPR44/DP2 axis mechanism of action in islets. Glucose and IL-1beta induced *PTGS2* (COX-2) expression in stellate cells which stimulates the production of PGD₂ synthesis. PGD₂ is secreted and may act in a paracrine fashion on GPR44/DP2 on beta-cells, inhibiting cAMP and thereby reducing GSIS.

<https://doi.org/10.1371/journal.pone.0208998.g005>

We explored if we could identify responder sub-populations with high PGD₂ tone by determining circulating surrogate biomarkers of local PGD₂ and to correlate that with PD response. However, it was not possible to establish a correlation between efficacy and such biomarkers (Fig I in S1 File).

Finally, there were no AZD1981 safety and tolerability findings that could have impacted the PD effects (Tables A and B in S1 File). All of the adverse events (AEs) reported were mild, none resulted in withdrawal from the study, and there were no serious adverse events (SAEs). The incidence of AEs was similar in the AZD1981 (n = 5, 25%) and placebo (n = 6, 30%) treatments. Two (2) of the AEs reported following AZD1981 treatment (diarrhoea) were considered related to the investigational product. There were no unusual or clinically significant vital sign changes observed, and no significant differences between treatments. No clinically significant abnormal findings were detected by ECG or physical examination.

Discussion

To our knowledge, this is the first study elucidating the effect of PGD₂ on human islet function, although it has been known for almost 30 years that PGE₂ inhibits glucose-stimulated insulin secretion [12]. Since then, PGE₂ control of both beta-cell function and mass has been extensively studied and its receptor EP3 is a potential drug target for treatment of diabetes (recently reviewed by Carboneau et al. [23]). Taking that into account, the unknown impact of




Fig 6. Mixed Meal Tolerance Test in T2DM patients. Glucose assessments at baseline (a), and after 3 days treatment (b) or Insulin Secretion Rate (ISR) at baseline (c) and after 3 days treatment (d). Circles connected with dotted lines represent placebo and squares connected with full lines represent AZD1981. Data are presented as mean \pm SD.

<https://doi.org/10.1371/journal.pone.0208998.g006>

PGD₂ on GSIS and the finding that GPR44/DP2 is highly expressed in human beta-cells [7, 8], motivated us to explore the role of PGD₂ in human beta-cell function.

We hypothesized that GPR44/DP2 modulates beta-cell function through an impact on the insulin secretory machinery, based on tissue expression of the GPR44/DP2 pathway genes and robust functional *in vitro* human islet/beta-cell data. Surprisingly, we demonstrate that GPR44/DP2 antagonism do not acutely improve insulin secretion in T2DM patients irrespective of the endogenous incretin response, *i.e.* neither during MMTT nor GGI. Furthermore, the GPR44/DP2 antagonist AZD1981 had no significant effect on the paracetamol profile and thus gastric emptying in T2DM patients, which could have confounded the interpretation of MMTT data. At a dose of 200 mg per day for three days, AZD1981 was safe and well tolerated in T2DM patients treated with metformin. The AZD1981 PK profile was adequate and in line with previous reports, and thus, allowed for a clinically and therapeutically relevant testing of

Table 2. Results of mixed effects model for treatment differences in pharmacodynamic parameters before and after GPR44/DP2 antagonism treatment.

Pharmacodynamic variable	AZD1981 LS mean (SE) # (n = 20)	Placebo LS mean (SE)# (n = 20)	AZD1981/placebo Mean ratio\$ (80% CI) (n = 20)	P-value\$
MMTT AUC _{Glucose 0-4h} *	0.92 (0.02)	0.98 (0.02)	0.94 (0.90; 0.98)	0.12
MMTT C _{max} Glucose *	0.93 (0.02)	0.97 (0.02)	0.96 (0.92; 0.99)	0.06
GGI AUC _{C-peptide 1-2h} *	1.05 (0.04)	1.06 (0.04)	0.99 (0.96; 1.03)	0.41
MMTT AUC _{C-peptide 0-4h}	1.01 (0.03)	1.09 (0.03)	0.93 (0.88; 0.98)	0.04
MMTT C _{max} C-peptide 0-4h	1.06 (0.04)	1.11 (0.04)	0.95 (0.89; 1.02)	0.17
MMTT AUC _{Insulin 0-4h}	0.99 (0.04)	1.08 (0.04)	0.92 (0.85; 0.99)	0.06
MMTT AUC _{Glucagon 0-4h}	1.02 (0.05)	0.97 (0.05)	1.05 (0.97; 1.14)	0.28
MMTT AUC _{total GLP1 0-4h}	1.06 (0.06)	0.99 (0.05)	1.07 (0.97; 1.19)	0.27
GGI AUC _{C-peptide 0-1h}	1.02 (0.04)	1.03 (0.04)	0.99 (0.94; 1.05)	0.45
GGI AUC _{Glucose0-1h}	1.01 (0.02)	0.97 (0.02)	1.04 (1.01; 1.06)	0.045
GGI AUC _{Glucose 1-2h}	1.01 (0.01)	0.99 (0.01)	1.02 (1.00; 1.04)	0.06
GGI AUC _{Insulin 0-1h}	1.04 (0.06)	1.07 (0.06)	0.97 (0.91; 1.04)	0.29
GGI AUC _{Insulin 1-2h}	1.04 (0.06)	1.04 (0.06)	0.99 (0.91; 1.06)	0.37
GGI AUC _{total GLP1 0-1h}	1.19 (0.06)	1.06 (0.05)	1.12 (1.00; 1.26)	0.09
AUC _{Glucose 0-24h}	0.96 (0.02)	0.97 (0.02)	0.99 (0.97; 1.02)	0.34

* Primary study variables

Ratio of Baseline to Endpoint. Data are presented as least squares geometric means with standard error.

\$ The difference in primary variables between AZD1981 and placebo were analysed using mixed-effects models with treatment, treatment sequence, and period as fixed effects, baseline of the primary variable as a covariate, and subject-within-sequence as a random effect. Significance was established at 1-sided at 10%. Treatment estimates were provided as estimated mean ratios with 80% 2-sided CI.

Abbreviations: AUC = Area Under Curve, CI = confidence interval, CGGI = Graded Glucose Infusion, LS = Least square, MMTT = Mixed Meal Tolerance Test, PD = Pharmacodynamic.

<https://doi.org/10.1371/journal.pone.0208998.t002>

the hypothesis. Thereby, we cannot conclude that the PGD₂-GPR44/DP2 axis is a major therapeutically relevant negative modulator of insulin secretion in T2DM patients.

The patient population recruited to the study was adequately representative of a therapeutically relevant target population e.g. patients with T2DM who have failed to achieve glycemic control on first line standard of care (oral metformin with a balanced gender, ethnic, BMI, and HbA1c distribution). However, it is noteworthy that the study population presented with comparatively long-standing disease as indicated by 9.5 years disease duration and HbA1c of 79 mmol/mol. In line with that, there are several typical pathophysiological features that accompany extended T2DM duration such as the observed glycemic profile during MMTT and the comparatively delayed gastric emptying as demonstrated by the delayed C_{max} observed in our study (See Fig F in S1 File and, for comparison, see Fig 2 in [24]). A potential limitation in the clinical study might be the chosen insulinotropic detection level of 8% in MMTT AUC_{Glc 0-4h} for which the study was powered (N = 20 evaluable subjects for 80% power at the 5% level). The 8% change in MMTT AUC_{Glc 0-4h} is the observed GSIS effect by DPPIV inhibitors [19]. It can be debated if this detection level was set too high given that we did not observe any effects. However, given that DPPIV inhibitors have comparatively modest chronic glycemic efficacy in T2DM, averaging 0.3–0.5% HbA1c reduction, when detecting and validating a novel insulin secretory regulatory pathway of broad clinical relevance and interest we found it hard to argue for a lower efficacy threshold.

A protocol deviation with GLP1 entrapment due to a microfilter incompatibility led to a less than planned GLP1 exposure during the GGI PD test and affected the ability to assess potential effects of GPR44/DP2 antagonism on GLP1 responsiveness during the GGI test. However, it was still possible to draw overall conclusions on GLP1 responsiveness due to the complete lack of effect on ISR after both the MMTT and GGI. The rationale for this is that we observed the expected increase in endogenous GLP1 during the MMTT but it did not result in increased insulinotropic action. This conclusion is further supported by the observation that the endogenous total GLP1 concentration after the MMTT was only about one third lower than the expected GLP1 exposure during the GGI/GLP1 PD test, and furthermore, by the knowledge that the GLP1-insulin secretion dose response relationship in T2DM is linear [25].

The underlying hypothesis for the clinical study, *i.e.* the insulinotropic action of GPR44/DP2 antagonism *in vitro*, was based on our data demonstrating that AZD1981 restores PGD₂-inhibited GSIS by potentiation of intracellular cAMP. That is, we observed that a PGD₂-analogue acutely reduces GSIS in the human beta-cell line EndoC-betaH1 through inhibition of cAMP. Moreover, high glucose with or without IL-1beta induced significant PGD₂ production in human islets, as previously seen in rodent islets [11]. Therefore, we hypothesized that the PGD₂ tone in islets is high in diabetics and that GPR44/DP2 antagonism consequently would improve the insulin response to glucose in humans with T2DM. However, irrespective of endogenous GLP1 secretion after both MMTT and GGI, AZD1981 for 3 days did not have any significant acute insulinotropic efficacy, and we concluded that the *in vitro* data did not translate to a therapeutically relevant *in vivo* disease setting. The possible reasons underlying the lack of translatability are speculative, but the primary hypotheses are [1] an absence of a PGD₂ tone in T2DM islets *in vivo* and/or [2] few/absent GPR44/DP2 receptors in T2DM islets. We analyzed metabolites for PGD₂ both in plasma and in urine but did not find any correlation between insulinotropic effects and PGD₂ metabolite levels. Unfortunately, PGD₂ has a very short half-life [26] and it is produced in multiple cell types and tissues [27]. Since islets comprise only a very small portion of the total cell pool that produce PGD₂, it is likely that global PGD₂ metabolite concentrations do not accurately reflect the PGD₂ production in islets. The lack of specific islet PGD₂ metabolites makes it impossible to conclude if a low islet PGD₂ tone was the reason for lack of efficacy. Obviously, it was not possible to measure GPR44/DP2 expression in the islets of the trial subjects. However, gene expression data from isolated islets do not support downregulation of GPR44/DP2 in diabetes (see Fig 2) and previously published data comparing GPR44/DP2 protein expression demonstrate no distinct difference between healthy and T2DM islets [8]. GPCR activity is, however, dependent on protein expression on the cell surface, and it is possible that long term diabetes leads to reduced GPR44/DP2 activity via receptor internalization or other mechanisms, which could have contributed to the poor *in vitro-in vivo* translation. In contrast to traditional insulin secretagogues, which have all demonstrated excellent *in vitro-in vivo* translation, the GPR44/DP2 mechanism depends on antagonism of a presumed negative feed-back loop. If this feed-back loop is not present in diabetics the antagonistic action of AZD1981 would have no acute effect, which seems to be the case.

This study demonstrates the complexity of islet function *in vivo*, where many different GPCRs work in concert to modulate insulin secretion to match the demand and achieve glucose control. The role of prostaglandins for beta-cell function was recently reviewed [23], further illustrating the complexity of prostaglandin signaling through different pathways that modulate beta-cell function, both stimulating and inhibiting insulin secretion. It has long been known that the enzymes needed for prostaglandin synthesis are expressed in rodent islets and that prostaglandins are produced in response to high glucose and IL-1beta [12, 13]. Also in human islets, activation of the prostaglandin synthesis pathway has been observed by increased expression of COX-2 after treatment with cytokines and high glucose [28, 29]. We confirm

this mechanism in intact human islet preparations. However, by mining a human islet single cell sequencing data set [16] we found, to our surprise, that the prostaglandin machinery in human islets is not expressed in endocrine cells but in the stellate cells. This is a novel finding and is not in line with the current consensus for prostaglandin synthesis in islets [28]. To our knowledge the expression pattern has so far only been studied in whole islet preparations and this is the first-time single cell sequencing of human islet has been utilized, allowing analysis of the expression of genes at the cell level. The pancreatic stellate cells are few and quiescent under normal conditions and they surround the acinar, ductal and peri-islet areas [30]. The single cell sequencing data clearly demonstrate that stellate cells are present in human islet preparations [16]. During development of pancreatitis and pancreatic cancer, the stellate cells activate into a myofibroblast phenotype, deteriorating the tissue by rapid growth and secretion of cytokines [31, 32]. Recent studies in animal models have demonstrated activation of stellate cells during development of islet fibrosis and beta-cell dysfunction in T2DM [33, 34], which we confirm, albeit with a small number of cells. The role of stellate cells in the development of human T2DM is currently not known and it could only be speculated that inhibition of PGD₂ signaling through GPR44/DP2 may require longer treatment periods than performed in the present study to alter beta-cell function and have an impact on glycemic control.

Supporting information

S1 File. Supplementary material vs 11.docx: **Fig A. Expression of PGD₂-related genes in all human islet endocrine cell types.** GPR44 (a), DP1 (b), L-PGDS (c), COX1 (d), and COX2 (e). Alpha-cell, n = 886. Beta-cell, n = 309. Delta-cells, n = 114. PP-cells, n = 197. **Fig B. Gene expression of PGD₂ receptors in rodent and human islets and beta cells.** (a) The mRNA expression of GPR44 in rodent islet and beta cell lines were significantly lower compared to human islet and beta cells. (b) mRNA expression of GPR44 and DP1 in rat and human primary islets shows almost undetectable levels of DP1 in human islets. The data were normalized against the expression of the housekeeping gene HPRT for each sample. **Fig C. PGD₂ potently activates signalling in human beta cells.** The natural ligand for GPR44 is PGD₂. Since PGD₂ is unstable in water solution, 15(R)-15-methyl PGD₂ was used in all experiments. The potency for 15(R)-15-methyl PGD₂ at the GPR44 receptor was determined using DMR, cAMP and insulin assays. The agonist showed strong potency and good correlation between the assays. A) The DMR signal (Epic) for the PGD₂ effect on human beta cells (Endo-βH1) ($EC_{50} = 0.8 \times 10^{-10} M$), B) the intracellular cAMP levels ($EC_{50} = 0.6 \times 10^{-10} M$) and C) effects on glucose-stimulated insulin secretion ($EC_{50} = 1.2 \times 10^{-10} M$) were determined. The curves illustrate one representative experiment for each method. Number of data points/assay were for cAMP and insulin assays 6-plicates and 3-plicates for DMR (Epic). Values shown are mean ± SEM. **Fig D. Signaling pathway for GPR44 in human beta cells.** GPCRs couple to different G-proteins and it is known that GPR44 couple to the G_{oi} pathway, inhibiting cAMP levels. To establish the signaling pathway in human beta cells, pertussis toxin (PTX) was added to inhibit signaling through G_{oi}. Using the DMR assay, PGD₂ induced a potent effect on the human EndoC-beta cells, while addition of PTX completely blocked the total cell response. This indicate that the major signaling pathway for GPR44 in human beta cells occur through the G_{oi} pathway. **Fig E. AZD1981 dose-response in human beta cell-line (Endo-βH1).** AZD1981 in different concentrations induced no significant agonist response in human beta cells while 15(R)-15-methyl-PGD₂, which was used as positive control, produced a dose response curve, demonstrating the presence of active GPR44 receptors on the cells. **Fig F. Graded Glucose Infusion in T2DM patients.** C-peptide assessments at baseline (A), after 3 days treatment (B) or Insulin Secretion Rate (ISR) at baseline (C) and after 3 days treatment. Data are presented as

mean \pm SD. **Fig G. Mixed meal test in T2DM patients with plasma paracetamol assessments after 3 days treatment of AZD1981 or placebo.** Data are presented as mean \pm SD. **Fig H. PK/PD analysis of AZD1981 exposure vs selected PD variable after 3 days treatment.** Mixed Meal Tolerance Test and AUC_{Glucose (0-4h)} (A) and for Graded Glucose Infusion and AUC_{C-peptide (0-1h)} (B) in T2DM patients. Data are presented as linear regression. **Fig I. Analysis of exploratory surrogate biomarkers of PGD₂ vs AZD1981 PD response eg MMTT AUC_{Glc (0-4h)} after 3 days treatment.** Plasma 11-beta-PGF₂-alpha (A) Urine 11-beta-PGF₂-alpha /creatinine (B) Urine tetranor-PGD_M / creatinine (C) and plasma L-PGDS (D) in T2DM patients. Data are presented as linear regression. **Table A. Safety evaluation: Adverse events / Serious Adverse events Table. B. Safety evaluation: Clinical chemistry. Table C. Maximum Plasma AZD1981 Concentration at Steady-State, C_{ss},Max (DOCX)**

S2 File. CONSORT 2010 checklist.

(DOC)

S3 File. D6420C00001 clinical study protocol final.

(PDF)

Acknowledgments

The authors would like to thank Anne-Christine Carlsson, Bengt Kull and Gun-Britt Forsberg for excellent work on the exploratory biomarkers.

Author Contributions

Conceptualization: Stanko Skrtic, Björn Tyrberg, Malin Broberg, Hans Ericsson, Marcus Hompesch, Lars Löfgren, Linda Morrow, Tina Rydén-Bergsten, Maria Sörhede Winzell.

Data curation: Stanko Skrtic, Björn Tyrberg, Hans Ericsson, Volker Schneck, Magnus Kjaer, Eva-Marie Andersson, Erik Ryberg, Alexander Aivazidis, Charlotte Wennberg Hultdt, Lars Löfgren, Joanna Parkinson, Elaine Watkins, Maria Sörhede Winzell.

Formal analysis: Stanko Skrtic, Björn Tyrberg, Hans Ericsson, Volker Schneck, Magnus Kjaer, Lars Löfgren, Joanna Parkinson.

Funding acquisition: Stanko Skrtic, Tina Rydén-Bergsten, Maria Sörhede Winzell.

Investigation: Stanko Skrtic, Magnus Kjaer, Marcus Hompesch, Eva-Marie Andersson, Erik Ryberg, Alexander Aivazidis, Charlotte Wennberg Hultdt, Lars Löfgren, Linda Morrow, Joanna Parkinson, Elaine Watkins, Maria Sörhede Winzell.

Methodology: Stanko Skrtic, Björn Tyrberg, Malin Broberg, Hans Ericsson, Magnus Kjaer, Marcus Hompesch, Eva-Marie Andersson, Erik Ryberg, Alexander Aivazidis, Charlotte Wennberg Hultdt, Lars Löfgren, Linda Morrow, Joanna Parkinson, Elaine Watkins, Maria Sörhede Winzell.

Project administration: Stanko Skrtic, Malin Broberg, Linda Morrow, Tina Rydén-Bergsten, Elaine Watkins, Maria Sörhede Winzell.

Resources: Stanko Skrtic, Tina Rydén-Bergsten, Maria Sörhede Winzell.

Software: Volker Schneck, Magnus Kjaer, Joanna Parkinson.

Supervision: Stanko Skrtic, Malin Broberg, Marcus Hompesch, Maria Sörhede Winzell.

Validation: Magnus Kjaer.

Visualization: Stanko Skrtic, Hans Ericsson, Volker Schneck, Alexander Aivazidis, Joanna Parkinson, Maria Sörhede Winzell.

Writing – original draft: Stanko Skrtic, Björn Tyrberg.

Writing – review & editing: Stanko Skrtic, Björn Tyrberg, Malin Broberg, Hans Ericsson, Volker Schneck, Magnus Kjaer, Marcus Hompesch, Eva-Marie Andersson, Erik Ryberg, Alexander Aivazidis, Charlotte Wennberg Hultdt, Lars Löfgren, Linda Morrow, Joanna Parkinson, Tina Rydén-Bergsten, Maria Sörhede Winzell.

References

- White MG, Shaw JA, Taylor R. Type 2 Diabetes: The Pathologic Basis of Reversible beta-Cell Dysfunction. *Diabetes Care*. 2016; 39(11):2080–8. <https://doi.org/10.2337/dc16-0619> PMID: 27926891
- Song I, Muller C, Louw J, Bouwens L. Regulating the beta cell mass as a strategy for type-2 diabetes treatment. *Current drug targets*. 2015; 16(5):516–24. PMID: 25654737
- O'Byrne PM, Naji N, Gauvreau GM. Severe asthma: future treatments. *Clinical and experimental allergy: journal of the British Society for Allergy and Clinical Immunology*. 2012; 42(5):706–11.
- Norman P. Update on the status of DP2 receptor antagonists; from proof of concept through clinical failures to promising new drugs. *Expert Opin Investig Drugs*. 2013; 23(1):55–66. <https://doi.org/10.1517/13543784.2013.839658> PMID: 24073896
- Kuna P, Bjermer L, Tornling G. Two Phase II randomized trials on the CRTh2 antagonist AZD1981 in adults with asthma. *Drug design, development and therapy*. 2016; 10:2759–70. <https://doi.org/10.2147/DDDT.S105142> PMID: 27621597
- Luker T, Bonnert R, Brough S, Cook AR, Dickinson MR, Dougall I, et al. Substituted indole-1-acetic acids as potent and selective CRTh2 antagonists-discovery of AZD1981. *Bioorg Med Chem Lett*. 2011; 21(21):6288–92. <https://doi.org/10.1016/j.bmcl.2011.08.124> PMID: 21944852
- Hellström-Lindahl E, Danielsson A, Ponten F, Czernichow P, Korsgren O, Johansson L, et al. GPR44 is a pancreatic protein restricted to the human beta cell. *Acta Diabetol*. 2016; 53(3):413–21. <https://doi.org/10.1007/s00592-015-0811-3> PMID: 26467464
- Lindskog C, Korsgren O, Ponten F, Eriksson JW, Johansson L, Danielsson A. Novel pancreatic beta cell-specific proteins: antibody-based proteomics for identification of new biomarker candidates. *Journal of proteomics*. 2012; 75(9):2611–20. <https://doi.org/10.1016/j.jprot.2012.03.008> PMID: 22465717
- Amisten S, Salehi A, Rorsman P, Jones PM, Persaud SJ. An atlas and functional analysis of G-protein coupled receptors in human islets of Langerhans. *Pharmacology & therapeutics*. 2013; 139(3):359–91.
- Hata AN, Zent R, Breyer MD, Breyer RM. Expression and molecular pharmacology of the mouse CRTH2 receptor. *The Journal of pharmacology and experimental therapeutics*. 2003; 306(2):463–70. <https://doi.org/10.1124/jpet.103.050955> PMID: 12721327
- Evans MH, Pace CS, Clements RS Jr. Endogenous prostaglandin synthesis and glucose-induced insulin secretion from the adult rat pancreatic islet. *Diabetes*. 1983; 32(6):509–15. PMID: 6354779
- Robertson RP. Eicosanoids as pluripotential modulators of pancreatic islet function. *Diabetes*. 1988; 37(4):367–70. PMID: 2897939
- Kimple ME, Keller MP, Rabaglia MR, Pasker RL, Neuman JC, Truchan NA, et al. Prostaglandin E2 receptor, EP3, is induced in diabetic islets and negatively regulates glucose- and hormone-stimulated insulin secretion. *Diabetes*. 2013; 62(6):1904–12. <https://doi.org/10.2337/db12-0769> PMID: 23349487
- Schmidt JA, Bell FM, Akam E, Marshall C, Dainty IA, Heinemann A, et al. Biochemical and pharmacological characterization of AZD1981, an orally available selective DP2 antagonist in clinical development for asthma. *Br J Pharmacol*. 2013; 168(7):1626–38. <https://doi.org/10.1111/bph.12053> PMID: 23146091
- Ravassard P, Hazhouz Y, Pechberty S, Bricout-Neveu E, Armanet M, Czernichow P, et al. A genetically engineered human pancreatic β cell line exhibiting glucose-inducible insulin secretion. *J Clin Invest*. 2011; 121(9):3589–97. <https://doi.org/10.1172/JCI58447> PMID: 21865645
- Segerstolpe A, Palasantza A, Eliasson P, Andersson EM, Andreasson AC, Sun X, et al. Single-Cell Transcriptome Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes. *Cell Metab*. 2016; 24(4):593–607. <https://doi.org/10.1016/j.cmet.2016.08.020> PMID: 27667667
- Zaccaria M, De Palo E, Zago E, Siculo N, Erle G, Federspil G. Metabolic and endocrine responses to a standard mixed meal. A physiologic study. *Acta Diabetol Lat*. 1979; 16(1):45–53. PMID: 452819

18. Byrne MM, Sturis J, Polonsky KS. Insulin secretion and clearance during low-dose graded glucose infusion. *Am J Physiol*. 1995; 268(1 Pt 1):E21–7. <https://doi.org/10.1152/ajpendo.1995.268.1.E21> PMID: 7840177
19. Solis-Herrera C, Triplitt C, Garduno-Garcia JeJ, Adams J, DeFronzo RA, Cersosimo E. Mechanisms of glucose lowering of dipeptidyl peptidase-4 inhibitor sitagliptin when used alone or with metformin in type 2 diabetes: a double-tracer study. *Diabetes Care*. 2013; 36(9):2756–62. <https://doi.org/10.2337/dc12-2072> PMID: 23579178
20. Hovorka R, Chassin L, Luzio SD, Playle R, Owens DR. Pancreatic beta-cell responsiveness during meal tolerance test: model assessment in normal subjects and subjects with newly diagnosed noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab*. 1998; 83(3):744–50. <https://doi.org/10.1210/jcem.83.3.4646> PMID: 9506719
21. Baron M, Veres A, Wolock SL, Faust AL, Gaujoux R, Vetere A, et al. A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population Structure. *Cell systems*. 2016; 3(4):346–60 e4. <https://doi.org/10.1016/j.cels.2016.08.011> PMID: 27667365
22. Schröder R, Janssen N, Schmidt J, Kebig A, Merten N, Hennen S, et al. Deconvolution of complex G protein-coupled receptor signaling in live cells using dynamic mass redistribution measurements. *Nat Biotechnol*. 2010; 28(9):943–9. <https://doi.org/10.1038/nbt.1671> PMID: 20711173
23. Carboneau BA, Breyer RM, Gannon M. Regulation of pancreatic β -cell function and mass dynamics by prostaglandin signaling. *J Cell Commun Signal*. 2017.
24. Monnier L, Colette C, Dunseath GJ, Owens DR. The loss of postprandial glycemic control precedes stepwise deterioration of fasting with worsening diabetes. *Diabetes Care*. 2007; 30(2):263–9. <https://doi.org/10.2337/dc06-1612> PMID: 17259492
25. Kjems LL, Holst JJ, Vølund A, Madsbad S. The influence of GLP-1 on glucose-stimulated insulin secretion: effects on beta-cell sensitivity in type 2 and nondiabetic subjects. *Diabetes*. 2003; 52(2):380–6. PMID: 12540611
26. Schuligoi R, Schmidt R, Geisslinger G, Kollrosier M, Peskar BA, Heinemann A. PGD2 metabolism in plasma: kinetics and relationship with bioactivity on DP1 and CRTH2 receptors. *Biochem Pharmacol*. 2007; 74(1):107–17. <https://doi.org/10.1016/j.bcp.2007.03.023> PMID: 17452035
27. Murata T, Maehara T. Discovery of anti-inflammatory role of prostaglandin D2. *The Journal of veterinary medical science*. 2016; 78(11):1643–7. <https://doi.org/10.1292/jvms.16-0347> PMID: 27498997
28. Parazzoli S, Harmon JS, Vallerie SN, Zhang T, Zhou H, Robertson RP. Cyclooxygenase-2, not microsomal prostaglandin E synthase-1, is the mechanism for interleukin-1 β -induced prostaglandin E2 production and inhibition of insulin secretion in pancreatic islets. *J Biol Chem*. 2012; 287(38):32246–53. <https://doi.org/10.1074/jbc.M112.364612> PMID: 22822059
29. Shanmugam N, Todorov IT, Nair I, Omori K, Reddy MA, Natarajan R. Increased expression of cyclooxygenase-2 in human pancreatic islets treated with high glucose or ligands of the advanced glycation end-product-specific receptor (AGER), and in islets from diabetic mice. *Diabetologia*. 2006; 49(1):100–7. <https://doi.org/10.1007/s00125-005-0065-7> PMID: 16341840
30. Omary MB, Lugea A, Lowe AW, Pandol SJ. The pancreatic stellate cell: a star on the rise in pancreatic diseases. *J Clin Invest*. 2007; 117(1):50–9. <https://doi.org/10.1172/JCI30082> PMID: 17200706
31. Apte MV, Wilson JS. Dangerous liaisons: pancreatic stellate cells and pancreatic cancer cells. *J Gastroenterol Hepatol*. 2012; 27 Suppl 2:69–74.
32. Pandol S, Gukovskaya A, Edderkaoui M, Edderkoui M, Dawson D, Eibl G, et al. Epidemiology, risk factors, and the promotion of pancreatic cancer: role of the stellate cell. *J Gastroenterol Hepatol*. 2012; 27 Suppl 2:127–34.
33. Ryall CL, Vilorio K, Lhaf F, Walker AJ, King A, Jones P, et al. Novel role for matricellular proteins in the regulation of islet β cell survival: the effect of SPARC on survival, proliferation, and signaling. *J Biol Chem*. 2014; 289(44):30614–24. <https://doi.org/10.1074/jbc.M114.573980> PMID: 25204658
34. Li FF, Chen BJ, Li W, Li L, Zha M, Zhou S, et al. Islet Stellate Cells Isolated from Fibrotic Islet of Goto-Kakizaki Rats Affect Biological Behavior of Beta-Cell. *J Diabetes Res*. 2016; 2016:6924593. <https://doi.org/10.1155/2016/6924593> PMID: 26697502