



Review

Bouveret's Syndrome: A Case-Based Review, Clinical Presentation, Diagnostics and Treatment Approaches

Murat Ferhat Ferhatoğlu, Abdulcabbar Kartal

Department of General Surgery, Okan University, Faculty of Medicine, Istanbul, Turkey

Abstract

Gastric outlet obstruction that arises from gallstones impacted in the distal stomach or proximal duodenum after passing through a cholecystoduodenal, cholecystogastric or rarely choledocoduodenal fistula is called as Bouveret's syndrome and it accounts for approximately 1-3% of all the patients with gallstone ileus. Although treatment modalities, including stone removal or fragmentation with classical endoscopic devices, such as snares, and forceps or fragmentation of gallstones with new devices, such as electrohydraulic lithotripsy, laser, extracorporeal shockwave lithotripsy have been described. However, only 29% of the patients benefit from nonsurgical methods. Removal of the stone through a gastrotomy or enterotomy and performing cholecystectomy and fistula repair with a second operation is an approach recommended for older patients with comorbid diseases. In this paper, a case of Bouveret's syndrome was presented. The authors also aimed to review the diagnosis, management and treatment of this rare disease and to update the previous reviews.

Keywords: Bouveret's syndrome; gallstone; ileus.

Please cite this article as "Ferhatoğlu MF, Kartal A. Bouveret's Syndrome: A Case-Based Review, Clinical Presentation, Diagnostics and Treatment Approaches. Med Bull Sisli Etfal Hosp 2020;54(1):1-7".

Complications related to gallstones are not rare in daily medical practice, which are acute cholecystitis, pancreatitis, choledocolithiasis and gallstone ileus. Gallstone ileus is obstruction of the gastrointestinal system that arises from gallstones and occurs only approximately 0.3-0.5% of the patients having gallstones.^[1] Obstruction usually occurs in the terminal ileum and ileocecal valve (70%), rarely occurs in proximal ileum or jejunum (20-40%).^[2] Gastric outlet obstruction that arises from gallstones impacted in the distal stomach or proximal duodenum after passing through a cholecystoduodenal, cholecystogastric or rarely choledocoduodenal fistula is called as Bouveret's syndrome (BS). The first two cases were described for the first time in 1896 by a Parisian surgeon, Leon Bouveret.^[3] This rare complication is observed only 1-3% of patients having gallstone

ileus and more frequent in patients with advanced age and comorbid diseases.^[4] It is mostly presented as single case reports; the largest case series in the literature only consist of six cases.^[5] Because of the rarity of this disease, no standardized diagnostic or therapeutic technique is described, including endoscopic techniques or surgical approaches (open or laparoscopic). In this paper, a case of BS was presented. The authors also aimed to review the diagnosis, management and treatment of this rare disease and to update the previous reviews. The patient's written consent was obtained in this paper.

Methods and Case Presentation

Keywords "Bouveret", "Bouveret syndrome", "Bouveret's" and "Bouveret's syndrome" were searched in PubMed and

Address for correspondence: Murat Ferhat Ferhatoglu, MD. Okan Universitesi Tip Fakultesi, Genel Cerrahi Anabilim Dalı, Istanbul, Turkey

Phone: +90 555 321 47 93 **E-mail:** ferhatferhatoglu@yahoo.co.uk

Submitted Date: April 05, 2018 **Accepted Date:** July 18, 2018 **Available Online Date:** March 25, 2020

©Copyright 2020 by The Medical Bulletin of Sisli Etfal Hospital - Available online at www.sislietfaltip.org

OPEN ACCESS This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).



Google Scholar database. All related articles published in English between the years 1968-2018 were evaluated. The 152 cases were found in 146 articles. Gender, age, symptoms, signs, diagnostic tools, treatment options were analysed. Also, a new case in our clinic was presented below.

A 69-year-old female patient was evaluated due to complaints of loss of appetite, nausea began 10 days ago and complaints of vomiting and abdominal pain after eating began two days before the patient was admitted to our hospital. On examination, the blood pressure of the patient who was on salbutamol 200 mcg, valsartan 320 mg, hydrochlorothiazide 25 mg treatment due to hypertension and chronic obstructive pulmonary disease was 155/90 mmHg. Her heart rate was 88/bpm, and her body temperature was 37.8°C. She also had a history of type 2 diabetes mellitus under insulin treatment. On physical examination of the abdomen, she had tenderness in her right upper quadrant, and her intestinal sounds were hypoactive. At laboratory examinations, leukocyte was 12300/mm³(4600-10200/mm³), c-reactive protein was 27 mg/dL (0-5 mg/dL) and potassium was 2.8 mmol/dL (3.5-5.1 mmol/dL) aspartate aminotransferase was 154U/L (5-34 U/L), alanine aminotransferase was 205 U/L (0-55 U/L), and she had a slightly elevated alkaline phosphatase level of 177 U/L (40-150 U/L). In her blood gases analysis, she had a pH of 7.31. Abdominal x-ray showed no abnormal findings. Abdominal ultrasound examination revealed an image in size of 4 cm with acoustic shadowing in the duodenum. Computed tomography scan with intravenous and oral contrast of the abdomen demonstrated that the stomach was dilated, and there was a solid mass approximately in size of 4 cm in the duodenal bulb (Fig. 1). In consequence of gastroscopy performed, a gallstone impacted the duodenal bulb in size of 4 cm was observed (Fig. 2). Gallstone was tried to be split by using endoscopic forceps and snares, but since this procedure failed, it was decided to perform surgery. In exploration performed with a supraumbilical vertical midline incision, it was observed that the gallbladder, duodenum and surrounding fatty tissues were oedematous; the areas of inflammation were wrapped by the omentum. Because the wall of the duodenum is very oedematous and fragile, it was decided to reach the stones via gastrotomy. After performing gastrotomy from the pyloric area, the impacted gallstone has been extracted in two pieces through gastrotomy using sponge forceps, and subsequent Heineke-Mikulicz pyloroplasty was performed. A 7-mm silicone drain was inserted in the operation area. And after closing the abdominal midline incision, the operation was finished uneventfully. Since our patient's advanced age, poor general condition and her comorbid diseases, cholecystectomy was not performed in the same operative



Figure 1. Gallstone impacted in the duodenum (blue arrow) and dilated stomach stone (white arrow) on tomography.



Figure 2. Gallstone impacted in the duodenal bulb (red arrow).

session. After surgery, the patient's c-reactive protein level became normal on the 6th day of surgery. On the fourth day of the surgery, the liquid diet was given to the patient and she was discharged on the postoperative 7th day uneventfully. Since she did not come for follow-ups after discharge, cholecystectomy in the second session could not be performed.

Discussion

Fistulas between the biliary system and the upper gastrointestinal system are rare pathologies, but BS is an even rarer

phenomenon that has the potential to cause lethal complications. BS tends to occur more commonly in women (65%) with a median age of 74.1, and patients mostly have comorbid diseases.^[4] A higher incidence of BS in females can be explained by the higher rates of gallstones in females than males, which is related to cholestatic effects of oestrogen and progesterone.^[6] Ileus, due to gallstones, has female predominance related to the same factors. Advanced age and gender of our patient are typical for BS.

The size of gallstones is an important risk factor for BS. Large gallstones (>2.5 cm) have the potential to migrate into the duodenal bulb or distal part of the stomach, where they frequently lead to obstruction by eroding gallbladder or choledochal wall.^[7] The gallstone of the presented case was 4 cm in size, which is typical for gallstone ileus.

BS has nonspecific symptoms, physical and radiological findings. Knowledge of risk factors, signs, symptoms, radiological and endoscopic findings are important for rapid and accurate diagnosis. However, the most important key to diagnosing is having a high level of suspicion in cases with a history of gallstone and sings of pyloric obstruction. The disease mimics pyloric obstruction very well. It may cause nausea, vomiting, abdominal pain or discomfort, loss of appetite, weight loss and anorexia. Symptoms of dehydration also may be observed due to vomiting. Hematemesis or melena may develop respectively, 13.8% and 4.6% of BS cases due to mucosal erosion that arises from an impacted gallstone.^[1] In recently reported 152 cases, vomiting, nausea and abdominal pain were the most common symptoms (Table 1). Our patient presented all three symptoms.

Abdominal tenderness, distension due to obstruction of the distal stomach or duodenal bulb and absence or hypoactivation of intestinal sounds is the most observed signs (Table 2). Our patient presented all three major signs.

Table 1. Symptoms of the patients with the Bouveret's syndrome (A review of 152 cases in 138 reports)

| Symptoms | Number of the findings | % of the findings |
|-----------------|------------------------|-------------------|
| Nausea/Vomiting | 124 | 81.5 |
| Abdominal pain | 103 | 67.7 |
| Hematemesis | 21 | 13.8 |
| Weight loss | 18 | 11.8 |
| Anorexia | 17 | 11.1 |
| Constipation | 11 | 7.2 |
| Melena | 7 | 4.6 |
| Pale stool | 2 | 1.3 |
| Diarrhea | 3 | 1.9 |
| Dysphagia | 2 | |

Since the clinical presentations were not reported, three of 152 cases were excluded (Reference number 44, 45, 46 were excluded)

Table 2. Signs of the patients with the Bouveret's syndrome (A review of 152 cases in 138 reports)

| Signs | Number of the findings | % of the findings |
|--------------------------------------|------------------------|-------------------|
| Abdominal tenderness | 59 | 38.8 |
| Abdominal distention | 39 | 25.6 |
| Hypoactive or absent of bowel sounds | 7 | 4.6 |
| Jaundie | 3 | 1.97 |
| Succussion splash | 10 | 0.65 |

Since the clinical presentations were not reported, three of 152 cases were excluded (Reference number 44, 45, 46 were excluded)

Metabolic alkalosis, hypokalaemia secondary to severe vomiting and elevation in leukocyte and CRP values due to the inflammatory process in the gallbladder, fistula and surrounding tissues can be seen in patients. Our patient had an increase in CRP value over 50-fold and leucocytosis. Additionally, hypokalaemia was determined.

The following findings can be observed on an abdominal x-ray or upper gastrointestinal system fluoroscopy: intestinal obstruction, gallstone in the intestinal lumen and the presence of gas in the biliary system (pneumobilia). These findings are known as Rigler's triad. In addition to these, Balthazar and Schechter also added two findings, air-fluid levels in the right upper quadrant and air in the gallbladder and duodenal bulb on abdominal x-ray.^[8] However, this triad of x-ray findings is observed only in 30-35% of the cases (Table 3) (7,9). Since obstruction is in the proximal part in BS, classical gallstone ileus findings cannot be observed in the abdominal x-ray. In the presented case, no significant pathology was determined in the abdominal x-ray.

Ultrasonography and computed tomography are helpful for diagnosis; these non-invasive tests are preferred as confirmation of endoscopic diagnosis. Ultrasonography is useful in identifying gallstones in the gallbladder. It has limited ability to identify stones in duodenum or stomach

Table 3. Abdominal X-ray findings in 81 cases of the Bouveret's syndrome

| Findings | Number of the findings | % of the findings |
|-----------------------------|------------------------|-------------------|
| Pneumobilia | 30 | 37 |
| Calcified mass or gallstone | 29 | 35.8 |
| Dilated stomach | 17 | 20.9 |
| Dilated intestinal loops | 10 | 12.3 |
| Paucity of bowel gas | 2 | 2.5 |
| No significant findings | 11 | 13.5 |

The 81 cases of BS from references: 4-6, 17, 20, 28, 29-32

compared to computed tomography^[9] (Table 4). In our presented case, ultrasonography identified gallstone in the duodenal bulb. Computed tomography is helpful in identifying fistula between gallbladder and stomach or duodenum, also helpful to show the accurate size of the impacted stone. However, diagnosis can be made by using computed tomography in 60% of patients.^[8] Computed tomography also cannot show choledocus lumen clearly in 25% of the cases. However, magnetic resonance cholangiopancreatography clearly shows the whole biliary system and differentiates fluid from calculi.^[10, 11] Magnetic resonance cholangiopancreatography also identifies fistula between the biliary system and the gastrointestinal system.^[10] It can be used for evaluation if the pericholecystic fluid is found or not and evaluation of gallbladder, stomach and the intestinal wall.^[12, 13] Oral contrast usage in computed tomography or magnetic resonance imaging demonstrates contrast extravasation into gallbladder via the fistula. Analysis of 67 computed topographies of BS patients revealed that the fistula and pneumobilia diagnosis rate is 46.2% in com-

Table 4. Ultrasonography findings in 48 cases of the Bouveret's syndrome

| Findings | Number of the findings | % of the findings |
|-----------------------------------------------------|------------------------|-------------------|
| Gallstone in gallbladder | 24 | 50 |
| Gallstone in duodenum | 10 | 20.8 |
| Thickened gallbladder wall | 4 | 8.3 |
| Unspecified echogenic focus in right upper quadrant | 2 | 4.1 |
| Cholecystoduodenal fistula and pneumobilia | 18 | 37.5 |
| No significant findings | 6 | 12.5 |

The 48 cases of BS from references: 4, 5, 6, 25, 30, 47

Table 5. Computed tomography findings in 67 cases of the Bouveret's syndrome

| Findings | Number of the findings | % of the findings |
|-------------------------------------------|------------------------|-------------------|
| Cholecystoduodenal fistula or pneumobilia | 31 | 46.2 |
| Gallstone in duodenum | 24 | 35.8 |
| Gallstone in gallbladder | 13 | 19.4 |
| Gallstone in stomach | 3 | 4.4 |
| Mass in duodenum | 15 | 22.3 |
| Mass in pylorus | 1 | 1.5 |
| Mass in gallbladder | 1 | 1.5 |
| Dilatation in stomach | 12 | 17.9 |
| Dilatation in duodenum | 7 | 10.4 |

The 67 cases of BS from references: 1, 3-6, 22, 23, 28, 29-39, 42, 43, 47, 48

Table 6. Endoscopic findings in 90 cases of Bouveret's syndrome

| Findings | Number of the findings | % of the findings |
|-------------------------|------------------------|-------------------|
| Dilated stomach | 4 | 4.4 |
| Obstruction | | |
| Pylorus | 12 | 13.3 |
| Duodenal bulb | 59 | 65.5 |
| Postbulber duodenum | 19 | 21.1 |
| Fistula | | |
| With stone | 4 | 4.4 |
| Without stone | 7 | 7.7 |
| No significant findings | 2 | 2.2 |

The 90 cases of BS from references: 1, 4, 6, 20, 22-24, 34, 36, 37, 39-43, 47*

*ERCP was performed

puted tomography (Table 5). Despite all developments in diagnostic techniques, the diagnosis can be made only in 50% of the cases preoperatively.^[14]

The upper gastrointestinal system endoscopy shows the impacted stone in the distal stomach or duodenal bulb with a dilated stomach (Table 6). The duodenal orifice of the cholecystoduodenal or choledocoduodenal fistula may also be identified in upper gastrointestinal system endoscopy.^[15-17] Although treatment modalities, including stone removal with endoscopic techniques (nets/baskets) or fragmentation of gallstones with mechanical, electrohydraulic, laser or extracorporeal shockwave lithotripsy, have been described, only 29% of the patients benefit from endoscopic methods.^[18] Mechanical lithotripsy can be used to split stones into small pieces before extracting impacted stone by nets or baskets. Baskets, snares, forceps may be used as mechanical lithotripter.^[19] If suspicion of malignancy occurs, biopsies should be taken at the time of the endoscopic approach. Extracting all stone pieces is necessary to avoid postoperative ileus due to gallstones.^[20]

Electrohydraulic lithotripsy is another option as an endoscopic treatment modality. The first usage of electrohydraulic lithotripsy in endoscopy is described by Sethi et al. They used a double-channel gastroscope during the operation.^[21] This method may have the potential of complications as perforation and bleeding due to damage in surrounding tissues.

Neodymium, rhodamine, holmium: YAG lasers are also promising techniques for gallstone fragmentation. The advantage of laser using is splitting the stone with minimal tissue damage.^[22, 23] Holmium: YAG laser fragmentation has benefits over neodymium and rhodamine laser given the capability of applying high energy via flexible and small probes.^[24]

Holl et al. first reported the success of extracorporeal shockwave lithotripsy in gallstone impacted in duodenum.^[25] After Holl et al., many cases were reported about extracorporeal shockwave lithotripsy treatment. Most of these cases have shown to be ineffective or addition of other endoscopic techniques required. In a case reported by Lenz et al., five times of extracorporeal shockwave lithotripsy was used to split the stone, which impacted in the duodenal bulb.^[26]

Limitation of endoscopic therapy is the impossibility of closure of fistula between stomach/duodenum and gallbladder or choledocus. Another limitation is defragmented stones may cause gallstone ileus by obstructing distal parts of intestines. Beside all these limitations, endoscopic treatment methods need a high degree of experience. Endoscopic treatment methods may be insufficient for the stones that are larger than 2.5 cm.^[4] An attempt of endoscopic treatment for the stone in size of 4 cm determined in our case failed.

As the surgical treatment modality, removal of the stone through a gastrotomy or enterotomy and in addition to this operation performing cholecystectomy and fistula repair was described. Although the surgical approach is the most efficient method, due to comorbid factors in most of the patients, postoperative mortality rates may reach a level of 12%; in fact, when cholecystectomy is added,

these rates may increase much more.^[27] Removal of the stone through a gastrotomy or enterotomy and performing cholecystectomy and fistula repair in a second operation is an approach recommended for elderly patients with comorbid diseases (Table 7). The likelihood of occurrence of gallstone ileus before the second operation is 5%.^[28] Cholangitis and cholecystitis are other complications that may occur before the second operation. Since our patients advanced age, poor general condition and her comorbid diseases, cholecystectomy was not performed in the same operative session.

Conclusion

Gallstone ileus is an infrequent complication of gallstone disease, but BS is a scarce complication of gallstone disease. BS presents differently from classical gallstone ileus with symptoms and signs of gastric outlet obstruction. Recognizing risks, symptoms, signs and radiological findings that are pointed out in this review is important for early diagnosis. Gallstone disease, ileus and BS have a higher incidence in the female population due to cholestatic effects of oestrogens. Gender and advanced age of this reported patient is typical for this rare disease. Small gallstones pass through the cystic duct, whereas large gallstones cannot pass and erode gallbladder wall and lead to a fistula between gallbladder and duodenum. In recent years, diagnostic tools and treatment strategies evolve to less invasive methods.

Table 7. Treatment modalities and clinical outcomes of 129 patients of Bouveret's syndrome

| Treatment modality | Cases | Success number | Mortality | Major complication |
|------------------------------------------------------------|-------|----------------|-----------|--------------------------------------------------------------------------------|
| Endoscopic extraction | 39 | 4 | 0 | 0 |
| Laser lithotripsy | 7 | 5 | 0 | 0 |
| ESWL | 5 | 1 | 0 | 0 |
| Mechanical or electrohydraulic lithotripsy | 16 | 3 | 0 | 1 e.coli sepsis |
| Enterotomy/gastrotomy+stone extraction | 49 | 45 | 1 | 1 respiratory failure, 1 wound dehiscence, 1 duodenal leak, 1 fungal sepsis |
| Enterotomy+stone extraction+cholecystectomy+fistula repair | 36 | 30 | 1 | 1 sepsis, 1 renal failure, 1 upper GI bleed, 1 pulmonary embolism, 1 bile leak |
| Enterotomy+stone extraction+cholecystectomy | 3 | 2 | 0 | 1 bile leak |
| Enterotomy+stone extraction+fistula repair | 3 | 3 | 0 | 0 |
| Enterotomy+stone extraction+cholecystectomy+pyloroplasty | 3 | 3 | 0 | 0 |
| Antrectomy+gastrojejunostomy+cholecystectomy | 1 | 1 | 1 | 0 |
| Laparoscopic enterolithotomy | 2 | 2 | 0 | 0 |
| No therapy | 1 | 0 | 1 | 0 |

Since the outcome of the case was not reported, the 23 of 152 BS cases were excluded from this Table (Excluded references: 5, 37, 40, 42, 48)

The use of laser technology with endoscopic tools promises a new era in the treatment of this rare disease. Avoiding surgical intervention may decrease mortality in cases with comorbid diseases; endoscopic interventions may be tried in these cases. However, in cases having the necessity of surgery, two-stage surgical interventions should be preferred in patients with comorbid risk factors.

Disclosures

Ethics Committee Approval: The study was approved by the Ethics Committee of Istanbul Okan University Hospital (09.05.2018/94).

Peer-review: Externally peer-reviewed.

Conflict of Interest: None declared.

Authorship Contributions: Concept – M.F.F.; Design – M.F.F.; Supervision – A.K.; Materials – M.F.F.; Data collection &/or processing – M.F.F., A.K.; Analysis and/or interpretation – M.F.F.; Literature search – M.F.F., A.K.; Writing – M.F.F., A.K.; Critical review – M.F.F., A.K.

References

- Caldwell KM, Lee SJ, Leggett PL, Bajwa KS, Mehta SS, Shah SK. Bouveret syndrome: current management strategies. *Clin Exp Gastroenterol* 2018;11:69–75.
- Khaira HS, Thomas DR. Gallstone emesis and ileus caused by common hepatic duct-duodenal fistula. *Br J Surg* 1994;81:723.
- Puri V, Lee RW, Amirlak BA, Lanspa SJ, Fitzgibbons RJ Jr. Bouveret syndrome and gallstone ileus. *Surg Laparosc Endosc Percutan Tech* 2007;17:328–30.
- Cappell MS, Davis M. Characterization of Bouveret's syndrome: a comprehensive review of 128 cases. *Am J Gastroenterol* 2006;101:2139–46.
- Printen KJ, Safaie-Shirazi S. Duodenal obstruction caused by gallstones: report of six cases. *Am Surg* 1973;39:688–91.
- Bateson MC. Gallbladder disease and cholecystectomy rate are independently variable. *Lancet* 1984;2:621–4.
- Altinkaya N, Koç Z, Alkan Ö, Demir S, Belli S. Multidetector computed tomography diagnosis of ileal and antropyloric gallstone ileus. *Ulus Travma Acil Cerrahi Derg* 2011;17:461–3.
- Reimann AJ, Yeh BM, Breiman RS, Joe BN, Qayyum A, Coakley FV. Atypical cases of gallstone ileus evaluated with multidetector computed tomography. *J Comput Assist Tomogr* 2004;28:523–7.
- Delabrousse E, Bartholomot B, Sohm O, Wallerand H, Kastler B. Gallstone ileus: CT findings. *Eur Radiol* 2000;10:938–40.
- Pickhardt PJ, Friedland JA, Hruza DS, Fisher AJ. Case report. CT, MR cholangiopancreatography, and endoscopy findings in Bouveret's syndrome. *AJR Am J Roentgenol* 2003;180:1033–5.
- Algin O, Ozmen E, Metin MR, Ersoy PE, Karaoğlanoğlu M. Bouveret syndrome: evaluation with multidetector computed tomography and contrast-enhanced magnetic resonance cholangiopancreatography. *Ulus Travma Acil Cerrahi Derg* 2013;19:375–9.
- Algin O, Ozlem N, Kilic E, Karaoglanoglu M, Arslan H. Gd-BOP-TA-enhanced MR cholangiography findings in gall bladder perforation. *Emerg Radiol* 2010;17:487–91.
- Algin O, Ozmen E, Ersoy PE, Karaoglanoglu M. Periapillary localized pancreatic intraepithelial neoplasia-3 (PanIN-3): evaluation with contrast-enhanced MR cholangiography (MRCP). *Radiol Oncol* 2011;45:300–3.
- Istvan DM, Arpad T, Etele E, Kucserik L, Andras Z, Borz B, et al. Surgical strategy in Bouveret's syndrome: A case report. *Acta Medica Marisiensis* 2016;62:272–5.
- Beuran M, Venter MD, Ivanov I, Smarandache R, Iftimie-Nastase I, Venter DP. Gallstone ileus—still a problem with heart. *Annals of Academy of Romanian Scientists: Medical sciences* 2012;3:5–28.
- Koulaouzidis A, Moschos J. Bouveret's syndrome. Narrative review. *Ann Hepatol* 2007;6:89–91.
- Trubek S, Bhama JK, Lamki N. Radiological findings in Bouveret's syndrome. *Emergency Radiology* 2001;8:335–7.
- Bramson J, Topilow A, Matteotti R. Bouveret's syndrome: The rarest obstructing gallstone. *New Horizons in Clinica Case Reports* 2017;2:17–9.
- Afzal M, Ghosh D, Leigh T. Mechanical lithotripsy for Bouveret's syndrome. *Gut* 2007;56:733–4.
- Alsolaiman MM, Reitz C, Nawras AT, Rodgers JB, Maliakkal BJ. Bouveret's syndrome complicated by distal gallstone ileus after laser lithotripsy using Holmium: YAG laser. *BMC Gastroenterol* 2002;2:15.
- Sethi S, Kochar R, Kothari S, Thosani N, Banerjee S. Good Vibrations: Successful Endoscopic Electrohydraulic Lithotripsy for Bouveret's Syndrome. *Dig Dis Sci* 2015;60:2264–6.
- Saldaña Dueñas C, Fernández-Urien I, Rullán Iriarte M, Vila Costa JJ. Laser lithotripsy resolution for Bouveret syndrome. *Endoscopy* 2017;49:E101–2.
- Maiss J, Hochberger J, Hahn EG, Lederer R, Schneider HT, Muehldorfer S. Successful laserlithotripsy in Bouveret's syndrome using a new frequency doubled doublepulse Nd:YAG laser (FRED-DY). *Scand J Gastroenterol* 2004;39:791–4.
- Goldstein EB, Savel RH, Pachter HL, Cohen J, Shamamian P. Successful treatment of Bouveret syndrome using holmium: YAG laser lithotripsy. *Am Surg* 2005;71:882–5.
- Holl J, Sackmann M, Hoffmann R, Schüssler P, Sauerbruch T, Jüngst D, et al. Shock-wave therapy of gastric outlet syndrome caused by a gallstone. *Gastroenterology* 1989;97(2):472–4.
- Lenz P, Domschke W, Domagk D. Bouveret's syndrome: unusual case with unusual therapeutic approach. *Clin Gastroenterol Hepatol* 2009;7:e72.
- Gan S, Roy-Choudhury S, Agrawal S, Kumar H, Pallan A, Super P, et al. More than meets the eye: subtle but important CT findings in Bouveret's syndrome. *AJR Am J Roentgenol* 2008;191:182–5.
- Newton RC, Loizides S, Penney N, Singh KK. Laparoscopic management of Bouveret syndrome. *BMJ Case Rep* 2015;2015:bcr2015209869.
- O'Neill C, Colquhoun P, Schlachta CM, Etemad-Rezai R, Jayaraman

- S. Gastric outlet obstruction secondary to biliary calculi: 2 cases of Bouveret syndrome. *Can J Surg* 2009;52:16–8.
30. Thompson RJ, Gidwani A, Caddy G, McKenna E, McCallion K. Endoscopically assisted minimally invasive surgery for gallstones. *Ir J Med Sci* 2009;178:85–7.
31. Rossi D, Khan U, McNatt S, Vaughan R. Bouveret syndrome: a case report. *WV Med J* 2010;106:18–22.
32. Mumoli N, Cei M, Luschi R, Carmignani G, Orlandi F. Bouveret syndrome. *Emerg Med J* 2010;27:525.
33. Palomeque-Jiménez A, Calzado-Baeza S, Reyes-Moreno M. Bouveret syndrome: an infrequent presentation of gallstone ileus. *Rev Esp Enferm Dig* 2012;104:324–5.
34. Gallego Otaegui L, Sainz Lete A, Gutiérrez Ríos RD, Alkorta Zuloaga M, Arteaga Martín X, Jiménez Agüero R, et al. A rare presentation of gallstones: Bouveret's syndrome, a case report. *Rev Esp Enferm Dig* 2016;108:434–6.
35. Vieiro Medina MV, Gómez Sanz R, Bra Insa E, Domínguez Sánchez I, de la Fuente Bartolomé M, Díaz Pérez D, et al. First case reported of Bouveret's syndrome associated to duodenal and biliary perforation to retroperitoneum. *Rev Esp Enferm Dig* 2016;108:376–8.
36. Baharith H, Khan K. Bouveret syndrome: when there are no options. *Can J Gastroenterol Hepatol* 2015;29:17–8.
37. Smith Z, Totten J, Hughes A, Strote J. Delayed diagnosis of gastric outlet obstruction from bouveret syndrome in a young woman. *West J Emerg Med* 2015;16:151–3.
38. Englert ZP, Love K, Marilley MD, Bower CE. Bouveret syndrome: gallstone ileus of the duodenum. *Surg Laparosc Endosc Percutan Tech* 2012;22:e301–3.
39. Shah SK, Walker PA, Fischer UM, Karanjawala BE, Khan SA. Bouveret syndrome. *J Gastrointest Surg* 2013;17:1720–1.
40. Patel A, Agarwal S. The yellow brick road of Bouveret syndrome. *Clin Gastroenterol Hepatol* 2014;12:A24.
41. Makker J, Muthusamy VR, Watson R, Sedarat A. Electrohydraulic lithotripsy and removal of a gallstone obstructing the duodenum: Bouveret syndrome. *Gastrointest Endosc* 2015;81:1021–2.
42. Chiang AL, Ryou M. Combined Mirizzi syndrome with Bouveret syndrome. *Gastrointest Endosc* 2016;83:1034.
43. Lopes CV, Lima FK, Hartmann AA. Bouveret syndrome and pancreatic acinar cell carcinoma. *Endoscopy* 2017;49:E62–3.
44. Dumonceau JM, Delhay M, Devière J, Baize M, Cremer M. Endoscopic treatment of gastric outlet obstruction caused by a gallstone (Bouveret's syndrome) after extracorporeal shock-wave lithotripsy. *Endoscopy* 1997;29:319–21.
45. Katsinelos P, Dimiropoulos S, Tsolkas P, Baltagiannis S, Kapelidis P, Galanis I, et al. Successful treatment of duodenal bulb obstruction caused by a gallstone (Bouveret's syndrome) after endoscopic mechanical lithotripsy. *Surg Endosc* 2002;16:1363.
46. Malvaux P, Degolla R, De Saint-Hubert M, Farchakh E, Hauters P. Laparoscopic treatment of a gastric outlet obstruction caused by a gallstone (Bouveret's syndrome). *Surg Endosc* 2002;16:1108–9.
47. Fedidat R, Safadi W, Waksman I, Hadary A. Choledochoduodenal fistula: an unusual case of pneumobilia. *BMJ Case Rep* 2014;2014:bcr2014206798.
48. Chick JF, Chauhan NR, Mandell JC, de Souza DA, Bair RJ, Khurana B. Traffic jam in the duodenum: imaging and pathogenesis of Bouveret syndrome. *J Emerg Med* 2013;45:e135–7.