

Assessment of $\alpha 4$, αv , $\beta 1$ and $\beta 3$ integrins expression throughout the implantation window phase in endometrium of a mouse model of polycystic ovarian syndromes

Fatemeh Peyghambari¹ Ph.D., Mehri Fayazi² M.Sc., Saeid Amanpour³ D.V.Sc., Mahnaz Haddadi³ M.Sc., Samad Muhammadnejad⁴ Ph.D., Ahad Muhammadnejad⁴ D.V.Sc., Samira Abdolapour⁴ B.Sc., Mozghan Enkesari⁴ B.Sc., Zohreh Mazaheri² Ph.D.

1. Department of Anatomical Sciences, Faculty of Medical Sciences, Islamic Azad University, Yazd Branch, Yazd, Iran.

2. Department of Anatomical sciences, Medical Sciences Faculty, Tarbiat Modares University, Tehran, Iran.

3. Vali-e-Asr Reproductive Health Research Centre, Tehran University of Medical Sciences, Tehran, Iran.

4. Tumor Model Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, Tehran, Iran.

Corresponding Author:

Zohreh Mazaheri, Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Jalale-Ale-Ahmad Highway, Tehran, Iran. P.O.Box. 14115-175,

Email: z_mazaheri@modares.ac.ir

Tel/Fax: (+98) 21 88011001

Received: 26 January 2014

Revised: 25 May 2014

Accepted: 9 July 2014

Abstract

Background: Endometrial integrin expression changes might be a reason for implantation failure in polycystic ovarian syndromes (PCOS).

Objective: Assessment of integrin genes and proteins expression upon endometrium in the PCOS experimental mouse model was the main goal of this study.

Materials and Methods: 30 NMRI female mice were equally divided into control, experimental (PCOS; received estradiol valerate (40 mg/kg)) and sham group (received; olive oil). After 8 weeks, each group was hyper stimulated by 7 IU PMSG and then, after 48hrs, 7 IU HCG was injected. Vaginal plaque was checked. After 5 days, Progesterone and estradiol levels and endometrial tissues were investigated to evaluate of $\alpha 4$, αv , $\beta 1$ and $\beta 3$ integrins gene and protein by qPCR method and immunohistochemistry, respectively.

Results: Tissue samples were assessed and showed that level of progesterone was significantly decreased in PCOS group. Results of molecular part in the amount of αv , $\beta 3$, $\beta 1$ and $\alpha 4$ gene expressions showed a great difference in $\beta 3$ and αv genes expressions between experimental groups. αv , $\beta 3$, $\alpha 4$ and $\beta 1$ proteins in the endometrial stroma in the control group were expressed, but they were not detected in PCOS group.

Conclusion: According to the results, integrins had different expression patterns in different areas of the endometrium; such as epithelial and stromal. It seems that in PCOS, this pattern has changed and the results might have a great influence on implantation failure. Therefore, this study suggests that a great attention to this problem may be essential in patients who are involved.

Key words: Integrin, Endometrium, Implantation, Polycystic ovarian syndromes.

Introduction

Polycystic ovary syndrome (PCOS) was the first illustrated by Stein and Leventhal in 1935. The major features of women with PCOS are polycystic ovaries and chronic anovulation, insulin resistance, obesity, irregular menstruation, infertility, and hirsutism (1-3). There is some evidences that the endocrinologic and metabolic imbalances in PCOS may have complex effects on the endometrium, take parting to the infertility and endometrial disorders observed in women with this syndrome (2, 4). The uterine receptivity disorder looks to be one of the most important factors in spontaneous abrasion cases such

as PCOS patients, and cure to develop implantation rates is probable to be taken to this process (5).

Researchers indicate that in women with PCOS, the $\alpha v \beta 3$ -integrin, HOXA-10, HOXA-11 and insulin-like growth factor (IGFBP-I) binding protein is down regulated throughout the secretory phase (3, 6-9). This is belief that integrins, cell adhesion molecules (CAMs), are good markers of endometrial receptivity and essential to cell contact. They represent a family of glycoproteins, formed by the two different non-covalent bonds: α and β subunits Integrins take part in cell matrix and cell-to-cell adhesion in many physiological events, such as embryological development, wound healing, immune and non-immune defense

mechanisms and oncogenic transformation. A range of integrins has been expressed in the lumen of the glandular endometrial epithelium (10). Up regulation of $\alpha 1\beta 1$, $\alpha 4\beta 1$ and $\alpha v\beta 3$ integrines has been explained in the endometrium in the mid-luteal phase, and the expression of subunit $\beta 3$ demonstrated no increase before the 19th day, which has been suggested as an endometrial blastocyst receptor (11).

They undergo hormonal effects, where high estrogen levels reduce the expression of integrins and increase the progesterone. Integrins have been expressed by the trophoblast at the time of implantation, demonstrating a sandwich model in embryonic attachment (12, 13). Despite technological progresses in assisted reproduction techniques, allowing the selection of top-quality embryos, the implantation rate is still low and has not increased adequately in recent decades (3, 14). Uterine receptivity has a key character in establishing the success of pregnancy and when changes, it may limit the success of assisted reproduction techniques and supply to infertility in certain gynecological diseases, such as PCOS (15).

Literature concerns on the expression and regulation of $\alpha 4\beta 1$ and $\alpha v\beta 3$ -integrin show that this protein may symbolize a marker for the human implantation process. The existing literature data are not very obvious if there is an endometrial dysfunction in PCOS, despite integrin action. Some studies show that the endometrial process, including cell proliferation, differentiation and response to biological effects, could be involved by the expression of integrins such as $\alpha 4\beta 1$ and $\alpha v\beta 3$ -integrin, explaining in part the poor reproductive outcomes in this group of women (15).

We think that there is a lower expression of integrins in implantation window period in mice with PCOS than in normal mice. To understand this propose, we investigated gene and protein expression of $\alpha 4\beta 1$ and $\alpha v\beta 3$ -integrin as some endometrial receptivity markers in the implantation window period of the experimental PCOS model, which were different from normal ones.

Materials and methods

Animals

Thirty mature female mice were randomly divided to estradiol valerate (EV)-induced PCOS, sham and control groups as an

experimental study (16). The mice were housed in a room under standard laboratory conditions (12 hr light/dark at 22°C) with free access to water and food at the cancer research center of Imam Khomeini hospital. The current study was conducted under the protocol approved by the animal experimentation committee at Islamic Azad University of Yazd animals' laboratory. The EV group received 40 mg/kg body weight estradiol valerate by intramuscular [IM] (17). Sham group also received 100 μ l of olive oil. All the groups were evaluated 60 days after the injection.

Design of the study

Body weight measurement

During the study, the body weight was registered, weekly.

Gonadal hormones assessment

Serum was isolated from heart blood sample by centrifugation (2000 Rounds per min/10 min) in 3 mice from each group, 8 weeks post PCOS induction. 17- β estradiol and progesterone concentrations were evaluated by RIA method (a very sensitive in vitro assay technique used to measure concentrations of hormone levels) in Vali-e-asr hospital lab in the Imam Khomeini hospital.

Polycystic ovaries evaluation

Ovaries of each group were picked up after the treatment period (n: 3 mice), cleaned from adherent fat and connective tissue, and fixed in 10% formaldehyde buffer. Tissue processing and sectioning was down by routine histological techniques, then sections 5-6 μ m were stained with hematoxylin-eosin stain as described by Guyer, and observed under a light microscope [Olympus BX51, Germany] (18).

Sampling of endometrial tissue during the implantation window period

Each group was hyper stimulated by 7 IU PMSG (Folligon, Intervet, Australia) and after 48hr, 7 IU HCG (Serenio, Switzerland) was injected. Vaginal plaques were checked, and the pregnant mice were dislocated 5 days after the test. Endometrial tissue was picked up and divided into two parts. One part was put in 10% formalin buffer, and another part

was kept in -70°C in order to extract RNA in RNA later for assessment of gene expression.

Molecular assessment by quantitative polymerase chain reaction (qPCR)

Total RNA was extracted from embryos in each group using QIAzol (QIAGEN Germany) according to the manufacturer's recommendations. To eliminate genomic contamination, RNA was treated with DNase I using a kit (EN0521; Fermentas). Concentrations of RNA were determined using a UV spectrophotometer (Eppendorf, Germany). The cDNAs were synthesized from 500 ng DNase-treated RNA samples with a RevertAid™ First Strand cDNA Synthesis kit (K1622; Fermentas, Germany) using oligo (dT) primers. For PCR reactions, primers were designed by the NCBI website and gene runner software and synthesized by Pishgam (Table I). PCRs were performed using Master Mix and SYBR Green in an Applied Biosystems, StepOne™ thermal cycler (Applied Biosystems, USA). The PCR program started with an initial melting cycle for 5 min at 95°C to activate the polymerase, followed by 40 cycles of melting (15 sec at 95°C), annealing (30 sec at 58°C) and extension (15 sec at 72°C). The quality of the PCR reactions was confirmed by melting curve analyses. Efficiency was determined for each gene using a standard curve (logarithmic dilution series). For each sample, the reference gene (β -actin) and target gene were amplified in the same run. Reference gene was approximately equal. The relative quantification of target genes, was normalized to a reference gene and determined using the method (19).

Protein assessment by immunohistochemistry method

Endometrial samples were fixed in the formalin buffer. The tissues were sectioned for immunostaining analysis. We used antibodies specific for αv (1:100; Chemicon, UK), $\alpha 4$ (1:100; Abcam, UK), $\beta 1$ (1:250; Abcam, UK) and $\beta 3$ (1:250; Abcam, UK). HRP-labeled secondary rabbit antibody (BIO-IDEA, Iran) was used as a secondary antibody and DAB (Diaminobenzidine tetra hydrochloride hydrate, BIO-IDEA; Iran). Counter staining was done with hematoxylin dye. All of the methods for immunohistochemistry were done according to the manufacturer's instructions. Images

were captured using a light microscope equipped Olympus U-TV0.5 XC-3 camera.

Statistical analysis

Data have been presented as the mean \pm SD. The results were analyzed by SPSS software (version 16.0) using One-way repeated measures analysis of variance (ANOVA) test between the groups. $p < 0.05$ were considered statistically significant, and it was followed by the Tukey post hoc test multiple.

Results

Gaining weight in all groups was observed at the end of study, after 2 months of PCOS induction. Anyway among all groups, there was a significant difference between the first days of study and the last weeks. However, there was not any significant difference between groups (Figure 1). The amount of 17- β estradiol in the sera of control, sham and PCOS groups was 49.14 ± 3.8 , 41.17 ± 4.1 and 133.11 ± 5.6 pg/ml and that the progesterone level was 28 ± 2.01 , 35.12 ± 1.72 , 18.63 ± 2.11 ng/ml, respectively. The level of 17- β estradiol increased significantly in EV-induced mice compared with the control and sham mice ($p < 0.01$). Also, the level of progesterone decreased significantly in EV-induced mice compared with the other groups ($p < 0.04$).

In the study of tissue sections, the presence of many cysts in ovaries full of cyst tissue in the PCOS induced groups was confirmed (Figure 2). Evaluations of molecular part in the amount of αv , $\beta 3$, $\beta 1$ and $\alpha 4$ gene expression showed that there was a great difference in gene expression between $\beta 3$, αv and other groups. The results showed that $\beta 1$ and $\alpha 4$ expressions in the endometrium of PCOS-induced mice were significantly ($p \leq 0.001$ and $p \leq 0.046$, respectively) increased whereas that of $\beta 3$ and αv expression was significantly ($p \leq 0.001$) decreased compared with other groups; however, the expressions observed in the control and sham groups showed no significant difference in none of the integrin genes (Figure 3).

The αv , $\alpha 4$, $\beta 1$ and $\beta 3$ integrins were detected in the endometrium of all groups during the implantation window period. The expression of integrin molecules was as a diffuse pattern in apical and basal part of surface and glandular endometrial epithelium

(Figure 4 & 5 A-D). The $\alpha 4$ subunit was most intense in the apical and basal membranes of the glandular uterine epithelial cells (Figure 4 & 5 C) and the αv subunit showed a diffuse pattern that was intense in the apical and basal membranes in both surface and glandular uterine epithelial cells (Figure 4 & 5 B). These two types of α subunit were not detected in the endometrial stroma of PCOS-induced mice during of the implantation window period. $\beta 1$ integrin subunit exhibited a

diffuse staining pattern, intensely positive in the membrane of glandular epithelial cells (Figure 4D). $\beta 3$ integrin subunit was small intense in both surface and glandular epithelial cells of the endometrium of PCOS-induced mice, and it was not detected in the stroma (Figure 5A). All of integrin molecules and expression were restricted to the surface and glandular epithelium and was not detected in the stroma in PCOS-induced mice (Figure 5A-D).

Table I. Primer Used for real- time PCR

Genes	Primer sequence	GenBank code	Tm (°C)
$\beta 1$ integrin	FOR: 5'- TGCCTACAACCTCTTTCTTC-3'	NM_010578.2	58
	REV: 5'- TGGTTTCAGACTCCTTATTG-3'		
β -actin	FOR: 5'- TCCCTGGAGAAGAGCTACG-3'	NM_001101	66.6
	REV: 5'- GTAGTTTCGTGGATGCCACA-3'		
$\beta 3$ integrin	FOR: 5'- TGGAAAGAGCCTGAGTGTC-3'	NM_016780	60
	REV: 5'- CCGTAGGTGATATTGGTGAAG-3'		
αv integrin	FOR: 5'- GGAACAACGAAGCCTTAG-3'	NM_008402	55
	REV: 5'- GTATCCATCTCTGACTGC-3'		
$\alpha 4$ integrin	FOR: 5'- GAATCTCCTCCACCTACTCACAG -3'	NM_010576	64
	REV: 5'- CCAACGGCTACATCAACATATCC-3'		




Figure 1. Mean of body weight changes after 8 weeks of PCOs induction in the experimental groups.




Figure 2. In the ovarian tissue, A; the cysts were mainly appeared by a single intramuscular dose of estradiol valerate, 40 mg/kg (H&E). B; ovarian tissue in the control group.




Figure 3. Real Time-PCR analysis. mRNA levels were normalized with respect to β -actin, chosen as an internal control and calibrated to control group. Profile of Integrin genes expression of endometrium after using a single intramuscular dose of estradiol valerate, 40 mg/kg. Histograms show mean expression values (\pm SD; $p < 0.05$). α : significant difference with other groups in the same genes.




Figure 4. Immunostaining reactivity in endometrial tissue of normal mice. A; $\beta 3$ Integrin expression, B; αV Integrin expression, C; $\alpha 4$ Integrin expression, D; $\beta 1$ Integrin expression. Red arrow: stromal expression, Black arrow: Epithelial expression.




Figure 5. Immunostaining reactivity in endometrial tissue of PCOs model. A; $\beta 3$ Integrin expression, B; αV Integrin expression, C; $\alpha 4$ Integrin expression, D; $\beta 1$ Integrin expression. Red arrow: Basal epithelium expression, Black arrow: Luminal (apical) epithelium expression.

Discussion

Understanding endometrial receptivity is an invasive subject in the fertility field in human. Thus, in this study, we used an animal PCOS model. An single dose injection of EV in the mouse produces an PCO condition. PCOS

signs in mice, received 40 mg/kg estradiol valerate, were made after 8 weeks. The effect of estradiol valerate on hypophysis-hypothalamus pathway was observed after measuring of sexual hormone secretion. This plasma gonadotropin pattern indirectly was related to ovules, too and makes multiple

small cysts. The cystic ovary produced by EV is similar in appearance to that seen in the human Stein-Leventhal condition, and thus provides a good model for the study of cystic ovarian disease (20-22).

Thus 8 weeks later, injection was used to assess hormone levels. Results have shown that a significant decrease in the progesterone level and a significant increase in the estrogen level in experimental groups. Luza *et al.* in 1995 and Brawer in 1978 made a hypothalamus-hypophysis pathway impairment by EV injection. This impairment could be a reason for change in the secretion of hypothalamic stimulators and consequently hypophysis secretion. This new pathway can affect the sexual hormone amount in blood serum (23-24).

We have demonstrated, for the first time, the expression of several integrin molecules in the endometrium of PCOS-induced mice, during the implantation window period using real-time RT-PCR in combination with immunohistochemistry techniques. Our observations, obtained by two techniques, demonstrated that there was not any sign of expression for integrins in the endometrial stroma of PCOS mice. These patterns of expression were correlated and synchronous well with the period when the serum progesterone concentration was increasing at the metestrus phase of the estrous cycle. The metestrus phase is like the secretory phase of the endometrium in human, and in this phase, the endometrium under the effect of circulatory progesterone undergoes some changes for reception of the embryo, and the implantation occurs in this phase (25).

It seems that the expression of these molecules in the endometrial stroma was regulated by progesterone while the estrogen had negligible effect in this regard. Regulation of integrin molecules expression is very complex and is hard to realize what regulates the expression of these during the implantation period. The expressions of epithelial $\alpha\beta3$ integrin 6-8 days after ovulation, occurs with serum progesterone level increase, simultaneously (26). It has described that estrogen down-regulated $\alpha\beta3$ expression, but no effect was observed for $\alpha4$ or $\beta1$ integrin subunit (27).

Down-regulation of $\alpha4$ by progesterone and its up regulation by estradiol, also reported by some researcher in contrast to our study, in

the blastocysts and the endometrium by a delayed-implantation mouse model (28). In this regards, some of studies showed that the expression of $\alpha\beta3$ integrin heterodimers not influenced by sex steroids in vivo (29-32). Lin *et al.* revealed that there is no correlation between the serum levels of steroid hormones and the $\alpha\beta3$ integrins expression (33). These controversially outcomes might be due to some reasons like various types of integrin molecules, different protocols or experimental designs and types of species.

Integrin cell adhesion molecules are one of the usual receptors upon endometrium surface in all mammals, and express in a cyclic pattern (27). We examined the profile of $\alpha\beta3$, $\beta3$, $\beta1$ and $\alpha4$ expression in epithelial level and endometrial stroma in PCOS mice model. $\alpha4$ subunit up regulated in the endometrium of the experimental mice groups than other molecules, and was expressed intense in the epithelium and stroma of the PCOS group. These findings can be a sign of roles and value of different types of integrins heterodimers like $\alpha4\beta3$, $\alpha4\beta1$, $\alpha\beta1$, $\alpha\beta3$ in different site of endometrium for making of contact between the embryo and endometrium during the implantation time. In this subject, Basak *et al* described the $\alpha4$ integrin expression on the basement membrane of the endometrium might aid in adherent of the embryo to the uterine lumen (28).

They have coated the $\alpha4$ integrin by monoclonal antibodies in the endometrium of pregnant mice during implantation window and revealed the implantation failure in both normal as well as the delayed implantation mice (28). $\alpha4$ epithelial subunit as a fibronectin receptor ($\alpha4\beta1$) has a high expression in fertile and infertile women in the whole menstrual cycle. Epithelial $\alpha4$ in fertile women increased in an implantation window period as similar as epithelial $\alpha4\beta1$ expression in those who used anti-pregnancy pills (34-35). Stromal $\alpha4$ is also seen in infertile women during the last proliferation period and the beginning of secretion (36). In the fertile women, the belief in molecular adherence expression pattern changed in the endometrial stroma is identified as unexplained infertility.

Epithelial $\beta3$ expression in fertile women is significantly increased in the implantation window period, while in infertile women this level is weakly expressed in the mid-secretion

phase (37). Thus, a deficiency in the epithelial $\beta 3$ expression can be a reason for infertility. In fertile women in the last part of the secretion, αv , $\alpha 4$, $\alpha 1$ and $\beta 3$ are secreted. Molecular importance of this change in infertile women is not known (34).

αv expression, which is a vitronectin receptor ($\alpha v\beta 3$), has a different expression pattern related to its molecule, i.e. $\beta 3$ (38). Integrin αv only expresses in last secretion phase in the epithelium of infertile women (36). This protein in women's epithelium and in Ishikawa cells is also shown. αv in addition to $\beta 3$ can be expressed along other β subunits such as $\beta 1$, $\beta 5$ and $\beta 8$ (34-35).

A decrease in stromal $\beta 1$ and $\alpha 1$ in unexplained infertile women is seen that can cause a deficiency in trophoblast linking with stromal cells. Regulation of $\beta 1$ and $\alpha 1$ and their function during the implantation period in human is actually unidentified. The function of the endometrial steroid hormones by some growth factors and cytokines may result in integrin expression depended on the cycle (39). So, based on other studies, it was revealed that hormonal fluctuations due to cystic ovarian syndrome and can regulate Integrin cell adhesion molecules involved in endometrium and embryo attachment. Furthermore, expression out of the cycle in these molecules can directly affect fertility in patients who are being treated.

Conclusion

The present study shows that in mouse with PCOS, αv and $\beta 3$ integrin expression in the endometrial tissue was significantly lower than normal mouse. Additionally, in these animal model, a high expression of $\alpha 4$ and $\beta 1$, involved integrins in the implantation process in a cyclic way during the whole period, could exist. Although this expression in different areas of the endometrium such as the epithelial and stromal, had a different expression pattern in investigated groups, but it seems that in PCOS, this pattern had changed and the results had a great effect on implantation failure.

Acknowledgments

This work was financially supported by Yazd Branch of Islamic Azad University. We gratefully thank Dr. Zargarzadeh and Dr. Dini for excellent technical assistance.

Conflict of interest

There is no conflict of interest in this study.

References

1. Speroff L, Fritz MA. Chronic anovulation and the polycystic ovary syndrome. In: Clinical gynecologic endocrinology and infertility. Philadelphia; 2011: 495-531.
2. Qiao J, Wang L, Li R, Zhang X. Microarray evaluation of endometrial receptivity in Chinese women with polycystic ovary syndrome. *Reprod Biomed Online* 2008; 17: 425-435.
3. Cakmak H, Taylor HS. Human implantation failure: molecular mechanisms and clinical treatment. *Hum Reprod Update* 2011; 17: 242-253.
4. Donaghy M, Lessey BA. Uterine receptivity: alterations associated with benign gynecological disease. *Semin Reprod Med* 2007; 25: 461-475.
5. Lopes IM, Baracat MC, Simões Mde J, Simões RS, Baracat EC, Soares Jr JM. Endometrium in women with polycystic ovary syndrome during the window of implantation. *Rev Assoc Med Bras* 2011; 57: 702-709.
6. Bergeron C, Ferenczy A, Shyamala G. Distribution of estrogen receptors in various cell types of normal, hyperplastic, and neoplastic human endometrial tissues. *Lab Invest* 1988; 58: 338-345.
7. Apparao KB, Lovely LP, Gui Y, Lininger RA, Lessey BA. Elevated endometrial androgen receptor expression in women with polycystic ovarian syndrome. *Biol Reprod* 2002; 66: 297-304.
8. Cermik D, Selam B, Taylor HS. Regulation of HOXA-10 expression by testosterone in vitro and in the endometrium of patients with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003; 88: 238-243.
9. Daftary GS, Kayisli U, Seli E, Bukulmez O, Arici A, Taylor HS. Salpingectomy increases peri-implantation endometrial HOXA10 expression in women with hydrosalpinx. *Fertil Steril* 2007; 87: 367-372.
10. Zhao Y, Garcia J, Kolp L, Cheadle C, Rodriguez A, Vlahos NF. The impact of luteal phase support on gene expression of extracellular matrix protein and adhesion molecules in the human endometrium during the window of implantation following controlled ovarian stimulation with a GnRH antagonist protocol. *Fertil Steril* 2010; 94: 2264-2271.
11. Lessey BA. Two pathways of progesterone action in the human endometrium: implications for implantation and contraception. *Steroids* 2003; 68: 809-815.
12. Wang J, Armant DR. Integrin-mediated adhesion and signalling during blastocyst implantation. *Cells Tissues Organs* 2002; 172: 190-201.
13. Achace H, Revel A. Endometrial receptivity markers, the journal to successful embryo implantation. *Hum Reprod Update* 2006; 12: 731-746.
14. Andersen AN, Gianaroli L, Felberbaum R, de Mouzon J, Nygren KG. Assisted reproductive technology in Europe, 2001. Results generated from European registers by ESHRE. *Hum Reprod* 2005; 20: 1158-1176.

15. Donaghy M, Lessey BA. Uterine receptivity: alterations associated with benign gynecological disease. *Semin Reprod Med* 2007; 25: 461-475.
16. Dikmen A, Ergenoglu AM, Yeniel AO, Dilisiz OY, Ercan G, Yilmaz H. Evaluation of glyceemic and oxidative/antioxidative status in the estradiol valerate-induced PCOS model of rats. *Eur J Obstet Gynecol Reprod Biol* 2012; 160: 55-59.
17. Mahood RAH. Effects of Pimpinella anisum oil Extract on Some Biochemical Parameters in Mice experimentally induced for human Polycystic Ovary Syndrome. *J Biotech Research Cent* 2012; 6: 67-73.
18. Guyer M. Animal microbiology. 5th Ed. The University of Chicago press, Chicago; 1993.
19. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001; 29: e45.
20. Stener-Victorin E, Kobayashi R, Watanabe O, Lundeberg T, Kurosawa M. Effect of electro-acupuncture stimulation of different frequencies and intensities on ovarian blood flow in anaesthetized rats with steroid-induced polycystic ovaries. *Reprod Biol Endocrinol* 2004; 2: 16.
21. Schulster A, Farookhi R, Brawer J R. Polycystic ovarian condition in Estradiol Valerate-treated rats: spontaneous changes in characteristic endocrine features. *Biol Reprod* 1984; 31: 587-593.
22. Shirwalker H, Deepak N M, Anurupa M. Exposure of adult rats to estradiol valerate induces ovarian cyst with early senescence of follicles. *Mol Cell Endocrinol* 2007; 272: 22-37.
23. Schulster A, Farookhi R, Brawer J R. Polycystic ovarian condition in Estradiol Valerate-treated rats: spontaneous changes in characteristic endocrine features. *Biol Reprod* 1984; 31: 587-593.
24. Brawer JR, Naftolin F, Martin J, Sonnenschein C. Effects of a single injection of estradiol valerate on the hypothalamic arcuate nucleus and on reproductive function in the female rat. *Endocrinology* 1978; 103: 501-512.
25. Peyghambari F, Salehnia M, Forouzandeh Moghadam M, Rezazadeh Valujerdi M, Hajizadeh E. The correlation between the endometrial integrins and osteopontin expression with pinopodes development in ovariectomized mice in response to exogenous steroids hormones. *Iran Biomed J* 2010; 14: 109-119.
26. Lessey BA, Damjanovich L, Coutifaris C, Castelbaum A, Albelda SM, Buck CA. Integrin adhesion molecules in the human endometrium. Correlation with the normal and abnormal menstrual cycle. *J Clin Invest* 1992; 90: 188-195.
27. Widra EA, Weeraratna A, Stepp MA, Stillman RJ, Patierno SR. Modulation of implantation-associated integrin expression but not uteroglobin by steroid hormones in an endometrial cell line. *Mol Hum Reprod* 1997; 3: 563-568.
28. Basak S, Dhar R, Das C. Steroids modulate the expression of alpha4 integrin in mouse blastocysts and uterus during implantation. *Biol Reprod* 2002; 66: 1784-1789.
29. Kimmins S, MacLaren LA. Cyclic Modulation of integrin expression in bovine endometrium. *Biol Reprod* 1999; 61: 1267-1274.
30. Lessey BA, Arnold JT. Paracrine signaling in the endometrium: integrins and the establishment of uterine receptivity. *J Reprod Immunol* 1998; 39: 105-116.
31. Apparao KB, Murray MJ, Fritz MA, Meyer WR, Chambers AF, Truong PR, Lessey BA. Osteopontin and its receptor alphavbeta (3) integrin are coexpressed in the human endometrium during the menstrual cycle but regulated differentially. *J Clin Endocrinol Metab* 2001; 86: 4991-5000.
32. Bowen JA, Bazer FW, Burghardt RC. Spatial and temporal analyses of integrin and Muc-1 expression in porcine uterine epithelium and trophectoderm in vivo. *Biol Reprod* 1996; 55: 1098-1110.
33. Lin H, Wang X, Liu G, Fu J, Wang A. Expression of alphaV and beta3 integrin subunits during implantation in pig. *Mol Reprod Dev* 2007; 74: 1379-1385.
34. Lim KJ. Molecular factors in human implantation: adhesion molecules, proteases and cytokines. *Malays J Pathol* 2003; 25: 1-13.
35. Reddy KV, Mangale SS. Integrin receptors: the dynamic modulators of endometrial function. *Tissue Cell* 2003; 35: 260-273.
36. Gonzalez RR, Palomino A, Boric A, Vega M, Devoto L. A quantitative evaluation of alpha1, alpha4, alphaV and beta3 endometrial integrins of fertile and unexplained infertile women during the menstrual cycle. A flow cytometric appraisal. *Hum Reprod* 1999; 14: 2485-2492.
37. Lessey BA, Castelbaum AJ, Sawin SW, Sun J. Integrins as markers of uterine receptivity in women with primary unexplained infertility. *Fertil Steril* 1995; 63: 535-542.
38. Lim JJ, Lee DR, Song HS, Kim KS, Yoon TK, Gye MC, Kim MK. Heparin-binding epidermal growth factor (HB-EGF) may improve embryonic development and implantation by increasing vitronectin receptor (integrin alphanubeta3) expression in peri-implantation mouse embryos. *J Assist Reprod Genet* 2006; 23: 111-119.
39. Singh H, Aplin JD. Adhesion molecules in endometrial epithelium: tissue integrity and embryo implantation. *J Anat* 2009; 215: 3-13.