



HHS Public Access

Author manuscript

Longhua Chin Med. Author manuscript; available in PMC 2024 May 31.

Published in final edited form as:

Longhua Chin Med. 2023 October 31; 6: . doi:10.21037/lcm-23-7.

Associations between traditional Chinese medicine body constitution and obesity risk among US adults

Xiangzhu Zhu^{1, #}, Xiaolin Yin^{2, #}, Xinqing Deng³, Yevheniy Eugene Shubin⁴, Harvey J. Murff⁵, Reid M. Ness⁶, Chang Yu⁷, Martha J. Shrubsole¹, Qi Dai¹

¹Division of Epidemiology, Department of Medicine, Vanderbilt University School of Medicine, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, TN, USA;

²Shanghai Municipal Hospital of Traditional Chinese Medicine Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai, China;

³Department of Medicine, Vanderbilt University School of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA;

⁴Saint Joseph Health System, South Bend, IN, USA;

⁵Department of Medicine, Division of Geriatric Medicine, Vanderbilt University Medical Center, Nashville, TN, USA;

⁶Department of Medicine, Division of Gastroenterology, Hepatology, and Nutrition, Vanderbilt University Medical Center, Nashville, TN, USA;

⁷Department of Population Health, Division of Biostatistics, NYU Grossman School of Medicine, New York, NY, USA

Abstract

Background: Traditional Chinese medicine (TCM) body constitution (BC), primarily determined by physiological and clinical characteristics, is an important process for clinical diagnosis and treatment and play a critical role in precision medicine in TCM. The purpose of the study was to explore whether the distributions of BC types differed by obesity status.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the noncommercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Correspondence to: Xiangzhu Zhu, MD, MPH. Division of Epidemiology, Department of Medicine, Vanderbilt University Medical Center, 2525 West End Ave, Nashville, TN 37203, USA. xiangzhu.zhu@vanderbilt.edu.

Contributions: (I) Conception and design: X Zhu, Q Dai; (II) Administrative support: Q Dai, MJ Shrubsole, C Yu, X Zhu; (III) Provision of study materials or patients: Q Dai, MJ Shrubsole, HJ Murff, RM Ness, X Zhu, X Yin; (IV) Collection and assembly of data: X Zhu, X Yin, X Deng, YE Shubin; (V) Data analysis and interpretation: X Zhu, Q Dai; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work as co-first authors.

Peer Review File: Available at <https://lcm.amegroups.com/article/view/10.21037/lcm-23-7/prf>

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study was approved by the Vanderbilt Institutional Review Board (IRB) (No. 100106) and informed consent was obtained from all individual participants.

Methods: We conducted a study to evaluate BC type in US population during 2012–2016. A total of 191 White participants from Personalized Prevention of Colorectal Cancer Trial (PPCCT) completed a self-administered Traditional Chinese Medicine Questionnaire (TCMQ, English version). In this study, we further compared the distribution of major types of TCM BC in the PPCCT to those Chinese populations stratified by obesity status.

Results: We found the Blood-stasis frequency was higher in US White adults, 22.6% for individuals with BMI <30 and 11.2% for obese individuals, compared to 1.4% and 1.8%, respectively, in Chinese populations. We also found the percentages Inherited-special and Qi-stagnation were higher in US White adults than those in Chinese populations regardless of obesity status. However, the proportions of Yang-deficiency were higher in Chinese populations than those in our study conducted in US White adults regardless of obesity status.

Conclusions: These new findings indicate the difference in distribution of BC types we observed between US and Chinese populations cannot be explained by the differences in prevalence of obesity. Further studies are needed to confirm our findings and understand the potential mechanism including genetic background and/or environmental factors.

Keywords

Traditional Chinese medicine (TCM); body constitution (BC); overweight; obesity

Introduction

In recent decades, obesity has surged to epidemic proportions and has been a major contributor to the global burden of chronic noncommunicable diseases and the associated disability. According to the World Health Organization (WHO) in 2016, over 1.9 billion aged 18 years and older were overweight and 650 million adults were obese. Obesity is not only a condition of increased body weight but also a complex disorder that affects glucose, lipid, and protein metabolism. Furthermore, it is closely related to metabolic disorders such as insulin resistance (IR), type 2 diabetes mellitus (T2DM), and liver steatosis, leading to significant morbidity, mortality, and societal burden (1).

Body constitution (BC), the foundational concept of traditional Chinese medicine (TCM), classifies individuals into nine types: Balanced constitution, Qi-deficiency constitution, Yang-deficiency constitution, Yin-deficiency constitution, Phlegm-dampness constitution, Damp-heat constitution, Blood-stasis constitution, Qi-stagnation constitution and Inherited-special constitution (2). This classification is based on physical, psychological, and physiological characteristics, as well as susceptibility to illnesses and adaptability to the environment. These constitutions are dynamic and can be influenced by acquired factors including lifestyles and environment (3). BC is an important process for clinical diagnosis and treatment and play a critical role in precision medicine and precision prevention.

However, TCM plays a very limited role in Western medical practice. Almost all these previous studies using TCM BC have been conducted in Chinese populations except for one study of 400 White college students attending three Beijing universities in China (4).

Very recently, in the first conducted in an American population (5), we found the distribution of TCM BC types in a US White population were different from the distribution among Chinese individuals in China. TCM BC has been previously associated with obesity in China (6) and the US population has higher prevalence of obesity than the Chinese populations (7). Thus, it is possible that the difference in distribution of major types of TCM BC may be due to higher obesity prevalence in our study population. In this study, we compared the distribution of major types of TCM BC between our study population and Chinese populations according to obesity status.

Methods

Study population

The participants in this study were from the Personalized Prevention of Colorectal Cancer Trial (PPCCT, [NCT01105169](https://clinicaltrials.gov/ct2/show/study/NCT01105169) at [ClinicalTrials.gov](https://clinicaltrials.gov)) which was a double-blind, placebo-controlled, randomized precision-based magnesium trial conducted between March 2011 and January 2016. Aged 40 to 85 years with history of colorectal polyp or high risk of colorectal cancer (CRC) participants were enrolled from Vanderbilt University Medical Center. Detailed inclusion and exclusion criteria have been reported (8,9). In the parent study, 250 participants enrolled and 239 completed the study with 11 participants finishing part of the study before withdrawal. One participant completed questionnaires and provided samples at baseline and at the end of the trial after withdrawal. In total, 240 participants were included (shown in Table 1). Traditional Chinese Medicine Questionnaire (TCMQ, English version: Health Questionnaire) as a component of the study was approved by IRB on May 31, 2012, which was a year after the trial started. A total of 191 participants (80% of 240) completed the self-administered TCMQ from May 31, 2012 to Jan 30, 2016 (5). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study was approved by the Vanderbilt Institutional Review Board (IRB) (No. 100106) and informed consent was obtained from all individual participants.

Constitution in TCMQ

TCMQ is an English version of the standardized Constitution in Chinese Medicine Questionnaire (CCMQ) was developed by Wang *et al.* in China (10). As reported in our study (5), the construct validity of TCMQ was confirmed with scaling success rates that ranged from 75.0% to 100%, and confirmatory factor analysis indicate an acceptable model fit. The results of reliability for test-retest reliability (intra-class correlation coefficients) were 0.7–0.8, the Cronbach's alphas ranged from 0.44 to 0.72 during three-month period. The TCMQ consists of 60 questions and each item (question) was graded on a 5-point Likert scale, ranging from 1 (not at all) to 5 (very much). Each of the nine subscales within the TCMQ assessed one type of the TCM BC individually. A total score of each subscale was obtained by summing relevant item scores and the transformed score was created for each type by using the following equation.

$$\text{Transform score} = 100\% \times \frac{\text{Original score} - \text{Num. of item}}{\text{Num of item} \times 4}$$

[1]

Following the criteria, a higher score in a specific TCMQ BC subscale indicates a higher likelihood of the corresponding BC type, with a score of 30 being a “threshold” for case definition. The following rule was applied to ultimately classify a participant within a specific TCM BC: when (I) the score for the Balanced subscale was greater than or equal to 60 and other BC type scores were less than 30, the study participant was diagnosed as “Balanced” which was considered the balanced BC; when (II) the score for an imbalanced BC type (all other BCs) was greater than or equal to 40, then the participant was regarded as one or more of eight imbalanced BC types; and when (III) a BC type score was between 30–40, the diagnosis was made by a well-trained Chinese Medicine Practitioner.

Anthropometric measurements

Anthropometry or body measures for the participants were conducted during their clinic visits by trained personnel. Weight was taken on a digital scale and measured in kilograms. When the participant was properly positioned and the digital readout was stable, the study nurse recorded the number on the screen. Standing height was measured in centimeter with a fixed stadiometer with a vertical backboard and a moveable headboard. They were asked to stand on the floor with the heels of both feet together and the toes pointed slightly outward at approximately a 60° angle. When the participant was properly positioned, the study nurse recorded the height. At least two measurements were required, the 3rd measurement was made only if the difference between the first two was greater than the difference threshold (0.1 kg for weigh and 1 cm for height). The means of two or three measurements (including weight, height, waist circumference, hip circumference) separated by 30 seconds were used for the analysis. BMI values are calculated for participants using measured height and weight values as follows: weight (kilograms)/height (meters squared). BMI criteria are used to for weight categories: normal or desirable weight (BMI values <25.0), overweight (BMI values 25.0–29.9), obese (BMI values ≥ 30.0).

Statistical analyses

Mean ± standard deviation for continuous demographic variables and percentage for categorical demographic variables were presented in Table 1. We performed general linear model for continuous variables or Pearson chi-squared tests, and Fisher’s exactly tests for categorical variables. We compared the distribution of nine TCM BC types by obesity status and population. All P values are two sided and statistical significance was determined using an alpha level of 0.05. The data analyses used software SAS Enterprise Guide 7.1.

Results

We compared baseline demographic characteristics between 191 participants who completed TCMQ and all participants (n=240) in the parent study (Table 1). No significant difference was found for baseline characteristics between these two groups. Table 2 showed the distributions of the nine-body constitution by body weight and no difference was found between normal, overweight, and obese participants.

We found that the Blood-stasis proportion was higher in US White adults, 22.6% for individuals with BMI <30 and 11.2% for obese individuals, compared to 1.4% and 1.8%, respectively, in Chinese populations (Table 3). Also, the percentages of Inherited-special were higher in US White adults, 12.8% for individuals with BMI <30 and 7.9% for obese individuals, than 1.7% and 1.6%, respectively, in Chinese populations (11). Likewise, we observed that the proportions of Qi-stagnation were higher in US White adults, 4.9% for individuals with BMI <30 and 7.9% for obese individuals, than 1.0% and 1.6%, respectively, in Chinese populations. Conversely, the proportions of Yang-deficiency were higher in Chinese populations, 18.8% for individuals with BMI <30 and 10.6% for obese individuals, than 4.9% and 4.5%, respectively in our study conducted in US White adults. ($P_1 < 0.0001$ among BMI <30 and $P_2 < 0.0001$ among obesity adults).

In our recent report (5), the distribution of BC types varied greatly between our study conducted in US White adults and those conducted in Chinese populations. Balanced (29.8%) was the predominant BC type in our study, while the three most common pathologic BC types were Blood-stasis (17.3%), Qi-deficiency (13.6%), and Inherited-special (10.5%). In contrast, the most common pathologic BC types found in Chinese populations were Qi-deficiency, Yang-deficiency, Yin-deficiency, and Phlegm-Dampness. In this study, the differences of the three BC proportions between US population and Chinese population remained after considering obesity status.

Discussion

These new findings indicate the difference in distribution of BC types we observed between populations cannot be explained by the differences in prevalence of obesity. However, our finding was not conclusive. The different distribution between our study population in US and that conducted in previous studies may be caused by selection biases. In our study, over 46% of the study participants were obese whereas only 10.1% of participants in studies conducted in China were obese. Thus, the difference could be due to the different prevalence rates in obesity. In addition, our study participants had a personal history of colorectal polyp or were at high risk of CRC. Another possibility is that individuals diagnosed with colorectal polyp or at high risk of CRC had different BC types. Tao *et al.* recently reported that the observed differences in the distribution of Blood-Stasis cannot be attributable to this selection bias, as well as different distributions in sex and age (12). Thus, it is possible that the observed differences are due to the different genetic background or environmental factors, including different dietary patterns.

Conclusions

The dissimilarity in distribution of BC types between US population and Chinese population cannot be solely attributed to the variations in obesity prevalence. Additional research is necessary to validate and comprehend the potential reasons for the observed disparity.

Acknowledgments

The authors are grateful to the participants for participating in PPCCT. Authors thank the study staffs for their participation.

Funding:

This study was supported by R01 CA149633 from the National Cancer Institute, R01DK110166 from National Institute of Diabetes and Digestive and Kidney Diseases, Department of Health and Human Services as well as the Ingram Cancer Center Endowment Fund. Data collection, sample storage, and processing for this study were partially conducted by the Survey and Biospecimen Shared Resource, which is supported in part by P30CA68485. Clinical visits to the Vanderbilt at the Clinical Research Center were supported in part by the Vanderbilt CTSA grant UL1 RR024975 from NCRR/NIH. The parent study data were stored in Research Electronic Data Capture (REDCap), and data analyses (VR12960) were supported in part by the Vanderbilt Institute for Clinical and Translational Research (UL1TR000445).

Conflicts of Interest:

All authors have completed the ICMJE uniform disclosure form (available at <https://lcm.amegroups.com/article/view/10.21037/lcm-23-7/coif>). The series “Different Distribution of Traditional Chinese Medicine Body Constitution Across Population, Mechanism and Implications” was commissioned by the editorial office without any funding or sponsorship. QD served as the unpaid Guest Editor of the series. RMN is the Chairperson for the Comprehensive Cancer Network (NCCN) Colorectal Cancer Screening Guidelines Committee. MJS received funding support R01DK110166 from National Institute of Diabetes and Digestive and Kidney Diseases and R03 CA189455 from the National Cancer Institute. QD received funding support (R01 CA149633 and R01 CA202936) from the National Cancer Institute, Department of Health and Human Services as well as the Ingram Cancer Center Endowment Fund. The authors have no other conflicts of interest to declare.

Data Sharing Statement:

Available at <https://lcm.amegroups.com/article/view/10.21037/lcm-23-7/dss>

References

- Ghaben AL, Scherer PE. Adipogenesis and metabolic health. *Nat Rev Mol Cell Biol* 2019;20:242–58. [PubMed: 30610207]
- Wang J, Li YS, Wang Q. Identification of Chinese Medicine Constitution in Public Health Services. *Chin J Integr Med* 2019;25:550–3. [PubMed: 28028719]
- Yap SY, Foo CN, Lim YM, et al. Traditional Chinese Medicine Body Constitutions and Psychological Determinants of Depression among University Students in Malaysia: A Pilot Study. *Int J Environ Res Public Health* 2021;18:5366. [PubMed: 34069915]
- Bai M Applying Constitution in Chinese Medicine Questionnaire, designed by WANG Qi(English version) to survey TCM constitutions of the American and Canadian Caucasian in Beijing. *China Journal of Traditional Chinese Medicine and Pharmacy* [Internet]. 2012 [cited 2023 Jan 22]; Available online: <https://www.semanticscholar.org/paper/Applying-Constitution-in-Chinese-Medicine-by-WANG-Ming-hua/bf5170facd2cf11f78abf50bfdc61dea7a5d4922>
- Shu L, Yin X, Zhu X, et al. Associations between Traditional Chinese Medicine Body Constitution and Cardiovascular Disease Risk in a White population [Internet]. *medRxiv*; 2022 [cited 2023 Jan 22]. p. 2022.12.13.22283433. Available online: <https://www.medrxiv.org/content/10.1101/2022.12.13.22283433v1>
- Sun Y, Liu P, Zhao Y, et al. Characteristics of TCM constitutions of adult Chinese women in Hong Kong and identification of related influencing factors: a cross-sectional survey. *J Transl Med* 2014;12:140. [PubMed: 24886055]
- Obesity Rates by Country 2023 [Internet]. [cited 2023 Jan 22]. Available online: <https://worldpopulationreview.com/country-rankings/obesity-rates-by-country>
- Zhu X, Borenstein AR, Zheng Y, et al. Ca:Mg Ratio, APOE Cytosine Modifications, and Cognitive Function: Results from a Randomized Trial. *J Alzheimers Dis* 2020;75:85–98. [PubMed: 32280092]
- Dai Q, Zhu X, Manson JE, et al. Magnesium status and supplementation influence vitamin D status and metabolism: results from a randomized trial. *Am J Clin Nutr* 2018;108:1249–58. [PubMed: 30541089]
- Wong W, Lam CL, Wong VT, et al. Validation of the constitution in chinese medicine questionnaire: does the traditional chinese medicine concept of body constitution exist? *Evid Based Complement Alternat Med* 2013;2013:481491. [PubMed: 23710222]

11. Li M, Mo S, Lv Y, et al. A Study of Traditional Chinese Medicine Body Constitution Associated with Overweight, Obesity, and Underweight. *Evid Based Complement Alternat Med* 2017;2017:7361896. [PubMed: 29234426]
12. Tao M, Zhu X, Yin X. Different Traditional Chinese Medicine Body Constitution in White and Chinese Patients with Colorectal Polyp. *Longhua Chinese Medicine*, 2023. In press.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlight box**Key findings**

- Blood-stasis, Inherited-special and Qi-stagnation proportions were higher in US White adults compared to Chinese populations despite obesity status.

What is known and what is new?

- This is first study to compare the distribution of Traditional Chinese medicine body constitution in White population to that in Chinese population stratified by obesity status.

What is the implication, and what should change now?

- Further large studies are needed to confirm the findings.

Table 1
Baseline characteristics of participants with or without TCMQ, results from PPCCT

Characteristic	Overall (n=240)	With TCMQ (n=191)	Without TCMQ (n=49)	P value
BMI (kg/m ²)	30.3±6.3	30.3±6.7	30.1±5.0	0.95
<25	47 (19.6)	40 (20.9)	7 (14.4)	
25–29.9	78 (32.5)	62 (32.5)	16 (32.6)	0.93
30	115 (47.9)	89 (46.6)	26 (53.0)	
Waist circumference (cm)	106.9±15.7	106.8±16.7	107.4±11.1	0.92
Age (year)	60.8±8.0	60.0±7.8	63.8±7.8	0.33
Gender male	127 (52.9)	100 (52.4)	27 (55.1)	0.93
Education college or beyond	216 (90.0)	173 (90.6)	43 (87.8)	0.84
Smoking status				0.87
Never	131 (54.6)	102 (53.4)	29 (59.2)	
Ever	91 (37.9)	72 (46.6)	19 (38.8)	
Current	18 (7.5)	17 (8.9)	1 (2.0)	
Drinking status				0.92
Never	90 (37.5)	70 (36.7)	20 (40.8)	
Ever	49 (20.4)	42 (21.9)	7 (14.3)	
Current	101 (42.1)	79 (41.4)	22 (44.9)	
Physically active 2 days/week	195 (81.2)	162 (84.8)	33 (67.3)	0.33
SBP (mmHg)	127.3±14.8	127.5±15.3	126.8±13.1	0.92
DBP (mmHg)	7.05±8.7	75.3±8.8	73.7±8.2	0.71
Total cholesterol (mg/dL)	202.9±56.1	204.7±57.7	195.9±49.4	0.74
Triglycerides (mg/dL)	130.1±73.7	124.8±65.6	151.0±97.2	0.43
LDL (mg/dL)	117.5±41.2	118.5±41.9	113.5±38.4	0.80
HDL (mg/dL)	59.4±20.4	61.2±21.3	52.2±14.8	0.36
C-reactive protein (mg/dL)	3.1±4.3	3.1±3.9	3.1±5.1	0.97
Antihypertensive medication use	106 (44.1)	78 (40.8)	28 (57.1)	0.48
Lipid control medication use	88 (36.7)	68 (35.6)	20 (40.8)	0.82
Glycemic control medication use	22 (9.2)	22 (11.5)	0 (0.0)	0.42

Continuous variables were presented as mean \pm SD. P values (the comparison between overall and participants who completed TCMQ) were calculated using GLM model; categorical variables presented as n (%). P values were calculated using chi-square test or fisher's exact test. TCMQ, Traditional Chinese Medicine Questionnaire; PPCCT, Personalized Prevention of Colorectal Cancer Trial; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; HDL, high-density lipoproteins; LDL, low-density lipoproteins.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Comparison of distribution of different body mass index among the TCMQ constitution groups, results from PPCCT

TCM constitution types	BMI (kg/m ²)			P value
	<25 (n=40)	25–29.9 (n=62)	30 (n=89)	
Balanced	16	18	23	0.27
Qi-deficiency	3	8	15	
Yang-deficiency	1	4	4	
Yin-deficiency	2	1	9	
Phlegm-dampness	1	4	11	
Damp-heat	2	1	3	
Blood-stasis	8	15	10	
Qi-stagnation	2	3	7	
Inherited-special	5	8	7	

Categorical variables presented as n (%), P values were calculated using chi-square test. TCM, traditional Chinese medicine; TCMQ, Traditional Chinese Medicine Questionnaire; PPCCT, Personalized Prevention of Colorectal Cancer Trial.

Table 3
Comparison of distribution of different body mass index among the TCMQ constitution groups

TCM constitution types	BMI (kg/m ²)				P ₁	P ₂
	PPCCT		China (Li et al. 2017) (11)			
	<30 (n=102)	30 (n=89)	<30 (n=4,513)	30 (n=510)		
Balanced	34 (33.3)	23 (25.8)	1,776 (39.4)	224 (44.4)	<0.0001	<0.0001
Qi-deficiency	11 (10.8)	15 (16.8)	1,050 (23.3)	103 (20.2)		
Yang-deficiency	5 (4.9)	4 (4.5)	848 (18.8)	54 (10.6)		
Yin-deficiency	3 (2.9)	9 (10.1)	298 (6.6)	32 (6.3)		
Phlegm-dampness	5 (4.9)	11 (12.4)	247 (5.5)	55 (10.8)		
Damp-heat	3 (2.9)	3 (3.4)	107 (2.4)	17 (3.3)		
Blood-stasis	23 (22.6)	10 (11.2)	65 (1.4)	9 (1.8)		
Qi-stagnation	5 (4.9)	7 (7.9)	45 (1.0)	8 (1.6)		
Inherited-special	13 (12.8)	7 (7.9)	77 (1.7)	8 (1.6)		

Categorical variables presented as n (%). P₁: the comparison of US and Chinese population among BMI <30; P₂: the comparison of US and Chinese population among BMI ≥30. TCMQ, Traditional Chinese Medicine Questionnaire; PPCCT, Personalized Prevention of Colorectal Cancer Trial.