

## Original Article

# Evaluation of the characteristics of infection prevention and control programs and infection control committees in Brazilian hospitals: A countrywide cross-sectional study

Beatriz Arns MD<sup>1,2</sup> , Crepin Aziz Jose Oluwafoumi Agani MSc<sup>1</sup>, Guilherme Prates Sesin MSc<sup>1</sup> , Jaqueline Driemeyer C. Horvath PhD<sup>1</sup> , Débora Vacaro Fogazzi MSc<sup>1</sup>, Fernanda Kelly Romeiro Silva BSc<sup>1</sup>, Lauren Sezera Costa MSc<sup>1</sup>, Adriano Jose Pereira PhD<sup>3</sup>, Antônio Paulo Nassar Junior PhD<sup>3</sup> , Bruno Tomazini Cavalcanti PhD<sup>4</sup>, Camila Dietrich PhD<sup>5</sup>, Viviane Cordeiro Veiga PhD<sup>6</sup>, Daniela G.M. Catarino PhD<sup>7</sup>, Maysa Yukari Cheno RN<sup>7</sup>, Alexandre Biasi PhD<sup>4</sup> , Bianca Ramos Ferronato MD<sup>8</sup>, Bil Randerson Bassetti MSc<sup>9</sup> , Caio Cesar Ferreira Fernandes MD<sup>10</sup>, Caroline Deutschendorf MD<sup>11</sup> , Cintia Magalhães Carvalho Grion PhD<sup>12</sup>, Claudia Fernanda de Lacerda Vidal PhD<sup>13</sup>, Cláudio Dornas de Oliveira MSc<sup>14</sup>, Eliana Bernadete Caser PhD<sup>15</sup>, Emerson Boschi MD<sup>16</sup>, Everton Macêdo Silva PhD<sup>17</sup>, Felipe Dal Pizzol PhD<sup>18</sup> , Hugo Correa de Andrade Urbano MD<sup>19</sup>, Iany Silva RN<sup>20</sup>, Israel Silva Maia MSc<sup>4,21</sup>, Leila Rezegue de Moraes Rego MSc<sup>22</sup>, Luana Pontes Oliveira MD<sup>23</sup>, Maria Brandão Tavares PhD<sup>24</sup>, Marianna Dewey Andrade Dracoulakis PhD<sup>25</sup>, Marina Peres Bainy MSc<sup>26</sup>, Nicole Alberti Golin MD<sup>27</sup>, Pablo Oscar Tomba MD<sup>28</sup>, Pedro Martins Pereira Kurtz PhD<sup>29</sup>, Rafael Botelho Foernges MD<sup>30</sup>, Rejane Martins Prestes MD<sup>31</sup>, Rodrigo Morel Vieira de Melo PhD<sup>32</sup>, Rodrigo Reghini Da Silva MG<sup>33</sup>, Tatiana Gozzi Pancev Toledo MD<sup>34</sup>, Valéria Paes Lima PhD<sup>35</sup> , Vanildes de Fátima Fernandes RN<sup>36</sup>, Wilson José Lovato MD<sup>37</sup> and Alexandre Prehn Zavascki PhD<sup>2,38</sup> 

<sup>1</sup>Hospital Moinhos de Vento, Porto Alegre, Rio Grande do Sul, Brazil, <sup>2</sup>Infectious Diseases Service, Hospital Moinhos de Vento, Porto Alegre, Rio Grande do Sul, Brazil, <sup>3</sup>Hospital Israelita Albert Einstein, São Paulo, Brazil, <sup>4</sup>HCor Research Institute, São Paulo, Brazil, <sup>5</sup>Hospital Sírio-Libanês, São Paulo, Brazil, <sup>6</sup>Hospital Beneficência Portuguesa, São Paulo, Brazil, <sup>7</sup>Hospital Alemão Oswaldo Cruz, São Paulo, Brazil, <sup>8</sup>Hospital Erasto Gaertner, Curitiba, Paraná, Brazil, <sup>9</sup>Hospital Estadual Central, Vitória, Espírito Santo, Brazil, <sup>10</sup>Hospital Estadual Mário Covas, Santo André, São Paulo, Brazil, <sup>11</sup>Hospital de Clínicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil, <sup>12</sup>Hospital Universitário da Universidade Estadual de Londrina, Londrina, Paraná, Brazil, <sup>13</sup>Hospital das Clínicas da Universidade Federal de Pernambuco, Recife, Pernambuco, Brazil, <sup>14</sup>Santa Casa de Misericórdia de Belo Horizonte, Belo Horizonte, Minas Gerais, Brazil, <sup>15</sup>Hospital Unimed Vitória, Vitória, Espírito Santo, Brazil, <sup>16</sup>Hospital Geral de Caxias do Sul, Caxias do Sul, Rio Grande do Sul, Brazil, <sup>17</sup>Hospital Base do Distrito Federal, Brasília, Federal District, Brazil, <sup>18</sup>Hospital São José, Criciúma, Santa Catarina, Brazil, <sup>19</sup>Hospital Vila da Serra, Nova Lima, Minas Gerais, Brazil, <sup>20</sup>Santa Casa de Misericórdia de São João Del Rei, São João del Rei, Minas Gerais, Brazil, <sup>21</sup>Hospital Nereu Ramos, Florianópolis, Santa Catarina, Brazil, <sup>22</sup>Hospital Jean Bitar, Belém, Pará, Brazil, <sup>23</sup>Hospital Presidente Vargas, São Luís, Maranhão, Brazil, <sup>24</sup>Hospital do Subúrbio, Salvador, Bahia, Brazil, <sup>25</sup>Hospital da Bahia, Salvador, Bahia, Brazil, <sup>26</sup>Hospital Escola Universidade Federal de Pelotas, Pelotas, Rio Grande do Sul, Brazil, <sup>27</sup>Hospital Tacchini, Bento Gonçalves, Rio Grande do Sul, Brazil, <sup>28</sup>Hospital do Cancer Barretos, Unidade Jales, Jales, São Paulo, Brazil, <sup>29</sup>Instituto Estadual do Cérebro Paulo Niemeyer, Rio de Janeiro, Rio de Janeiro, Brazil, <sup>30</sup>Hospital Santa Cruz, Santa Cruz, Rio Grande do Sul, Brazil, <sup>31</sup>Hospital Universitário da Universidade Federal do Piauí, Teresina, Piauí, Brazil, <sup>32</sup>Hospital Ana Nery, Salvador, Bahia, Brazil, <sup>33</sup>Hospital A.C. Camargo Cancer Center, São Paulo, Brazil, <sup>34</sup>Hospital e Maternidade Brasil, Santo André, São Paulo, Brazil, <sup>35</sup>Hospital Universitário de Brasília, Brasília, Federal District, Brazil, <sup>36</sup>Hospital Santa Casa de Misericórdia de Passos, Passos, Minas Gerais, Brazil, <sup>37</sup>Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto, Ribeirão Preto, São Paulo, Brazil and <sup>38</sup>Department of Internal Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

## Abstract

**Objective:** Data are scarce regarding hospital infection control committees and compliance with infection prevention and control (IPC) recommendations in Brazil, a country of continental dimensions. We assessed the main characteristics of infection control committees (ICCs) on healthcare-associated infections (HAIs) in Brazilian hospitals.

**Author for correspondence:** Beatriz Arns, MD, Rua Ramiro Barcelos, 630, sala 815. Bairro Moinhos de Vento, Porto Alegre - RS, Brazil. E-mail: [beatriz.arns@hmv.org.br](mailto:beatriz.arns@hmv.org.br)

**Cite this article:** Arns B, Agani CAJO, Sesin GP, *et al.* Evaluation of the characteristics of infection prevention and control programs and infection control committees in Brazilian hospitals: A countrywide cross-sectional study. *Antimicrob Steward Healthc Epidemiol* 2023. doi: [10.1017/ash.2023.136](https://doi.org/10.1017/ash.2023.136)

**Methods:** This cross-sectional study was conducted in ICCs of public and private hospitals distributed across all Brazilian regions. Data were collected directly from the ICC staff by completing an online questionnaire and during on-site visits through face-to-face interviews.

**Results:** In total, 53 Brazilian hospitals were evaluated from October 2019 to December 2020. All hospitals had implemented the IPC core components in their programs. All centers had protocols for the prevention and control of ventilator-associated pneumonia as well as bloodstream, surgical site, and catheter-associated urinary tract infections. Most hospitals (80%) had no budget specifically allocated to the IPC program; 34% of the laundry staff had received specific IPC training; and only 7.5% of hospitals reported occupational infections in healthcare workers.

**Conclusions:** In this sample, most ICCs complied with the minimum requirements for IPC programs. The main limitation regarding ICCs was the lack of financial support. The findings of this survey support the development of strategic plans to improve IPCs in Brazilian hospitals.

(Received 8 February 2023; accepted 11 February 2023)

Healthcare-associated infections (HAIs) affect millions of patients worldwide annually and represent a major challenge to patient care within hospitals.<sup>1</sup> The World Health Organization (WHO) estimates that ~1.7 million patients are affected by HAIs in the United States and 4 million in Europe every year.<sup>2</sup> HAIs affect patient outcomes by increasing mortality, promoting antimicrobial resistance and increasing healthcare-associated costs.<sup>1,3</sup> The burden of HAI is even higher in low- and middle-income countries, representing a greater epidemiological problem than in high-income countries.<sup>4</sup> The pooled prevalence of HAI in resource-limited settings (~15.5 per 100 patients) is 2–3 times higher than the rates observed in Europe and the United States. These infections also determine a marked economic impact at the societal level.<sup>3,4</sup>

Infection prevention and control (IPC) programs are a key strategy in preventing HAIs and containing antimicrobial resistance in hospitals.<sup>1–3</sup> Adequate conditions for the development and implementation of IPC programs have been considered a key parameter of the quality of healthcare services.<sup>1–3,5</sup> The insufficiency of these conditions has been a challenge for low- and middle-income countries, resulting in the aforementioned consequences of high HAI rates.<sup>3,4</sup> Evidence-based guidance for national IPC programs is needed to support national and global capacity building for the reduction of HAI and antimicrobial resistance.<sup>1,3,5</sup>

In 2018, the WHO launched the IPC Assessment Framework (IPCAF) with 8 core components, with the main objective of providing an orientation to assess the situation of IPC at the individual healthcare facility level and to monitor the development and improvement of IPC activities over time.<sup>6</sup> The components are as follows: IPC program; IPC guidelines; IPC education and training; HAI surveillance; multimodal strategies for implementation of IPC interventions; monitoring and audit of IPC practices and feedback; workload, staffing, and bed occupancy; and built environment, materials, and equipment for IPC at the facility level.<sup>6</sup>

In the last decade, several studies have evaluated IPC programs worldwide, and virtually all from high-income countries.<sup>3,7–17</sup> The results of countrywide surveys of IPC programs are relevant nationally to drive public health policies and internationally to identify global inequalities in both quality and implementation processes of IPC programs among countries.<sup>18</sup> These surveys are also important because they contribute to continued awareness of the importance of IPC and alert participants and policy makers to review gaps and priorities for action, allowing them to set targets to improve national IPC programs.<sup>18</sup>

Brazil is a country of continental dimensions with a population of >215 million.<sup>19</sup> IPC program content and implementation are an attribution of infection control committees (ICCs), whose existence in Brazil has been mandatory by law since 1997,<sup>20</sup> whereas the

management of IPC programs at the federal level has been assigned to the Brazilian Health Regulatory Agency (ANVISA for short, in Portuguese).<sup>21</sup> Although some studies have evaluated IPC programs in Brazil, they have usually been restricted to a single city or state.<sup>11,22–24</sup> Owing to the high heterogeneity of the economic, social, and healthcare structure in Brazil, Padoveze *et al*<sup>25</sup> highlighted the fragility of IPC and ICC quality assessments in the country and the urgent need to overcome this limitation. Only 1 has study assessed the structures for prevention of HAI in Brazilian hospitals, and it was conducted between 2011 and 2013.<sup>11</sup> Since then, however, several global and national actions have been implemented to increase awareness of HAI and to promote improvements in IPC programs. In this study, we assessed IPC programs and ICCs from Brazil, including hospitals from all 5 regions of the country, after the implementation of national and international actions to increase awareness of HAI.

## Methods

### Study design and data collection

We conducted a cross-sectional study in Brazilian public and private hospitals from October 2019 to December 2020. The ICCs included in this study were those from hospitals participating in the Impact of Infections by Antimicrobial-Resistant Microorganisms in Patients Admitted to Adult Intensive Care Units in Brazil: Platform of Projects to Support the National Action Plan for the Prevention and Control of Antimicrobial Resistance (IMPACTO-MR) program. This multicenter, country-wide platform is supported by the Brazilian Ministry of Health and was created to develop studies related to HAIs and antimicrobial resistance.<sup>26</sup> The centers were selected by convenience sampling ensuring they encompass all 5 Brazilian federative regions.

The questionnaire used in this study, containing the IPC program elements and specific structural characteristics of the hospital, was developed and adapted from the ANVISA guidelines for completing the assessment of patient safety practices.<sup>27</sup> The questionnaire shares similarities with most of the variables in each WHO IPCAF core component (Supplementary Material), and for this reason, no scoring system was used.

The data collection process started with an online questionnaire sent to the professional in charge of the ICC in all institutions, to be answered using the REDCap platform. Once completed, a team of 2–3 investigators conducted an on-site visit to the participating institutions to check documents and validate the answers received online. The audit visit focused on the documents, processes, and structures that were reported on the questionnaire. In addition, we collected monthly epidemiological data through a standardized form sent to the ICC for information on adherence to the guidelines for HAI prevention.

### Statistical analysis

In the study, we used descriptive analysis. Categorical variables (yes or available) were presented as proportions, and continuous variables have been presented as means (SD) or medians (IQR) according to their distribution. Comparisons between Brazilian federative regions were performed using frequency analysis, cross tabulation, and  $\chi^2$  test for categorical variables and the Kruskal-Wallis test for continuous variables, assuming the statistical significance of  $P < .05$ . The analyses were performed in R version 4.0.2 software (R Foundation for Statistical Computing, Vienna, Austria).<sup>28</sup>

### Ethics

The research was conducted in accordance with the ethical and legal principles contained in Resolution No. 466/2012 of the Brazilian National Health Council and was approved by the Research Ethics Committee of Hospital Moinhos de Vento (coordinating center) and the hospitals that participate in the IMPACTO-MR program (no. 3.385.438 and CAAE no. 02139418.3.2003.5330) (Supplementary Material).

### Results


In total, 53 hospitals randomly distributed across the country were evaluated through their respective ICCs (Fig. 1). All included hospitals had the following features: an ICC, at least 6 intensive care unit (ICU) beds, and access to a microbiology laboratory that follows the guidelines and interpretative criteria for antimicrobial susceptibility testing of the European Committee on Antimicrobial Susceptibility Testing (EUCAST),<sup>29</sup> the Clinical and Laboratory Standards Institute (CLSI),<sup>30</sup> or the Brazilian Committee on Antimicrobial Susceptibility Testing (BrCAST, affiliated with EUCAST).<sup>31</sup>

All ICCs had as members doctors (100%), nurses (100%), nursing technicians (43.4%), pharmacists (32.1%), and administrators (43.4%) (Table 1). Among other health professional categories involved in IPC practices, there were differences in the compositions of IPC departments among the analyzed regions (Table 2).

The analysis of data collected via electronic form and on-site visit to the centers revealed that 52 (98%) hospitals had an IPC program (Table 1), 50 (96%) of which had a clearly defined objective for the program. The mean time of existence of ICC was 21.6 years ( $\pm 11.9$ ). We detected a statistically significant difference in the time of existence of the ICCs between the South and North regions ( $P = .012$ ), with no significant differences between the other regions in the adjusted analysis (Table 2). Only 9 hospitals had a budget allocated to the IPC program, mostly localization in the Southeast region. Only 2 of the 25 public hospitals with the IPC program had a budget allocated annually to the program (8%), whereas 7 (25%) of the 28 private hospitals had this budget.

The guidelines available from the IPC programs are described in Table 1. Among them, the guidelines for HAI prevention and hand hygiene were evaluated in more detail regarding their recommendations. All centers (100%) had either protocols or guidelines for the prevention and control of ventilator-associated pneumonia, bloodstream infection, surgical site infection, and catheter-associated urinary tract infection. Also, most centers had guidelines for medical waste management and hand hygiene (98.1% and 100%, respectively).

The surveillance indicators are also described in Table 1. All hospitals (100%) reported conducting nosocomial infection surveillance and epidemiological surveillance to detect outbreaks in



**Fig. 1.** Distribution of the 53 hospitals located in the 5 federative regions of Brazil. Hospitals in each region were distributed as follows: North (Acre, Amapá, Amazonas, Pará, Rondônia, Roraima, and Tocantins); Northeast (Maranhão, Piauí, Ceará, Rio Grande do Norte, Paraíba, Pernambuco, Alagoas, Sergipe, and Bahia); Midwest (Goiás, Mato Grosso, and Mato Grosso do Sul; along with Distrito Federal, the Federal District), Southeast (Espírito Santo, Minas Gerais, Rio de Janeiro, and São Paulo), and South (Paraná, Santa Catarina, and Rio Grande do Sul).

a timely manner. Within the HAI surveillance component, only 7.5% of hospitals reported occupational infections in healthcare workers. All hospitals had access to a microbiology laboratory to support HAI surveillance, and 38% of laboratory services were outsourced. Within the IPC education and training component, 88.7% of centers provided specific, systematic, and periodic IPC training for hospital staff. Adherence to specific training provided by the ICC was lower among the laundry staff (34.0%) than among other professionals (Table 1).

Within the monitoring and audit of IPC practices and feedback component, nearly all ICCs reported performing routine data collection on nosocomial infections, usually monthly, and using these indicators to evaluate services and plan future actions (100%). Antimicrobial use per patient day was the least-monitored indicator (22.6%) (Table 1).

Nearly all centers had an adequate structure and policy for cleaning, with regular assessment of the policies and structures that guide routine hand hygiene practices in the hospital environment. All hospitals (100%) had personal protective equipment for all professionals and specific containers for disposal of hospital waste, and 98.1% had routine bacteriological control of water (Table 1). There was no statistically significant difference between the Brazilian regions in most of the characteristics described.

### Discussion

To support countries in their efforts to strengthen IPC, the WHO released evidence-based guidelines addressing the key components of IPC programs.<sup>2,32</sup> These guidelines cover 8 core components of IPC and include recommendations and best practice statements. Some publications provide an evaluation of ICCs using these core components. In Georgia, for example, 41 hospitals underperformed in the implementation of WHO recommendations, especially those for IPC practices.<sup>15</sup>

In view of the various problems that occurred during the COVID-19 pandemic and the health-system collapse in Brazil and worldwide, the existence of an ICC proved to be indispensable. ICCs were particularly useful in rebuilding care pathways and protocols for IPC, enabling immediate and appropriate decisions to

**Table 1.** Indicators to Evaluate Infection Prevention and Control programs in Brazilian Hospitals<sup>a</sup>

<b>Component 1: IPC program</b>	No. (%) or Mean (SD)
IPC program has been implemented	52 (98.1)
Regular meetings with the ICC	51 (96.2)
Records of the minutes of ICC meetings	53 (100)
Medical doctor (member of ICC team)	53 (100)
Nurse (member of ICC team)	53 (100)
Nursing technician (member of ICC team)	23 (43.4)
Pharmacist (member of ICC team)	17 (32.1)
Administrator (member of ICC team)	23 (43.4)
Other professionals (members of ICC team)	23 (43.4)
<b>Component 2: IPC guidelines</b>	
Standard precautions	53 (100)
Antimicrobial control (stewardship)	42 (79.2)
Prevention of surgical site infections	53 (100)
Prevention of bloodstream infections	53 (100)
Prevention of ventilator-associated pneumonia	53 (100)
Prevention of catheter-associated urinary tract infections	53 (100)
Prevention of transmission of multidrug-resistant (MDR) pathogens	50 (94.3)
Basic guideline written by ICC for cleaning services	44 (83.0)
Basic guideline written by ICC for laundry services	26 (49.1)
Disinfection, surface cleaning, and sterilization	52 (98.1)
Hand hygiene	53 (100)
Waste management	52 (98.1)
<b>Component 3: IPC education and training</b>	
ICC establishes continuing education measures for the medical team in relation to the prescription of antimicrobials	37 (69.8)
ICC establishes continuing education measures for the medical team in relation to the prescription of antimicrobials	37 (69.8)
ICC promotes debates with the hospital community about IPC	48 (90.6)
Orientation program for professionals entering the institution	48 (90.6)
Specific, systematic and periodic IPC training for hospital staff	47 (88.7)
<b>Component 3: IPC education and training</b>	
ICC establishes continuing education measures for the medical team in relation to the prescription of antimicrobials	37 (69.8)
<b>ICC promotes debates with the hospital community about IPC</b>	48 (90.6)
Orientation program for professionals entering the institution	48 (90.6)
Specific, systematic and periodic IPC training for hospital staff	47 (88.7)
Periodicity of IPC training for hospital health professionals	16 (30.2)
Monthly	14 (26.4)
Semiannually	13 (24.5)
Annually	11 (20.8)
Other	
Specific training for hygiene professionals by the ICC	41 (77.4)
Specific training for laundry professionals by the ICC	18 (34.0)
<b>Component 4: HAI surveillance</b>	
Surveillance conducted for infections	53 (100)
Epidemiological surveillance to detect outbreaks in a timely manner	53 (100)
Active prospective surveillance	53 (100)
Passive prospective surveillance (notification form/medical)	36 (67.8)
Catheter-associated bloodstream infections	53 (100)

(Continued)

**Table 1.** (Continued)

Catheter-associated urinary tract infections	53 (100)
Skin and soft tissue infections	34 (64.2)
Pneumonia	53 (100)
Reported occupational infections in healthcare workers	4 (7.5)
<b>Component 5: Multimodal strategies</b>	
ICC participates in the technical committee for specification of products and related products to be acquired	46 (86.8)
Own microbiology laboratory	33 (62.3)
<b>Component 6: Monitoring/audit of IPC practices and feedback</b>	
Monitoring the rate of healthcare-associated infections	53 (100)
Monitoring the case fatality rate of healthcare-associated infections	32 (60.4)
Monitoring antimicrobial use per patient day	12 (22.6)
Monitoring the coefficient of sensitivity or resistance of microorganisms to antimicrobials	38 (71.7)
Monitoring the records of puncture-cutting accidents in employees	41 (77.4)
ICC reports informational data on IPC indicators	53 (100)
ICC shares the hospital infection control indicators with the hospital management	52 (98.1)
<b>Component 7: Weekly workload of ICC team, median h (IQR)</b>	
Medical doctor	20 (20–40)
Nurse	44.5 (40–117)
Pharmacist	40 (36–44)
Administrator	40 (35–44)
<b>Component 8: Built environment, materials, and equipment for IPC at the facility level</b>	
Own cleaning service	29 (54.7)
Own laundry service	18 (34.0)
All hospital units or wards have washbasins with running water, soap or antiseptic and paper towels for hand hygiene	52 (98.1)
Personal protective equipment available for all professionals	53 (100)
Routine bacteriological control of the water that supplies the hospital	52 (98.1)
Routine cleaning of the water tank that supplies the hospital	53 (100)
Routine control of air quality management	39 (73.6)
Specific containers for disposal of hospital waste	53 (100)

Note. IPC, infection prevention and control; ICC, infection control committee.

\*This table shows the number of Brazilian hospitals that meet some indicators in the ICC assessment. The indicators are grouped by components. Data were available from 53 hospitals.

control the spread of the pathogen and to protect health professionals.<sup>33</sup>

A well-structured IPC program can reduce infection rates, and some recommendations result from the economic and social burden generated by these events. HAI is preventable and can be reduced by up to 70% through effective IPC measures.<sup>34</sup> Although the implementation of the ICC was not supported in all assessed hospitals,<sup>24</sup> most hospitals evaluated in this study had an IPC program. Our data show that all types of infections occurring in Brazilian hospitals are generally reported by the ICC. This reporting practice has a major impact on public health by contributing significantly to infection control. Therefore, simple IPC measures and policies frequently implemented by the ICC, such as hand hygiene, are initial steps for the future success of the IPC program and the effective implementation of IPC practices to prevent the transmission of pathogens.<sup>35</sup>

The ICCs evaluated in our study reported low infection rates in employees. Underreporting of infection rates among health

professionals may not only contribute to the failure of IPC actions but also present a threat to the functioning of hospitals due to the lack of preparedness of professionals.<sup>36</sup> This assessment proved to be very important during the COVID-19 pandemic, and several studies have highlighted, among other indicators, the infection rate in health professionals as a critical factor for facility level improvement.<sup>37,38</sup>

Based on the results of our survey, the major obstacle seems to be the allocation of a dedicated budget for IPC implementation. Only 17% of hospitals had a budget specifically allocated to the IPC program. A dedicated budget for implementing IPC strategies and plans is a minimum requirement according to the WHO.<sup>32</sup> This gap was more pronounced in public hospitals; most of the hospitals that had a dedicated IPC budget were private. The ICC needs economic and administrative support to implement the IPC program, otherwise it might be inefficient and yield unsatisfactory results. The simultaneous availability of qualified personnel, infection control policies, and adequate structures



**Table 2.** Characteristics of Hospital Infection Prevention and Control Programs in Brazilian Hospitals, for Federative Regions

Characteristic	Brazilian Federative Regions, No. (%) or mean ( $\pm$ SD)				
	South	Southeast	Midwest	North	Northeast
<b>Hospital management</b>					
Public	5 (45.0)	8 (36.3)	4 (100)	2 (50.0)	6 (50.0)
Private	6 (55.0)	14 (63.7)	0 (0.0)	2 (50.0)	6 (50.0)
<b>Existence of infection control committees, mean years</b>					
	31 ( $\pm$ 15)	21 ( $\pm$ 8)	22 ( $\pm$ 11)	9 ( $\pm$ 6)	18 ( $\pm$ 12)
<b>Budget specifically to the IPC program</b>					
	1 (11.1)	5 (21.7)	0 (0.0)	1 (25.0)	2 (16.7)
<b>Internal audits to assess compliance with the IPC program</b>					
	7 (77.8)	22 (100)	4 (100)	3 (75.0)	7 (58.3)
<b>Consortium with other hospitals for the reciprocal use of technical, material, and human resources in the implementation of the IPC program</b>					
	2 (22.2)	3 (13.0)	1 (25.0)	0 (0.0)	2 (16.7)
<b>Professionals that compose the IPC department</b>					
Medical doctor	11 (100)	22 (100)	4 (100)	4 (100)	12 (100)
Nurse	11 (100)	22 (100)	4 (100)	4 (100)	12 (100)
Nursing technician	9 (81.8)	4 (17.4)	1 (25.0)	3 (75.0)	6 (50.0)
Pharmaceutical	7 (63.6)	4 (17.4)	0	0	6 (50.0)
Administrator	3 (27.3)	6 (26.1)	0	0	5 (41.7)
Interns	4 (36.7)	4 (17.4)	0	1 (25.0)	0 (0.0)

Note. IPC, infection prevention and control. Data available from 53 hospitals.

(equipment and materials) is crucial for the implementation of optimal IPC measures. Data from the WHO global report on IPC, with surveys conducted in 62 countries (including Brazil), show a significant increase in the proportion of countries having a dedicated budget for IPC programs between 2017–2018 (25.8%) and 2021–2022 (48.4%;  $P = .02$ ).<sup>39</sup> However, consistent with the present data, there is still considerable potential for improvement. An Australian study showed that implementing a budget for a cleaning bundle (audit, communication, technique, training, and product) returned a positive net monetary benefit and incremental cost-effectiveness ratio.<sup>40</sup>

Regarding IPC education and training of staff, most ICCs offered IPC training to employees and staff members, but not all human resources had the same level of training in the different areas of the facility. Despite the good connection with the cleaning staff, which reported appropriate training, adherence to specific training provided by the ICC was low (34%) among the laundry staff in most hospitals. Specific training for all health professionals is crucial for the success of the IPC program. Therefore, training should not be limited to frontline health workers or IPC specialists but rather encompass all staff regardless of position or function, as providing technical and specific training for the entire hospital personnel is essential to prevent and reduce HAI.<sup>17</sup>

Previous studies conducted in Brazil have pointed out deficiencies in HAI prevention and control practices in outsourced services, such as laundry and laboratory.<sup>23,41</sup> Outsourcing microbiology services, as reported by many centers in the present study, may create obstacles to the actions of the ICC, which depends on the turnover of microbiological results for timely IPC actions.<sup>11,42</sup>

A strength of our study is the analysis of IPC programs from all 5 Brazilian regions, which provided a broad view of the country's reality that had not been possible in most previous publications. Limitations of the study include the eligibility criteria for inclusion of hospitals in the IMPACTO-MR program, based on which a sample of hospitals with better conditions was selected, which may have led to improved indicators in our survey. In addition, we modified the assessment form designed by the WHO and other national forms used in Brazilian studies to measure adherence to recommendations for ICC and IPC programs. For this reason, no scoring system was used to evaluate the hospitals or to measure and compare their respective ICCs.

The results of our analysis of IPC programs and ICCs in Brazilian hospitals showed great adherence to IPC guidelines and recommendations, particularly those on hand hygiene, sterilization, and catheter use. The profile of the ICCs of Brazilian hospitals described in the present study indicates the implementation of a range of strategies and policies to combat antimicrobial resistance and prevent HAI. These data are crucial for the establishment of new goals aiming to improve the evaluation of IPC indicators and, consequently, to reduce the rate of infections due to multi-drug-resistant organisms in the country. We were able to identify relevant aspects of IPC that may help guide public policies aiming to improve the quality of healthcare services in low- and middle-income countries.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/ash.2023.136>

**Acknowledgments.**

**Financial support.** This research was funded by the Brazilian Ministry of Health through the Institutional Development Support Program of the Brazilian Unified Health System (PROADI-SUS). The funding body has no role in coordinating the study.

**Conflict of interest.** A.P.Z. is a research fellow of the National Council for Scientific and Technological Development, Ministry of Science and Technology, Brazil. A.P.Z. received a research grant from Pfizer and was a member of the advisory board for Spero Therapeutics and Eurofarma. All other authors have no competing interests to declare.

## References

- Pittet D, Donaldson L. Clean Care is Safer Care: the first global challenge of the WHO World Alliance for Patient Safety. *Infect Control Hosp Epidemiol* 2005;26:891–4.
- Guidelines on core components of infection prevention and control programmes at the national and acute health care facility level. World Health Organization website. <https://www.who.int/publications-detail-redirect/9789241549929>. Published 2016. Accessed August 23, 2022.
- Price L, MacDonald J, Melone L, et al. Effectiveness of national and subnational infection prevention and control interventions in high-income and upper-middle-income countries: a systematic review. *Lancet Infect Dis* 2018;18:e159–e171.
- Allegranzi B, Nejad SB, Combescure C, et al. Burden of endemic healthcare-associated infection in developing countries: systematic review and meta-analysis. *Lancet* 2011;377:228–241.
- Storr J, Twyman A, Zingg W, et al. Core components for effective infection prevention and control programmes: new WHO evidence-based recommendations. *Antimicrob Resist Infect Control* 2017;6:6.
- Infection prevention and control assessment framework at the facility level. World Health Organization website. <https://www.who.int/publications-detail-redirect/WHO-HIS-SDS-2018>. Published 2018. Accessed August 23, 2022.
- Stone PW, Pogorzelska-Maziarz M, Herzig CTA, et al. State of infection prevention in US hospitals enrolled in the National Health and Safety Network. *Am J Infect Control* 2014 ;42:94–99.
- Oh HS, Uhm DC. Current status of infection prevention and control programs for emergency medical personnel in the Republic of Korea. *J Prev Med Public Health Yebang Uihakhoe Chi* 2015;48:330–341.
- Hansen S, Zingg W, Ahmad R, et al. Organization of infection control in European hospitals. *J Hosp Infect* 2015;91:338–345.
- Yoon YK, Lee SE, Seo BS, et al. Current status of personnel and infrastructure resources for infection prevention and control programs in the Republic of Korea: a national survey. *Am J Infect Control* 2016;44:e189–e193.
- Padoveze MC, Fortaleza CMCB, Kiffer C, et al. Structure for prevention of health care-associated infections in Brazilian hospitals: a countrywide study. *Am J Infect Control* 2016;44:74–79.
- Hansen S, Schwab F, Zingg W, Gastmeier P, Group the P study. Process and outcome indicators for infection control and prevention in European acute-care hospitals in 2011 to 2012: results of the PROHIBIT study. *Eurosurveillance* 2018;23:1700513.
- Aghdassi SJS, Hansen S, Bischoff P, Behnke M, Gastmeier P. A national survey on the implementation of key infection prevention and control structures in German hospitals: results from 736 hospitals conducting the WHO Infection Prevention and Control Assessment Framework (IPCAF). *Antimicrob Resist Infect Control* 2019;8:73.
- Aghdassi SJS, Grisold A, Wechsler-Fördös A, et al. Evaluating infection prevention and control programs in Austrian acute-care hospitals using the WHO Infection Prevention and Control Assessment Framework. *Antimicrob Resist Infect Control* 2020;9:92.
- Deryabina A, Lyman M, Yee D, et al. Core components of infection prevention and control programs at the facility level in Georgia: key challenges and opportunities. *Antimicrob Resist Infect Control* 2021;10:39.
- Ni K, Jin D, Wu Z, Sun L, Lu Q. The status of infection prevention and control structures in Eastern China based on the IPCAF tool of the World Health Organization. *Antimicrob Resist Infect Control* 2022;11:46.
- Jeong Y, Joo H, Bahk H, Koo H, Lee H, Kim K. A nationwide survey on the implementation of infection prevention and control components in 1,442 hospitals in the Republic of Korea: comparison to the WHO Infection Prevention and Control Assessment Framework (IPCAF). *Antimicrob Resist Infect Control* 2022;11:71.
- Tartari E, Tomczyk S, Pires D, et al. Implementation of the infection prevention and control core components at the national level: a global situational analysis. *J Hosp Infect* 2021;108:94–103.
- Projeção da população do Brasil e das Unidades da Federação. Instituto Brasileiro de Geografia e Estatística-IBGE website. [https://www.ibge.gov.br/apps/populacao/projecao/index.html?utm\\_source=portal&utm\\_medium=popclock&utm\\_campaign=novo\\_popclock](https://www.ibge.gov.br/apps/populacao/projecao/index.html?utm_source=portal&utm_medium=popclock&utm_campaign=novo_popclock). Published 2022. Accessed August 23, 2022.
- Portaria no. 2.616, de 12 de maio de 1998. Dispõe sobre diretrizes e normas para a prevenção e o controle das infecções hospitalares. Diário Oficial da União, Brasília, 13 maio 1998. Brasil Ministério da Saúde website. [https://bvsms.saude.gov.br/bvs/saudelegis/gm/1998/prt2616\\_12\\_05\\_1998.html](https://bvsms.saude.gov.br/bvs/saudelegis/gm/1998/prt2616_12_05_1998.html). Published 1998. Accessed September 1, 2022.
- Programa Nacional de Controle de Infecção Hospitalar. Anvisa Tecnologia em Serviços de Saúde website. <https://www.anvisa.gov.br/servicosade/infec.htm>. Published 2003. Accessed August 23, 2022.
- Meneguetti MG, Canini SRM da S, Bellissimo-Rodrigues F, Laus AM. Evaluación de los Programas de Control de Infección Hospitalaria en servicios de salud. *Rev Lat Am Enfermagem* 2015;23:98–105.
- Alves DCI, Lacerda RA. Evaluation of programs of infection control related to healthcare assistance in hospitals. *Rev Esc Enferm USP* 2015;49:65–73.
- Santos PLC, Padoveze MC, Lacerda RA. Performance of infection prevention and control programs in small hospitals. *Rev Esc Enferm USP* 2020. doi: 10.1590/S1980-220X2019002103617.
- Padoveze MC, Fortaleza CMCB. Healthcare-associated infections: challenges to public health in Brazil. *Rev Saúde Pública* 2014;48:995–1001.
- Página 118 da Seção 3 do Diário Oficial da União (DOU) de 16 de Abril de 2021. Jusbrasil website. <https://www.jusbrasil.com.br/diarios/1121922021/dou-secao-3-16-04-2021-pg-118>. Published 2021. Accessed August 23, 2022.
- Orientações-Avaliação-Práticas-Seg. Paciente-2022-HOSPITAIS COM UTI\_14.04.2022.pdf. Agência Nacional de Vigilância Sanitária (ANVISA) website. [https://www.gov.br/anvisa/pt-br/assuntos/servicosdesaude/seguranca-do-paciente/orientacoes-avaliacao-praticas-seg-paciente-2022-hospitais-com-uti\\_14-04-2022.pdf/view](https://www.gov.br/anvisa/pt-br/assuntos/servicosdesaude/seguranca-do-paciente/orientacoes-avaliacao-praticas-seg-paciente-2022-hospitais-com-uti_14-04-2022.pdf/view). Published 2022. Accessed August 23, 2022.
- R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, version 4.0.2. Vienna, Austria. <https://www.R-project.org/>. Published 2022. Accessed August 23, 2022.
- Clinical breakpoints and dosing of antibiotics. EUCAST website. [https://www.eucast.org/clinical\\_breakpoints/](https://www.eucast.org/clinical_breakpoints/). Published 2022. Accessed August 23, 2022.
- CLSI guidelines. Clinical & Laboratory Standards Institute website. <https://clsi.org/>. Published 2022. Accessed August 23, 2022.
- BR Cast. Brazilian Committee on Antimicrobial Susceptibility Testing website. <http://brcast.org.br/documentos/>. Published 2022. Accessed August 23, 2022.
- Minimum requirements for infection prevention and control programmes. World Health Organization website. <https://www.who.int/publications-detail-redirect/9789241516945>. Published 2019. Accessed August 23, 2022.
- Oliveira EC da S, Silva FP da, Pereira EBF e, Oliveira RC de. Actions of the hospital infection control committee in front of the new coronavirus. *Rev Baiana Enferm* 2020;e37259–e37259.
- Umscheid CA, Mitchell MD, Doshi JA, Agarwal R, Williams K, Brennan PJ. Estimating the proportion of healthcare-associated infections that are reasonably preventable and the related mortality and costs. *Infect Control Hosp Epidemiol* 2011;32:101–114.
- Alhumaid S, Al Mutair A, Al Alawi Z, et al. Knowledge of infection prevention and control among healthcare workers and factors influencing compliance: a systematic review. *Antimicrob Resist Infect Control* 2021;10:86.
- Atnafie SA, Anteneh DA, Yimenu DK, Kifle ZD. Assessment of exposure risks to COVID-19 among frontline health care workers in Amhara Region, Ethiopia: a cross-sectional survey. *PLoS One* 2021;16:e0251000.

37. Bandyopadhyay S, Baticulon RE, Kadhum M, *et al*. Infection and mortality of healthcare workers worldwide from COVID-19: a systematic review. *BMJ Glob Health* 2020;5:e003097.
38. Albuquerque M de FPM de, Souza WV de, Montarroyos UR, *et al*. Risk of SARS-CoV-2 infection among frontline healthcare workers in Northeast Brazil: a respondent-driven sampling approach. *BMJ Open* 2022;12:e058369.
39. Global report on infection prevention and control. World Health Organization website. <https://www.who.int/publications-detail-redirect/9789240051164>. Published 2022. Accessed August 23, 2022.
40. White NM, Barnett AG, Hall L, *et al*. Cost-effectiveness of an environmental cleaning bundle for reducing healthcare-associated infections. *Clin Infect Dis* 2020;70:2461–2468.
41. Meneguetti MG, Canini SRM da S, Bellissimo-Rodrigues F, Laus AM. Evaluation of nosocomial infection control programs in health services. *Rev Lat Am Enfermagem* 2015;23:98–105.
42. Costa LB, Cardoso MRA, Ferreira CG, *et al*. National prevalence survey in Brazil to evaluate the quality of microbiology laboratories: the importance of defining priorities to allocate limited resources. *Rev Panam Salud Pública* 2013;33:73–78.