

RESEARCH ARTICLE

Factors associated with postmenstrual age at full oral feeding in very preterm infants

Grégoire Brun¹, Céline J. Fischer Fumeaux², Eric Giannoni², Myriam Bickle Graz^{3*}

1 Emergency Care, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland, **2** Clinic of Neonatology, Department Mother-Woman-Child, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland, **3** Follow-up Unit, Department Mother-Woman-Child, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

* Myriam.Bickle-Graz@chuv.ch

Abstract

Aim

We aimed to identify variables associated with gestational age at full oral feeding in a cohort of very preterm infants.

Methods

In this retrospective study, all infants born below 32 weeks of gestation and admitted to a level III neonatal unit in 2015 were included. We dichotomized our population of 122 infants through the median age at full oral feeding, and explored which variables were statistically different between the two groups. We then used linear regression analysis to study the association between variables known from the literature and variables we had identified and age at full oral feeding.

Results

The median postnatal age at full oral feeding was 36 6/7 weeks post menstrual age (Q1-Q3 35 6/7-392/7), and was associated with the duration of hospital of stay. In the univariable linear regression, the variables significantly associated with full oral feeding were gestational age, socioeconomic status, sepsis, patent ductus arteriosus, duration of supplementary oxygen, of non-invasive and invasive ventilation, and bronchopulmonary dysplasia. In the multivariable regression analysis, duration of non-invasive ventilation and oxygen therapy, bronchopulmonary dysplasia, and patent ductus arteriosus were associated with an older age at full oral feeding, with bronchopulmonary dysplasia the single most potent predictor.

Discussion

Lung disease severity is a major determinant of age at full oral feeding and thus length of stay in this population. Other factors associated with FOF include socioeconomic status and patent ductus arteriosus. There is a need for research addressing evidence-based bundles of care for these infants at risk of long-lasting feeding and neurodevelopmental impairments.

OPEN ACCESS

Citation: Brun G, Fischer Fumeaux CJ, Giannoni E, Bickle Graz M (2020) Factors associated with postmenstrual age at full oral feeding in very preterm infants. *PLoS ONE* 15(11): e0241769. <https://doi.org/10.1371/journal.pone.0241769>

Editor: Umberto Simeoni, Centre Hospitalier Universitaire Vaudois, FRANCE

Received: April 14, 2020

Accepted: October 21, 2020

Published: November 11, 2020

Copyright: © 2020 Brun et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its [Supporting Information](#) files.

Funding: Yes, The Leenaards Foundation supports E.G.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: BPD, Bronchopulmonary dysplasia; CPAP, continuous positive airway pressure; FOF, Full oral feeding; PMA, post menstrual age; SGA, small for gestational age.

Introduction

Worldwide, 10% of children are born preterm [1] and are at risk of lifelong disabilities, including neurodevelopmental impairments. Brain growth and maturation is of paramount importance during these last weeks of gestation, and depends crucially on adequate nutrition [2]. Very preterm neonates generally require parenteral nutrition, while enteral feeding is provided in increasing doses. Once full enteral feeding is achieved, the transition period from tube to full oral feeds (FOF) is a major determinant of the length of stay in the neonatal unit [3] and depends notably on the maturation and efficiency of the coordination of sucking-swallowing and breathing. Prolonged tube feeding is associated with long-term feeding difficulties [4], long-term oral sensitivity and speech problems [5]. Finally, feeding difficulties may have a negative impact on nutrition in the neonatal period and thereafter, which may lead to stunted growth, altered neurodevelopment and lower academic achievement, and potentially adult onset metabolic disease [6]. Among the many complications of preterm birth, feeding and nutritional issues thus carry a heavy burden in terms of financial and social cost [7].

Different factors are known to be associated with age at tube weaning, such as gestational age, need for respiratory support and bronchopulmonary dysplasia (BPD) [8], as well as parental involvement [3]. The association of socio-economic status with several neurodevelopmental outcomes in preterm children has been demonstrated [9], but the association with feeding is underreported [10, 11]. Moreover, studies of oral motor interventions to enhance feeding skills, such as non-nutritive sucking and oral stimulation, do not show unanimous results [12–15], which could be explained by the heterogeneity of the populations and of the interventions.

The aim of this study was thus to identify and characterize, in infants born before 32 weeks of gestation, the variables associated with postmenstrual age (PMA) at full oral feeding.

Patients and methods

The "Commission cantonale d'éthique de la recherche sur l'être humain" approved the study protocol, and all patients gave written consent.

This was a retrospective cohort study of infants born before 32 weeks of gestation and admitted to a level III neonatal unit admitting 800 infants per year, as the reference center for a region counting 15 000 births/year [16]. All infants born between 01.01.2015 and 31.12.2015 at a gestational age less than 32 weeks were considered for this study. Parents were asked for written informed consent for the use of neonatal and follow-up data, according to the Swiss regulation and local ethics committee. Exclusion criteria were consent refusal, death before full oral feeding, and major congenital abnormalities.

Feeding practice

In the setting of our neonatal unit in 2015, enteral feeds were started on the first day of life with mother's own milk when available, fresh or frozen, or preterm formula (BEBA Alprem or BEBA Aliment pour Prématurés Etape 1; Nestlé, Vevey, Switzerland). The milk was administered every 2 hours for preterm infants weighing < 1500g and every 3 hours for those weighing \geq 1500g, with a targeted average daily volume increase of 10–20 ml/kg/day, adapted according to enteral tolerance until 160 ml/kg/day. Breastfeeding was encouraged as early as possible, with no lower age limit, whereas bottle-feeding was started from 34 weeks PMA. Donor milk was not available. A standard fortifier (Aptamil Frauen-Milch-Supplement 4%; Milupa SA, Domdidier, Switzerland) was introduced when 100 mL/kg/day of human milk was tolerated for all preterm infants < 32 weeks of GA. To assess feeding tolerance, gastric residuals were checked before every feed as a part of enteral feeding indicator; when the volumes of aspirates were superior to volumes of feeds, and/or in the presence of severe abdominal

distension, it was possible to suspend enteral feeding for a couple of hours until resolution of symptoms.

Probiotics were not administered in 2015, due to concerns regarding safety of available products [17]. Oral stimulation and non-nutritive sucking were systematically offered to all preterm infants as part of general developmental care, following the procedures described in Pfister et al [18], which include the use of pacifiers, exposure to the smell and taste of milk, massage, and stimulation of the rooting reflex. Parental presence and skin-to-skin practice were also strongly encouraged, but these interventions were not monitored. The feeding tube was removed when the infant was able to breast or bottle-feed more than 60% of their total milk intake.

Variables

Neonatal physiological and treatment variables were prospectively recorded in a widely available computerized patient Clinical Information System (Metavision® iMDsoft, Massachusetts, USA) [19, 20], which allows the extraction of selected data. Social and diagnostic variables were prospectively recorded in a specific *ad hoc* database. The selection of independent variables for this study was based on the literature and on an exploratory analysis of our population. Variables used for the study were thus: gestational age, assessed with early first trimester ultrasound (before 14 weeks) and last menstrual period, gender, birthweight and birthweight z-score [21], small for gestational age (SGA) defined as birthweight z-score < 2, mode of delivery. Socioeconomic status was assessed with the score described by Largo [22], which entails mother's education (scored from 1 to 6) and father's occupation (scored from 1 to 6), which is used in neonatal research in Switzerland [23] (total score from 2 to 12, 2 corresponding to higher parental education or occupation). Blood culture proven sepsis [24], necrotising enterocolitis (Bell stage ≥ 2), major brain injury such as cystic leukomalacia or grade III-IV intraventricular haemorrhage according to Papile [25], In 2015, cardiac echocardiography to detect PDA was made on the base of suggestive clinical signs. Hemodynamically significant PDA was treated medically first (indomethacin) if there were no contraindications, surgically in the case of a failure of the medical treatment or in the presence of contraindications. Variables were thus treated PDA, including both medically and surgically treated PDA, as usually reported, and surgically treated PDA.

To assess the impact of lung disease, we used the diagnoses of bronchopulmonary dysplasia, defined as a requirement for more than 28 days of supplemental oxygen between birth and 36 weeks PMA [26], duration of invasive ventilation, of continuous positive airway pressure (CPAP), and supplementary oxygen corresponding to the total number of days for which the patient received more than 21% oxygen, whatever the mode of administration (low or high flow nasal cannula, non-invasive or invasive ventilation). Nasal CPAP using the Medijet® generator [27] was the primary mode of respiratory support in spontaneously breathing infants. Respiratory distress syndrome was managed according to the most recent guidelines [28], to ensure continuity of care between the different attending physicians. Finally studied outcomes were post menstrual age (PMA) at enteral and oral feeding milestones, type of milk at discharge (any or no mother's own milk), and length of stay.

Statistical analysis

Analyses were performed with STATA® 13 (StataCorp. 2013. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP). Continuous variables were described with median and first to third quartiles, categorical variables with proportions. As a preliminary exploration, we divided our cohort in 2 subgroups on the basis of the median PMA at FOF




Fig 1. Flowchart of the study population.

<https://doi.org/10.1371/journal.pone.0241769.g001>

(early PMA at FOF and late PMA). We tested the difference between the 2 groups for the above mentioned variables, with t-tests or Kruskal-Wallis test for continuous variables and chi2 test for categorical variables.

For the linear regression analysis, we chose variables which were significantly different in our preliminary exploration, as well as variables known from the literature to be associated with age at FOF. The association of the variables with PMA at FOF was analysed with unilinear regression first. Secondly, we used multilinear regression including variables which were significant at the $p < 0.2$ level in the univariable analysis.

Results

The study population consisted of 122 patients (64 girls, 52%), with a median gestational age of 29.7 weeks (Q1-Q3 28–30.8, range 23 6/7–31 6/7) and a median birthweight of 1138 g (Q1-Q3 900–1425, range 510–1930g) (Fig 1), 88.5% of whom had received any antenatal steroids.

Patient characteristics

Enteral feeding was initiated on the first day of life for most patients ($n = 112$, 9%). Weaning of the feeding tube occurred after a median of 55 days (Q1-Q3 38–74), at a median of 36 6/7 weeks PMA (Q1-Q3 35 6/7–39 2/7) with a wide range between 34 and 141 weeks. Patients were discharged home at a median postmenstrual age of 37 4/7 (Q1-Q3 36 3/7–40 3/7, range 34 4/7–109 weeks PMA). The age at discharge home was associated with age at FOF (coef. (IC95%) 4.06 (3.45; 4.67), $p < 0.001$). At discharge home 70% of infants were fed with some mother's milk, and 30% with formula only.

The main patient characteristics are described in Table 1, with the characteristics of the groups of early (\leq median FOF) and late age at FOF ($>$ median FOF).

Table 1. Population characteristics.

	All	Feeding tube weaned	Feeding tube weaned	P
	n = 122	≤ 36 6/7 weeks PMA n = 60	> 36 6/7 weeks PMA n = 62	
Socioeconomic status (median, Q1-Q3)	6 (4–7)	6 (4–7)	6 (4–7)	NS
Antenatal steroids (n, %)	108 (88.5)	54 (90)	54(87)	NS
Gestational age, weeks (median, Q1-Q3)	29.7 (28; 30.8)	30.4 (29.7; 31.4)	28.4 (27.4; 29.7)	0.001
Gender, female (n, %)	64 (52.4)	27 (45)	37 (59.7)	0.047
Birthweight (g, median, Q1-Q3)	1138 (900;1425)	1350 (1089; 1580)	1010 (770; 1155)	0.001
Birthweight z-score (mean, SD)	-0.38 (0.72)	-0.12 (0.64)	-0.55 (0.76)	0.007
SGA (z-score < -2DS, %)	3 (2.5)	0	3 (4.8)	NS
Caesarean section (%)	98 (80)	41 (68)	57 (92)	0.001
Blood culture-proven sepsis (%)	13 (10)	0 (0)	13 (21)	0.001
Necrotizing enterocolitis (%)	4 (3)	0 (0)	4 (6)	0.001
Treated patent ductus arteriosus (n, %)	22 (18)	6 (10)	16 (25.8)	0.011
Surgical PDA	6 (5)	1(1.7)	5(8.0)	NS
Major brain lesions ¹ (n, %)	10 (8.2)	2 (3.3)	8 (12.9)	0.048
Supplementary oxygen (days, median, Q1-Q3)	2 (0; 28)	0 (0; 1)	25 (5; 44)	0.0001
Range	0–284	0–45	0–284	
Duration of CPAP ² (days, median, Q1-Q3)	23 (6; 41)	7 (2; 19)	38 (26; 51)	0.0001
Range	0–130	0–55	0–130	
Duration of invasive ventilation (days, median, Q1-Q3)	0.5 (0; 2)	0.5 (0; 2)	1 (0; 6)	0.0001
Range	0–10	0–52	0–52	
Length of stay (days, median, Q1-Q3)	62 (42–86)	43 (36–51)	82 (66–107)	0.0001

p values were computed with student's t-test or Kruskal-Wallis test for normally distributed and non-normally distributed continuous variables and with chi2 tests for categorical variables.

1: Major brain lesions: Cystic leucomalacia and/or grade III-IV intraventricular hemorrhage.

2: CPAP: continuous positive airway pressure.

<https://doi.org/10.1371/journal.pone.0241769.t001>

Univariable linear regression analyses

Variables significantly associated with PMA at FOF were BPD (+6 days), parental socioeconomic status, with a better status (i.e. lower score) associated with younger PMA at FOF as the maximum difference of 10 points in SES could add 11 days of FOF, gestational age, blood culture proven sepsis (+ 12 days) and treated PDA (+ 5.7 days), surgically treated PDA (+19 days) (Table 2). Respiratory support variables were strongly associated with the outcome, as each week of supplementary oxygen added 1.6 days of feeding tube, each week of CPAP 1.7 day of feeding tube and each day of invasive ventilation added 0.9 days of feeding tube. No association could be shown for gender, birthweight z-score, SGA, necrotizing enterocolitis, mode of delivery and major brain lesions with age at FOF.

Multivariable linear regression analyses

The variables for the multivariable regression were socioeconomic status, gestational age, sepsis, necrotizing enterocolitis, treated PDA and surgically treated PDA and the 4 respiratory variables (supplementary oxygen and CPAP, invasive ventilation, and bronchopulmonary dysplasia). The final analysis is shown in Table 3.

Table 2. Univariable regression.

	Coefficient (CI 95%)	Beta coefficient	R ²	P
Socioeconomic status	1.13 (0.27 ; 2)	0.24	0.06	0.011
Gestational age (weeks)	-2.05 (-3.14; -0.95)	-0.32	0.1	<0.001
Female gender	1.16 (-3.11 ; 5.44)	0.049	0.0024	0.59
Birthweight z-score	-1.28 (-4.25 ; 1.68)	-0.079	0.0061	0.394
SGA ¹	0.7 (-13.09 ; 14.49)	0.0091	0.0001	0.92
Vaginal delivery	-2.03 (-7.39 ; 3.33)	-0.068	0.0047	0.45
Sepsis	12.64 (6.11; 19.17)	0.33	0.11	<0.001
Necrotizing enterocolitis	10.73 (-1.1 ; 22.57)	0.16	0.026	0.075
Treated PDA	5.75 (-0.45; 10.8)	0.19	0.036	0.036
Surgically treated PDA	19.24 (9.99; 28.48)	0.35	0.12	<0.001
Major brain lesion	1.98 (-5.8 ; 9.75)	0.073	0.0021	0.616
Mother's milk at discharge (any)	1.81 (-2.64; 6.27)	0.046	0.0054	0.422
Oxygen treatment(days)	0.23 (0.18 ; 0.28)	0.67	0.44	<0.001
CPAP³(days)	0.24 (0.16 ; 0.32)	0.47	0.23	<0.001
Invasive ventilation (days)	0.92 (0.71 ; 1.13)	0.62	0.38	<0.001
Bronchopulmonary dysplasia	6.41 (1.65; 11.2)	0.23	0.05	0.008

1. Small for gestational age

2. PDA, patent ductus arteriosus

3 CPAP: continuous positive airway pressure

<https://doi.org/10.1371/journal.pone.0241769.t002>

Discussion

This study examined the association of main neonatal characteristics with the postmenstrual age at which our cohort of very preterm infants were weaned from the feeding tube. FOF was achieved at a median of 36 6/7 weeks PMA (Q1-Q3 35 6/7-39 2/7) within the upper reported range of 35 1/7–36 6/7 weeks PMA [8, 29, 30].

The univariable regression showed a weak but significant association of socioeconomic status, gestational age, sepsis and PDA with the outcome, whereas all variables linked with respiratory support (supplementary oxygen, duration of CPAP, of invasive ventilation, and, BPD)

Table 3. Multivariable linear regression.

PMA at full oral feeding	Coefficient (CI 95%)	P
Socioeconomic status	0.29 (-0.24 , 0.82)	0.282
Gestational age (days)	0.85 (-0.26; 1.96)	0.133
Sepsis	2.02 (-3.74; 7.78)	0.488
Necrotizing enterocolitis	-4.00 (-13.8; 5.81)	0.420
Treated patent ductus arteriosus	-5.60 (-9.82; -1.38)	0.010
Surgically treated PDA	6.52 (-0.83; 13.8)	0.082
Invasive ventilation (days)	0.36 (-0.11 ; 0.73)	0.058
CPAP ¹ (days)	0.12 (0.01; 0.23)	0.032
Oxygen treatment (days)	0.23 (0.14 ; 0.32)	<0.001
Bronchopulmonary dysplasia	-11.23 (-15.5; -6.9)	<0.001

In the multivariable linear regression ($p < 0.0001$, $r^2 0.67$), bronchopulmonary dysplasia was the most important single predictor of delayed FOF, followed by oxygen and CPAP duration, as well as treated PDA (any treatment).

PMID: 30974433

<https://doi.org/10.1371/journal.pone.0241769.t003>

and surgically treated PDA were associated with the outcome with moderate to large effect sizes. Not surprisingly, the multivariable analysis also showed that supplementary oxygen, CPAP, invasive ventilation, BPD and PDA were associated with PMA at FOF. BPD, which affected 25% of our population, was the single most potent predictor, followed by treated PDA.

Our findings are similar to other studies for some variables such as gestational age [29], with each supplementary week decreasing the age at FOF by 2 days, or treated patent ductus arteriosus [8, 30, 31]. The mechanism through which PDA is linked with FOF is not clear, PDA may be seen as a marker of the severity of the infant's condition. Contrary to most published studies, there was no association with SGA, which could be explained by the varying definition of SGA, based on a birthweight z score < -2 SD in our cohort, which included very few patients (3%), which were all in the late FOF group, nor with NEC, which was a rare event in our cohort.

Better socio-economic status was associated with earlier FOF in the univariable model, although this result needs to be interpreted with caution, as multiple testing may impact the significance of some results. To our knowledge, this association has not been studied before, but the association of socioeconomic status with parental presence and skin to skin care [32], and breastfeeding during NICU [11, 33], could have mediated this outcome. The association of socio-economic status with later feeding difficulties in preterm children has also been shown [34].

Respiratory support was a risk factor for delayed FOF in several studies [8, 29–31, 35] with several possible explanations. In most neonatal units, infants are not offered oral feeds as long as they are treated for respiratory distress (mainly by fear of aspiration). The transition period from tube to oral feeding thus starts later for these infants. Moreover, infants treated by invasive or non-invasive ventilation experience nociceptive stimuli in the naso- and orofacial region, which can lead to altered processing of sensory information [36], as well as with olfactory stimuli that are known to be very important in feeding processes particularly at this age. Moreover, in a population of very preterm infants, Neubauer showed that BPD and delayed FOF were independently associated with delayed brain maturation at term-equivalent age [37]. Delayed age at FOF may thus be seen as an early marker of immature sensorimotor development and delayed or altered brain maturation, a risk factor for long-lasting neurodevelopmental impairments [38–40].

The major strengths of this study are the prospective collection of data entered and retrieved from our clinical information system Metavision® and in our study database. The analysis of respiratory support was based on objective measures of duration of support, as well as the diagnoses of BPD, more subject to interpretation, [26].

The major limitation of this study is its single-centre retrospective design, with some variables of interest not available, such as time of parental presence, quantitative assessment of skin to skin care or oral stimulation which have been shown to be associated with FOF [3]. Other obstacles described by Tubbs- Cooley, the “missed opportunities”, where infants are tube fed for convenience, thus depriving them of feeding stimulations, were not recorded either [41]. The feeding protocol differed slightly between infants fed with mother's milk, who could breastfeed as early as possible, and infants fed formula (30% at discharge), who were offered bottle feeding from 34 weeks on. The data about timing of these milestones, and about the moment mothers switched from breast to bottle was not recorded in our data set, so the impact of these variables was not analyzed. However, the type of milk at discharge was not associated with FOF. Finally, the risk of collinearity is high among the studied variables, such as the variables based on respiratory support, PDA, and gestational age, which warrants caution in the interpretation of the results.

Perspectives

The feeding protocol has changed in our unit since this period and infants are offered a bundle of care including administration of oropharyngeal colostrum even for mechanically ventilated neonates, breastfeeding peer support, early bottle feeding when needed, systematic skin-to-skin care, and oral feeding under CPAP. The impact of these interventions need to be monitored. Recent studies also seem to show benefits in starting cue-based feeding preterm infants at a very early age [42]. Research on oral stimulation in unselected populations of preterm infants have shown inconclusive results according to a recent Cochrane review [14], due to methodological flaws such as varying inclusion criteria. We suggest that future studies of bundles including oral and olfactory stimulation, parental education and enhanced skin to skin care should target this specific group of infants with severe lung disease at risk of long-lasting feeding and neurodevelopmental difficulties.

Conclusion

Lung disease was a major independent determinant of delayed feeding skills in this cohort of very preterm infants, whereas gestational age, socio-economic status, and sepsis, were associated with age at FOF in an univariable linear analysis only. These findings which replicate studies in different contexts, ought to guide research on interventions aimed at shortening this learning period to FOF and thus the length of stay.

Supporting information

S1 Data.
(XLSX)

Acknowledgments

To Jocelyne Urfer and Corinne Stadelmann, responsible for maintaining our clinical information system Metavision®, and who diligently provided us with data extractions as needed.

Author Contributions

Conceptualization: Myriam Bickle Graz.

Data curation: Grégoire Brun, Myriam Bickle Graz.

Formal analysis: Grégoire Brun, Myriam Bickle Graz.

Investigation: Grégoire Brun, Eric Giannoni, Myriam Bickle Graz.

Methodology: Grégoire Brun, Céline J. Fischer Fumeaux, Eric Giannoni, Myriam Bickle Graz.

Supervision: Céline J. Fischer Fumeaux, Eric Giannoni.

Writing – original draft: Grégoire Brun.

Writing – review & editing: Céline J. Fischer Fumeaux, Eric Giannoni, Myriam Bickle Graz.

References

1. Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller A-B, et al. Born Too Soon: The global epidemiology of 15 million preterm births. *Reproductive Health*. 2013; 10(Suppl 1):S2. <https://doi.org/10.1186/1742-4755-10-S1-S2> PMID: 24625129
2. Schneider J, Fischer Fumeaux CJ, Duerden EG, Guo T, Foong J, Bickle Graz M, et al. Nutrient Intake in the First Two Weeks of Life and Brain Growth in Preterm Neonates. 2018; 141(3):e20172169.

3. Gianni ML, Sannino P, Bezze E, Comito C, Plevani L, Roggero P, et al. Does parental involvement affect the development of feeding skills in preterm infants? A prospective study. *Early Hum Dev.* 2016; 103:123–8. <https://doi.org/10.1016/j.earlhumdev.2016.08.006> PMID: 27591506
4. Adams-Chapman I, Bann CM, Vaucher YE, Stoll BJ, Eunice Kennedy Shriver National Institute of Child H, Human Development Neonatal Research N. Association between feeding difficulties and language delay in preterm infants using Bayley Scales of Infant Development-Third Edition. *J Pediatr.* 2013; 163(3):680–5 e1-3. <https://doi.org/10.1016/j.jpeds.2013.03.006> PMID: 23582139
5. Adams-Chapman I, Bann C, Carter SL, Stoll BJ, Network NNR. Language outcomes among ELBW infants in early childhood. *Early Hum Dev.* 2015; 91(6):373–9. <https://doi.org/10.1016/j.earlhumdev.2015.03.011> PMID: 25955535
6. Guellec I, Lapillonne A, Marret S, Picaud JC, Mitanchez D, Charkaluk ML, et al. Effect of Intra- and Extrauterine Growth on Long-Term Neurologic Outcomes of Very Preterm Infants. *J Pediatr.* 2016; 175:93–9 e1. <https://doi.org/10.1016/j.jpeds.2016.05.027> PMID: 27318373
7. Lau C, Bhat K, Potak D, Schanler RJ. Oral Feeding Assessment Predicts Length of Hospital Stay in Late Preterm Infants. *J Pediatr Mother Care.* 2015; 1(1). PMID: 27042698
8. Hwang YS, Ma MC, Tseng YM, Tsai WH. Associations among perinatal factors and age of achievement of full oral feeding in very preterm infants. *Pediatr Neonatol.* 2013; 54(5):309–14. <https://doi.org/10.1016/j.pedneo.2013.03.013> PMID: 23660538
9. Burnett AC, Cheong JLY, Doyle LW. Biological and Social Influences on the Neurodevelopmental Outcomes of Preterm Infants. *Clin Perinatol.* 2018; 45(3):485–500. <https://doi.org/10.1016/j.clp.2018.05.005> PMID: 30144851
10. Pineda R, Bender J, Hall B, Shabosky L, Annecca A, Smith J. Parent participation in the neonatal intensive care unit: Predictors and relationships to neurobehavior and developmental outcomes. *Early Human Development.* 2018; 117:32–8. <https://doi.org/10.1016/j.earlhumdev.2017.12.008> PMID: 29275070
11. Gertz B, DeFranco E. Predictors of breastfeeding non-initiation in the NICU. *Matern Child Nutr.* 2019; 15(3):e12797. <https://doi.org/10.1111/mcn.12797> PMID: 30767426
12. Fontana C, Menis C, Pesenti N, Passera S, Liotto N, Mosca F, et al. Effects of early intervention on feeding behavior in preterm infants: A randomized controlled trial. *Early Hum Dev.* 2018; 121:15–20. <https://doi.org/10.1016/j.earlhumdev.2018.04.016> PMID: 29730130
13. Say B, Simsek G, Canpolat F, Oguz S. Effects of Pacifier Use on Transition Time from Gavage to Breastfeeding in Preterm Infants: A Randomized Controlled Trial. *Breastfeeding Medicine.* 2018; 13. <https://doi.org/10.1089/bfm.2018.0031> PMID: 29912580
14. Greene Z, O'Donnell CP, Walshe M. Oral stimulation for promoting oral feeding in preterm infants. *Cochrane Database Syst Rev.* 2016; 9:CD009720. <https://doi.org/10.1002/14651858.CD009720.pub2> PMID: 27644167
15. Tian X, Yi LJ, Zhang L, Zhou JG, Ma L, Ou YX, et al. Oral Motor Intervention Improved the Oral Feeding in Preterm Infants: Evidence Based on a Meta-Analysis With Trial Sequential Analysis. *Medicine (Baltimore).* 2015; 94(31):e1310. <https://doi.org/10.1097/MD.0000000000001310> PMID: 26252313
16. <https://www.bfs.admin.ch/bfs/fr/home/statistiques/population/naissances-deces/naissances.assetdetail.13187376.html>
17. Bertelli C, Pillonel T, Torregrossa A, Prod'hom G, Fischer CJ, Greub G, et al. Bifidobacterium longum bacteremia in preterm infants receiving probiotics. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America.* 2015; 60(6):924–7.
18. Pfister R, Launoy V, Vassant C, Martinet M, Picard C, Bianchi JE, et al. Transition de l'alimentation passive à l'alimentation active chez le bébé prématuré. 2008; 60(4):317–35.
19. Levesque E, Hoti E, Azoulay D, Ichai P, Samuel D, Saliba F. The implementation of an Intensive Care Information System allows shortening the ICU length of stay. *Journal of clinical monitoring and computing.* 2015; 29(2):263–9. <https://doi.org/10.1007/s10877-014-9592-4> PMID: 24973014
20. Duvoisin G, Fischer C, Maucourt-Boulch D, Giannoni E. Reduction in the use of diagnostic tests in infants with risk factors for early-onset neonatal sepsis does not delay antibiotic treatment. *Swiss medical weekly.* 2014; 144:w13981. <https://doi.org/10.4414/smw.2014.13981> PMID: 24964177
21. Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr.* 2013; 13:59. <https://doi.org/10.1186/1471-2431-13-59> PMID: 23601190
22. Largo RH, Pfister D, Molinari L, Kundu S, Lipp A, Duc G. Significance of prenatal, perinatal, and postnatal factors in the development of AGA preterm infants at five to seven years. *Dev Med Child Neurol.* 1989; 31:440–56. <https://doi.org/10.1111/j.1469-8749.1989.tb04022.x> PMID: 2680687
23. Natalucci G, Bucher HU, Von Rhein M, Borradori Tolsa C, Latal B, Adams M. Population based report on health related quality of life in adolescents born very preterm. *Early Hum Dev.* 2017; 104:7–12. <https://doi.org/10.1016/j.earlhumdev.2016.11.002> PMID: 27936395

24. Giannoni E, Agyeman PKA, Stocker M, Posfay-Barbe KM, Heining U, Spycher BD, et al. Neonatal Sepsis of Early Onset, and Hospital-Acquired and Community-Acquired Late Onset: A Prospective Population-Based Cohort Study. *J Pediatr*. 2018; 201:106–14.e4. <https://doi.org/10.1016/j.jpeds.2018.05.048> PMID: 30054165
25. Papile LA, Burstein J, Burstein R, Keffler H. Incidence and evolution of the subependymal intraventricular hemorrhage: a study of infants with weights less than 1500 grams. *J Pediatr*. 1978; 92:529–34. [https://doi.org/10.1016/s0022-3476\(78\)80282-0](https://doi.org/10.1016/s0022-3476(78)80282-0) PMID: 305471
26. Hines D, Modi N, Lee SK, Isayama T, Sjörs G, Gagliardi L, et al. Scoping review shows wide variation in the definitions of bronchopulmonary dysplasia in preterm infants and calls for a consensus. *Acta Paediatr*. 2017; 106(3):366–74. <https://doi.org/10.1111/apa.13672> PMID: 27862302
27. <https://www.medin-medical.com/en/Products/Patient-interfaces/Medijet.html>
28. Sweet DG, Carnielli V, Greisen G, Hallman M, Ozek E, Plavka R, et al. European consensus guidelines on the management of neonatal respiratory distress syndrome in preterm infants—2013 update. *Neonatology*. 2013; 103(4):353–68. <https://doi.org/10.1159/000349928> PMID: 23736015
29. Gianni ML, Sannino P, Bezze E, Plevani L, di Cugno N, Roggero P, et al. Effect of co-morbidities on the development of oral feeding ability in pre-term infants: a retrospective study. *Sci Rep*. 2015; 5:16603. <https://doi.org/10.1038/srep16603> PMID: 26558841
30. Van Nostrand SM, Bennett LN, Coraglio VJ, Guo R, Muraskas JK. Factors influencing independent oral feeding in preterm infants. *J Neonatal Perinatal Med*. 2015. <https://doi.org/10.3233/NPM-15814045> PMID: 25766199
31. Park J, Knaf G, Thoyre S, Brandon D. Factors associated with feeding progression in extremely pre-term infants. *Nurs Res*. 2015; 64(3):159–67. <https://doi.org/10.1097/NNR.000000000000093> PMID: 25932696
32. Pineda R, Bender J, Hall B, Shabosky L, Annecca A, Smith J. Parent participation in the neonatal intensive care unit: Predictors and relationships to neurobehavior and developmental outcomes. *Early Hum Dev*. 2018; 117:32–8. <https://doi.org/10.1016/j.earlhumdev.2017.12.008> PMID: 29275070
33. Herich LC, Cuttini M, Croci I, Franco F, Di Lallo D, Baronciani D, et al. Maternal Education Is Associated with Disparities in Breastfeeding at Time of Discharge but Not at Initiation of Enteral Feeding in the Neonatal Intensive Care Unit. *J Pediatr*. 2017; 182:59–65 e7. <https://doi.org/10.1016/j.jpeds.2016.10.046> PMID: 27865429
34. Crapnell TL, Rogers CE, Neil JJ, Inder TE, Woodward LJ, Pineda RG. Factors associated with feeding difficulties in the very preterm infant. *Acta Paediatr*. 2013; 102(12):e539–45. <https://doi.org/10.1111/apa.12393> PMID: 23952198
35. Jadcherla SR, Wang M, Vijayapal AS, Leuthner SR. Impact of prematurity and co-morbidities on feeding milestones in neonates: a retrospective study. *J Perinatol*. 2010; 30(3):201–8. <https://doi.org/10.1038/jp.2009.149> PMID: 19812589
36. Dodrill P, McMahon S, Ward E, Weir K, Donovan T, Riddle B. Long-term oral sensitivity and feeding skills of low-risk pre-term infants. *Early Hum Dev*. 2004; 76(1):23–37. <https://doi.org/10.1016/j.earlhumdev.2003.10.001> PMID: 14729160
37. Neubauer V, Junker D, Griesmaier E, Schocke M, Kiechl-Kohlendorfer U. Bronchopulmonary dysplasia is associated with delayed structural brain maturation in preterm infants. *Neonatology*. 2015; 107(3):179–84. <https://doi.org/10.1159/000369199> PMID: 25632975
38. Jadcherla SR, Khot T, Moore R, Malkar M, Gulati IK, Slaughter JL. Feeding Methods at Discharge Predict Long-Term Feeding and Neurodevelopmental Outcomes in Preterm Infants Referred for Gastrostomy Evaluation. *J Pediatr*. 2017; 181:125–30 e1. <https://doi.org/10.1016/j.jpeds.2016.10.065> PMID: 27939123
39. Johnson S, Matthews R, Draper ES, Field DJ, Manktelow BN, Marlow N, et al. Eating difficulties in children born late and moderately preterm at 2 y of age: a prospective population-based cohort study. *Am J Clin Nutr*. 2016; 103(2):406–14. <https://doi.org/10.3945/ajcn.115.121061> PMID: 26718420
40. Lainwala S, Kosyakova N, Power K, Hussain N, Moore JE, Hagadorn JI, et al. Delayed Achievement of Oral Feedings Is Associated with Adverse Neurodevelopmental Outcomes at 18 to 26 Months Follow-up in Preterm Infants. *Am J Perinatol*. 2019. <https://doi.org/10.1055/s-0039-1681059> PMID: 30822799
41. Tubbs-Cooley HL, Pickler RH, Meinen-Derr JK. Missed oral feeding opportunities and preterm infants' time to achieve full oral feedings and neonatal intensive care unit discharge. *Am J Perinatol*. 2015; 32(1):1–8. <https://doi.org/10.1055/s-0034-1372426> PMID: 24683073
42. Fry TJ, Marfurt S, Wengier S. Systematic Review of Quality Improvement Initiatives Related to Cue-Based Feeding in Preterm Infants. *Nursing for women's health*. 2018; 22(5):401–10. <https://doi.org/10.1016/j.nwh.2018.07.006> PMID: 30138603