



A two-dimensional Zn coordination polymer with a three-dimensional supramolecular architecture

Fuhong Liu,* Yan Ding, Qiuyu Li and Liping Zhang

Basis Department, Jilin Business and Technology College, Changchun, Jilin, People's Republic of China.

*Correspondence e-mail: liuhf563@nenu.edu.cn

Received 27 July 2017

Accepted 29 August 2017

Edited by A. Van der Lee, Université de Montpellier II, France

Keywords: crystal structure; coordination polymer; zinc complex; two-dimensional layer.**CCDC reference:** 1564369**Supporting information:** this article has supporting information at journals.iucr.org/e


The title compound, poly[bis{ μ_2 -4,4'-bis[(1,2,4-triazol-1-yl)methyl]biphenyl- $\kappa^2N^4:N^4$ }bis(nitrato- κO)zinc(II)], $[\text{Zn}(\text{NO}_3)_2(\text{C}_{18}\text{H}_{16}\text{N}_6)_2]_n$, is a two-dimensional zinc coordination polymer constructed from 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl units. It was synthesized and characterized by elemental analysis and single-crystal X-ray diffraction. The Zn^{II} cation is located on an inversion centre and is coordinated by two O atoms from two symmetry-related nitrate groups and four N atoms from four symmetry-related 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligands, forming a distorted octahedral $\{\text{ZnN}_4\text{O}_2\}$ coordination geometry. The linear 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligand links two Zn^{II} cations, generating two-dimensional layers parallel to the crystallographic (132) plane. The parallel layers are connected by C—H...O, C—H...N, C—H... π and π — π stacking interactions, resulting in a three-dimensional supramolecular architecture.

1. Chemical context


Over the past few decades, the self-assembly of coordination polymers (CPs) or metal–organic frameworks (MOFs) based on metal ions or clusters and organic ligands has attracted much attention, owing to their intriguing molecular topologies and potential applications. Multidentate ligands derived from 1,2,4-triazole that contain an aromatic core have been used for this purpose, examples being 1,4-bis(1*H*-1,2,4-triazol-1-ylmethyl)benzene (Wang *et al.*, 2007; Ding & Zou, 2010; Zhu *et al.*, 2010), 1,3-bis(1*H*-1,2,4-triazol-1-ylmethyl)benzene (Zhang *et al.*, 2012; Ge *et al.*, 2008; Zhu *et al.*, 2015), 1,2-bis(1*H*-1,2,4-triazol-1-ylmethyl)benzene (Yang *et al.*, 2009; Zhao *et al.*, 2017; Zhang *et al.*, 2013), 1,3,5-tris(1*H*-1,2,4-triazol-1-ylmethyl)benzene (Li *et al.*, 2012; Yin *et al.*, 2009; Shi *et al.*, 2011), 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl (Mu *et al.*, 2011; Ren *et al.*, 2010; Ni *et al.*, 2010). Hydrothermal synthesis has been proved to be an effective method for the construction of these new coordination polymers. In this study, a new two-dimensional CP, *viz.* poly[bis{ μ_2 -4,4'-bis[(1,2,4-triazol-1-yl)methyl]biphenyl- $\kappa^2N^4:N^4$ }bis(nitrato- κO)zinc], $[\text{Zn}(\text{NO}_3)_2(\text{C}_{18}\text{H}_{16}\text{N}_6)_2]_n$, was synthesized under hydrothermal conditions by the reaction of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ and 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl at 313 K for 48 h. We report here its crystal structure and its elemental analysis.

2. Structural commentary

The title complex crystallizes in the triclinic space group $P\bar{1}$; the asymmetric unit of the structure consists of one Zn^{II} cation



(site symmetry $\bar{1}$), one nitrate anion and one 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligand.



As shown in Fig. 1, each Zn^{II} cation exhibits a slightly distorted octahedral [ZnN₄O₂] coordination geometry and is coordinated by four N atoms (N1, N4, N1ⁱ and N4ⁱ) from four symmetry-related organic ligands and two O atoms (O3 and O3ⁱ) from two symmetry-related nitrate groups (see Fig. 1 for symmetry code). The Zn–O [2.191 (2) Å] and Zn–N bond lengths [2.124 (3)–2.168 (2) Å] are in agreement with corresponding bond lengths found in previously reported Zn^{II} coordination polymers. For the title coordination polymer, the Zn^{II} cation is coordinated by four 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligands and two nitrate anions, and each organic ligand in turn connects two Zn^{II} cations to generate a two-dimensional layer parallel to the crystallographic (132) plane. The organic ligand adopts a *cis,cis* substituent conformation. The two distinct Zn···Zn distances are 18.397 (3) and 18.964 (3) Å (see Fig. 2). The two benzene rings of the 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligand lie nearly in one plane [dihedral angle = 0.00 (2)°]. The two triazole groups of the 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl ligand are inclined to the plane of the central biphenyl groups, with dihedral angles of 80.050 (2)° (C1/C2/N1/N2/N3) and 85.511 (2)° (C10/C11/N4/N5/N6). Four adjacent Zn^{II} cations are connected by four linear organic ligands and form a 72-membered macrocyclic




Figure 1
The asymmetric unit of (I), showing the atom-numbering scheme. Displacement ellipsoids drawn at the 25% probability level. [Symmetry code: (i) $-x, 2 - y, -z$.]

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C3–H3 <i>B</i> ···O1 ⁱ	0.97	2.31	3.2728 (7)	170
C3–H3 <i>B</i> ···O1 <i>A</i> ⁱ	0.97	2.33	3.2765 (7)	165
C10–H10···O2 <i>A</i> ⁱⁱ	0.93	2.53	3.0888 (6)	115
C14–H14···O2 ⁱⁱ	0.93	2.46	3.5454 (7)	158
C15–H15···N6 ⁱⁱⁱ	0.93	2.58	3.482 (16)	162

Symmetry codes: (i) $x - 1, y + 1, z$; (ii) $-x + 1, -y, -z$; (iii) $x - 1, y, z$.

ring in the above-mentioned two-dimensional layer (see Fig. 2).

3. Supramolecular features

Neighbouring layers are linked to each other by weak interactions (Table 1), including C–H···O, C–H···N, C–H··· π [C11–H11···C*g*1ⁱⁱ = 3.6756 (8) Å and C12–H12···C*g*2ⁱⁱⁱ = 3.5252 (7) Å; C*g*1 and C*g*2 are the centroids of the triazole (C1/C2/N1/N2/N3) and phenyl (C4–C9) rings, respectively; symmetry codes: (ii) $2 - x, -y, -z$; (iii) $1 - x, 1 - y, -z$]. These interactions, together with the covalent interactions in the infinite two-dimensional polymeric-like layer, make up a three-dimensional supramolecular structure.

4. Database survey

A search in the Cambridge Structural Database (Groom *et al.*, 2016) for zinc and the 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl moiety gave eight hits. Seven of them are constructed by 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl units and different carboxylate ligands. One example is a chain structure based on Zn and 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl (PUQWAA; Ni *et al.*, 2010).




Figure 2
The two-dimensional layer parallel to the crystallographic (132) plane.

Table 2
Experimental details.

Crystal data	
Chemical formula	[Zn(NO ₃) ₂ (C ₁₈ H ₁₆ N ₆) ₂]
<i>M</i> _r	822.12
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	293
<i>a</i> , <i>b</i> , <i>c</i> (Å)	7.3257 (15), 9.0188 (18), 15.578 (3)
α , β , γ (°)	81.70 (3), 77.64 (3), 68.90 (3)
<i>V</i> (Å ³)	935.4 (4)
<i>Z</i>	1
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.72
Crystal size (mm)	0.24 × 0.22 × 0.20
Data collection	
Diffractometer	Bruker APEXII Quazar
Absorption correction	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)
<i>T</i> _{min} , <i>T</i> _{max}	0.84, 0.86
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	7292, 3271, 2589
<i>R</i> _{int}	0.042
(sin θ/λ) _{max} (Å ⁻¹)	0.595
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.051, 0.143, 1.05
No. of reflections	3271
No. of parameters	278
No. of restraints	85
H-atom treatment	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.34, -0.54

Computer programs: *APEX3* (Bruker, 2016), *SAINT-Plus* (Bruker, 2016), *SHELXT* (Sheldrick, 2008), *SHELXTL-Plus* (Sheldrick, 2008) and *OLEX2* (Dolomanov *et al.*, 2009).

5. Synthesis and crystallization

Zn(NO₃)₂·6H₂O (0.1 mmol), 4,4'-bis[(1*H*-1,2,4-triazol-1-yl)methyl]-1,1'-biphenyl (0.1 mmol) and water (6 ml) were mixed and placed in a thick Pyrex tube, which was sealed and heated to 413 K for 72 h. After cooling to room temperature, colourless block-shaped crystals (53% yield, based on Zn) suitable for X-ray analysis were obtained. Elemental analysis calculated for C₃₆H₃₂N₁₄O₆Zn: C 52.59, H 3.92, N 23.85%; found: C 52.23, H 3.74, N 23.49%.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms bonded to C atoms were

positioned geometrically and allowed to ride on their parent atoms, with C—H = 0.93 Å with *U*_{iso}(H) = 1.2*U*_{eq}(C) for other H atoms. Atoms O1 and O2 of the nitrate group are disordered over two orientations, with occupancies of 0.511 (11) and 0.489 (11), and were refined through the use of SADI, RIGU and SIMU commands.

Funding information

Funding for this research was provided by: Science and Technology Research Projects of Jilin Province Department of Education (JJKH20170186KJ) .

References

- Bruker (2016). *APEX3*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ding, B. & Zou, H.-A. (2010). *Acta Cryst.* **E66**, m932.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Ge, H., Liu, K., Yang, Y., Li, B. & Zhang, Y. (2008). *Inorg. Chem. Commun.* **11**, 260–264.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.
- Li, Q.-X., Shi, X.-J. & Chen, L.-C. (2012). *Acta Cryst.* **E68**, m1299.
- Mu, Y., Song, Y., Wang, C., Hou, H. & Fan, Y. (2011). *Inorg. Chim. Acta*, **365**, 167–176.
- Ni, T., Shao, M., Zhu, S., Zhao, Y., Xing, F. & Li, M. (2010). *Cryst. Growth Des.* **10**, 943–951.
- Ren, C., Liu, P., Wang, Y.-Y., Huang, W.-H. & Shi, Q.-Z. (2010). *Eur. J. Inorg. Chem.* pp. 5545–5555.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shi, X., Li, Q., Zhang, Y., Chang, X. & Hou, H. (2011). *J. Coord. Chem.* **64**, 3918–3927.
- Wang, W.-B., Wang, L.-Y., Li, B.-L. & Zhang, Y. (2007). *Acta Cryst.* **E63**, m2416–m2417.
- Yang, Y., Feng, Y.-F., Liang, N., Li, B.-L. & Zhang, Y. (2009). *J. Coord. Chem.* **62**, 3819–3827.
- Yin, X.-J., Zhou, X.-H., Gu, Z.-G., Zuo, J.-L. & You, X.-Z. (2009). *Inorg. Chem. Commun.* **12**, 548–551.
- Zhang, Z., Ma, J.-F., Liu, Y.-Y., Kan, W.-Q. & Yang, J. (2013). *CrystEngComm*, **15**, 2009–2018.
- Zhang, H.-K., Wang, X., Wang, S. & Wang, X.-D. (2012). *Acta Cryst.* **E68**, m856.
- Zhao, S., Zheng, T.-R., Zhang, Y.-Q., Lv, X.-X., Li, B.-L. & Zhang, Y. (2017). *Polyhedron*, **121**, 61–69.
- Zhu, X., Guo, Y. & Zou, Y.-L. (2010). *Acta Cryst.* **E66**, m85.
- Zhu, X., Yang, Y., Jiang, N., Li, B., Zhou, D., Fu, H. & Wang, N. (2015). *Z. Anorg. Allg. Chem.* **641**, 699–703.

supporting information

Acta Cryst. (2017). E73, 1402-1404 [https://doi.org/10.1107/S2056989017012452]

A two-dimensional Zn coordination polymer with a three-dimensional supramolecular architecture

Fuhong Liu, Yan Ding, Qiuyu Li and Liping Zhang

Computing details

Data collection: *APEX3* (Bruker, 2016); cell refinement: *S SAINT-Plus* (Bruker, 2016); data reduction: *S SAINT-Plus* (Bruker, 2016); program(s) used to solve structure: *SHELXT* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL-Plus* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

Poly[bis(μ_2 -4,4'-bis([1,2,4-triazol-1-yl)methyl]biphenyl- κ^2 N⁴:N^{4'})bis(nitrato- κ O)zinc(II)]

Crystal data

[Zn(NO₃)₂(C₁₈H₁₆N₆)₂]

$M_r = 822.12$

Triclinic, *P1*

$a = 7.3257$ (15) Å

$b = 9.0188$ (18) Å

$c = 15.578$ (3) Å

$\alpha = 81.70$ (3)°

$\beta = 77.64$ (3)°

$\gamma = 68.90$ (3)°

$V = 935.4$ (4) Å³

$Z = 1$

$F(000) = 424$

$D_x = 1.459$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7292 reflections

$\theta = 1.6$ – 25.1 °

$\mu = 0.72$ mm⁻¹

$T = 293$ K

Block, colorless

$0.24 \times 0.22 \times 0.2$ mm

Data collection

Bruker APEXII Quazar
diffractometer

Radiation source: microfocus sealed X-ray tube,
Incoatec I μ s

Mirror optics monochromator

Detector resolution: 7.9 pixels mm⁻¹

0.5° ω and 0.5° ϕ scans

Absorption correction: multi-scan
(SADABS; Krause *et al.*, 2015)

$T_{\min} = 0.84$, $T_{\max} = 0.86$

7292 measured reflections

3271 independent reflections

2589 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\max} = 25.0$ °, $\theta_{\min} = 3.0$ °

$h = -8 \rightarrow 8$

$k = -10 \rightarrow 9$

$l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.143$

$S = 1.05$

3271 reflections

278 parameters

85 restraints

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.079P)^2 + 0.2956P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.34$ e Å⁻³

$\Delta\rho_{\min} = -0.54$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.000000	1.000000	0.000000	0.0406 (2)	
O3	0.1631 (4)	1.1607 (3)	0.00189 (17)	0.0574 (7)	
N1	0.2322 (4)	0.8123 (3)	0.05733 (18)	0.0441 (7)	
N2	0.4790 (5)	0.6937 (4)	0.1274 (2)	0.0482 (7)	
N3	0.4061 (6)	0.5766 (4)	0.1198 (2)	0.0602 (9)	
N4	0.1701 (4)	0.9540 (4)	-0.12813 (18)	0.0457 (7)	
N5	0.3961 (5)	0.8456 (4)	-0.23645 (19)	0.0466 (7)	
N6	0.2787 (6)	0.9865 (5)	-0.2713 (2)	0.0729 (11)	
N7	0.1398 (5)	1.2530 (4)	0.0601 (2)	0.0570 (8)	
C1	0.2590 (6)	0.6550 (5)	0.0771 (3)	0.0554 (10)	
H1	0.179295	0.606514	0.061538	0.066*	
C2	0.3743 (6)	0.8301 (4)	0.0904 (2)	0.0474 (8)	
H2	0.397217	0.925660	0.087958	0.057*	
C3	0.6466 (6)	0.6585 (5)	0.1731 (3)	0.0601 (10)	
H3A	0.687840	0.751186	0.165279	0.072*	
H3B	0.757323	0.571442	0.146296	0.072*	
C4	0.5997 (6)	0.6146 (4)	0.2700 (3)	0.0513 (9)	
C5	0.7455 (7)	0.5120 (7)	0.3132 (4)	0.0902 (17)	
H5	0.873179	0.468413	0.281685	0.108*	
C6	0.7082 (8)	0.4711 (8)	0.4028 (4)	0.0927 (17)	
H6	0.812884	0.404978	0.430318	0.111*	
C7	0.5224 (6)	0.5251 (4)	0.4518 (3)	0.0532 (9)	
C8	0.3756 (8)	0.6273 (6)	0.4079 (3)	0.0813 (16)	
H8	0.246945	0.667746	0.439147	0.098*	
C9	0.4126 (8)	0.6717 (6)	0.3195 (3)	0.0823 (16)	
H9	0.309014	0.741721	0.292602	0.099*	
C10	0.3284 (6)	0.8298 (5)	-0.1520 (2)	0.0515 (9)	
H10	0.384162	0.743198	-0.114162	0.062*	
C11	0.1465 (6)	1.0486 (5)	-0.2035 (2)	0.0588 (10)	
H11	0.046686	1.147437	-0.206804	0.071*	
C12	0.5625 (7)	0.7311 (5)	-0.2909 (3)	0.0658 (13)	
H12A	0.509806	0.678916	-0.324861	0.079*	
H12B	0.641231	0.649898	-0.252615	0.079*	
C13	0.6941 (6)	0.8103 (4)	-0.3527 (3)	0.0530 (10)	
C14	0.8175 (7)	0.8650 (6)	-0.3228 (3)	0.0722 (13)	
H14	0.821652	0.853057	-0.262825	0.087*	
C15	0.9375 (7)	0.9384 (6)	-0.3803 (3)	0.0684 (13)	
H15	1.021917	0.973475	-0.358088	0.082*	
C16	0.9348 (5)	0.9608 (4)	-0.4696 (2)	0.0431 (8)	

C17	0.8101 (7)	0.9040 (6)	-0.4990 (3)	0.0738 (14)	
H17	0.804965	0.915631	-0.558860	0.089*	
C18	0.6915 (8)	0.8296 (6)	-0.4409 (3)	0.0750 (14)	
H18	0.608813	0.792084	-0.462688	0.090*	
O1	-0.0075 (14)	1.3759 (11)	0.0613 (8)	0.0890 (19)	0.500 (12)
O2	0.2754 (14)	1.2075 (12)	0.1048 (6)	0.0792 (17)	0.500 (12)
O1A	-0.0249 (12)	1.3242 (11)	0.1028 (7)	0.0807 (18)	0.500 (12)
O2A	0.2775 (13)	1.2743 (14)	0.0828 (7)	0.0812 (17)	0.500 (12)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0365 (3)	0.0492 (4)	0.0287 (3)	-0.0098 (2)	-0.0025 (2)	0.0033 (2)
O3	0.0658 (17)	0.0635 (16)	0.0492 (15)	-0.0325 (14)	-0.0020 (13)	-0.0084 (13)
N1	0.0444 (16)	0.0499 (17)	0.0331 (15)	-0.0123 (13)	-0.0074 (13)	0.0027 (13)
N2	0.0514 (18)	0.0509 (17)	0.0384 (16)	-0.0136 (14)	-0.0123 (14)	0.0052 (13)
N3	0.077 (2)	0.0481 (18)	0.060 (2)	-0.0216 (17)	-0.0251 (19)	0.0036 (16)
N4	0.0463 (17)	0.0559 (18)	0.0310 (15)	-0.0158 (14)	-0.0048 (13)	0.0013 (13)
N5	0.0506 (17)	0.0545 (17)	0.0347 (16)	-0.0245 (15)	0.0054 (14)	-0.0049 (14)
N6	0.072 (3)	0.088 (3)	0.042 (2)	-0.019 (2)	0.0022 (19)	0.0074 (19)
N7	0.0616 (17)	0.0610 (18)	0.0580 (19)	-0.0299 (14)	-0.0085 (15)	-0.0136 (15)
C1	0.065 (3)	0.052 (2)	0.052 (2)	-0.0185 (19)	-0.022 (2)	-0.0003 (18)
C2	0.052 (2)	0.0469 (19)	0.0385 (19)	-0.0165 (17)	-0.0040 (17)	0.0047 (15)
C3	0.052 (2)	0.071 (3)	0.057 (3)	-0.020 (2)	-0.020 (2)	0.009 (2)
C4	0.059 (2)	0.048 (2)	0.046 (2)	-0.0146 (17)	-0.0201 (18)	0.0056 (17)
C5	0.044 (3)	0.133 (5)	0.070 (3)	-0.017 (3)	-0.018 (2)	0.044 (3)
C6	0.057 (3)	0.133 (5)	0.070 (3)	-0.021 (3)	-0.028 (3)	0.047 (3)
C7	0.071 (3)	0.043 (2)	0.047 (2)	-0.0121 (18)	-0.027 (2)	0.0009 (16)
C8	0.083 (3)	0.077 (3)	0.041 (2)	0.025 (2)	-0.010 (2)	-0.006 (2)
C9	0.091 (4)	0.075 (3)	0.045 (2)	0.021 (3)	-0.027 (2)	0.002 (2)
C10	0.052 (2)	0.055 (2)	0.0356 (19)	-0.0112 (17)	0.0015 (17)	0.0035 (16)
C11	0.049 (2)	0.068 (2)	0.036 (2)	-0.0013 (19)	0.0002 (17)	0.0047 (18)
C12	0.079 (3)	0.055 (2)	0.057 (3)	-0.033 (2)	0.028 (2)	-0.017 (2)
C13	0.058 (2)	0.047 (2)	0.045 (2)	-0.0194 (18)	0.0153 (18)	-0.0088 (17)
C14	0.087 (3)	0.100 (3)	0.034 (2)	-0.048 (3)	0.002 (2)	0.000 (2)
C15	0.075 (3)	0.108 (4)	0.039 (2)	-0.054 (3)	-0.004 (2)	-0.002 (2)
C16	0.0386 (18)	0.0427 (18)	0.0388 (18)	-0.0073 (14)	0.0024 (15)	-0.0055 (15)
C17	0.087 (3)	0.114 (4)	0.038 (2)	-0.060 (3)	-0.007 (2)	0.003 (2)
C18	0.085 (3)	0.111 (4)	0.051 (3)	-0.066 (3)	-0.002 (2)	-0.005 (2)
O1	0.091 (3)	0.077 (3)	0.080 (4)	-0.006 (3)	-0.008 (3)	-0.016 (3)
O2	0.094 (3)	0.080 (4)	0.075 (3)	-0.035 (3)	-0.032 (3)	-0.003 (3)
O1A	0.080 (3)	0.075 (3)	0.074 (4)	-0.020 (3)	0.009 (3)	-0.016 (3)
O2A	0.085 (3)	0.088 (4)	0.087 (4)	-0.041 (3)	-0.030 (3)	-0.008 (3)

Geometric parameters (Å, °)

Zn1—O3	2.191 (2)	C4—C5	1.367 (6)
Zn1—O3 ⁱ	2.191 (2)	C4—C9	1.376 (6)

Zn1—N1 ⁱ	2.167 (3)	C5—H5	0.9300
Zn1—N1	2.167 (3)	C5—C6	1.385 (7)
Zn1—N4 ⁱ	2.124 (3)	C6—H6	0.9300
Zn1—N4	2.124 (3)	C6—C7	1.363 (7)
O3—N7	1.263 (4)	C7—C7 ⁱⁱ	1.505 (8)
N1—C1	1.359 (5)	C7—C8	1.376 (6)
N1—C2	1.322 (5)	C8—H8	0.9300
N2—N3	1.372 (4)	C8—C9	1.375 (7)
N2—C2	1.318 (5)	C9—H9	0.9300
N2—C3	1.465 (5)	C10—H10	0.9300
N3—C1	1.314 (5)	C11—H11	0.9300
N4—C10	1.318 (5)	C12—H12A	0.9700
N4—C11	1.353 (5)	C12—H12B	0.9700
N5—N6	1.365 (5)	C12—C13	1.506 (5)
N5—C10	1.310 (5)	C13—C14	1.359 (6)
N5—C12	1.475 (5)	C13—C18	1.364 (6)
N6—C11	1.307 (5)	C14—H14	0.9300
N7—O1	1.237 (8)	C14—C15	1.387 (6)
N7—O2	1.249 (8)	C15—H15	0.9300
N7—O1A	1.237 (7)	C15—C16	1.381 (5)
N7—O2A	1.221 (8)	C16—C16 ⁱⁱⁱ	1.487 (7)
C1—H1	0.9300	C16—C17	1.375 (5)
C2—H2	0.9300	C17—H17	0.9300
C3—H3A	0.9700	C17—C18	1.390 (6)
C3—H3B	0.9700	C18—H18	0.9300
C3—C4	1.501 (6)		
O3—Zn1—O3 ⁱ	180.0	C5—C4—C3	120.3 (4)
N1—Zn1—O3 ⁱ	92.22 (11)	C5—C4—C9	116.5 (4)
N1 ⁱ —Zn1—O3	92.22 (11)	C9—C4—C3	123.2 (4)
N1 ⁱ —Zn1—O3 ⁱ	87.78 (11)	C4—C5—H5	119.1
N1—Zn1—O3	87.78 (11)	C4—C5—C6	121.8 (5)
N1 ⁱ —Zn1—N1	180.0	C6—C5—H5	119.1
N4—Zn1—O3 ⁱ	94.65 (10)	C5—C6—H6	119.1
N4 ⁱ —Zn1—O3	94.65 (10)	C7—C6—C5	121.9 (5)
N4—Zn1—O3	85.35 (10)	C7—C6—H6	119.1
N4 ⁱ —Zn1—O3 ⁱ	85.35 (10)	C6—C7—C7 ⁱⁱ	122.4 (5)
N4 ⁱ —Zn1—N1 ⁱ	90.36 (12)	C6—C7—C8	116.2 (4)
N4 ⁱ —Zn1—N1	89.64 (12)	C8—C7—C7 ⁱⁱ	121.4 (5)
N4—Zn1—N1	90.36 (12)	C7—C8—H8	118.9
N4—Zn1—N1 ⁱ	89.64 (12)	C9—C8—C7	122.2 (4)
N4 ⁱ —Zn1—N4	180.0	C9—C8—H8	118.9
N7—O3—Zn1	128.5 (2)	C4—C9—H9	119.3
C1—N1—Zn1	130.5 (3)	C8—C9—C4	121.4 (4)
C2—N1—Zn1	126.5 (2)	C8—C9—H9	119.3
C2—N1—C1	102.7 (3)	N4—C10—H10	124.8
N3—N2—C3	120.7 (3)	N5—C10—N4	110.5 (4)
C2—N2—N3	109.9 (3)	N5—C10—H10	124.8

C2—N2—C3	129.4 (3)	N4—C11—H11	123.6
C1—N3—N2	102.0 (3)	N6—C11—N4	112.9 (4)
C10—N4—Zn1	128.1 (3)	N6—C11—H11	123.6
C10—N4—C11	103.9 (3)	N5—C12—H12A	109.2
C11—N4—Zn1	128.0 (3)	N5—C12—H12B	109.2
N6—N5—C12	122.6 (3)	N5—C12—C13	112.2 (3)
C10—N5—N6	109.0 (3)	H12A—C12—H12B	107.9
C10—N5—C12	128.3 (4)	C13—C12—H12A	109.2
C11—N6—N5	103.8 (3)	C13—C12—H12B	109.2
O1—N7—O3	115.4 (6)	C14—C13—C12	121.5 (4)
O1—N7—O2	130.6 (7)	C14—C13—C18	118.1 (4)
O2—N7—O3	114.0 (5)	C18—C13—C12	120.4 (4)
O1A—N7—O3	122.8 (5)	C13—C14—H14	119.5
O2A—N7—O3	123.5 (6)	C13—C14—C15	121.0 (4)
O2A—N7—O1A	113.6 (7)	C15—C14—H14	119.5
N1—C1—H1	122.6	C14—C15—H15	119.2
N3—C1—N1	114.8 (3)	C16—C15—C14	121.7 (4)
N3—C1—H1	122.6	C16—C15—H15	119.2
N1—C2—H2	124.7	C15—C16—C16 ⁱⁱⁱ	120.9 (4)
N2—C2—N1	110.7 (3)	C17—C16—C15	116.7 (3)
N2—C2—H2	124.7	C17—C16—C16 ⁱⁱⁱ	122.4 (4)
N2—C3—H3A	108.9	C16—C17—H17	119.4
N2—C3—H3B	108.9	C16—C17—C18	121.2 (4)
N2—C3—C4	113.5 (3)	C18—C17—H17	119.4
H3A—C3—H3B	107.7	C13—C18—C17	121.4 (4)
C4—C3—H3A	108.9	C13—C18—H18	119.3
C4—C3—H3B	108.9	C17—C18—H18	119.3

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x+2, -y+2, -z-1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3B \cdots O1 ^{iv}	0.97	2.31	3.2728 (7)	170
C3—H3B \cdots O1A ^{iv}	0.97	2.33	3.2765 (7)	165
C10—H10 \cdots O2A ^v	0.93	2.53	3.0888 (6)	115
C14—H14 \cdots O2 ^v	0.93	2.46	3.5454 (7)	158
C15—H15 \cdots N6 ^{vi}	0.93	2.58	3.482 (16)	162

Symmetry codes: (iv) $x-1, y+1, z$; (v) $-x+1, -y, -z$; (vi) $x-1, y, z$.