CORRECTION


Monoallelic and bi-allelic variants in *NCDN* cause neurodevelopmental delay, intellectual disability, and epilepsy

Ambrin Fatima, Jan Hoeber, Jens Schuster, Eriko Koshimizu, Carolina Maya-Gonzalez, Boris Keren, Cyril Mignot, Talia Akram, Zafar Ali, Satoko Miyatake, Junpei Tanigawa, Takayoshi Koike, Mitsuhiro Kato, Yoshiko Murakami, Uzma Abdullah, Muhammad Akhtar Ali, Rein Fadoul, Loora Laan, Casimiro Castillejo-López, Maarika Liik, Zhe Jin, Bryndis Birnir, Naomichi Matsumoto, Shahid M. Baig, Joakim Klar, and Niklas Dahl*

(The American Journal of Human Genetics 108, 739–748; April 1, 2021)

In the originally published version of this article, family 2 was assigned an incorrect as substitution as a result of the G>A transition in codon 478 of the *NCDN* gene. The as substitution in individual F2: II.1 should read p.Arg478Gln, not p.Arg478Glu. The error appeared consistently throughout the text, in Table 1, and in Figures 1, 2, and 3; the error has been corrected online and Figures 1, 2, and 3 appear correctly here as well. The authors regret this error.










Figure 1. Segregation of rare NCDN missense variants in families with neurodevelopmental phenotypes (corrected)




Figure 1. Segregation of rare NCDN missense variants in families with neurodevelopmental phenotypes (original)




Figure 2. Missense variants in NCDN alter length and number of neurites in SH-SY5Y cells (corrected)




Figure 2. Missense variants in NCDN alter length and number of neurites in SH-SY5Y cells (original)




Figure 3. NCDN variants interfere with mGluR5 signaling and alter electrophysiological properties of SH-SY5Y cells (corrected)




Figure 3. NCDN variants interfere with mGluR5 signaling and alter electrophysiological properties of SH-SY5Y cells (original)