



ELSEVIER

Contents lists available at ScienceDirect

Data in brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Liver proteome dataset of *Sparus aurata* exposed to low temperaturesS. Ghisaura^a, R. Melis^a, G. Biosa^a, D. Pagnozzi^a, H. Slavski^b,
S. Uzzau^{a, c}, R. Anedda^{a, **}, M.F. Addis^{a, d, *}^a Porto Conte Ricerche, SP 55 Porto Conte/Capo Caccia, Km 8.400, Loc. Tramarglio, 07041 Alghero, Italy^b Aller Aqua, Allervej 130, DK-6070, Christiansfeld, Denmark^c Università degli Studi di Sassari, Dipartimento di Scienze Biomediche, Viale S. Pietro 43/B, 07100 Sassari, Italy^d Università degli Studi di Milano, Dipartimento di Medicina Veterinaria, Via Celoria 10, 20133 Milano, Italy

ARTICLE INFO

Article history:

Received 16 April 2019

Received in revised form 29 July 2019

Accepted 12 August 2019

Available online 19 August 2019

Keywords:

Cold stress

Gilthead sea bream

Low temperature exposure

Liver proteins

Shotgun proteomics

Liver proteomic dataset

ABSTRACT

We report the proteomic dataset of livers from *Sparus aurata* exposed to low temperature during growth. Gilthead sea bream juveniles were reared in Recirculating Aquaculture Systems (RAS) and exposed to a temperature ramp made of two phases of four weeks each: a Cooling phase from 18 °C (t0) to 11 °C (t1) and a Cold Maintenance phase at 11 °C (t1-t2) in a 8 week feeding trial. At the end of the experiment, sea bream livers were collected and analyzed with a shotgun proteomics approach based on filter-aided sample preparation followed by tandem mass spectrometry, peptide identification carried out using Sequest-HT as search engine within the Proteome Discoverer informatic platform, and label-free differential analysis.

The mass spectrometry data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD011059 (Vizcaíno et al., 2016; Deutsch et al., 2017; Perez-Riverol et al., 2016). The dataset described here is also related to the research article entitled "Liver proteomics of gilthead sea bream (*Sparus aurata*) exposed to cold stress" (Ghisaura et al., 2019).

© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <https://doi.org/10.1016/j.jtherbio.2019.04.005>.

* Corresponding author. Porto Conte Ricerche, SP 55 Porto Conte/Capo Caccia, Km 8Loc. Tramarglio, 07041 Alghero, Italy.

** Corresponding author.

E-mail addresses: anedda@portocontericerche.it (R. Anedda), filippa.addis@unimi.it (M.F. Addis).

<https://doi.org/10.1016/j.dib.2019.104419>

2352-3409/© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	<i>Biology</i>
More specific subject area	<i>Proteomics</i>
Type of data	Tables
How data was acquired	Q-TOF hybrid mass spectrometer equipped with a nano lock Z spray source and coupled on-line with a NanoAcquity chromatography system (Waters)
Data format	Raw, processed
Experimental factors	Proteome analysis of gilthead sea bream livers during exposure to low temperatures: Cooling phase from 18 °C (t0) to 11 °C (t1); Cold maintenance Phase at 11 °C (t2)
Experimental features	1) Protein extraction (mechanical disruption in TUC-based buffer) 2) Filter-aided sample preparation (FASP) 3) LC-MS/MS analysis
Data source location	Tramariglio, Alghero (SS), Italy; Torregrande (OR), Italy
Data accessibility	Data is within this article and available via the ProteomeXchange Consortium, dataset identifier PXD011059
Related research article	Ghisaura S, Pagnozzi D, Melis R, Biosia G, Slawski H, Uzzau S, Anedda R, Addis MF. Liver proteomics of gilthead sea bream (<i>Sparus aurata</i>) exposed to cold stress. J Therm Biol 2019; 82:234–41. https://doi.org/10.1016/j.jtherbio.2019.04.005 . Author names: S.Ghisaura; D.Pagnozzi; R.Melis; G.Biosia; H.Slawski; S.Uzzau; R.Anedda; M.F.Addis; Title: Liver proteomics of gilthead sea bream (<i>Sparus aurata</i>) exposed to cold stress; Journal: Journal of Thermal Biology

Value of the data

- Detailed proteomic dataset of gilthead sea bream livers in fish exposed to low temperature during growth;
- Differential protein abundances between the two temperature phases: Cooling phase (t0-t1, from 18 °C to 11 °C) and Cold Maintenance phase (t1-t2, at 11 °C) are useful to understand the dynamics and metabolic shifts occurring in sea bream liver with decreasing water temperature;
- Shotgun proteomics dataset improves previous data on fish hepatic metabolism during cold exposure;
- The proteomic dataset might be advantageous to other research groups working on the development of feeds designed to compensate the thermal stress encountered by fish in offshore farming conditions.

1. Data

Sea bream livers exposed to low temperatures (Cooling phase and Cold maintenance phase) were characterized with a shotgun proteomic approach. A summary of protein identifications obtained in all samples is provided in [Table 1](#). All protein identifications obtained with the Proteome Discoverer software in gilthead seabream livers exposed to three temperature phases (t0, t1, t2) are listed in [Supplementary Table 1](#).

The differential analysis was carried out with a label-free approach by comparing all different groups according to temperature variations: Cooling phase (t0-t1), Cold Maintenance phase (t1-t2) and Overall changes (t0-t2). The differential proteins passing the significance thresholds ($R_{NSAF} > 0.5$ or < -0.5 ; P value < 0.05 ; $FDR < 0.1$) are summarized in detail in the related research article [\[4\]](#). Detailed protein identification and abundance data for the three comparisons (t0 vs t1; t1 vs t2 and t0 vs t2) are provided in [Supplementary Table 1](#).

2. Experimental design, materials, and methods**2.1. Sparus aurata liver samples**

Gilthead sea bream specimens with an average weight of 82.0 ± 4.5 g were selected for the experimental feeding trial. A total of 60 juveniles were transferred in three 550 L tanks and an

Table 1

Summary of protein identifications obtained in all samples.

	t0	t1	t2
#Proteins	649	620	654
#Proteins with PSM ≥ 2	510	458	487
#Peptides	1491	1328	1448
# PSM(s) ^a	5176	4274	4837
#Search inputs ^b	19391	19025	20644
#Total Spectra ^c	22036	21607	23302

^a The number of peptide spectrum matches obtained for protein identification.^b The number of preprocessed spectra by the Spectrum Selector node in the workflow used in Proteome Discoverer software (precursor mass range: 350–5000 Da; Signal/Noise Threshold: 1.5).^c The total number of spectra obtained by LC-MS/MS run.

acclimation phase of two weeks from 20 °C to 18 °C was carried out. Water temperature was gradually lowered as described in Ghisaura et al., 2019 [4]. During the trial, fish were fed with an experimental feed formulation (Aller Aqua, Christiansfeld, Denmark) by hand, once a day. After fish anesthetization with 1,1,1-trichloro-2-methylpropan-2-ol (2% in marine water) and transfer in a mixture of marine water and ice, liver tissues were collected at each time point (t0, t1, t2). The complete procedure of tissue excision and storage is described by Melis et al., 2017 [5].

2.2. Protein extraction and digestion

Liver tissues were subjected to protein extraction and quantification according to Ghisaura et al., 2016 [6]. All protein extracts were then subjected to on-filter reduction, alkylation, and trypsin digestion according to the filter-aided sample preparation (FASP) protocol [7], with some modifications [8]. Peptide mixture concentration was estimated by using the BCA protein assay kit (Thermo Scientific - Rockford, IL).

2.3. LC-MS/MS analysis

A Q-TOF hybrid mass spectrometer with a nano lock Z spray source, coupled with a NanoAcquity chromatography system (Waters) on-line, was used for LC-MS/MS analyses as described in Pagnozzi et al., 2014 [9]. LC-MS/MS procedures are fully described in Ghisaura et al., 2019 [4].

2.4. Data analysis

Proteome Discoverer software (version 1.4.0.288; Thermo Scientific) was used to analyze the peak lists from the Q-TOF instrument, after conversion into an MGF file. The workflow was as described in Ghisaura et al., 2019 [4]. Gene ontology and protein annotations were retrieved from UniProtKB (<http://www.uniprot.org>). The uncharacterized sequences were identified by homology through blasting on NCBI as another non-redundant database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Differential protein abundances of different functional categories (Cooling phase; Cold maintenance phase; and Overall changes) were estimated by the Normalized Spectral Abundance Factor (NSAF) according to Zybailov et al., 2006 [10]. The significance threshold $RNSAF > 0.5$ or < -0.5 was applied for analysis. To evaluate the statistical significance of differential protein abundance between logarithmized (normally distributed) NSAF values, a student's t-test (two-sample comparison, $p < 0.05$) was applied. Logarithmized NSAF values were furthermore corrected by using a false discovery rate (FDR) as a multiple hypothesis testing, with $FDR < 0.1$ as a threshold limit. The dataset was then deposited in the ProteomeXchange Consortium via the PRIDE partner repository (identifier PXD011059) [1,2,3,11].

Acknowledgments

The PRIDE team is acknowledged for the support for MS data deposition into ProteomeXchange (identifier PXD011059). This work was funded by the Sardinia Regional Government by means of Sardegna Ricerche (art. 26 L.R. 37/98).

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.dib.2019.104419>.

References

- [1] J.A. Vizcaíno, A. Csordas, N. Del-Toro, J.A. Dianes, J. Griss, I. Lavidas, et al., 2016 Update of the PRIDE database and its related tools, *Nucleic Acids Res.* 44 (2016) D447–D456, <https://doi.org/10.1093/nar/gkv1145>.
- [2] E.W. Deutsch, A. Csordas, Z. Sun, A. Jarnuczak, Y. Perez-Riverol, T. Ternent, et al., The ProteomeXchange consortium in 2017: supporting the cultural change in proteomics public data deposition, *Nucleic Acids Res.* 45 (2017) D1100–D1106, <https://doi.org/10.1093/nar/gkw936>.
- [3] Y. Perez-Riverol, Q.-W. Xu, R. Wang, J. Uszkoreit, J. Griss, A. Sanchez, et al., PRIDE inspector toolsuite: moving toward a universal visualization tool for proteomics data standard formats and quality assessment of ProteomeXchange datasets, *Mol. Cell. Proteom.* 15 (2016) 305–317, <https://doi.org/10.1074/mcp.O115.050229>.
- [4] S. Ghisaura, D. Pagnozzi, R. Melis, G. Biosia, H. Slawski, S. Uzzau, et al., Liver proteomics of gilthead sea bream (*Sparus aurata*) exposed to cold stress, *J. Therm. Biol.* 82 (2019) 234–241, <https://doi.org/10.1016/j.jtherbio.2019.04.005>.
- [5] R. Melis, R. Sanna, A. Braca, E. Bonaglini, R. Cappuccinelli, H. Slawski, et al., Molecular details on gilthead sea bream (*Sparus aurata*) sensitivity to low water temperatures from 1H NMR metabolomics, *Comp. Biochem. Physiol. Part A Mol Integr Physiol* 204 (2017) 129–136, <https://doi.org/10.1016/j.cbpa.2016.11.010>.
- [6] S. Ghisaura, B. Loi, G. Biosia, M. Baroli, D. Pagnozzi, T. Roggio, et al., Proteomic changes occurring along gonad maturation in the edible sea urchin *Paracentrotus lividus*, *J. Proteomics* 144 (2016) 63–72, <https://doi.org/10.1016/j.jprot.2016.05.035>.
- [7] J.R. Wiśniewski, A. Zougman, N. Nagaraj, M. Mann, Universal sample preparation method for proteome analysis, *Nat. Methods* 6 (2009) 359–362, <https://doi.org/10.1038/nmeth.1322>.
- [8] A. Tanca, G. Biosia, D. Pagnozzi, M.F. Addis, S. Uzzau, Comparison of detergent-based sample preparation workflows for LTQ-Orbitrap analysis of the *Escherichia coli* proteome, *Proteomics* 13 (2013) 2597–2607, <https://doi.org/10.1002/pmic.201200478>.
- [9] D. Pagnozzi, G. Biosia, M.F. Addis, S. Mastrandrea, G. Masala, S. Uzzau, An easy and efficient method for native and immunoreactive *Echinococcus granulosus* antigen 5 enrichment from hydatid cyst fluid, *PLoS One* 9 (2014) e104962, <https://doi.org/10.1371/journal.pone.0104962>.
- [10] B. Zybailov, A.L. Mosley, M.E. Sardu, M.K. Coleman, L. Florens, M.P. Washburn, Statistical analysis of membrane proteome expression changes in *Saccharomyces cerevisiae*, *J. Proteome Res.* 5 (2006) 2339–2347, <https://doi.org/10.1021/pr060161n>.
- [11] J.A. Vizcaíno, E.W. Deutsch, R. Wang, A. Csordas, F. Reisinger, D. Ríos, et al., ProteomeXchange provides globally coordinated proteomics data submission and dissemination, *Nat. Biotechnol.* 32 (2014) 223–226, <https://doi.org/10.1038/nbt.2839>.