


ORIGINAL ARTICLE

Beyond the kidney biopsy: genomic approach to undetermined kidney diseases

Thomas Robert ^{1,2}, Laure Raymond³, Marine Dancer³, Julia Torrents⁴, Noémie Jourde-Chiche^{1,5}, Stéphane Burtey^{1,5}, Christophe Bérout² and Laurent Mesnard⁶

¹Centre of Nephrology and Renal Transplantation, Hôpital de la Conception, CHU de Marseille, Marseille, France, ²Marseille Medical Genetics, Bioinformatics & Genetics, INSERM U1251, Aix-Marseille Université, Marseille, France, ³Genetics Department, Laboratoire Eurofins Biomnis, Lyon, France, ⁴Department of Renal Pathology, CHU Timone, AP-HM, Marseille, France, ⁵Aix-Marseille Univ, INSERM, INRAE, C2VN, Marseille, France and ⁶Urgences Néphrologiques et Transplantation Rénale, Sorbonne Université, APHP, Hôpital Tenon, Paris, France

Correspondence to: Thomas Robert; E-mail: thomas.robert@ap-hm.fr

ABSTRACT

Background. According to data from large national registries, almost 20%–25% of patients with end-stage kidney disease have an undetermined kidney disease (UKD). Recent data have shown that monogenic disease-causing variants are under-diagnosed. We performed exome sequencing (ES) on UKD patients in our center to improve the diagnosis rate.

Methods. ES was proposed in routine practice for patients with UKD including kidney biopsy from January 2019 to December 2021. Mutations were detected using a targeted bioinformatic customized kidney gene panel (675 genes). The pathogenicity was assessed using American College of Medical Genetics guidelines.


Results. We included 230 adult patients, median age 47.5 years. Consanguinity was reported by 25 patients. A family history of kidney disease was documented in 115 patients (50%). Kidney biopsies were either inconclusive in 69 patients (30.1%) or impossible in 71 (30.9%). We detected 28 monogenic renal disorders in 75 (32.6%) patients. Collagenopathies was the most common genetic kidney diagnosis (46.7%), with *COL4A3* and *COL4A4* accounting for 80% of these diagnoses. Tubulopathies (16%) and ciliopathies (14.7%) yielded, respectively, the second and third genetic kidney diagnosis category and *UMOD*-associated nephropathy as the main genetic findings for tubulopathies (7/11). Ten of the 22 patients having ES “first” eventually received a positive diagnosis, thereby avoiding 11 biopsies. Among the 44 patients with glomerular, tubulo-interstitial or vascular nephropathy, 13 (29.5%) were phenocopies. The diagnostic yield of ES was higher in female patients ($P = .02$) and in patients with a family history of kidney disease ($P < .0001$), reaching 56.8% when the patient had both first- and second-degree family history of renal disease.

Conclusion. Genetic diagnosis has provided new clinical insights by clarifying or reclassifying kidney disease etiology in over a third of UKD patients. Exome “first” may have a significant positive diagnostic yield, thus avoiding invasive kidney biopsy; moreover, the diagnostic yield remains elevated even when biopsy is impossible or inconclusive. ES provides a clinical benefit for routine nephrological healthcare in patients with UKD.

Received: 19.12.2022; Editorial decision: 1.4.2023

© The Author(s) 2023. Published by Oxford University Press on behalf of the ERA. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

GRAPHICAL ABSTRACT



Keywords: chronic kidney disease, exome sequencing, undetermined nephropathy

INTRODUCTION

Chronic kidney disease (CKD) is a public health problem that is associated with high costs and morbidity. Despite kidney biopsy, different national registries have established that 20%–25% of patients with end-stage kidney disease (ESKD) have an undetermined kidney disease (UKD) [1–3]. In this category, several studies have already suggest an enrichment of monogenic/Mendelian kidney diseases with causative variants [4]. Despite rare Mendelian diseases beginning to appear more frequently than expected in the CKD population, their molecular diagnosis remains challenging, especially given the paucity of the phenotype. In France, as in many developed countries, a national effort to facilitate whole-genome sequencing as part of the routine healthcare process for different disease entities is moving forward. Despite this, few institutions can currently provide exome sequencing (ES) as a first-line clinical diagnostic tool in nephrology practice.

Patients with UKD represent a significant proportion of CKD patients despite there being no clear definition of UKD at present. Since 2012, the national Renal Epidemiology and Information Network (REIN) registry in France has provided an overview of incidental and prevalent patients with ESKD. REIN data show that 20% of registry patients have a UKD, while 13% are still labelled as “other.” A recent Kidney Disease: Improving Global Outcomes (KDIGO) conference suggested that UKD could be an indication for genetic screening. This would be

especially beneficial if a kidney biopsy is neither informative nor possible because the patient is at too advanced a stage and there are no indications of an hereditary disease. This approach could reduce costly diagnostic workups for those patients [4, 5]. Genetic screening may even be the only available tool when newly diagnosed ESKD is discovered in adult patients, sometimes resolving atypical cases. And last, but not least, the analysis of a local registry of kidney biopsies (NéphroMIC², NCT03305211) revealed that around 20% of patients still have inconclusive kidney biopsies, thus remaining within the UKD category (data not published). Since 2019, ES has been proposed for all UKD patients in our center.

A small number of studies support the implementation of ES in the diagnostic workup of UKD, particularly if the patient is waiting for kidney transplantation, or if early onset or familial nephropathy is observed. Whether those findings can be extended to the UKD population or, more especially, to sporadic cases, is unclear. Our strategy is based on an in-silico panel analysis of known genes related to kidney diseases (Supplementary data, Table S1). To address the usefulness of ES to resolve UKD in routine care, we reported our experience.

MATERIALS AND METHODS

Patient cohort and phenotype characterization

ES was proposed as part of routine clinical care in patients with UKD from January 2019 to December 2021. UKD is defined as




Figure 1: Diagnostic workflow for undetermined kidney diseases selection.

the absence of any of the following criteria: biopsy-proven diagnosis [e.g. immunoglobulin A (IgA) nephropathy], absence of a specific morphological renal diagnosis (e.g. polycystic kidney disease) or absence of a specific or plausible renal diagnosis (such as, history of long-term insulin-dependent diabetes mellitus before the onset of CKD, cyclosporin-induced nephropathy). Patients with kidney disease who underwent ES before kidney biopsy were included in the analysis and considered as the exome “first” approach. Since hypertensive nephropathy is a non-specific diagnosis, and hypertension is also a very common consequence of CKD, patients with hypertensive nephropathy in the absence of a clear underlying disorder, such as renal artery stenosis, are considered to have unexplained CKD. Patients with

renal hypoplasia, renal atrophy and nonspecific histological conditions (such as secondary focal segmental glomerulosclerosis, glomerulonephritis of unknown origin or interstitial nephritis) are also considered to have UKD. We excluded patients with familial IgA nephropathy, patients with typical presentation of Gitelman or Bartter syndrome, or an established kidney-related genetic diagnosis in the family. None of the 230 index cases had a genetic diagnosis and therefore all were considered as unsolved (Fig. 1). If prior genetic analyses were performed, they were at the discretion of the referring clinician. Patients with UKD both from our center and other tertiary care centers were referred for a nephrogenomic consultation with one of our adult nephrologists. Phenotypes were acquired by using a standardized questionnaire and review of medical reports during

Table 1: Demographic characteristics of the whole cohort.

Variable ^a	Patients (n = 230)
Age at ES (years)	47.5 [36.8–62]
Age at kidney disease onset or discovery	35 [23–50]
Male	133 (57.8)
Geographic origin	
Europe	173 (75.2)
North Africa	31 (13.5)
Sub-Saharan Africa	12 (5.2)
French Antilles	7 (3.1)
Asia	2 (0.01)
Others	5 (2.2)
Consanguinity	25 (10.9)
Kidney disease onset before 35 years old	126 (54.8)
With familial history	50 (21.7)
Familial history of kidney disease	115 (50)
First degree only	58 (25.2)
Second degree only	19 (8.3)
Both first and second degree	37 (16.2)
Prior negative genetic exploration with gene panel	34 (14.7)
ES first to avoid kidney biopsy	22 (9.5)
Prior inconclusive kidney biopsy	69 (30.1)
Kidney disease discovered at ESKD	43 (18.7)
Impossible kidney biopsy	71 (30.9)
Undetermined clinical nephropathy subgroup	
Unclassified	88 (38.3)
Glomerular	64 (27.8)
Tubulointerstitial	30 (13)
Cyst	32 (13.9)
Vascular	16 (7)
Stage of kidney disease at ES	
I	32 (13.9)
II	13 (5.7)
III	39 (17)
IV	32 (13.9)
V	22 (9.5)
V (D)	54 (23.5)
V (T)	38 (16.5)
Status of patient transplantation	80 (34.5)
Active in transplant waiting list	16 (6.9)

^aFor quantitative variables, values are expressed as median [interquartile ranges]. For qualitative variables, values are expressed as n (%).

consultation. Consanguinity was also established during consultation either as reported by the patient or as suspected by the clinician. Patients were assigned to four subgroups of undetermined nephropathy: unclassified nephropathy; undetermined glomerular nephropathy; undetermined tubulointerstitial nephropathy; and undetermined vascular nephropathy. We defined a phenocopy as a patient having a phenotype that corresponded to a specific clinical nephropathy (such as glomerular nephropathy) without detection of the expected genotype (e.g. a glomerular gene such as *TRPC6*) but with detection of a different genotype (for example, a tubulointerstitial gene such as *CLCN5*).

Blood samples were collected after written informed consent from the patients or their legal guardians during the consultation. All patients gave their written informed consent for genetic testing.

Exome sequencing

DNA was extracted from peripheral blood using the QIAAsymphony[®] DSP DNA Mini Kit on a QIAAsymphony instrument following the manufacturer's (QIAGEN) guidelines.

From 50 ng of fragmented DNA, indexed libraries were prepared and hybridized with a biotinylated probe from Twist Human Core Exome (33 Mb) and, from April 2019, Twist Human Comprehensive Exome (37 Mb). ES was performed on the Illumina NextSeq 500 in paired-end mode (2 × 75 bp reads) then, from March 2021, NextSeq 2000 platform in paired-end mode (2 × 150 bp reads) on FlowCell P3. Raw data (bcl format) were converted to FASTQ format using the Dragen software sequencer (Illumina). Reads were aligned to the human reference genome (UCSC Genome Browser build hg19). Sequences were analyzed according to GATK Broad Institute good practice with two pipelines: Intern pipeline (BWA-MEM, GATK v3.6–44ge7d1cd2) and SeqOne pipeline (v1.2, 2018). Copy number variants (CNVs) calls were performed using the GATK4 CNV calling module and were validated using Multiplex Ligation-dependent Probe Amplification [6].

To identify diagnostic variants, we assessed the pathogenicity of the variant using American College of Medical Genetics (ACMG) guidelines [7]. Variants were filtered according to coverage level (DP > 10), allelic frequency (>20%) and protein effect. Frequency of variants in GnomAd was also considered: for the analysis of *de novo*, autosomal dominant, autosomal recessive and X-linked variants, only variants with a minor allele frequency (MAF) <1% in the GnomAd database were considered. For the analysis of autosomal recessive and X-linked variants (homozygous, hemizygous or putative compound heterozygous) in unsolved cases, additional research with an MAF up to 3% was considered. In addition, we assessed the *APOL1* genotype [G1 (rs73885139 and rs60910145) and G2 (rs71785313)] as Mendelian forms of nephropathy when two copies were present as follows: G1/G1, G1/G2 or G2/G2 [8]. Identified variants were compared with available databases for pathogenic variants such ClinVar, the Human Gene Mutation Database (HGMD) [9] and the Leiden Open Variation Database (LOVD), and databases for pathogenic CNVs such as DECIPHER. Only variants rated as “likely pathogenic” or “pathogenic” according to the ACMG classification, and with a genotype in agreement with the mode of inheritance and the phenotype, led to a positive ES result. Patients with variants classified as benign, likely benign or of unknown significance according to ACMG led to a negative exome result. ES results were communicated to the patients by the same nephrologist with whom they had had the initial nephrogenomic consultation.

Statistical analysis

Baseline characteristics were expressed as frequencies (n, %), means [standard deviations (SDs)] and medians (range). Fisher's exact test was performed for categorical data. Diagnostic yield was calculated based on counts of variants classified as “pathogenic” or “likely pathogenic.” To compare two continuous variables, we used normality tests and then an unpaired t-test or unpaired Mann–Whitney nonparametric test if values were not sampled from Gaussian distribution. P-values <.05 were considered statistically significant. Two-tailed P-values <.05 were regarded as statistically significant. Statistical analyses were performed using Prism 9 (GraphPad Software) software.

RESULTS

Characteristics of the population

Two hundred and thirty-two adult patients were included (133 males) with a median age of 47.5 years, and 173 were




Figure 2: Gene-associated kidney disease according to the category of genetic kidney disease. The inner circle represents the categories of genetic kidney disease and in the outer circle, gene reported with a pathogenic variant in patients; numbers in brackets represent the number of patient's carrier. Genes have been clustered according to their associated categories of genetic kidney disease.

self-declared European. Of these, 54 patients were treated with hemodialysis and 38 patients had a kidney transplantation. Consanguinity was reported or suspected in 25 patients, and 126 had kidney disease onset before the age of 35 years. A family history of kidney disease was present in 115 patients and 37 patients had both first and second degree with history of kidney disease. Kidney biopsies were inconclusive in 69 patients and impossible in 71 patients. An ES was performed first to avoid kidney biopsy in 22 patients. A negative genetic testing with gene panel prior to the ES was performed in 34 patients. Finally, 88 patients had unclassified kidney disease (Table 1).

Diagnostic yield of the ES

We identified 28 monogenic renal disorders in 75 patients carrying either pathogenic or likely pathogenic variants [single nucleotide variants (SNV)/small indels; $n = 70$] or CNVs ($n = 5$) among 230 patients (32.6%) (Supplementary data, Table S2). Collagenopathies were the major genetic kidney disease category with 46.7% (COL4A4 $n = 15$; COL4A3 $n = 13$; COL4A5 $n = 7$), followed by tubulopathies with 16% (UMOD $n = 7$; CLCN5 $n = 2$; HNF1B $n = 1$; KCNJ1 $n = 1$; SLC5A2 $n = 1$). Ciliopathies (TTC21B $n = 3$; NPHP1 $n = 2$; DNAJB11 $n = 2$; NPHP3 $n = 1$; TULP3 $n = 1$) and podocytopathies (INF2 $n = 2$; PAX2 $n = 2$; APOL1 $n = 2$; TRPC6 $n = 1$; NPHS1 $n = 1$) represent, respectively, the third and fourth majors

genetic kidney disease category with 14.7% and 10.7% (Fig. 2A). Finally, when ES was performed first to avoid renal biopsy, 11 of the 22 patients (50%) had a positive genetic finding [9 collagenopathies, 1 APOL1 nephropathy and 1 ciliopathy-gene associated disease (TTC21B)] (Table 2).

Relationship between genotype and nephropathy category in ES solved cases

Glomerular and unclassified nephropathy had the highest ES-solved case rates in, respectively, 39.1% and 35.2%, whilst tubulointerstitial and vascular nephropathy had the lowest ES-solved case rates (respectively, 23.3% and 18.8%) (Fig. 3A). Among the 44 patients who had their nephropathy categorized, 13 (29.5%) were phenocopies (Fig. 3B and Supplementary data, Fig. 1). Vascular (2/3) and tubulointerstitial nephropathy (4/7) had the highest phenocopy rates, at 66.3% and 42.9%, respectively. Among glomerular nephropathy, 16% (4/24) were phenocopies. Cystic kidney disease and congenital anomalies of the kidneys and urinary tract (CAKUT) had a phenocopy rate of 22.2% (2/9) (Fig. 3B). Collagenopathies (COL4A3 and COL4A4), ciliopathy category (TTC21B, TULP3) and tubulopathies (UMOD, CLCN5) show the highest phenocopy rates (Supplementary data, Fig. S2). Of the unclassified nephropathies, the largest proportions were due to

Table 2: Comparison of characteristics between exome-solved and -unsolved cases.

Variable ^a	Unsolved (n = 155)	Solved (n = 75)	P-value
Age at ES (years)	46 [36–62]	50 [40–63]	.25
Age at kidney disease onset or discovery	37 [24–50]	32 [20–49]	.26
Male	98 (63.2)	35 (46.7)	.02
Geographic origin			.2
Europe	116 (74.8)	57 (76)	
North Africa	17 (11)	14 (18.6)	
Sub-Saharan Africa	10 (6.5)	2 (2.7)	
French Antilles	5 (3.2)	2 (2.7)	
Asia	2 (1.3)	0	
Others	5 (3.2)	0	
Consanguinity	13 (8.5)	12 (16)	.1
Kidney disease onset before 35 years old	81 (52.3)	45 (60)	.3
With familial history	26 (16.8)	24 (32)	.01
Familial history of kidney disease	63 (40.6)	52 (69.3)	.0001
First degree only	33 (21.3)	25 (33.3)	.05
Second degree only	13 (8.4)	6 (8)	.9
Both first and second degree	16 (10.3)	21 (28)	.001
Prior negative genetic exploration with gene panel	22 (14.2)	12 (16)	.7
ES first to avoid kidney biopsy	11 (7.1)	11 (14.7)	.09
Prior inconclusive kidney biopsy	45 (29.2)	24 (32)	.7
Impossible kidney biopsy	48 (31)	23 (30.7)	1
Undetermined clinical nephropathy subgroup			
Unclassified	57 (36.8)	31 (41.4)	
Glomerular	39 (25.2)	25 (33.3)	
Tubulointerstitial	23 (14.8)	7 (9.3)	.3
Cyst	23 (14.8)	9 (12)	
Vascular	13 (8.4)	3 (4)	
Stage of kidney disease at ES			
I	22 (14.2)	10 (13.3)	
II	10 (6.5)	3 (4)	
III	28 (18.1)	11 (14.7)	
IV	20 (12.9)	12 (16)	.3
V	16 (10.3)	6 (8)	
V (D)	39 (25.2)	15 (20)	
V (T)	20 (12.9)	18 (24)	

collagenopathies (61.3%) and tubulopathies (22.6%) (Fig. 3B and Supplementary data, Fig. 3).

In biopsy-associated UKD patients, the pathologist's conclusion about the main histological lesion was consistent with the genetic findings for all glomerular and tubulo-interstitial diseases. In patients whose pathologist concluded that nephroangiosclerosis was the primary histological lesion, ES revealed two tubulopathies—one collagenopathy and one vasculopathy associated with the *TREX1* gene (Fig. 4). The most frequent causes of global fibrosis in the kidneys were found to be either a collagenopathy (33.3%) or a ciliopathy (33.3%) (Fig. 4).

Clinical predictors of positive exome diagnosis for kidney disease

The ES diagnostic yield was higher in female patients (54.5% vs 47%, $P = .01$) and those with a familial history of kidney disease (69.2% vs 40.4%, $P < .0001$) (Table 2 and Fig. 5A). Out of patients with a family history of kidney disease in both first and second degree, 56.8% had a positive ES (Fig. 5B). Younger age at onset of the disease (<35 years) was not significantly associated with a positive ES ($P = .3$) except for those with positive family history of kidney disease (32% vs 16.8%, $P = .01$) (Table 2).

DISCUSSION

In the present UKD cohort, implementation of ES in routine healthcare allowed us to resolve diagnosis in one-third of patients (32.6%). Previous studies provided a diagnostic yield of genetic testing in patients with CKD, in the absence of a specific suspicion of monogenic kidney disease, of between 10% and 30% in adults [4, 5, 10] and between 30% and 50% for pediatric cohorts [11–14]. Our study illustrates the underestimation of the prevalence of inherited case. CNV accounts for 6.7% of genetic findings, suggesting that a combination of SNV and CNV detection is suitable for genomic analysis prior to concluding that ES is negative and that whole-genome sequencing is necessary [6]. Interestingly, the Groopman study was unable to detect CNVs, suggesting that the proportion of inherited kidney disease among its participants is likely greater than was reported [4]. CKD affects 11%–13% of the world population [15], but UKD accounts for approximately 20% of CKD patients in developed countries. The application of ES to nephrology, and especially to UKD, has the potential to revolutionize the way kidney diseases are classified, taking into account not only clinical features and kidney biopsy, but also the underlying pathogenesis of the disease. This could lead to a much more accurate and comprehensive reclassification and ontology of kidney diseases.




Figure 3: Genetic findings according to nephropathy category. (A) The diagnostic yielded according to the clinical nephropathy category. (B) The distribution of the genetic kidney disease category according to the clinical nephropathy category.

In selecting a panel of genes for testing, geneticists traditionally take a phenotypically driven approach. This is referred to as the “candidate gene genetics” method and is responsible for most of the human gene–disease associations that have so far been discovered with penetrant phenotypes. In recent years, and with the ease of access to NGS, a gene-oriented approach has emerged which consists of beginning with the variant in a gene and looking back to the patient’s phenotype. This method is called “reverse phenotyping” and seems particularly interesting in the context of UKD patients. This approach is especially valuable in genetically related kidney diseases that may remain completely undiagnosed until ESKD [16].

Glomerular diseases, most particularly collagenopathies, are the major genetic findings in our cohort, which differs from the pediatric and young adult population where CAKUT constitutes the major clinical diagnostic group (50% of cases) [17]. In our elderly population (most of whom were over 30 years old), nearly half of the genetic findings were a collagen type IV nephropathy, demonstrating wide variability in phenotype among patients with mutations affecting type IV collagen. Interestingly, mutations in the *COL4A3* and *COL4A4* genes accounted for 80% of the collagenopathies, which is much higher than previously reported [18]. This suggests that collagen IV nephropathy associated with these two genes is underdiagnosed. Of the patients

with a pathogenic variation in either *COL4A3* or *COL4A4*, 78.6% had a stage of chronic kidney disease below stage 3 chronic kidney disease and may have been classified as having “thin basement membrane disease,” which does not accurately reflect this mild condition. This finding supports the recent consensus report which proposed that all individuals with mutations in either the *COL4A3* or *COL4A4* gene should be diagnosed as having Alport Syndrome [19]. This suggests that the accurate clinical diagnosis of monogenic kidney diseases in elderly CKD patients is challenging due to their high phenotypic overlap, extensive genetic heterogeneity and the presence of comorbidities. This result also confirms that collagenopathies are the major causative genes of monogenic CKD in a UKD population [4]. Tubulopathies was the second genetic kidney disease category with *UMOD* as the next main genetic finding in our study and ciliopathy accounted for the third genetic kidney disease category. This confirms that this locus is one of those most strongly associated with CKD [20], representing one of the most common monogenic kidney diseases with an overall prevalence of ~2% in patients with renal failure [21, 22].

Among patients with unclassified nephropathy, collagenopathies and tubulopathies are the major findings, highlighting the clinical heterogeneity and variable expression of gene-associated phenotype. Our result confirms Groopman’s




Figure 4: Distribution of genetic kidney disease categories based on kidney biopsy findings.

findings, with 45% and 29.4% of glomerular and TI-gene associated disease, respectively, in patients with nephropathy of unknown origin. The former should be more accurately termed unclassified nephropathy as Gropman *et al.* classified patients by clinical diagnosis subgroup (glomerulopathy, congenital or cystic kidney disease, hypertensive nephropathy, diabetic nephropathy, tubulointerstitial disease and nephropathy of unknown origin) [4]. As recommended by KDIGO, there is an urgent need to clarify UKD definition, which is referred to using different terms: unexplained kidney disease, undetermined kidney disease, nephropathy of unknown origin. We propose that UKD is a clinical diagnostic group that should be based on clinical findings, renal morphology, biological parameters and, if available, renal histology. Undetermined kidney disease can thus be classified as being glomerular, tubulointerstitial, vascular, cystic or related to a renal development abnormality. If there is not a strong enough argument bundle, the undetermined kidney disease remains unclassifiable.

Our study reveals that nearly one-quarter of the classifiable UKDs are phenocopies with a wide range of clinical presentations. This suggests that whole-exome sequencing is more suitable than gene panels in UKD [23]. However, the pathologist's conclusion can be used to help select the most suitable gene panel when kidney biopsy is possible. Ottlewski *et al.* observed a 12% genetic diagnostic yield using a 200-gene panel in a UKD cohort [5]. In comparison, Schrezenmeier and colleagues reported a higher yield of 20% using a larger 600-gene panel [24]. All these results suggest that, given the regular discovery of new genes associated with UKD and the need for rapid and cost-effective sequencing, whole-exome sequencing with a bioinformatic gene panel is advantageous. In our center, we follow the ACMG guidelines in reanalyzing ES data for each newly discovered gene-

associated kidney disease. To this end, we conduct periodically a literature review to identify newly published genes and incorporate them into our bioinformatics panel and we then reanalyze all unsolved ES. As an example, index case 240 displaying *TULP3* variants was initially classified as negative. Upon the discovery of *TULP3* as a ciliopathy-associated disease gene, we were able to reanalyze all sample displaying rare variants in *TULP3* with their phenotype, leading to the reclassification of the original case as positive for ciliopathy [25, 26]. The *TULP3* gene was incorporated into our custom kidney gene panel. We have included *APOL1* in our gene panel but the current clinical use of *APOL1* genotyping has been widely discussed, and there is still no clear agreement among clinicians. Identifying the triggers of *APOL1* kidney disease can be challenging. Possible triggers include infection, interferon activation, environmental kidney damage, superimposed lupus, diabetes and arterial hypertension. However, determining the direction of causality between *APOL1* kidney disease and diabetes or arterial hypertension is particularly difficult. Given that many biotechnology and academic groups are now looking into potential therapies for *APOL1* kidney disease, we have chosen to consider *APOL1*-related nephropathy as resolved in our study and as suggested by a recent review paper [27].

In our experience, ES avoided renal biopsy in 12 of 23 selected patients (52.2% diagnostic yield rate) at high suspicion of hereditary kidney disease. Alport disease represented the major finding. Renal biopsy is the traditional diagnostic approach, but it may either reveal nonspecific findings or be completely contraindicated. In addition, unlike in the USA or UK, electron microscopy is not readily available in routine care in France, and its use is highly restricted to a few indications. ES represents a non-invasive new tool that should be included in the diagnostic workup of nephropathy, either when biopsy




Figure 5: ES yielded according family history of kidney disease. (A) Comparison of the prevalence of kidney disease in relatives of positive and negative ES patients; panel (B) shows in positive ES patients the proportion of familial history of kidney disease by degree in the relatives.

features are nonspecific or when hereditary kidney disease is suspected and renal biopsy has not been helpful [28]. ES could replace renal biopsy as the gold standard in challenging situations such as pregnancy, in patients with atrophic kidney, or in the case of a high suspicion of inherited kidney disease (glomerular presentation and familial history, for example). With expanding indications of genetic tests, registries will be essential for improving our understanding of genetic kidney disease [29].

We observed that among patients with classified nephropathy, almost one-third were phenocopies, with collagenopathies, tubulopathies and ciliopathies being the most common contributors among all nephropathy categories. This proportion is slightly higher than that reported by Riedhammer *et al.* [30]. ES in patient known as vascular or hypertensive nephropathy based on histological finding could lead to recategorization of the kidney disease and suggest that a diagnosis of unclassifiable UKD would be more accurate before ES. Patients with suspected vascular or hypertensive nephropathy, who are young and have few comorbidities, a family history of kidney disease and/or extrarenal features such as hearing impairment, should be

classified as vascular kidney-NOS, a condition similar to autosomal dominant tubulointerstitial kidney disease not otherwise specified (ADTKD-NOS) [31].

The literature reports a positive family history of CKD in 25%–44% of cases in patients with CKD, and familial clustering is a common phenomenon in patients with ESKD, confirming that inherited kidney disease is a common cause of CKD [3, 32–35]. In our study, the segregation of kidney disease in a family was the main predictive factor associated with a high rate of ES resolution in patients with UKD, demonstrating that inherited kidney disease should not be overlooked in the adult population. In clinical nephrology practice, the systematic investigation of family history should be a standard element of the consultation, especially when the renal disease is of undetermined origin. History of kidney disease in a relative can be defined as requiring renal replacement therapy, either with dialysis or kidney transplant, or kidney disease that warrants repeated outpatient follow-up with a nephrology service. Depending on the patient, this may require a thorough survey of both the patient's history and that of their relatives prior to the nephrology consultation. Consanguinity was also associated with a slightly higher likelihood of

ES-resolved case rates, confirming what is already known in the literature.

Female sex was also reportedly associated with a positive family history of kidney disease. Being female has a protective effect on the progression of chronic renal disease [36]. Nowadays, X-linked kidney disorders, including Alport syndrome, Fabry disease, nephrogenic diabetes insipidus, X-linked hypophosphatemic rickets, Dent disease, oral facial digital syndrome and Lowe syndrome, are widely recognized [37]. Males have single X chromosome whereas females have two X chromosomes, and thus 100 Mb more DNA than males (X: 155 Mb, Y: 55 Mb), and this leads to a dosage imbalance between males and females for the 1000 X-linked genes. Most of these diseases predominantly affect males and have variable penetrance in female carriers. In our cohort, only two X-linked inherited kidney diseases (*COL4A5* and *CLCN5*) are present and do not explain this difference. X inactivation may contribute to the difference by cellular mosaicism because most of the kidney is formed after the onset of X inactivation. The distribution of the two mosaic populations is expected to be random, following a bell-shaped curve around a mean of 50%. It is possible that women with inherited kidney disease may have very skewed patterns of X inactivation, leading to major expression of the maternal autosomal chromosome.

Early-onset CKD is frequently associated with inherited kidney disease and may even be the leading cause of CKD in children. Interestingly, in our older cohort, early-onset disease was not the characteristic criterion associated with cases resolved by ES, as classically described. Our results indicate that late-onset disease or late age of end-stage renal failure should not negate the value of a genetic test such as ES.

Combining SNV and CNV using an exome-wide CNV detection pipeline raises the diagnostic yield of Mendelian kidney disease in our experience (5/75 patients, 6.7%) without adding any direct laboratory cost. It confirmed the result of previous studies [6]. This requires a bioinformatics team to set up CNV analysis in exome sequencing data-analysis pipelines with method validation by control samples. A careful analysis with updated CNV calling should be considered before performed a whole-genome sequencing in patients with high suspicion of inherited kidney disease.

An important limitation of our study is the analysis of variable number tandem repeat of the *MUC1* gene (GC-rich sequence), which is not really feasible despite it being the dominant gene associated with tubulointerstitial disease. In the future, long-read single molecule technology could directly assess many difficult or even previously unsequenceable regions of the genome by ES, such as repetitive elements of the *MUC1* gene, non-targeted structural variant breakpoints at base-pair resolution or discrimination of pseudogenes, and this would increase the diagnostic yield in patients with genetic CKD. However, variant interpretation remains a challenge with both short- and long-read technologies.

Most studies of genetic testing in CKD are limited to a research setting. Our study highlighted the importance of opening up genetic testing in routine nephrological care to address the problem of UKD, which accounts for nearly 20% of the CKD population. In addition, the clinical impact is significant because of 73 actionable genes proposed by the ACMG, many of which are associated with phenotypes relevant to nephrology. Furthermore, the most recent KDIGO conference, which centered on the controversies surrounding the topic of genetics in CKD, named additional kidney-specific actionable genes, selected on the

basis of the availability of interventions, which could prevent renal morbidity [29].

CONCLUSION

Genetic diagnosis has provided new clinical insights in UKD patients by clarifying or reclassifying kidney disease etiology or informing prognosis, treatment or transplant decisions. The exome first approach can have a high diagnostic yield, avoiding an invasive procedure such as renal biopsy or in cases where a biopsy is not possible. ES appears to be a compelling tool in a routine healthcare setting for patients with UKD.

SUPPLEMENTARY DATA

Supplementary data are available at [ckj](#) online.

ACKNOWLEDGEMENTS

The authors would like to thank CHU de Marseille for its investment in the genomic diagnosis process. The authors thank the affected individuals and their families for agreeing to participate in this study. The authors would also like to thank all nephrologists who referred samples from patients.

CONFLICT OF INTEREST STATEMENT

All the authors declared no competing interests.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

1. Titze S, Schmid M, Köttgen A et al. Disease burden and risk profile in referred patients with moderate chronic kidney disease: composition of the German Chronic Kidney Disease (GCKD) cohort. *Nephrol Dial Transplant* 2015;30:441–51. <https://doi.org/10.1093/ndt/gfu294>
2. Johansen KL, Chertow GM, Foley RN et al. US Renal Data System 2020 Annual Data Report: epidemiology of kidney disease in the United States. *Am J Kidney Dis* 2021;77:A7–8. <https://doi.org/10.1053/j.ajkd.2021.01.002>
3. Connaughton DM, Kennedy C, Shril S et al. Monogenic causes of chronic kidney disease in adults. *Kidney Int* 2019;95:914–28. <https://doi.org/10.1016/j.kint.2018.10.031>
4. Groopman EE, Marasa M, Cameron-Christie S et al. Diagnostic utility of exome sequencing for kidney disease. *N Engl J Med* 2019;380:142–51. <https://doi.org/10.1056/NEJMoa1806891>
5. Ottlewski I, Münch J, Wagner T et al. Value of renal gene panel diagnostics in adults waiting for kidney transplantation due to undetermined end-stage renal disease. *Kidney Int* 2019;96:222–30. <https://doi.org/10.1016/j.kint.2019.01.038>
6. Testard Q, Vanhoye X, Yauy K et al. Exome sequencing as a first-tier test for copy number variant detection: retrospective evaluation and prospective screening in 2418 cases. *J Med Genet* 2022;59:1234–40. <https://doi.org/10.1136/jmg-2022-108439>

7. Richards S, Aziz N, Bale S et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 2015;17:405–24. <https://doi.org/10.1038/gim.2015.30>
8. Kopp JB, Nelson GW, Sampath K et al. APOL1 genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy. *J Am Soc Nephrol* 2011;22:2129–37. <https://doi.org/10.1681/ASN.2011040388>
9. Stenson PD, Mort M, Ball EV et al. The Human Gene Mutation Database: towards a comprehensive repository of inherited mutation data for medical research, genetic diagnosis and next-generation sequencing studies. *Hum Genet* 2017;136:665–77. <https://doi.org/10.1007/s00439-017-1779-6>
10. Yao T, Udwan K, John R et al. Integration of genetic testing and pathology for the diagnosis of adults with FSGS. *Clin J Am Soc Nephrol* 2019;14:213–23. <https://doi.org/10.2215/CJN.08750718>
11. van der Ven AT, Connaughton DM, Ityel H et al. Whole-exome sequencing identifies causative mutations in families with congenital anomalies of the kidney and urinary tract. *J Am Soc Nephrol* 2018;29:2348–61. <https://doi.org/10.1681/ASN.2017121265>
12. Domingo-Gallego A, Pybus M, Bullich G et al. Clinical utility of genetic testing in early-onset kidney disease: seven genes are the main players. *Nephrol Dial Transplant* 2022;37:687–96. <https://doi.org/10.1093/ndt/gfab019>
13. Bullich G, Domingo-Gallego A, Vargas I et al. A kidney-disease gene panel allows a comprehensive genetic diagnosis of cystic and glomerular inherited kidney diseases. *Kidney Int* 2018;94:363–71. <https://doi.org/10.1016/j.kint.2018.02.027>
14. Mann N, Braun DA, Amann K et al. Whole-exome sequencing enables a precision medicine approach for kidney transplant recipients. *J Am Soc Nephrol* 2019;30:201–15. <https://doi.org/10.1681/ASN.2018060575>
15. Hill NR, Fatoba ST, Oke JL et al. Global Prevalence of chronic kidney disease - a systematic review and meta-analysis. *PLoS One* 2016;11:e0158765. <https://doi.org/10.1371/journal.pone.0158765>
16. Dryja TP. Gene-based approach to human gene-phenotype correlations. *Proc Natl Acad Sci USA* 1997;94:12117–21. <https://doi.org/10.1073/pnas.94.22.12117>
17. Smith JM, Stablein DM, Munoz R et al. Contributions of the transplant registry: the 2006 annual report of the North American Pediatric Renal Trials and Collaborative Studies (NAPRTCS). *Pediatr Transplant* 2007;11:366–73. <https://doi.org/10.1111/j.1399-3046.2007.00704.x>
18. Quinlan C, Rheault MN. Genetic basis of type IV collagen disorders of the kidney. *Clin J Am Soc Nephrol* 2021;16:1101–9. <https://doi.org/10.2215/CJN.19171220>
19. Kashtan CE, Ding J, Garosi G et al. Alport syndrome: a unified classification of genetic disorders of collagen IV α 345: a position paper of the Alport Syndrome Classification Working Group. *Kidney Int* 2018;93:1045–51. <https://doi.org/10.1016/j.kint.2017.12.018>
20. Köttgen A, Glazer NL, Dehghan A et al. Multiple loci associated with indices of renal function and chronic kidney disease. *Nat Genet* 2009;41:712–7. <https://doi.org/10.1038/ng.377>
21. Olinger E, Schaeffer C, Kidd K et al. An intermediate-effect size variant in. *Proc Natl Acad Sci USA* 2022;119:e2114734119. <https://doi.org/10.1073/pnas.2114734119>
22. Gast C, Marinaki A, Arenas-Hernandez M et al. Autosomal dominant tubulointerstitial kidney disease-UMOD is the most frequent non polycystic genetic kidney disease. *BMC Nephrol* 2018;19:301. <https://doi.org/10.1186/s12882-018-1107-y>
23. Ahram DF, Aggarwal VS, Sanna-Cherchi S, Phenocopies, phenotypic expansion, and coincidental diagnoses: time to abandon targeted gene panels? *Am J Kidney Dis* 2020;76:451–3. <https://doi.org/10.1053/j.ajkd.2020.07.003>
24. Schrezenmeier E, Kremerskothen E, Halleck F et al. The underestimated burden of monogenic kidney disease in adults waitlisted for kidney transplantation. *Genet Med* 2021;23:1219–24. <https://doi.org/10.1038/s41436-021-01127-8>
25. Robert T, Savenkoff B, Legris T et al. Exome sequencing re-analysis solved case in undetermined nephropathy with detection of TULP3-truncating variant. *Nephrol Dial Transplant* 2023;38:1057–60. <https://doi.org/10.1093/ndt/gfac319>
26. Devane J, Ott E, Olinger EG et al. Progressive liver, kidney, and heart degeneration in children and adults affected by TULP3 mutations. *Am J Hum Genet* 2022;109:928–43. <https://doi.org/10.1016/j.ajhg.2022.03.015>
27. Friedman DJ, Pollak MR. APOL1 nephropathy: from genetics to clinical applications. *Clin J Am Soc Nephrol* 2021;16:294–303. <https://doi.org/10.2215/CJN.15161219>
28. Kalatharan V, Lemaire M, Lanktree MB. Opportunities and challenges for genetic studies of end-stage renal disease in Canada. *Can J Kidney Health Dis* 2018;5:205435811878936. <https://doi.org/10.1177/2054358118789368>
29. KDIGO Conference Participants. Genetics in chronic kidney disease: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) controversies conference. *Kidney Int* 2022;101:1126–41.
30. Riedhammer KM, Braunisch MC, Günthner R et al. Exome sequencing and identification of phenocopies in patients with clinically presumed hereditary nephropathies. *Am J Kidney Dis* 2020;76:460–70. <https://doi.org/10.1053/j.ajkd.2019.12.008>
31. Eckardt KU, Alper SL, Antignac C et al. Autosomal dominant tubulointerstitial kidney disease: diagnosis, classification, and management—A KDIGO consensus report. *Kidney Int* 2015;88:676–83. <https://doi.org/10.1038/ki.2015.28>
32. Shang N, Khan A, Polubriaginof F et al. Medical records-based chronic kidney disease phenotype for clinical care and “big data” observational and genetic studies. *NPJ Digit Med* 2021;4:70. <https://doi.org/10.1038/s41746-021-00428-1>
33. Akrawi DS, Li X, Sundquist J et al. Familial risks of kidney failure in Sweden: a nationwide family study. *PLoS One* 2014;9:e113353. <https://doi.org/10.1371/journal.pone.0113353>
34. Skrunes R, Svarstad E, Reisæter AV. et al. Familial clustering of ESRD in the Norwegian population. *Clin J Am Soc Nephrol* 2014;9:1692–700. <https://doi.org/10.2215/CJN.01680214>
35. Freedman BI, Robinson TW. Risk factors: familial aggregation of ESRD in Europeans—is it in the genes? *Nat Rev Nephrol* 2014;10:677–8. <https://doi.org/10.1038/nrneph.2014.181>
36. Connaughton DM, Bukhari S, Conlon P et al. The Irish kidney gene project—prevalence of family history in patients with kidney disease in Ireland. *Nephron* 2015;130:293–301. <https://doi.org/10.1159/000436983>
37. Quinlan C, Rheault MN. X-linked kidney disorders in women. *Semin Nephrol* 2022;42:114–21. <https://doi.org/10.1016/j.semnephrol.2022.04.002>