

Knowledge, Attitude, and Practices regarding Leptospirosis among Visitors to a Recreational Forest in Malaysia

Nur Juliani Shafie,^{1*} Najma Syahmin Abdul Halim,¹ Mohamed Nor Zalipah,¹ Nur Amalin Zahirah Mohd Amin,¹ Sharifah Masit'ah Syed Esa,¹ Shukor Md-Nor,² Arnau Casanovas-Massana,³ Albert I. Ko,^{3,5} Fabiana Palma,⁴ Fabio Neves Souza,⁴ and Federico Costa^{3,4,5}

¹Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, Terengganu, Malaysia; ²School of Environmental and Natural Resource Sciences, Universiti Kebangsaan Malaysia, Bangi, Malaysia; ³Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, Connecticut; ⁴Institute of Collective Health, Federal University of Bahia, UFBA, Salvador, Brazil; ⁵Oswaldo Cruz Foundation, Brazilian Ministry of Health, Salvador, Bahia, Brazil

Abstract. Leptospirosis is a zoonotic disease and a worldwide public health problem that affects mainly high-risk groups. Characterizing knowledge, attitude, and practice (KAP) among high-risk groups is important to develop appropriate prevention programs. Here, we performed a cross-sectional study among 300 visitors of a recreational forest in Malaysia to examine leptospirosis KAP and demographics. These variables were integrated to create knowledge and practice scores for each respondent. All respondents had heard about leptospirosis, and 87% of them correctly identified it as a disease. The majority of respondents had high knowledge (63%), positive attitude, and good practice (68%) toward prevention of the disease. However, there were gaps in knowledge, with 78% of the respondents indicating eating without washing hands as the major cause of leptospirosis transmission. Our final model identified that higher knowledge score was associated with higher practice score. Our results indicate that it is important to increase knowledge, especially on transmission routes of leptospirosis, among visitors in recreational areas. Moreover, more attention needs to be paid to promote good practice habits among visitors, targeting those at higher risk of being infected by leptospirosis to prevent potential outbreaks in the recreational areas.

INTRODUCTION

Leptospirosis is a globally distributed zoonotic disease with more than one million cases and 50,000 deaths annually.¹ Leptospirosis shows a range of symptoms in humans such as high fever, headache, jaundice, renal failure, meningitis, hemorrhagic fever, abdominal pain, and muscle aches; however, most infections are subclinical.² Leptospirosis is caused by pathogenic spirochetes from the genus *Leptospira*. Pathogenic *Leptospira* spp. infect and colonize the kidneys of almost all mammalian species³ and are excreted at high concentrations^{4,5} into the environment where they can survive for extended time.⁶ Small mammals, and specifically rodents, are chronic carriers of *Leptospira* that act as the major reservoir host in the transmission of leptospirosis.³ Infection occurs either through direct contact of broken skin, abrasion, or mucous membranes with urine or tissue of infected animal, or indirect contact with contaminated soil and water.⁷ Leptospirosis is highly prevalent in the Asia Pacific region, and specifically in Southeast Asia and Oceania.^{1,3} The estimated incidence of leptospirosis ranges from 0.1 to 1 for every 100,000 per year in temperate climates, but it can be as high as 10–100 per 100,000 in tropical regions, and more than 100 per 100,000 among high-risk groups during outbreaks.⁸ The high prevalence of leptospirosis in Southeast Asia has been related to flooding, recreational activities, occupational exposure, travel to endemic regions, poor sanitation, and waste disposal.³

Leptospirosis is endemic in Malaysia, and the first case was reported in 1925.⁹ A total of 12,325 cases with 338 deaths were reported between 2004 and 2012.¹⁰ According to the Ministry of Health Malaysia,¹¹ Terengganu was among the five states in Malaysia with the highest number of leptospirosis

cases (≈ 8.00 per 100,000 population), and recreational areas were categorized as one of the important hotspots of leptospirosis outbreaks. In Malaysia, most recreational areas are equipped with many infrastructures such as community halls, chalets, public toilets, and food stalls.¹² However, poor sanitation and waste disposal make recreational areas favorable environments for the transmission of leptospirosis. The presence of pathogenic *Leptospira* has been previously reported in two recreational areas in Terengganu, Lata Tembakah Waterfall and Lata Belatan Waterfall, with 5% (1/20 water and soil samples) and 10% (2/20 water and soil samples) positive samples, respectively.¹³ Several factors have been identified as contributing to the increasing number of outbreak, including water pH, temperature, soil moisture, and the presence of wild animals which act as reservoirs.^{13,14} *Rattus tiomanicus* was found to be the dominant carrier with 88.1% prevalence (37/42 leptospiral isolates) for serovars Icterohaemorrhagiae, Canicola, Ballum, Pyrogenes, and Hebdomadis.¹⁵ Serovar hebdomadis was also reported as the dominant serovar in Terengganu.¹⁴

Despite the increasing incidence, leptospirosis is often neglected, and there is still a lack of awareness and basic knowledge on the disease, especially among high-risk groups.¹⁶ Understanding the knowledge of individual subjects and their attitudes with respect to risky behaviors is key to implementing better prevention practices. Similarly, it is also important to identify the sociodemographic characteristics that influence individual behaviors and their prevention practices of leptospirosis.¹⁷ Knowledge, attitude, and practice (KAP) studies on leptospirosis conducted in Malaysia have focused mainly on describing the KAP among urban and rural communities,^{18,19} residents in flood-prone areas,^{20,21} town service workers,²² wet market workers,²³ non-high-risk group,²⁴ and university students.²⁵ Yet, there are no KAP studies on leptospirosis conducted among visitors of recreational areas, which limits our ability to understand their level of awareness toward the prevention of leptospirosis. Here, we aimed to identify KAP related to leptospirosis among visitors of

* Address correspondence to Nur Juliani Shafie, Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia. E-mail: nur.shafie@umt.edu.my

Lata Belatan, a recreational forest in Besut (Terengganu, Malaysia), and to determine the factors that influence leptospirosis prevention practices.

MATERIALS AND METHODS

Study area. This study was conducted in Lata Belatan, a recreational forest in Terengganu, Malaysia (5°63'90"N, 102°58'78"E). This recreational forest is located at the base of Gunung Tebu in Besut, Terengganu (Figure 1). It consists of fast-flowing cascades and clear water that attracts visitors from Terengganu, other states in Malaysia, and foreigners. This recreational area is characterized by lowland riparian forest and low hill dipterocarp forest.²⁶ There are also many facilities provided for the visitors such as food stalls, public toilets, camping sites, rest areas, and a prayer room.

Study design. We conducted a cross-sectional study among visitors of Lata Belatan Recreational Forest. A list of 345 respondents were selected randomly from a database from the Department of Forestry Terengganu ($N = 13,043$) using IBM SPSS Statistics for Windows version 21 (IBM Corp.,

Armonk, NY). The sample size was calculated based on 95% CI, 5% margin of error, and 50% positive response rate. However, 13% ($N = 45$) of nonresponse was detected because of failure to reach. Inclusion criteria were > 16 years old, visited the recreational forest between January–June 2018, and agree to participate in the study. The questionnaire was prepared in Malay language (local dialect) because the local residents were predominantly Malay native speakers. All data and visitors' personal information were confidential. Parental consent letters were given for respondents younger than 18 years, and informed consent was also obtained from adult respondents before they participated in this study. Data collection was conducted using printed questionnaires from September 2018 until March 2019. The approval to conduct this study was obtained from Ethical Review Committee Board, Universiti Malaysia Terengganu (UMT/JKEPHT/2019/30).

Data collection. A questionnaire was developed using previous KAP studies on leptospirosis.^{27–29} The questionnaire consisted of 42 questions categorized into two sections: 1) sociodemographic characteristics (age, gender, educational





FIGURE 1. Lata Belatan Recreational Forest in Terengganu, Malaysia. (A) Location of the study site in Peninsular Malaysia and within the state of Terengganu; (B) visitors who are involved with recreational or water activities. This figure appears in color at www.ajtmh.org.

level, occupation, and monthly income) and 2) KAP of leptospirosis. Knowledge related to leptospirosis was assessed by asking whether they had ever heard of “leptospirosis” or also known as “rat-urine disease” among local residents. The respondents were also asked questions about the 1) causative organism and disease transmission, 2) common symptoms, and 3) prevention practices regarding leptospirosis. Respondents were given three answer options including “true,” “false,” and “do not know.” Each affirmative answer was given as one score, whereas negative answer and “do not know” were given as zero score. The attitude of the respondents toward leptospirosis was evaluated by asking questions on their reactions to statements about leptospirosis and other attitudinal issues based on a five-level Likert scale (one = strongly disagree, two = disagree, three = not sure, four = agree, and five = strongly agree).³⁰ Respondents who agreed or strongly agreed to the statements were considered to have a positive attitude, whereas respondents who were not sure, disagreed, or strongly disagreed were considered to have a negative attitude. The preventive practices were evaluated by asking them “yes or no” questions. Each “yes” answer was given one score, whereas “no” answer was given zero score. Scores for each individual were calculated and summed up to obtain the total score.

Data analysis. The data were coded, entered into Microsoft Excel 2013 (Microsoft Corp., Redmond, WA), and then imported to R program for further analysis.³¹ Descriptive statistics including means and frequencies were calculated for knowledge and practice scores. Knowledge and practice scores were used as continuous variables. Sociodemographic characteristics and knowledge score were considered as independent variables and the practice score as the dependent variable. Generalized linear models (GLMs) with Poisson error family was used to determine the predictors and associations between dependent and independent variables. Subsequently, these variables were evaluated one by one in the univariate analyses, and determinants having a screening significance of $P < 0.20$ were included in multivariate analyses.³² Model simplification was carried, and we used the *dredge* () function of the “MuMIn” package.³³ This function performs a set of models and valid combinations between the fixed variables of the multiple model, with all possible models ranked by the value of Akaike’s information criterion (AIC). This process was continued by removing the least significant terms from the model until the minimum adequate model was achieved. When comparing two models, the smaller AIC was accepted as indicating the better fit.³⁴ All P -values were considered statistically significant at $P < 0.05$.

RESULTS

Sociodemographics. Of 300 respondents, there was an equal number of men and women (49.3% and 50.7%, respectively). Most respondents were aged between 21 and 30 years (47.0%), and more than half of the respondents had completed tertiary education (70.3%) (Table 1). The main occupations were students (48.7%) and private sector workers (22.6%). The majority of participants did not have a monthly income (52.3%), and the mean household income was Ringgit Malaysia 1,279 per month (\approx United States Dollar 303). This was expected considering most of respondents lived close to the recreational forest, where this study was conducted, which consists of low-income residential communities.

TABLE 1
Sociodemographic characteristics of the respondents ($N = 300$)

Category	Profile of respondents	Number	Percentage
Gender	Male	148	49.3
	Female	152	50.7
Age-group (years)	16–20	89	29.7
	21–30	141	47.0
	31–40	44	14.7
	> 41	26	8.7
	Education	Primary school	0
	Secondary school	85	28.3
	Tertiary education	211	70.3
Occupation	No formal education	4	1.3
	Student	146	48.7
	Private sector	68	22.6
	Government sector	51	17.0
	Housewife	19	6.3
	Unemployed	11	3.7
	Retired	5	1.7
Household income	< RM 500	8	2.7
	RM 500–RM 1499	38	12.7
	RM 1500–RM 2499	39	13.0
	RM 2500–RM 3499	17	5.7
	RM 3500–RM 4999	20	6.7
	> RM 5000	21	7.0
	No income	157	52.3

Knowledge. All respondents had heard about leptospirosis, and 87.0% of them identified leptospirosis as a disease (Table 2). When asked about the causative agent of leptospirosis, 57.3% of respondents answered bacteria and 22.0% of respondents thought that parasites or worms were the causal organisms. Regarding the transmission of leptospirosis to humans, 78.0% answered eating without washing hands as the main risk factor. Only 47.3% of respondents knew contact with contaminated water and soil is one of the main factors of transmission. With regard to symptoms of leptospirosis, most of the respondents were aware of high fever (73.0%), headache (72.0%), and jaundice (51.0%). The analysis of the knowledge score showed that of a maximum of 20 points, the mean score of the respondents was $11.1 \pm \text{SD } 5.1$ (range: 0–19). A total of 189 respondents (63.0%) were ranked above the mean score, indicating high knowledge among respondents.

When asked about prevention, 74.0% of respondents reported that the disease could be detected by blood tests and could be treated early (71.3%). However, 22.0% of respondents thought that the disease could be prevented by taking a bath after water activities. In addition, 222 (74.0%) respondents agreed that leptospirosis could cause death. The main source of information on leptospirosis was media (82.7%), followed by community associations (76.7%) and healthcare providers (41.7%).

Attitude and prevention practices. In all the five statements used to measure attitude, high percentage of respondents had positive attitudes toward prevention and control of leptospirosis (Table 3). The majority of respondents believed that leptospirosis is a serious disease (87.7%). They also had positive attitudes on the statement, “I must know about leptospirosis” (85.3%) and “I should make sure that the recreational area is clean” (78.0%). However, only 59.7% of respondents had positive attitude on the statement, “I believe that leptospirosis has a cure.” A total of 204 respondents (68%) scored above the mean score ($3.0 \pm \text{SD } 1.5$, range: 0–6), which indicates good practices against leptospirosis among

TABLE 2

Proportion of respondents with affirmative answers for knowledge on leptospirosis ($N = 300$)

Knowledge regarding leptospirosis	Number	Percentage
Leptospirosis		
Leptospirosis is a disease	261	87.0
Bacteria are the causative agent of leptospirosis	172	57.3
Viruses are the causative agent of leptospirosis	165	55.0
Parasites or worms are the causative agent of leptospirosis	66	22.0
Transmission		
Eating without washing hands is a risk to get leptospirosis	234	78.0
Leptospirosis is transmitted through rat urine	221	73.7
Walking without shoes is a risk to get leptospirosis	204	68.0
Leptospirosis can be transmitted through cut skin wounds	176	58.7
Water activities are risky for leptospirosis infection	165	55.0
Contaminated food is a risk for leptospirosis infection	145	48.3
Leptospirosis can be transmitted through contaminated water and soil	142	47.3
Symptoms		
Leptospirosis will cause high fever	219	73.0
Infected individuals will develop headache	216	72.0
Leptospirosis will cause jaundice	153	51.0
Leptospirosis will cause liver damage	120	40.0
Treatment and prevention		
The disease can be detected by a blood test	222	74.0
This disease can be treated early	214	71.3
The disease can be prevented by maintaining cleanliness of recreational area	190	63.3
The disease can be prevented by taking a bath after water activities	66	22.0
Leptospirosis can cause death	222	74.0
Source of knowledge		
Media (newspaper, television, radio, Internet)	248	82.7
Community associations (family, friends, neighbor)	230	76.7
Health care	125	41.7
School	78	26.0
Work	44	14.7

respondents. When asked about prevention practices, 90.3% of them maintained the cleanliness of the recreational area and 89.3% of respondents said that they always drank clean water supplies. However, only 171 (57.0%) and 154 (51.3%) of respondents mentioned wearing shoes in the recreational area and avoiding water activities, respectively, as prevention practices toward leptospirosis.

Regarding association of the respondent's sociodemographic characteristics with practice score, the univariate multiple Poisson regression (GLM) analysis (Table 4) showed that knowledge was significantly associated with practice against leptospirosis, females were more likely to have higher practice score than males (Odds ratio [OR] = 1.1; 95% confidence interval [CI] = 1.0–1.2), and respondents > 30 years old have higher practice score than respondents < 30 years old (OR = 1.1; 95% CI = 1.0–1.2). In the final adjusted model, only knowledge score was associated with the practice against leptospirosis score (aOR = 1.0; 95% CI = 1.00–1.01). No

significant differences were observed between practice score and other sociodemographic characteristics.

DISCUSSION

In this study, we performed a cross-sectional evaluation of the KAP related to leptospirosis among visitors of Lata Belatan, a recreational forest in Besut (Terengganu, Malaysia), and to determine the factors that influence leptospirosis prevention practices. Our results showed that all respondents had heard about leptospirosis, and 87.0% of them identified leptospirosis as a disease. The majority of respondents had high knowledge (63%) and good practices (68%) regarding leptospirosis based on their overall scores. However, a large proportion of respondents had a poor understanding on the transmission factors of the disease, especially on the role of contaminated water and soil in the transmission of *Leptospira*. Moreover, higher knowledge score was significantly associated with higher leptospirosis prevention practices.

All of the respondents had heard of leptospirosis, and 87.0% of them identified it as a disease, with an overall knowledge score of 63%. The majority of the respondents had heard about leptospirosis from media (82.7%) and also community associations (76.7%) that involved knowledge sharing among families, friends, and neighbors. These findings are in agreement with previous studies that have shown that mass media are the main source of knowledge about leptospirosis.^{18,19,22,24} Media can be an effective channel to deliver messages to the public in a fastest way because face-to-face communications often require costly human resources.³⁵ Public health social media campaigns are viewed as powerful tools to enhance health behaviors, attitudes, and awareness at a population level.³⁶ However, the relevance of media and community associations need to be taken into account as not all information and knowledge received from the Internet and social networks are accurate and reliable.

Regarding transmission of leptospirosis, less than 50% of the respondents knew that contaminated water and soil are relevant risk factors for leptospirosis infection. Moreover, 22.0% of respondents thought that the disease could be prevented by rinsing their body after bathing in the waterfall. These findings are consistent with a previous study that found there is a gap in the knowledge regarding leptospirosis, especially on transmission routes and prevention measures.¹⁹ In our study, most of the respondents had misconceptions regarding leptospirosis because of the use of the layman term "rat-urine disease," which focuses only on rodents. However, human infection can occur from exposure to infected urine from a wide variety of non-rodent animals including bats, ruminants, carnivores, primates, wild boars, deers, and dogs.^{37–39} Leptospirosis can be prevented by avoiding direct or indirect contact with infected animals or possible contaminated water and soil, wearing protective clothing and shoes, maintaining personal hygiene, protecting any wounds or scars with medical plaster, and drinking clean water.⁸ Therefore, to improve prevention measures in recreational areas, it is important to provide this correct information and increase visitors' health education on leptospirosis.

Overall, most of the respondents had positive attitudes toward prevention and control of leptospirosis. Most of them strongly agreed and believe that leptospirosis is a serious disease and they must know about it. Leptospirosis

TABLE 3

Proportion of respondents with positive answers for attitude and prevention practices related to leptospirosis ($N = 300$)

Item	Number	Percentage
Attitude		
I believe leptospirosis is a serious disease	263	87.7
I must know about leptospirosis	256	85.3
I should make sure that recreational area is clean	234	78.0
It is important to control rat population	226	75.3
I believe that leptospirosis has a cure	179	59.7
Practices		
Maintain cleanliness of the recreational area	271	90.3
Drink clean water supply	268	89.3
Wash hands with soap and water	241	80.3
Scrutinize any broken wounds and scars	209	69.7
Wear shoes in the recreational area	171	57.0
Avoid water activities	154	51.3

was reported to be a silent killer disease as the infected person will be asymptomatic or experience mild symptoms that can be misinterpreted as a typical flu-like illness until it becomes fatal.⁴⁰ It is important for them to know that leptospirosis is an acute disease, in order for them to understand the consequences of this disease. Most of the respondents strongly agreed that they will make sure the recreational area is clean. Even though this attitude might not be directly related to leptospirosis, it is still an important precautionary measure because poor hygiene will increase the population of *Leptospira* reservoirs and create favorable environments for its transmission. The high percentage of positive responses might be due to the phrasing of the questions for the attitudes. However, to minimize bias in this section, respondents were provided with equally balanced answer scales between positive and negative options. The respondents were given five-level answer scales: strongly disagree, disagree, not sure, agree, and strongly agree.

Our final model identified that respondents with higher knowledge score showed higher practice score than those with lower knowledge score. Our result is consistent with previous studies in the Philippines²⁹ and Argentina⁴¹ that showed higher knowledge score on leptospirosis may influence a higher adoption of preventive practices, but no other studies have identified this association. Our finding suggests that increasing knowledge on leptospirosis is important to promote positive preventive practices in the community. According to the theoretical constructs in the

Health Belief Model (HBM), perceived severity and perceived susceptibility to a disease depend on individual knowledge about the disease.⁴² The HBM suggests that higher perceived severity and perceived susceptibility (perceived threat) lead to higher likelihood of positive behavior toward the disease. Increasing knowledge on leptospirosis among visitors may promote more concern about this disease, which in turn may affect leptospirosis prevention practices. Leptospirosis information through signage in the recreational area could help to convey important information about leptospirosis to the visitors. It was found that interpretive signage has successfully increased knowledge, awareness, and positive behavior among visitors at Penang National Park, Malaysia, on conservation aspects.⁴³ In addition, the development and implementation of gamification is a promising and interactive approach to increase knowledge on health information by involving the participants in various activities.⁴⁴⁻⁴⁶ Gamification modules such as role play, puzzles, goal-oriented activities, and social media engagement have been applied successfully to increase knowledge among university students in Malaysia regarding leptospirosis.⁴⁶

There were no associations between demographic factors (gender and age) with leptospirosis prevention practices. However, our univariate results identified that males and younger respondents were more likely to have lower practice scores against leptospirosis. Majority of the visitors in the recreational forest in this study were aged between 21 and 30 years. Previous studies in Malaysia showed that most of the positive cases on leptospirosis were reported to occur among males and those aged between 20 and 29 years as they are usually more involved with riskier occupational and outdoor activities.^{10,47,48} This tendency could be due to the frequent outdoor recreational activities among those groups without knowing or acknowledging the risk factors of leptospirosis.⁴⁹ Thus, it is crucial to increase the knowledge and awareness among male and younger visitors in recreational areas. In addition, more comprehensive efforts should be established to educate these groups to change their behavior and to avoid infection of this disease.

This study has limitations. First, the respondents were not asked why they did or did not take precautionary measures. This question needs to be included to determine in details the reasons for respondents' responses. Second, this study was conducted in only one recreational area in Malaysia which may

TABLE 4

Multiple poisson regression (Generalized linear model [GLM]) analysis of the association between socio-demographic characteristics, knowledge score, and preventive practice score against leptospirosis ($N = 300$)

Variables	Categories	Odds ratio (95% CI)	Adjusted odds ratio (95% CI)	P-value
Gender	Female	1.08 (0.98–1.20)	–	–
	Male	1	1	–
Education	Has formal education	1.05 (0.69–1.72)	–	–
	No formal education	1	1	–
Age (years)	> 30	1.09 (0.97–1.23)	–	–
	< 30	1	1	–
Occupation	Employed	0.98 (0.88–1.08)	–	–
	Unemployed	1	1	–
Income	No monthly income	1.14 (0.63–2.08)	–	–
	Has monthly income	1	1	–
Knowledge score		1.01 (1.00–1.02)	1.01 (1.00–1.01)	0.018

The bold values show that female was more likely to have higher practice score than male, respondents > 30 years old have higher practice score than respondents < 30 years old and knowledge was significantly ($P < 0.05$) associated with practice.

affect the generalizability of our findings. However, we believe that our initial findings provide reliable information on the level of KAP among individual on the specific setting of recreational areas.

In conclusion, this study demonstrated that most of the respondents had high knowledge, positive attitudes, and good practices toward prevention of leptospirosis. However, high percentage of them still had limited knowledge on its route of transmission and prevention measures. Our results suggest that appropriate education programs targeting the high-risk groups should be conducted to deliver and increase their knowledge on leptospirosis. Male and young respondents who perform recreational and outdoor activities were also found to have higher risk of being infected by leptospirosis in this area. The number of cases and deaths will continue to increase if there are no efficient measures taken to mitigate this situation. Public health workers, researchers, and policy-makers should improve prevention measures and treatments for this disease. Because community involvement is a key in public health, local communities themselves should also be involved in these approaches to reduce the burden of leptospirosis.

Received April 17, 2020. Accepted for publication December 6, 2020.

Published online January 18, 2021.

Acknowledgments: We thank Forestry Department of Peninsular Malaysia and Forestry Department of Terengganu for the approval and support to conduct this study. We also thank the visitors of Lata Belatan Recreational Forest, Terengganu, for participating in the study.

Financial support: The work presented in this study is funded by the Ministry of Higher Education, Malaysia, through UKM-UMT collaboration research grant (Vot 53138; 53269), the University of Putra Malaysia via Long-Term Research Grant Scheme (LRGS Phase 2/2014, UPM/700-2/7/LRGS/55264000), as well as the University of Malaya Research University Grant (RU005-2017), the Wellcome Trust (102330/Z/13/Z and 218987/Z/19/Z), and the Brazilian National Council for Scientific and Technological Development (CNPq).

Authors' addresses: Nur Juliani Shafie, Najma Syahmin Abdul Halim, Mohamed Nor Zalipah, Nur Amalin Zahirah Mohd Amin, and Sharifah Masit'ah Syed Esa, Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, Terengganu, Malaysia, E-mails: nur.shafie@umt.edu.my, najmasyahmin@gmail.com, zalipah@umt.edu.my, zahirahnuramalin@gmail.com, and masitah97cetah@gmail.com. Shukor Md-Nor, School of Environmental and Natural Resource Science, Universiti Kebangsaan Malaysia, Bangi, Malaysia, E-mail: shukor@ukm.edu.my. Amau Casanovas-Massana and Albert I. Ko, Yale School of Public Health, New Haven, CT, E-mails: amau.casanovas@yale.edu and albert.ko@yale.edu. Fabiana Palma, Fabio Neves Souza, and Federico Costa, Institute of Collective Health, Federal University of Bahia, UFBA, Salvador, Brazil, E-mails: fabiana.palma@yahoo.com.br, fabionevesouza@gmail.com, and fcosta2001@gmail.com.

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC-BY) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

- Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS, Stein C, Abela-Ridder B, Ko AI, 2015. Global morbidity and mortality of leptospirosis: a systematic review. *PLoS Negl Trop Dis* 9: e0003898.
- Bharti AR et al., 2003. Leptospirosis: a zoonotic disease of global importance. *Lancet Infect Dis* 3: 757-771.
- Victoriano AFB et al., 2009. Leptospirosis in the Asia Pacific region. *BMC Infect Dis* 9: 147.
- Costa F, Wunder EA, De Oliveira D, Bisht V, Rodrigues G, Reis MG, Ko AI, Begon M, Childs JE, 2015. Patterns in *Leptospira* shedding in Norway Rats (*Rattus norvegicus*) from Brazilian slum communities at high risk of disease transmission. *PLoS Negl Trop Dis* 9: e0003819.
- Barragan V et al., 2016. High *Leptospira* diversity in animals and humans complicates the search for common reservoirs of human disease in rural Ecuador. *PLoS Negl Trop Dis* 10: e0004990.
- Casanovas-Massana A, Pedra GG, Wunder EA, Diggle PJ, Begon M, Ko AI, 2018. Quantification of *Leptospira interrogans* survival in soil and water microcosms. *Appl Environ Microbiol* 84: e00507.
- Ko AI, Goarant C, Picardeau M, 2009. Leptospira: the dawn of the molecular genetics era for an emerging zoonotic pathogen. *Nat Rev Microbiol* 7: 736-747.
- WHO, 2007. *Leptospirosis*. Geneva, Switzerland: World Health Organization. Available at: http://apps.searo.who.int/PDS_DOCS/B2147.pdf. Accessed April 2, 2019.
- El Jalii IM, Bahaman AR, 2004. A review of human leptospirosis in Malaysia. *J Trop Biomed* 21: 113-119.
- Benacer D, Thong KL, Min NC, Verasahib KB, Galloway RL, Hartskeerl RA, Souris M, Zain SNM, 2016. Epidemiology of human leptospirosis in Malaysia, 2004-2012. *Acta Trop* 157: 162-168.
- MOH, 2015. *Epidemiology and Current Situation of Leptospirosis in Malaysia*. Available at: <http://moh.gov.my>. Accessed April 2, 2019.
- Adriansyah D, Busu I, Eva H, Muqtada M, 2015. Geoheritage as the basis of geotourism development: a case study in Jeli district, Kelantan, Malaysia. *Geoj Tour Geosites* 15: 25-43.
- Ridzlan FR, Bahaman AR, Khairani-Bejo S, Mutalib AR, 2010. Detection of pathogenic *Leptospira* from selected environment in Kelantan and Terengganu, Malaysia. *Trop Biomed* 27: 632-638.
- Ismail S et al., 2014. A study on the presence of pathogenic *Leptospira* spp. in environmental water samples obtained from selected recreational areas in Terengganu, Malaysia. *Res J Pharm Tech* 7: 1153-1157.
- Mohamed-Hassan SN, Bahaman AR, Mutalib AR, Khairani-Bejo S, 2012. Prevalence of pathogenic leptospires in rats from selected locations in Peninsular Malaysia. *Res J Anim Sci* 6: 12-25.
- Schneider MC, Jancloes M, Buss DF, Aldighieri S, Bertherat E, Najera P, Galan DI, Durski K, Espinal MA, 2013. Leptospirosis: a silent epidemic disease. *Int J Environ Res Public Health* 10: 7229-7234.
- Azimullah AZ, Aziah BD, Fauziah BD, 2016. The rise of leptospirosis in Kelantan 2014: characteristics, geographical pattern and associated factors. *Int J Public Health Clin Sci* 3: 52-62.
- Nozmi N et al., 2018. Low levels of knowledge, attitudes and preventive practices on leptospirosis among a rural community in Hulu Langat District, Selangor, Malaysia. *Int J Environ Res Public Health* 15: 693.
- Sukeri S, Idris Z, Zahiruddin WM, Shafei MN, Idris N, Hamat RA, Jamaluddin TZT, Osman M, Wahab ZA, Daud A, 2018. A qualitative exploration of the misconceptions, knowledge gaps and constructs of leptospirosis among rural and urban communities in Malaysia. *PLoS One* 13: e0200871.
- Edre MA, Hayati KS, Salmiah MS, Sharifah Norkhadajah SI, 2015. A case control study on factors associated with leptospirosis infection among residents in flood-prone area, Kuantan: a geographical information system-based approach. *Int J Public Health Clin Sci* 2: 151-163.
- Edre MA, Hayati KS, Salmiah MS, Sharifah Norkhadajah SN, 2018. Determinants of leptospirosis preventive practices among the community in a flood-prone residential area in Kuantan, Malaysia. *Mal J Med Health Sci* 14: 27-33.
- Azfar ZM, Nazri SM, Rusli AM, Maizurah O, Zahiruddin WM, Azwany YN, Nabilah I, Asma HS, Aziah BD, 2018. Knowledge, attitude and practice about leptospirosis prevention among town service workers in northeastern Malaysia: a cross sectional study. *J Prev Med Hyg* 59: e92-e98.
- Rahman M et al., 2018. Leptospirosis health intervention module effect on knowledge, attitude, belief, and practice among wet

- market workers in northeastern Malaysia: an Intervention Study. *Int J Environ Res Public Health* 15: 1396.
24. Sakinah SNS, Suhailah S, Jamaluddin TZMT, Norbaya SM, Malina O, 2015. Seroprevalence of leptospiral antibodies and knowledge, attitude and practices of leptospirosis to non-high risk group in Selangor. *Int J Public Health Clin Sci* 2: 92–104.
 25. Bakar SMA, Rahman HA, 2015. Knowledge, attitude and practice on leptospirosis among undergraduate students in University Putra Malaysia. *Mal J Med Health Sci* 14: 104–111.
 26. Sumarli AX, Grismer LL, Anuar S, Muin MA, Quah ES, 2015. First report on the amphibians and reptiles of a remote mountain, Gunung Tebu in northeastern Peninsular Malaysia. *Check List* 11: 1679.
 27. Rahim MS, Aziah B, Nazri S, Azwany Y, Habsah H, Zahiruddin WM, Zahila I, Rusli MA, 2012. Town service workers' knowledge, attitude and practice towards leptospirosis. *Brunei Darussalam J Health* 5: 1–12.
 28. Quina CR, Almazan JU, Tagarino JB, 2014. Knowledge, attitudes, and practices of leptospirosis in Catbalogan city, samar, Philippines. *Am J Public Health Res* 2: 91–98.
 29. Arbiol J, Orencio P, Nomura H, Takahashi Y, Yabe M, 2016. Knowledge, attitude and practices towards leptospirosis among lakeshore communities of Calamba and Los Baños, Laguna, Philippines. *Agriculture* 6: 18.
 30. Likert RA, 1932. Technique for the measurement of attitudes. *Arch Psychol* 22: 1–55.
 31. R core Team, 2016. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at: <https://www.R-project.org/>.
 32. Koenraad CJ, Tuiten W, Sithiprasasna R, Kijchalao U, Jones JW, Scott TW, 2006. Dengue knowledge and practices and their impact on *Aedes aegypti* populations in Kamphaeng Phet, Thailand. *Am J Trop Med Hyg* 74: 692–700.
 33. Burnham KP, Anderson DR, 2007. *Model Selection and Multi-model Inference: A Practical Information-Theoretic Approach*. New York, NY: Springer.
 34. Bozdogan H, 1987. Model selection and Akaike's information criterion (AIC): the general theory and its analytical extensions. *Psychometrika* 52: 345–370.
 35. Rogala MB, 2017. Health promotion actions in mass-media for seniors in selected European countries. *Epidemiol Biostat Public Health* 14: 1–10.
 36. Randolph KA, Whitaker P, Arellano A, 2012. The unique effects of environmental strategies in health promotion campaigns: a review. *Eval Program Plann* 35: 344–353.
 37. Koizumi N, Muto M, Yamamoto S, Baba Y, Kudo M, Tamae Y, Soma H, 2008. Investigation of reservoir animals of *Leptospira* in the northern part of Miyazaki Prefecture. *Jpn J Infect Dis* 61: 465–468.
 38. Lagadec E, Gomard Y, Guernier V, Dietrich M, Pascalis H, Temmam S, Ramasindrazana B, Goodman SM, Tortosa P, Dellagi K, 2012. Pathogenic *Leptospira* spp. in bats, Madagascar and Union of the Comoros. *Emerg Infect Dis* 18: 1696–1698.
 39. Andersen-Ranberg EU, Phipper C, Jensen PM, 2016. Global patterns of *Leptospira* prevalence in vertebrate reservoir hosts. *J Wildl Dis* 52: 468–477.
 40. Haake DA, Levett PN, 2015. Leptospirosis in humans. *Curr Top Microbiol Immunol* 387: 65–97.
 41. Ricardo T, Bergero LC, Bulgarella EP, Previtali MA, 2018. Knowledge, attitudes and practices (KAP) regarding leptospirosis among residents of riverside settlements of Santa Fe, Argentina. *PLoS Negl Trop Dis* 12: e0006470.
 42. Rosenstock IM, 1974. Historical origins of the health belief model. *Health Educ Monogr* 2: 328–335.
 43. Ismail MH, 2008. *The Role of Interpretive Signage in Enhancing Conservation Knowledge, Awareness and Behaviour Among Visitors at the Penang National Park*, Masters dissertation, Universiti Putra Malaysia, Serdang, Malaysia.
 44. Zahiruddin WM et al., 2018. Development and validation of a new knowledge, attitude, belief and practice questionnaire on leptospirosis in Malaysia. *BMC Public Health* 18: 331.
 45. Kostenius C, Hallberg J, Lindqvist AK, 2018. Gamification of health education. *Health Educ* 118: 354–368.
 46. Azhari NN, Abdul Manaf R, Ng SW, Shakeeb Arsalaan Bajunid SFB, Mohd Gobil AR, Saad WZ, Amin Nordin S, 2019. Gamification, a successful method to foster leptospirosis knowledge among university students: a pilot study. *Int J Environ Res Public Health* 16: 2108.
 47. Thayaparan S, Robertson ID, Fairuz A, Suut L, Abdullah MT, 2013. Leptospirosis, an emerging zoonotic disease in Malaysia. *Mal J Pathol* 35: 123–132.
 48. El-Jalii IM, 2000. *Epidemiology and Diagnosis of Human Leptospirosis in Malaysia*, Doctoral dissertation, Universiti Putra Malaysia, Serdang, Malaysia.
 49. Tan WL, Soelar SA, Suan MAM, Hussin N, Cheah WK, Verasahib K, Goh PP, 2016. Leptospirosis incidence and mortality in Malaysia. *Southeast Asian J Trop Med Public Health* 47: 434–440.