

LECTURE PRESENTATION

Open Access

Tumours of the prostate

F Grillone

From de Senectute: Age and Health Forum
Catanzaro, Italy. 5-7 December 2009

Background

In the United States, Prostate cancer is the most common tumour in men, with an expected 192.280 new cases and 27.360 deaths estimated in 2009 [1]. About 81% of patients with prostate cancer are over 65 years of age. Prostate cancer is a biologically heterogeneous neoplasm in which, especially in the elderly, some forms remain silent and with risk of disease-treatment. Moreover, this population may not receive optimal therapies for their disease, if decisions are based only on their chronological age. Health status, more than age, is a major factor affecting individual life expectancy. Comorbidity is the key predictor of health status, and should weigh more on the treatment decision than age alone. Other important parameters to consider in the elderly are the degree of dependence in activities of daily living, the nutritional status and the presence or not of a geriatric syndrome. The androgen-deprivation therapy (ADT) has become an integral piece of the armamentarium for treating prostate cancer in neoadjuvant, adjuvant and palliative therapy [2]. The impact on health-related quality of life (HRQOL) following ADT induction has gained significant clinical attention. Several series documented an association between ADT and declining HRQOL [3]. Treatment of hormone refractory metastatic prostate cancer with Docetaxel 75 mg/m² every 3 weeks (3W) in combination with daily prednisone has been shown to prolong survival and to have the same efficacy in healthy elderly as in younger patients [4].

Conclusions

Geriatric assessment in elderly patients with prostate cancer is crucial. Each treatment has to be adapted to health status.

Published: 19 May 2010

Medical Oncology Unit, Magna Graecia University and Tommaso Campanella Cancer Center, Campus Salvatore Venuta, Catanzaro, Italy

References

1. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ: **Cancer Statistics, 2009.** *Ca Cancer J Clin* 2009, **59**:225-249.
2. Kumar S, Shelley M, Harrison C, Coles B, Wilt TJ, Mason MD: **Neo-adjuvant and adjuvant hormone therapy for localised and locally advanced prostate cancer.** *Cochrane Database Syst Rev* 2006, **18**(4):CD006019.
3. Dacal K, Sereika SM, Greenspan SL: **Quality of life in prostate cancer patients taking androgen deprivation therapy.** *J Am Geriatr Soc* 2006, **54**:85-90.
4. Tannock IF, de Wit R, Berry WR, Horti J, Pluzanska A, Chi KN, Oudard S, Théodore C, James ND, Turesson I, Rosenthal MA, Eisenberger MA: **Docetaxel and prednisone or mitoxantrone plus prednisone for advanced prostate cancer.** *N Engl J Med* 2004, **351**:1502-1512.

doi:10.1186/1471-2318-10-S1-L37

Cite this article as: Grillone: Tumours of the prostate. *BMC Geriatrics* 2010 **10**(Suppl 1):L37.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

BioMed Central