


# BMJ Open Multidisciplinary approach to functional somatic syndromes: study protocol for a population-based prospective cohort study

Peyman Adibi,<sup>1,2</sup> Alireza Ani,<sup>3,4</sup> Ahmad Vaez ,<sup>3,4</sup> Fatemeh Hadizadeh,<sup>4</sup> Harold Snieder,<sup>3</sup> Hamidreza Roohafza<sup>5</sup>

**To cite:** Adibi P, Ani A, Vaez A, *et al.* Multidisciplinary approach to functional somatic syndromes: study protocol for a population-based prospective cohort study. *BMJ Open* 2022;**12**:e048941. doi:10.1136/bmjopen-2021-048941

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-048941>).

Received 11 January 2021  
Accepted 16 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Isfahan Gastroenterology and Hepatology Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup>Department of Internal Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>3</sup>Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>4</sup>Department of Bioinformatics, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>5</sup>Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

## Correspondence to

Dr Ahmad Vaez;  
[a.vaez@umcg.nl](mailto:a.vaez@umcg.nl)

## ABSTRACT

**Introduction** Isfahan functional disorders (ISFUN) cohort study aims to describe the interplay of genetic and environmental factors in shaping the characteristics of functional somatic syndromes (FSS). This study is primarily intended to investigate the epidemiology, risk factors, course and prognosis of FSSs in a sample of adult Iranian population. The other aim is to develop a new delimitation of FSSs based on an integrated multidisciplinary approach comprising of phenotypic and multiomics data.

**Methods and analysis** ISFUN is a population-based prospective cohort study designed to follow a population of randomly selected seemingly healthy adults (18–65 years) through annual visits during a 4-year observation period. Structured questionnaires are used for data collection and clinical assessment of the participants. Questionnaire-based diagnosis of FSSs are validated in a medical interview. Human DNA genotyping, microbial amplicon sequencing and urine analysis is under progress for genomics, microbiota and metabolomics profiling, respectively. Enrolment began in September 2017, and study completion is expected in 2022. A total number of 1943 participants were initially recruited.

**Ethics and dissemination** Ethical approval for data collection was granted by the National Research Ethics Committee of the Iranian Ministry of Health and Medical Education and the Research Ethics Committee of Isfahan University of Medical Sciences (IR.MUI.REC.1395.1.149). Following the description of the study procedure, we obtained written informed consent from all study participants. Study findings will be disseminated through peer-reviewed publications and presentations at scientific meetings.

## INTRODUCTION

Among the most common reasons for which people seek medical care are the so-called bodily complaints without an apparent organic origin.<sup>1 2</sup> It is reported that the main ailments of almost one-third of patients referred to the family physician are non-specific, somatic symptoms with no precise pathology.<sup>3</sup> More often than not, these complaints are not attributable to a

## Strengths and limitations of this study

- ⇒ This multidisciplinary population-based longitudinal cohort study does not have a disease-oriented approach and a diverse range of data including somatic symptoms, biomarkers, psychological measures and lifestyle information in addition to genomic, faecal microbiota and urine metabolomics profiles of the participants is being collected.
- ⇒ Variations in chronicity and severity of the functional symptoms are checked over a 5-year period in an effort to further our understanding of the course, risk factors and prognosis of functional somatic syndromes (FSSs).
- ⇒ The longitudinal study design coupled with serial collection of clinical data, biological samples and omics profiles will provide a unique platform for personalised medicine and omics-based clinical management of patients suffering from FSSs.
- ⇒ This cohort might lack the power to study rare FSSs.

conventionally defined medical disease or mental disorder and no organic causes can be identified even after complementary diagnostic procedures.<sup>4</sup> Multiple terms are proposed for such symptoms from which the umbrella concept of functional somatic syndromes (FSSs) is the accepted choice in medical specialties.<sup>5 6</sup>

The aetiology or underlying mechanisms of FSSs are not yet fully understood. However, various elements like biological, psychological and social factors and possibly the complex interactions among them are all potentially important.<sup>7 8</sup> Patients frequently meet the diagnostic criteria for more than one syndrome and usually share other non-clinically relevant characteristics.<sup>9 10</sup> Also, occurrence of a new syndrome could be predicted based on antecedent syndromes which implies a common etiopathogenesis.<sup>11</sup> It is suggested that FSSs might be a family

of closely related disorders or expressions of the same underlying illness with various subtypes.<sup>12</sup>

It is difficult to deal with such patients in medical practice because a disease-based explanation cannot be made and the direction of diagnosis mainly hinges on the speciality of the medical consultant rather than the nature of the disorder itself. Experts in each field of medicine have defined and named the syndromes according to their organ of interest without a holistic multiorgan or system view.<sup>13</sup> Patients may receive diagnoses such as irritable bowel syndrome, functional dyspepsia, fibromyalgia, non-cardiac chest pain, chronic fatigue syndrome, etc depending on the physician's speciality.<sup>14</sup> Hence, a more precise syndrome/symptoms identification system is needed for better patient management.<sup>15</sup> Latent class analysis and cluster analysis of the somatic symptoms have been previously used for creating symptom profiles.<sup>9 16 17</sup> But we believe the best way to overcome this challenge is through an integrated multidisciplinary approach comprising of both phenotypic and multiomics data.

Therefore, in this cohort study, we are gathering a multifaceted longitudinal dataset including functional symptoms, biomarkers, sociodemographic data, psychological measures and lifestyle information in addition to genomic, faecal microbiota and urine metabolomics profiles of seemingly healthy participants. This will give us the basic requirements for creating a new quantitative definition of FSSs via data modelling and machine learning methods. To the best of our knowledge, this is the first cohort study in the Middle Eastern region aimed at investigating FSSs.

### Study aims

The Isfahan functional disorders (ISFUN) cohort study is a population-based longitudinal cohort designed to:

1. Investigate the epidemiology, risk factors, course and prognosis of FSSs in a sample of adult Iranian population. The questionnaire-based diagnoses will be validated against diagnoses made by a personal medical interview as the gold standard method.
2. Challenge the organ/system-oriented view of FSSs by creating a personalised profile according to the clinical, psychological and omics characteristics of each individual. This profile is an effort to plan a tailor-made medical strategy towards personalised medicine.

## METHODS AND ANALYSIS

### Study population and exclusion criteria

ISFUN study is centred in the Kerdabad neighbourhood of Isfahan, Iran. The main criteria for selecting this location were population diversity of the inhabitants and its generalisability to the main population of Isfahan city according to their mean household income and socioeconomic status. All households from the fourth and fifth district of this area were included using census-based sampling technique. Subsequently, one person per household was randomly selected and if this nominee

did not meet our eligibility criteria, another random individual from the same household was selected and invited to participate in the study. All seemingly healthy adults (aged between 18 and 65) were included except for non-Iranian nationals, women who were pregnant or lactating, close relatives of a previously selected individual (up to the third degree) and a small number who suffered physical or mental disability to the degree that could not participate in providing reliable data on themselves (eg, severe mental retardation or untreated psychosis). Caregivers in the province health centre were responsible for evaluating the selected population using official family and personal health records or a phone call in case of missing data. Cohort management team created the initial participants list, supervised the recruitment process and decided on the new invitations or when to exclude an individual.


### Recruitment and follow-up

Recruitment was planned over 6 months beginning on September 2017. First, the objectives of the study were explained by a trained interviewer in a household visit. A custom-made package including introductory brochure, questionnaires, writing paper and matching envelopes, urine and stool specimen containers and guidelines on how to collect biological samples were also offered. The invitee was given a few days to read the educational materials. A direct phone line to an informed caregiver was provided for any questions or concerns remained unanswered. After consenting to join the study, a visit at the local health centre was scheduled during which the participants underwent clinical examination and medical interviews. Questionnaires were completed at home and controlled in different working stations.

There is a natural variance in functional symptoms characteristics over time which affects patient-reported complaints. Thus, the participants are followed through three scheduled annual visits (2019–2021) after baseline data collection to check the chronicity and severity of their symptoms and search for possible responsible factors. Two rounds of annual follow-ups are done and the next is planned to commence in fall 2021. [Figure 1](#) illustrates a flow chart of the ISFUN cohort study.

### Outcomes and measurements

Questionnaire data and phenotypic measurements were gathered at baseline and the following annual visits. The cohort management team selected the current list of measurements after a rapid review of existing methods followed by extensive consultation with experts in this field of medicine including psychologists, neurologists, internal medicine specialists and gastroenterologists. Our aim was to use the most reliable, applicable and popular measurements that are currently applied in medical practice for diagnosing FSSs in each body organ. The questionnaire-based data including psychological evaluation, dietary intake and functional symptoms were validated in personal medical visits with a psychologist,



**Figure 1** Flow chart of the Isfahan functional disorders cohort study.

a dietician and a physician, respectively, as the gold-standard method. A fixed medical team interviewed all the individuals and were responsible for making sure whether the questions are correctly comprehended and answered. This was done in three separate work stations and could take up to an hour. Incomplete questions were filled during this visit and interview-based diagnosis of the symptoms was also recorded. Phenotype assessment tools and evaluation schedule are described in online supplemental tables 1 and 2.

#### Physical examination and anthropometric measurements

Clinical examination including measurements of blood pressure, anthropometric parameters (ie, weight, height, neck, chest, waist and hip circumferences), body composition analysis (by bioelectrical impedance), lung function (ie, spirometry), pain sensitivity (using a handheld algometer) and adipose tissue thickness (by skinfold calliper) are performed at baseline and during subsequent follow-ups.

#### Laboratory methods

Participants were asked to refer to the local health centre early in the morning after 8 hours of fasting. Participants were requested to collect morning midstream urine and a stool specimen in the provided containers at home just before leaving for the health centre. Biological samples were collected on arrival and transferred to the laboratory under cold chain conditions. Stool, urine and fasting blood samples are collected from all the participants. As a quality control indicator, 5% of samples were rechecked for all the metabolic markers. Additional samples of whole blood, serum, plasma, stool, urine and DNA samples are preserved in a biobank for future complementary studies. General laboratory evaluation was performed for all participants in the baseline phase of the study, but only the more important and more variable diagnostic tests were done in the consequent visits due to budget limitations. Laboratory tests and instruments are described in online supplemental table 1.

#### Biota

DNA has been extracted from blood samples of all participants to be genotyped on Illumina Global Screening Array. Faecal microbial DNA has been extracted for 16S rRNA amplicon sequencing. DNA samples were subjected to spectrophotometric assay for quality assurance.

#### Dietary intake

Dietary intake assessment is done using a dish-based, semiquantitative Food frequency questionnaire for the Iranian adult population.<sup>18</sup> A qualified dietician assessed the filled questionnaires. Missing or doubtful responses were completed and verified during a medical interview. Participants are reassessed at each annual visit.

#### Questionnaires

Questionnaires are used to obtain detailed information on functional symptoms, psychological assessment, life-style evaluation, quality of life and sociodemographic information. Considering that some health behaviours or psychological traits (such as eating attitude and coping mechanisms) are fairly stable over time and the high number of questionnaire items for each visit that could cause respondent fatigue, we decided that there was no need to include every questionnaire in all annual visits. However, symptoms profiles, psychological traits and behaviours that are expected to vary over time were checked annually. Four questionnaires were not included in the baseline survey and were added in the first follow-up visit. The list of questionnaires for each phase of the study is mentioned in online supplemental table 2.

#### Data analysis

##### Statistical analysis

All data are collected in a central database using SPSS Version 23 software according to a standardised protocol. The results of participants' demographics, questionnaire-based prevalence of functional symptoms and other outcome variables will be summarised using descriptive summary measures, expressed as mean±SD for continuous variables and percentages and frequencies for categorical variables. Student's t-test will be used to assess differences between two means.  $\chi^2$  test will be used to assess the degree of association between categorical variables. A two-sided  $p < 0.05$  will be considered statistically significant. The diagnostic agreement between questionnaire-based and interview-based diagnosis of syndromes will be assessed using kappa statistic. Exploratory factor analysis and latent class analysis will be performed to detect qualitatively different subgroups inside the study group. Random-effects regression models will be done for time-series analysis. The correlation between genomic and meta-genomic variations with FSSs will be explored in genome wide and meta-genome wide association study analyses, respectively.

#### Sample size

This is an exploratory observational study and the sample size was based on pragmatic considerations, given time

and facility constraints of the study along with local capacity. Taking the high prevalence of FSSs into account, it was predicted that a total number of 2000 participants would suffice. We estimate that we would have 90% power to detect the hypothetical associations between most of the functional symptoms and the biota/biological data.

### Ethics and dissemination

The study was approved by the National Research Ethics Committee of the Iranian Ministry of Health and Medical Education and the Research Ethics Committee of Isfahan University of Medical Sciences (IR.MUI.REC.1395.1.149). All participants agreed to participate and signed a written consent form after receiving written and oral information about the study. Subjects consented to undertake annual investigations as set out in the study schedule and for long-term access to their collected biological samples. Study findings will be published in peer-reviewed scientific journals and will be presented at national and international conferences.

### DISCUSSION

Functional somatic symptoms (FSSs) are common and are shown to be a risk factor for poor health status and high healthcare use. Despite the high prevalence of FSSs and the burden they put on public health, very few population-based cohorts, such as the Danish study of Functional Disorders have been carried out to investigate their epidemiology and clinical features.<sup>19–25</sup> A robust classification system of functional disorders is still lacking and although various approaches are suggested for their delimitation, each has its own strengths and weaknesses and hence, more studies are required for better definition and categorization of patient symptoms.<sup>26</sup> A major issue in this field is that symptoms could occur in multiple body organs belonging to various medical specialties which warrants the need for a holistic view when confronting such disorders. Moreover, co-occurrence of symptoms could play an important role in diagnosis and treatment of affected individuals. Symptoms clusters based on functional symptoms spread over body organs have shown promising results for studying the risk factors and prognosis of FSSs.<sup>10 16</sup> We believe this approach could be further improved when corrected for personalised traits such as psychological, life style and biota characteristics.

This longitudinal population-based cohort study is the first nationwide survey of epidemiological characteristics of FSSs in Iranian adult population. In this study, we collect a validated multidomain dataset containing medical, psychological, lifestyle and multiomics data. This will be used to investigate symptoms associations and to suggest a new delimitation of FSS by identifying similar symptom patterns in combination with personalised omics profile of the individuals.

Availability of omics data, repeated measurements of different phenotypes over time and a biobank of blood, serum, plasma, stool and urine samples are among the

other important strengths of this study. On the other hand, although our sample size is relatively large, it may still not be sufficient for evaluating rare FSSs.

**Acknowledgements** We wish to thank all participants and their families for their commitment to this study. We acknowledge the management of Isfahan province health centre and also Isfahan health centre #1 for their administrative supports. We also acknowledge the staff and health workers at Kerdabad health centre for their gracious role in recruiting and following the participants. We thank the Rome Foundation for supporting this publication. We appreciate the support from Ammar Hassanzadeh Keshteli and Hassan Shahoon for their time and effort in this project.

**Contributors** PA is the lead investigator of this cohort study. PA, AV and HR contributed to the conception and design of the study. PA and HR were responsible for the construction of the questionnaires and HR handled data entry and primary analyses. AV, FH, AA, HR and HS provided methodological and clinical consults on various aspects of the study design and participant recruitment. AA drafted the manuscript. All authors were involved in critical revision of this cohort profile and approved the final manuscript.

**Funding** This work has been supported by the Iranian Ministry of Health and Medical Education (700/144), Isfahan University of Medical Sciences (195149) and the Iranian National Institute for Medical Research Development (NIMAD) (972 953).

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iD

Ahmad Vaez <http://orcid.org/0000-0001-9048-3795>

### REFERENCES

- Henningsen P, Zipfel S, Sattel H, *et al*. Management of functional somatic syndromes and bodily distress. *Psychother Psychosom* 2018;87:12–31.
- Finset A. Why do doctors not learn how to explain "medically unexplained symptoms"? *Patient Educ Couns* 2018;101:763–4.
- Rosendal M, Carlsen AH, Rask MT. Symptoms as the main problem: a cross-sectional study of patient experience in primary care. *BMC Fam Pract* 2016;17:29.
- Harvey SB, Wessely S. How should functional somatic syndromes be diagnosed? *Int J Behav Med* 2013;20:239–41.
- Barsky AJ, Borus JF. Functional somatic syndromes. *Ann Intern Med* 1999;130:910–21.
- Rief W, Isaac M. The future of somatoform disorders: somatic symptom disorder, bodily distress disorder or functional syndromes? *Curr Opin Psychiatry* 2014;27:315–9.
- D'Souza RS, Hooten WM. Somatic syndrome disorders., 2019. Available: <https://www.ncbi.nlm.nih.gov/books/NBK532253/> [Accessed 4 Feb 2020].
- Kato K, Sullivan PF, Evengård B, *et al*. A population-based twin study of functional somatic syndromes. *Psychol Med* 2009;39:497–505.
- Lacourt T, Houtveen J, van Doornen L. "Functional somatic syndromes, one or many?" An answer by cluster analysis. *J Psychosom Res* 2013;74:6–11.

- 10 Eliassen M, Jørgensen T, Schröder A, *et al.* Somatic symptom profiles in the general population: a latent class analysis in a Danish population-based health survey. *CLEP* 2017;9:421–33.
- 11 Warren JW, Langenberg P, Clauw DJ. The number of existing functional somatic syndromes (FSSs) is an important risk factor for new, different FSSs. *J Psychosom Res* 2013;74:12–17.
- 12 Wessely S, Nimnuan C, Sharpe M. Functional somatic syndromes: one or many? *The Lancet* 1999;354:936–9.
- 13 Fantoni F, Salvetti G, Manfredini D, *et al.* Current concepts on the functional somatic syndromes and temporomandibular disorders. *Stomatologija* 2007;9:3–9.
- 14 Frølund Pedersen H, Holsting A, Frostholm L, *et al.* "Understand your illness and your needs": Assessment-informed patient education for people with multiple functional somatic syndromes. *Patient Educ Couns* 2019;102:1662–71.
- 15 Rief W, Burton C, Frostholm L, *et al.* Core outcome domains for clinical trials on somatic symptom disorder, bodily distress disorder, and functional somatic syndromes: European network on somatic symptom disorders recommendations. *Psychosom Med* 2017;79:1008–15.
- 16 Eliassen M, Schröder A, Fink P, *et al.* A step towards a new delimitation of functional somatic syndromes: a latent class analysis of symptoms in a population-based cohort study. *J Psychosom Res* 2018;108:102–17.
- 17 Witthöft M, Fischer S, Jasper F, *et al.* Clarifying the latent structure and correlates of somatic symptom distress: a bifactor model approach. *Psychol Assess* 2016;28:109–15.
- 18 Keshteli A, Esmailzadeh A, Rajaie S, *et al.* A Dish-based semi-quantitative food frequency questionnaire for assessment of dietary intakes in epidemiologic studies in Iran: design and development. *Int J Prev Med* 2014;5:29–36.
- 19 Dantoft TM, Ebstrup JF, Linneberg A, *et al.* Cohort description: the Danish study of functional disorders. *Clin Epidemiol* 2017;9:127–39. Volume.
- 20 Budtz-Lilly A, Vestergaard M, Fink P, *et al.* The prognosis of bodily distress syndrome: a cohort study in primary care. *Gen Hosp Psychiatry* 2015;37:560–6.
- 21 Ahmad O, Ahmad KE. Functional neurological disorders in outpatient practice: an Australian cohort. *J Clin Neurosci* 2016;28:93–6.
- 22 Markkula RA, Kalso EA, Kaprio JA. Predictors of fibromyalgia: a population-based twin cohort study. *BMC Musculoskelet Disord* 2016;17:29.
- 23 Krogsgaard LR, Engsbro AL, Jones MP, *et al.* The epidemiology of irritable bowel syndrome: symptom development over a 3-year period in Denmark. A prospective, population-based cohort study. *Neurogastroenterol Motil* 2017;29. doi:10.1111/nmo.12986. [Epub ahead of print: 14 Nov 2016].
- 24 Joustra ML, Janssens KAM, Bültmann U, *et al.* Functional limitations in functional somatic syndromes and well-defined medical diseases. results from the general population cohort lifelines. *J Psychosom Res* 2015;79:94–9.
- 25 Harvey SB, Wadsworth M, Wessely S, *et al.* Etiology of chronic fatigue syndrome: testing popular hypotheses using a national birth cohort study. *Psychosom Med* 2008;70:488–95.
- 26 Petersen MW, Schröder A, Eliassen MH, *et al.* Three different approaches to delimitation of functional somatic disorders: DanFunD. *J Psychosom Res* 2021;145:110475.