

SCIENTIFIC REPORTS



OPEN

Quantum Fisher information of atomic system interacting with a single cavity mode in the presence of Kerr medium

N. Zidan^{1,2}, H. F. Abdel-Hameed^{1,3} & N. Metwally^{4,5}

The quantum Fisher information of an atomic system interacting with a single cavity mode in the presence of Kerr medium is discussed. It is shown that quantum Fisher information for an initial separable atomic system is larger than that depicted for the initial entangled atomic system. For initial vacuum state of the cavity mode, the quantum Fisher information with respect to the Kerr medium and the phase decoherence parameter is larger than that displayed for the detuning parameter. Both phase decoherence and Kerr medium have the same effect on the decay of quantum Fisher information, while they have an opposite effect on its maximum values.

Fisher information is proposed to detect the precision of the parameter in estimation theory¹. Extending the estimation theory from the classical regime to the quantum regime, quantum Fisher information plays a paramount role in quantum information theory^{2–4} and quantum metrology^{5,6}. Quantum Fisher information has been widely studied theoretically and experimentally^{7–18}. Quantum Fisher information has become a highly studied subject because it measures the phase sensitivity of the systems^{19–23}. Moreover, quantum Fisher information in non-inertial frames has been investigated for different systems, such as the performance of quantum Fisher information under the Unruh-Hawking effect in the context of relativistic quantum information theory²⁴, the dynamic of the teleported quantum Fisher information²⁵, and the Unruh acceleration effect on the precision of parameter estimation for a general two-qubit system²⁶. The existence of the maximal quantum information matrix in multi-parameter quantum estimation²⁷ and entanglement detection have been discussed extensively^{28,29}. The dynamics of quantum Fisher information in the two-level system coupled to a single bosonic reservoir^{30,31} and multiple bosonic reservoirs^{32,33} has been discussed. Quantum Fisher information is used as a predictor of decoherence in the preparation of spin-cat states for quantum metrology³⁴. Also, quantum Fisher information of a single qubit has quantified^{35,36}. The dynamics of quantum Fisher information is studied and employed as a measure to quantify the precision of the estimation of two interacting qubits subject to decoherence³⁷. The weak measurement based pre- and post-flips is used to protect the average quantum Fisher information in the independent amplitude damping channel for N-qubit GHZ states³⁸. The energy-level crossing behavior in a two-dimensional well with the Rashba and Dresselhaus-orbit coupling is studied and the approximate ground state and its quantum Fisher information via performing a unitary transformation is obtained³⁹. Furthermore, enhancing parameter precision optimal quantum estimation by direct quantum feedback⁴⁰ and quantum screening were reported⁴¹. The studying how to estimate two parameters in a spin-boson dephasing system by periodical projective measurements has been investigated⁴². The quantum Fisher information of a two charged qubits system interact locally with a dephasing channel is quantified with respect to the charged qubits and the channel parameters⁴³. For the Jaynes-Cummings-Dicke model, enhanced spin squeezing and quantum entanglement have been explained⁴⁴. Also, it has been shown that the Kerr-like medium plays an important role in generating twin beams by parametric amplification four-wave-mixing beams and triplet beams from the parametric amplification six-wave mixing process, where it can be used as a control parameter⁴⁵. Therefore we are motivated to consider, an atomic

¹Mathematics Department, Faculty of Science, Sohag University, Sohag, Egypt. ²Mathematics Department, College of Science, Jouf University, Sakaka, Saudi Arabia. ³Mathematics Department, Khurma University College, Taif University, Al-Taif, Saudi Arabia. ⁴Mathematics Department, College of Science, Bahrain University, Zallaq, Bahrain. ⁵Mathematics Department, Faculty of Science, Aswan University, Aswan, Egypt. Correspondence and requests for materials should be addressed to N.Z. (email: nazidan@yahoo.com) or H.F.A.-H. (email: hf_elsheikh@yahoo.com) or N.M. (email: nmetwally@gmail.com)

system containing two-atoms interacting with a single cavity mode with Kerr medium in the presence of a phase decoherence. Our contribution focus on estimating the parameters of the system by evaluating the quantum Fisher information. We focus on three parameters, the Kerr medium (χ), phase decoherence (γ) and detuning (Δ) and study the effect of these parameters on the behavior of quantum Fisher information. As an application of this model is generating a three qubit entangled state between the two atoms and the cavity mode. Also, this system may be used to teleport information from one party to another one between the three qubits. Moreover, it can be used to perform entanglement swapping between the one atom of the atomic system and the cavity mode. Therefore, It is important to know the values of the parameters that maximize/minimize the efficiency to perform these applications.

This paper is organized as follows: We introduce the model and calculation the reduced density matrix of two-two level atoms in sect. 2. In sect. 3, some properties of the quantum Fisher information are given and then we estimate the parameters of the system. Numerical discussion of our results is presented in sect. 4. Finally, the conclusion is given in sect. 5.

The Suggested Model

It is assumed that an atomic system consists of two two-level atoms interacting with a single cavity mode field which is initially prepared in the Fock state $|n + 1\rangle$. In the rotating wave approximation, the Hamiltonian which describes this system is given by⁴⁶:

$$\mathcal{H} = \omega_f a^\dagger a + \frac{\omega_a}{2} \sum_{j=A,B} \sigma_z^{(j)} + \kappa \sum_{j=A,B} \{a\sigma_+^{(j)} + a^\dagger\sigma_-^{(j)}\} + \chi(a^\dagger a a^\dagger a), \tag{1}$$

where $a(a^\dagger)$ denotes to the annihilation (creation) operator of the single cavity mode, ω_a and ω_f are the atomic and field transition frequencies, respectively. Here χ represents the coupling of the field induced by the Kerr medium and κ is the coupling constant between the atoms and field. $\sigma_z^{(j)} = |1\rangle_j\langle 0| - |0\rangle_j\langle 1|$, $\sigma_+^{(j)} = |1\rangle_j\langle 0|$, $\sigma_-^{(j)} = |0\rangle_j\langle 1|$ with $|1\rangle_j$ and $|0\rangle_j$ being the excited and ground states of j th atom ($j=A, B$). By the basis, $|n, 11\rangle$, $|n + 1, 10\rangle$, $|n + 1, 01\rangle$ and $|n + 2, 00\rangle$ the eigenvalues of Hamiltonian (1) can be given by:

$$\begin{aligned} E_1 &= \Delta - 2(n + 1)\chi, \quad E_2 = -\chi, \\ E_{3,4} &= \frac{1}{2}(\Delta - (2n + 3)\chi \mp \delta), \end{aligned} \tag{2}$$

where $\delta = \sqrt{8\kappa^2(2n + 3) + (-\Delta + \chi + 2n\chi)^2}$ and $\Delta = \omega_a - \omega_f$ is the detuning of the cavity field. The corresponding eigenvectors are given by:

$$\begin{aligned} |\psi_1\rangle &= -\sqrt{\frac{n+2}{n+3}}|n,11\rangle + \sqrt{\frac{n+1}{n+3}}|n+2,00\rangle, \\ |\psi_2\rangle &= \frac{1}{\sqrt{2}}(-|n+1,10\rangle + |n+1,01\rangle), \\ |\psi_3\rangle &= \frac{2\kappa\sqrt{n+1}}{\sqrt{\eta_1}}|n,11\rangle + \frac{2\kappa\sqrt{n+2}}{\sqrt{\eta_1}}|n+2,00\rangle \\ &\quad - \frac{\sqrt{\eta_1}}{2\delta}(|n+1,10\rangle + |n+1,01\rangle), \\ |\psi_4\rangle &= \frac{2\kappa\sqrt{n+1}}{\sqrt{\eta_2}}|n,11\rangle + \frac{2\kappa\sqrt{n+2}}{\sqrt{\eta_2}}|n+2,00\rangle \\ &\quad + \frac{\sqrt{\eta_2}}{2\delta}(|n+1,10\rangle + |n+1,01\rangle). \end{aligned} \tag{3}$$

Here $\eta_1 = \delta(\delta + \Delta - (2n + 1)\chi)$, $\eta_2 = \delta(\delta - \Delta + (2n + 1)\chi)$.

The master equation which governs the time evolution of the system under the Markovian approximation is given by⁴⁷:

$$\frac{d\rho}{dt} = -i[\mathcal{H}, \rho] - \frac{\gamma}{2}[\mathcal{H}, [\mathcal{H}, \rho]], \tag{4}$$

where γ is the phase decoherence coefficient. The formal solution of the master Eq. (4) may be expressed in the following form⁴⁸:

$$\rho(t) = \sum_{k=0}^{\infty} \frac{(\gamma t)^k}{k!} \mathcal{M}^k(t)\rho(0)\mathcal{M}^{\dagger k}(t), \tag{5}$$

where $\rho(0)$ is the density operator of the initial atom-field system. $\mathcal{M}^k(t)$ is defined as:

$$\mathcal{M}^k(t) = \mathcal{H}^k \exp(-it\mathcal{H}) \exp\left(-\frac{\gamma t}{2}\mathcal{H}^2\right). \tag{6}$$

Let us assume that the cavity field is initially prepared in the Fock state $|\psi_f(0)\rangle = |n+1\rangle$ and the atomic system is initially prepared in the entangled states $|\psi_{AB}(0)\rangle = \cos\varphi|10\rangle + \sin\varphi|10\rangle$. Then the initial state of the total system may be written as:

$$|\psi_s(0)\rangle = |\psi_{AB}(0)\rangle \otimes |\psi_f(0)\rangle. \quad (7)$$

Using Eqs (5) and (7) we can obtain the reduced density matrix of the atomic system $\rho_{AB}(t)$ after taking the trace over the field in the form

$$\rho_{AB}(t) = \rho_{11}|11\rangle\langle 11| + \rho_{22}|10\rangle\langle 10| + \rho_{33}|01\rangle\langle 01| + \rho_{23}|10\rangle\langle 01| + \rho_{32}|01\rangle\langle 10| + \rho_{44}|00\rangle\langle 00|, \quad (8)$$

where

$$\begin{aligned} \rho_{11} &= \frac{2\kappa^2(n+1)}{\delta^2}X, & \rho_{44} &= \frac{2\kappa^2(n+2)}{\delta^2}X, \\ \rho_{22} &= \frac{1}{2} - \frac{\kappa^2(2n+3)}{\delta^2}X \\ &\quad + \frac{\cos(2\varphi)}{4\delta^2} \left(Y_1 \cos\left[t\left(\frac{\eta_1}{2\delta}\right)\right] + Y_2 \cos\left[t\left(\frac{\eta_2}{2\delta}\right)\right] \right), \\ \rho_{33} &= \frac{1}{2} - \frac{\kappa^2(2n+3)}{\delta^2}X \\ &\quad - \frac{\cos(2\varphi)}{4\delta^2} \left(Y_1 \cos\left[t\left(\frac{\eta_1}{2\delta}\right)\right] + Y_2 \cos\left[t\left(\frac{\eta_2}{2\delta}\right)\right] \right), \\ \rho_{23} &= \rho_{32}^* = \frac{\sin(2\varphi)}{2} - \frac{\kappa^2(2n+3)}{\delta^2}X \\ &\quad - \frac{i\cos(2\varphi)}{4\delta^2} \left(Y_1 \sin\left[t\left(\frac{\eta_1}{2\delta}\right)\right] - Y_2 \sin\left[t\left(\frac{\eta_2}{2\delta}\right)\right] \right). \end{aligned}$$

with

$$\begin{aligned} X &= (1 + \sin(2\varphi)) \left(1 - \exp\left[-\frac{\gamma t \delta^2}{2}\right] \cos(t\delta) \right), \\ Y_1 &= \eta_1 \exp\left[-\frac{\gamma t}{2} \left(\frac{\eta_1}{2\delta}\right)^2\right], & Y_2 &= \eta_2 \exp\left[-\frac{\gamma t}{2} \left(\frac{\eta_2}{2\delta}\right)^2\right] \end{aligned}$$

Quantum Fisher Information

Quantum Fisher information of any parameter characterizes the sensitivity of the state with respect to changes of this parameter. Let us assume that the parameter τ is coded in the quantum state ρ_τ . The quantum Fisher information with respect to the parameter τ is defined as^{2,49}:

$$\mathcal{F}_\tau = \text{Tr}(\rho_\tau \mathcal{E}_\tau^2) = \text{Tr}\left(\frac{\partial \rho_\tau}{\partial \tau} \mathcal{E}_\tau\right), \quad (9)$$

where \mathcal{E}_τ is so-called symmetric logarithmic derivative which is defined by:

$$\frac{\partial \rho_\tau}{\partial \tau} = (\mathcal{E}_\tau \rho_\tau + \rho_\tau \mathcal{E}_\tau)/2. \quad (10)$$

Typically there are three methods to calculate the quantum Fisher information²⁴. The most frequently method used is the diagonalizing of the matrix $\rho_\tau = \sum_{i=1}^n \lambda_i |V_i\rangle\langle V_i|$, where λ_i and $|V_i\rangle$ are the eigenvalues and eigenvectors of the density operator ρ_τ , respectively. Then the quantum Fisher information \mathcal{F}_τ with respect to the parameter τ is written as^{50,51}:

$$\begin{aligned} \mathcal{F}_\tau &= \sum_{i=1}^n \frac{1}{\lambda_i} \left(\frac{\partial \lambda_i}{\partial \tau} \right)^2 + 4 \sum_{i=1}^n \lambda_i \left(\left\langle \frac{\partial V_i}{\partial \tau} \middle| \frac{\partial V_i}{\partial \tau} \right\rangle - \left| \left\langle V_i \middle| \frac{\partial V_i}{\partial \tau} \right\rangle \right|^2 \right) \\ &\quad - 8 \sum_{i \neq j}^n \frac{\lambda_i \lambda_j}{\lambda_i + \lambda_j} \left| \left\langle V_i \middle| \frac{\partial V_j}{\partial \tau} \right\rangle \right|^2. \end{aligned} \quad (11)$$




Figure 1. Quantum Fisher information \mathcal{F}_χ as a function of time t with $\kappa = 1$, $\gamma = 0.02$ and $\chi = 0.1$ for different initial state settings. The dotted-black, solid-red, and dashed-blue curves represent \mathcal{F}_χ for a weight angle $\theta = 0, \pi/8$, and $\pi/3$, respectively. Here: (a) $n = 0, \Delta = 0$, (b) $n = 0, \Delta = 1$ (c) $n = 1, \Delta = 0$, (d) $n = 1, \Delta = 1$.

To calculate the quantum Fisher information of the suggested atomic system, one needs to find the eigenvalues and eigenvectors of the reduced density operator Eq. (8) of the atomic system. In this context, the eigenvalues are given explicitly by:

$$\lambda_1 = \rho_{11}, \quad \lambda_2 = \rho_{44}, \quad \lambda_{3,4} = \frac{1}{2} \left(\rho_{22} + \rho_{33} \pm \sqrt{(\rho_{22} - \rho_{33})^2 + 4|\rho_{23}|^2} \right), \quad (12)$$

with the corresponding eigenvectors:

$$\begin{aligned} |V_1\rangle &= (1, 0, 0, 0), \quad |V_2\rangle = (0, 0, 0, 1), \\ |V_3\rangle &= \frac{1}{|\rho_{23}| \sqrt{|\rho_{23}|^2 + (\lambda_3 - \rho_{33})^2}} (0, (\lambda_3 - \rho_{33})\rho_{23}, |\rho_{23}|^2, 0), \\ |V_4\rangle &= \frac{1}{|\rho_{23}| \sqrt{|\rho_{23}|^2 + (\lambda_4 - \rho_{33})^2}} (0, (\lambda_4 - \rho_{33})\rho_{23}, |\rho_{23}|^2, 0). \end{aligned} \quad (13)$$

Making use of Eq. (11), we can obtain the quantum Fisher information of the density matrix $\rho_{AB}(t)$ given in Eq. (8). As its expression is very complicated, we will display the results mainly by numerical simulations in the following section.

Numerical Discussion

Quantum Fisher information with respect to the Kerr medium. In this subsection, we estimate the parameter χ by evaluating the corresponding quantum Fisher information, namely \mathcal{F}_χ . The effect of the phase decoherence parameter γ , the mean photon number n , the detuning parameter Δ and the Kerr medium parameter χ on the quantum Fisher information \mathcal{F}_χ will be discussed.

Figure 1 describes the quantum Fisher information behavior with respect to the Kerr-like medium parameter for different cases. It is assumed that the atomic-field system is initially prepared in the resonance and non-resonance cases. The field is prepared in the vacuum and one photon Fock states, while the atomic system is prepared in different initial states; separable and partially entangled atomic states. As soon as the interaction is switched on \mathcal{F}_χ increases suddenly to reach its maximum values and then oscillates between its upper and lower bounds. As the time increases the upper bounds decrease to vanish completely at the further time. Moreover, the upper and lower bounds of \mathcal{F}_χ depend on the initial state settings of the atomic system, where the upper bounds of \mathcal{F}_χ decrease as the weight angle increases. This shows that the possibility of estimating the χ parameter increases if the initial atomic system is coded to classical information. Additionally, the resonance and non-resonance cases have a clear effect. At the resonance case, the upper bounds of \mathcal{F}_χ are larger than those displayed for the non-resonance case. The effect of photon number can be seen clearly when compare between




Figure 2. Quantum Fisher information \mathcal{F}_χ as a function of time t with $\kappa = 1$, $\gamma = 0.02$ and $\theta = \pi/3$ for different values of Kerr medium. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_χ at $\chi = 0.1, 1$ and $\chi = 2$, respectively. Here: (a) $n = 0$, $\Delta = 0$, (b) $n = 0$, $\Delta = 1$ (c) $n = 1$, $\Delta = 0$, (d) $n = 1$, $\Delta = 1$.

Fig. 1(a) with 1(c) (or 1(b) with 1(d)) where the upper bounds of \mathcal{F}_χ for one-photon are higher than that depicted for vacuum state. On the other hand, for the non-resonance case, the upper bounds of the \mathcal{F}_χ are smaller than those are displayed for the resonance case as shown from Fig. 1(c,d).

Figure 2 displays the effect of different initial values of the Kerr medium where the atomic system is prepared in a partial entangled state with a weight angle $\theta = \pi/3$. It is clear that at small values of χ the upper bounds of \mathcal{F}_χ are larger than those displayed for larger values of χ . On the other hand, for the further time, the quantum Fisher information decreases gradually to vanish completely at very high values of time t . The survival time of \mathcal{F}_χ increases as χ increases. The resonance, non-resonance, vacuum and one photon cases have a noticeable effect on the upper bounds of \mathcal{F}_χ and its survival time.

In Fig. 3 we investigate the effect of the phase decoherence parameter γ on the behavior of \mathcal{F}_χ . The general behavior shows that the quantum Fisher information decreases as the phase decoherence increases. The decay rate depends on the initial state of the cavity mode and the detuning between the atomic system and the cavity. It is clear that the upper bounds of \mathcal{F}_χ are large for the one-photon and non-resonance case. From Figs 1–3 one may conclude that the precision of estimating the Kerr medium χ parameter depends on the initial state settings of the field and the atomic systems. The larger initial values of χ , the larger upper bounds of the quantum Fisher information \mathcal{F}_χ . Although the maximum value of \mathcal{F}_χ decreases as the phase decoherence increases, one may increase these values by considering the detuning and the one-photon case.

Quantum Fisher information with respect to the detuning. In this section, the quantum Fisher information with respect to the detuning parameter is discussed. It is assumed that the atomic system is initially prepared in a partial entangled state and the cavity is prepared in a vacuum and one photon case.

Figure 4 shows the behavior of \mathcal{F}_Δ for different initial values of the χ parameter. The behavior is similar to that displayed in Fig. 2(a,b) but the maximum values of \mathcal{F}_Δ are smaller than those displayed for \mathcal{F}_χ . The one-photon cavity mode makes the oscillations of the quantum Fisher information \mathcal{F}_Δ collapse. Moreover, the survival time of \mathcal{F}_Δ is larger than that displayed for the vacuum case. The phase decoherence effect on the behavior of \mathcal{F}_Δ is described in Fig. 5, where different values of γ are considered. The general behavior is similar to that displayed for \mathcal{F}_χ but has smaller maximum values.

Quantum Fisher information with respect to the phase decoherence. Quantum Fisher information with respect to \mathcal{F}_γ is described in Figs 6 and 7. The upper bounds of \mathcal{F}_γ are much larger than those displayed for \mathcal{F}_χ and \mathcal{F}_Δ .




Figure 3. Quantum Fisher information \mathcal{F}_χ as a function of time t with $\kappa = 1$, $\chi = 1$ and $\theta = \pi/3$ for different values of phase decoherence. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_χ at $\gamma = 0.02, 0.03$ and $\gamma = 0.04$, respectively. Here: (a) $n = 0, \Delta = 0$, (b) $n = 1, \Delta = 1$, (c) $n = 1, \Delta = 0$, (d) $n = 1, \Delta = 1$.




Figure 4. Quantum Fisher information \mathcal{F}_Δ as a function of time t with $\kappa = 1$, $\gamma = 0.02$ and $\theta = \pi/3$ for different values of Kerr medium. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_Δ at $\chi = 0.1, 1$ and $\chi = 2$, respectively. Here: (a) $n = 0, \Delta = 1$, and (b) $n = 1, \Delta = 1$.

As it is shown in Fig. 6 the effect of the Kerr medium on \mathcal{F}_γ shows that the larger values of χ make the upper bounds of \mathcal{F}_γ increase. The one-photon case of the cavity causes a delay of vanishing of the quantum Fisher information \mathcal{F}_γ . The effect of the initial values of γ on \mathcal{F}_γ is displayed in Fig. 7.

The general behavior shows that \mathcal{F}_γ decays as γ increases. The decay rate depends on the initial state of the total system. At one photon cavity mode, the quantum Fisher information collapses and consequently, the minimum values are larger than those displayed at the vacuum state.

Conclusion

In this contribution, a two-atom system interacting with a cavity mode which is initially prepared in the Fock state in presence of Kerr medium under decoherence is considered. The estimation of the Kerr medium, detuning, and the phase decoherence parameters are discussed in different initial states. Four different cases are considered (namely, resonance, non-resonance case, vacuum and one-photon cavity). First, it is important to mention that before the interaction is switched on, the Fisher information of all the estimated parameters is zero. Because, these parameters don't appear on the atomic system before the interaction. However, it is shown that, quantum Fisher information is much larger than those depicted for the Kerr-medium and the detuning parameters. Also preparing the cavity mode in non-vacuum state causes an increase of the upper bounds of quantum Fisher information




Figure 5. Quantum Fisher information \mathcal{F}_Δ as a function of time t with $\kappa=1$, $\chi=1$ and $\theta = \pi/3$ for different values of phase decoherence. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_Δ at $\gamma=0.02$, 0.03 and $\gamma=0.04$, respectively. Here: (a) $n=0$, $\Delta=1$, and (b) $n=1$, $\Delta=1$.




Figure 6. Quantum Fisher information \mathcal{F}_γ as a function of time t with $\kappa=1$, $\gamma=0.02$ and $\theta = \pi/3$ for different values of Kerr medium. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_γ at $\chi=0.1$, 1 and $\chi=2$, respectively. Here: (a) $n=0$, $\Delta=1$, and (b) $n=1$, $\Delta=1$.




Figure 7. Quantum Fisher information \mathcal{F}_γ as a function of time t with $\kappa=1$, $\chi=1$ and $\theta = \pi/3$ for different values of phase decoherence. The dotted-black, solid-red and dashed-blue curves represent \mathcal{F}_γ at $\gamma=0.02$, 0.03 and $\gamma=0.04$, respectively. Here: (a) $n=0$, $\Delta=1$, and (b) $n=1$, $\Delta=1$.

and its survival time too. The initial states settings of the atomic and the field systems play an important role in the behavior of the three kinds of the quantum Fisher information. Quantum Fisher information for the separable atomic system is larger than that displayed for entangling atomic system. The non-vacuum state of the field causes an increase the maximum values of quantum Fisher information with respect to the Kerr medium and the phase decoherence parameter. The Kerr medium and the phase decoherence have a similar effect on all the three types of the quantum Fisher information.

References

1. Fisher, R. A. Theory of Statistical Estimation. *Proc. Camb. Phil. Soc.* **22**, 700 (1925).
2. Helstrom, C. W. Quantum Detection and Estimation Theory. (Academic Press, New York, 1976).

3. Holevo, A. S. *Statistical Structure of Quantum Theory*. (Springer, Berlin, 2001).
4. Nielsen, M. A. & Chuang, I. L. *Quantum computation and quantum information*. (Cambridge University Press, Cambridge, 2000).
5. Giovannetti, V., Lloyd, S. & Maccone, L. Quantum Metrology. *Phys. Rev. Lett.* **96**, 010401 (2006).
6. Giovannetti, V., Lloyd, S. & Maccone, L. Advances in Quantum Metrology. *Nature Photon* **5**, 222 (2011).
7. Ma, J., Huang, Y., Wang, X. & Sun, C. P. Quantum Fisher information of the Greenberger-Horne-Zeilinger state in decoherence channels. *C. P. Phys. Rev. A* **84**, 022302 (2011).
8. Escher, B. M., Filho, M. & Davidovich, L. General framework for estimating the ultimate precision limit in noisy quantum-enhanced metrology. *Nature Physics* **7**, 406 (2011).
9. Yi, X., Huang, G. & Wang, J. Quantum Fisher Information of a 3-Qubit State. *Int. J. Theor. Phys.* **51**, 3458 (2012).
10. Liu, Z. Spin Squeezing in Superposition of Four-Qubit Symmetric State and W States. *Int. J. Theor. Phys.* **52**, 820 (2013).
11. Ozaydin, F., Altintas, A. A., Bugu, S. & Yesilyurt, C. Quantum Fisher Information of N Particles in the Superposition of W and GHZ States. *Int. J. Theor. Phys.* **52**, 2977 (2013).
12. Ozaydin, F. *et al.* Quantum Fisher Information of Several Qubits in the Superposition of A GHZ and two W States with Arbitrary Relative Phase. *Int. J. Theor. Phys.* **53**, 3219 (2014).
13. Ozaydin, F. Phase damping destroys quantum Fisher information of W states. *Phys. Lett. A* **378**, 3161 (2014).
14. Strobel, H. *et al.* Fisher information and entanglement of non-Gaussian spin states. *Science* **345**, 424 (2014).
15. Jing, X. X., Liu, J., Xiong, H. & Wang, X. Maximal quantum Fisher information for general $su(2)$ parametrization processes. *Phys. Rev. A* **92**, 012312 (2015).
16. Liu, Z., Qiu, L. & Pan, F. Enhancing quantum coherence and quantum Fisher information by quantum partially collapsing measurements. *Quantum Inf. Process* **16**, 109 (2017).
17. Ozaydin, F., Altintas, A. A., Bugu, S. & Yesilyurt, C. Behavior of Quantum Fisher Information of Bell Pairs under Decoherence Channels. *Acta Physica Polonica A* **125**, 606 (2014).
18. Ozaydin, F. Quantum Fisher Information of a 3 Bound Entangled State and its Relation with Geometric Discord. *Int. J. Theor. Phys.* **54**, 3304 (2015).
19. Ang, S., Harris, G. I., Bowen, W. P. & Tsang, M. Optomechanical parameter estimation. *New J. Phys.* **15**, 103028 (2013).
20. Iwasawa, K. *et al.* Quantum-Limited Mirror-Motion Estimation. *Phys. Rev. Lett.* **111**, 163602 (2013).
21. Tsang, M. Quantum metrology with open dynamical systems. *New J. Phys.* **15**, 073005 (2013).
22. Altintas, A. A. Quantum Fisher information of an open and noisy system in the steady state. *Annals of Physics* **367**, 192 (2016).
23. Ziv-Zakai Error Bounds for Quantum Parameter Estimation. *Phys. Rev. Lett.* **108** 230401 (2012).
24. Yao, Y. *et al.* Quantum Fisher information in noninertial frames. *Phys. Rev. A* **89**, 042336 (2014).
25. Metwally, N. Estimation of teleported and gained parameters in a non-inertial frame. *Laser Phys. Lett.* **14**, 045202 (2017).
26. Metwally, N. Unruh acceleration effect on the precision of parameter estimation. arXiv.1609.02092[quant-ph] (2016).
27. Chen, Y. & Yuan, H. Maximal quantum Fisher information matrix. *New J. Phys.* **19**, 063023 (2017).
28. Mirkhalaf, S. S. & Smerzi, A. Entanglement detection in a coupled atom-field system via quantum Fisher information. *Phys. Rev. A* **95**, 022302 (2017).
29. Anwar, S. J., Ramzan, M. & Khan, M. K. Dynamics of entanglement and quantum Fisher information for N-level atomic system under intrinsic decoherence. *Quantum Inf. Process* **16**, 142 (2017).
30. Lu, X. M., Wang, X. G. & Sun, C. P. Quantum Fisher information flow and non-Markovian processes of open systems. *Phys. Rev. A* **82**, 042103 (2010).
31. Zhong, W. *et al.* Fisher information under decoherence in Bloch representation. *Phys. Rev. A* **87**, 022337 (2013).
32. Wang, G. Y., Guo, Y. N. & Zeng, K. Dynamics of quantum Fisher information in a two-level system coupled to multiple bosonic reservoirs. *Chin. Phys. B* **24**, 114201 (2015).
33. Yan, K., Xie, Y. Q., Haung, Y. M. & Hao, X. Quantum Fisher information of driven multi-particle systems in structured bosonic reservoirs. *Comm. Theor. Phys.* **67**, 261 (2017).
34. Nolan, S. P. & Haine, S. A. Quantum Fisher information as a predictor of decoherence in the preparation of spin-cat states for quantum metrology. *Phys. Rev. A* **95**, 043642 (2017).
35. Metwally, N. & Hassan, S. S. Estimation of pulsed driven qubit parameters via quantum Fisher information. *Laser Phys. Lett.* **14**, 115204 (2017).
36. Metwally, N. Fisher information of a single qubit interacts with a spin-qubit in the presence of a magnetic field. *Physica E* **100**, 14 (2018).
37. Zhang, Q., Yao, Y. & Li, Y. Optimal quantum channel estimation of two interacting qubits subject to decoherence. *Eur. Phys. J. D* **68**, 170 (2014).
38. Chen, Y., Zou, J., Long, Z.-W. & Shao, B. Protecting quantum Fisher information of N-qubit GHZ state by weak measurement with fips against dissipation. *Sci. Ret.* **7**, 6160 (2017).
39. Wang, Z. H., Zheng, Q., Wang, X. & Li, Y. The energy-level crossing behavior and quantum Fisher information in a quantum well with spin-orbit coupling. *Sci. Ret.* **6**, 22347 (2016).
40. Zheng, Q., Ge, L., Yao, Y. & Zhi, Q. J. Enhancing parameter precision of optimal quantum estimation by direct quantum feedback. *Phys. Rev. A* **91**, 033805 (2015).
41. Huang, J., Guo, Y. N. & Xie, Q. Enhancing parameter precision of optimal quantum estimation by quantum screening. *Chin. Phys. B* **25**, 020303 (2016).
42. Yang, L., Dai, H. Y. & Zhang, M. Quantum parameter estimation in a spin-boson dephasing quantum system by periodical projective measurements. *Chin. Phys. B* **27**, 040601 (2018).
43. Abdel-Hameed, H., Zidan, N. & Metwally, N. Quantum Fisher information of two superconducting charge qubits under dephasing noisy channel. *Int. J. Mod. Phys. B* **32**, 1850245 (2018).
44. Bhattacharjee, A. B. & Sharma, D. Enhanced spin squeezing and quantum entanglement near the critical point of the Jaynes-Cummings-Dicke model. *Int. J. Mod. Phys. B* **31**, 1750062 (2017).
45. Li, C. *et al.* Controlled Correlation and Squeezing in $Pr^{3+}Y_2SiO_5$ to Yield Correlated Light Beams. *Phys. Rev. Applied* **7**, 014023 (2017).
46. Bu, S. P., Zhang, G. F., Liu, J. & Chen, Z. Y. Berry phase in a two-atom Jaynes-Cummings model with Kerr medium. *Phys. Scr.* **78**, 065008 (2008).
47. Gardiner, C. W. & Zoller, P. *Quantum noise*. (Springer-Verlag, Berlin, 2000).
48. Xu, J. B. & Zou, X. B. Dynamic algebraic approach to the system of a three-level atom in the Λ configuration. *Phys. Rev. A* **60**, 4743 (1999).
49. Braunstein, S. L. & Caves, C. M. Statistical distance and the geometry of quantum states. *Phys. Rev. Lett.* **72**, 3439 (1994).
50. Knysh, S., Smelyanskiy, V. N. & Durkin, G. A. Scaling laws for precision in quantum interferometry and the bifurcation landscape of the optimal state. *Phys. Rev. A* **83**, 021804 (2011).
51. Liu, J., Jing, X. & Wang, X. Phase-matching condition for enhancement of phase sensitivity in quantum metrology. *Phys. Rev. A* **88**, 042316 (2013).

Author Contributions

N. Zidan and H. Abdel-Hameed initiated the research project and established the main results. N. Metwally discussed the results and provided suggestions.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019