

Bilateral renal involvement in diffuse large B-cell lymphoma on fluorodeoxyglucose positron emission tomography/computed tomography

ABSTRACT

Secondary renal involvement in patients with diffuse large B-cell lymphoma (DLBCL) is rare and associated with poor prognosis. We, hereby, described a case of a patient diagnosed with DLBCL, in whom bilateral renal involvement was detected on 18F-fluorodeoxyglucose positron emission tomography/computed tomography (¹⁸F-FDG PET/CT). The patient received 4 courses of chemotherapy, and follow-up ¹⁸F-FDG PET/CT revealed the complete resolution of the diffuse increased renal ¹⁸F-FDG uptake. Renal lymphoma is uncommon and ¹⁸F-FDG PET/CT is particularly useful for detecting extranodal involvement in DLBCL.

Keywords: 18F-fluorodeoxyglucose positron emission tomography/computed tomography, diffuse large B-cell lymphoma, renal 18F-FDG uptake, renal lymphoma

INTRODUCTION

Renal involvement in aggressive lymphoma is most often seen along with the dissemination of systemic disease.^[1] In this case report, we describe a 17-year-old female with diffuse large B-cell lymphoma (DLBCL) and kidney involvement at diagnosis visualized on 18F-fluorodeoxyglucose positron emission tomography/computed tomography (¹⁸F-FDG PET/CT).

CASE REPORT

A 17-year-old female, without medical history, presented with a 15-day history of diplopia, nasal obstruction, and left orbital pain, associated with fever and night sweats. The physical examination found a left exophthalmos and bilateral cervical lymphadenopathy. T2-weighted orbital magnetic resonance imaging showed a left soft-tissue mass slightly hyperintense compared to muscle involving intra- and extraconal space responsible for an exophthalmos, with an extension to the left ethmoid sinus and the left nasal fossa. Biopsy of the nasal mass

and histopathologic examination objectified a high-grade DLBCL (CD20+, Bcl2+, CD3-, and CD5-, with a Ki-67 proliferation index of 95%). In addition, the cerebrospinal fluid analysis was consistent with a central nervous system (CNS) involvement.

Baseline ¹⁸F-FDG PET/CT revealed intense ¹⁸F-FDG uptake throughout the bilaterally enlarged renal parenchyma, on the soft tissue in the left orbit extending into the left nasal cavity, and on cervical and abdominal lymph nodes.

**YASSIR BENAMEUR, ADNANE HAMMANI¹,
KAMAL DOGHMI¹, ABDERRAHIM DOUDOUH**

Departments of Nuclear Medicine and ¹Clinical Hematology, Mohammed V Military Teaching Hospital, Faculty of Medicine and Pharmacy of Rabat, Mohammed V University in Rabat, Morocco


Address for correspondence: Dr. Yassir Benameur, Department of Nuclear Medicine, Mohammed V Military Teaching Hospital, Mohammed V University in Rabat, Rabat, Morocco. E-mail: benameur.yassir@gmail.com

Submitted: 04-Aug-2020, **Revised:** 16-Sep-2020, **Accepted:** 04-Oct-2020, **Published:** 12-Feb-2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Benameur Y, Hammani A, Doghmi K, Doudouh A. Bilateral renal involvement in diffuse large B-cell lymphoma on fluorodeoxyglucose positron emission tomography/computed tomography. World J Nucl Med 2021;20:195-7.

Access this article online	
Website: www.wjnm.org	Quick Response Code 
DOI: 10.4103/wjnm.WJNM_111_20	

Peritoneal nodules with diffusely increased ^{18}F -FDG and multiple hypermetabolic bones foci, including humeri, ribs, and pelvis uptake were also noted [Figure 1]. After 4 cycles of dose-adjusted EPOCH-rituximab, the patient was referred for ^{18}F -FDG PET/CT for response evaluation. The ^{18}F -FDG PET/CT showed a total regression of renal lesions suggesting good response to therapy [Figure 2].

DISCUSSION

DLBCL is the most common aggressive lymphoma.^[2] The usual presentation of DLBCL is symptomatic enlarged lymph nodes; around 40% of patients have extranodal involvement, most commonly in the gastrointestinal tract (36%), ear, nose, throat (20%), and bone marrow (14%).^[3] Renal parenchymal involvement is uncommon in DLBCL, it appears to be a secondary process, either by direct extension from a retroperitoneal mass or through hematogenous spread in the setting of disseminated disease.^[4] In a study with 821 patients diagnosed with DLBCL, renal involvement was noted in 26 (3%) at the time of diagnosis, the majority of patients presented in advanced disease, 86% were Ann Arbor stage \geq III.^[5] Infiltrative lymphomatous proliferation within the renal tissue is manifested as nephromegaly with the preservation of renal contour. The diagnosis, therefore,

depends on the finding of global renal enlargement and poor renal function.^[6]

More uncommon, primary renal non-Hodgkin's lymphoma is an extremely rare disease and only a few cases have been reported, accounting for <1% of all lymphomas.^[7] The most common presentation is acute renal failure, proteinuria, microscopic hematuria, and renal enlargement.^[7] Reuter *et al.* described a primary bilateral renal B-cell lymphoma; in this case, the patient had urinary symptoms and was in nonoliguric acute renal failure.^[8] Renal failure was also found in secondary renal lymphoma, Navalkisoor *et al.* reported a patient with DLBCL and renal failure, with the particularity of being normalized after only one cycle of R-CHOP (Rituximab plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisolone).^[9] The same features were also noted in the case reported by Kara *et al.* with bilateral renal involvement and renal impairment.^[10] As opposed to our patient, who had no urinary symptoms and was not in renal failure at the time of diagnosis (glomerular filtration rate 85 mL/min, serum creatinine 13 μM).

Villa *et al.* demonstrated that patients with DLBCL and kidney involvement at diagnosis have a poor prognosis with CNS relapse incidence which occurs early.^[4] Thus, the assessment for CNS involvement is highly recommended in patients with




Figure 1: Maximum intensity projection (a), coronal ^{18}F -fluorodeoxyglucose positron emission tomography/computed tomography fusion (b) and transaxial images (c) demonstrated very intense diffuse ^{18}F -fluorodeoxyglucose uptake in bilaterally enlarged kidneys




Figure 2: Resolution of the renal ^{18}F -fluorodeoxyglucose uptake after 4 courses of dose-adjusted EPOCH-rituximab, on maximum intensity projection (a) image as well as on coronal (b) and transaxial images (c)

renal lymphomatous lesions at the time of diagnosis, even in the absence of neurological symptoms. In addition, the R-CHOP protocol is not a sufficient therapy in these patients with or without CNS disease.^[4] However, randomized controlled trials are required to establish a standard treatment approach. Our patient had CNS involvement and was treated by dose-adjusted EPOCH-rituximab protocol and high-dose methotrexate with the complete resolution of the diffuse increased renal ¹⁸F-FDG uptake after 4 courses of chemotherapy, but a partial response on FDG-avid bone lesions.

¹⁸F-FDG PET/CT imaging has revolutionized the management of patients with lymphoma. It is now essential for accurate staging and has become the foundation of widely adopted response criteria. Currently, ¹⁸F-FDG PET/CT is considered as the gold standard for response assessment.^[11] Imaging procedures alone cannot differentiate between primary and secondary renal lymphoma as ¹⁸F-FDG is excreted by the kidneys, this physiologic uptake in the renal collecting system can lead to false-positive and false-negative interpretations on PET-CT.^[12] Renal involvement in lymphoma is seen as multiple focal areas of increased ¹⁸F-FDG uptake mostly in renal cortices that may be unilateral or less commonly, bilateral.^[13] As in our case, where multiple focal areas of increased FDG uptake were noted in both kidneys.

It is necessary to point out that when diffuse renal FDG uptake is observed on ¹⁸F-FDG PET-CT, one of the most frequent differential diagnoses is renal sarcoidosis, as in the case described by Toyonaga *et al.* with the particularity of homogenous increased FDG uptake in both kidneys.^[14]

CONCLUSION

Renal involvement in DLBCL is rare; however, it is associated with poor prognosis, and it includes an increased risk for CNS involvement.

This case supports the important clinical value of the ¹⁸F-FDG PET/CT as a whole-body imaging modality in detecting unusual sites of involvement in DLBCL.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Erdogmus S, Akturk S, Kendi Celebi Z, Kiremitçi S, Kaygusuz G, Kemal Altınbaş N, *et al.* Diffuse large B-cell lymphoma presenting with bilateral renal masses and hematuria: A case report. *Turk J Haematol* 2016;33:159-62.
2. Delaby G, Hubaut M-A, Morschhauser F, Besson A, Huglo D, Herbaux C, *et al.* Prognostic value of the metabolic bulk volume in patients with diffuse large B-cell lymphoma on baseline ¹⁸F-FDG PET-CT. *Leuk Lymphoma*. 2020:1-8.
3. Kaur A, Abughanim O, Zafar Y, Pluard T. Diffuse large B-cell lymphoma presenting as bilateral renal masses: Successful treatment with dose-adjusted REPOCH (Rituximab, Etoposide, Prednisone, Vincristine, Cyclophosphamide, Doxorubicin) chemotherapy regimen. *Cureus* 2019;11:e3814.
4. Villa D, Connors JM, Sehn LH, Gascoyne RD, Savage KJ. Diffuse large B-cell lymphoma with involvement of the kidney: Outcome and risk of central nervous system relapse. *Haematologica* 2011;96:1002-7.
5. Lehnert N, Kramer I, Schwarzbich MA, Ho AD, Witzens Harig M. Analysis of clinical characteristics and outcome of patients with previously untreated diffuse large B-cell lymphoma and renal involvement in the rituximab era. *Leuk Lymphoma*. 2016;57:2619-25.
6. Basu S, Ramani SK, Lad S. Unusual involvement of scalp and bilateral kidneys in an aggressive mediastinal diffuse large B cell lymphoma: Documentation by FDG-PET imaging. *Clin Nucl Med* 2009;34:638-41.
7. Tan YZ, Yilmaz S, Ozhan M, Halaç M. FDG PET-CT finding in bilateral renal and bone involvement of diffuse large B-cell lymphoma. *Mol Imaging Radionucl Ther* 2014;23:104-6.
8. Reuter S, Rahbar K, Busch V, Hillebrand U, Velden J, Pavenstädt H, *et al.* Acute renal failure due to primary bilateral renal large B-cell lymphoma: Diagnostics and follow-up by FDG-PET/CT. *Clin Nucl Med* 2009;34:722-4.
9. Navalkissoor S, Szyszko T, Gnanasegaran G, Nunan T. Diffuse FDG renal uptake in lymphoma. *Clin Nucl Med* 2010;35:813-5.
10. Kara PO, Koc ZP, Ozgur ES, Ayan E, Tombak A. A rare case of bilateral renal lymphoma on FDG PET/CT Imaging. *ARC J Radiol Med Imaging* 2017;2:14-6.
11. Cheson BD. PET/CT in lymphoma: Current overview and future directions. *Semin Nucl Med* 2018;48:76-81.
12. Even-Sapir E, Lievshitz G, Perry C, Herishanu Y, Lerman H, Metser U. Fluorine-18 fluorodeoxyglucose PET/CT patterns of extranodal involvement in patients with non-Hodgkin lymphoma and Hodgkin's disease [vii.]. *Radiol Clin North Am* 2007;45:697-709.
13. Dhull VS, Sharma P, Singla S, Faizi NA, Thulkar S, Bal C, *et al.* Extensive extranodal involvement of rare sites in non Hodgkin's lymphoma detected on ¹⁸F-FDG PET-CT: A case report. *Nucl Med Mol Imaging* 2013;47:125-9.
14. Toyonaga T, Manabe O, Gaertner FC, Nakagaki T, Nishio S, Suzuki A, *et al.* Diffuse renal ¹⁸F-FDG uptake of a patient with fever of unknown origin revealed sarcoidosis. *Clin Nucl Med* 2014;39:648-9.