
ERRATUM

Erratum to: Evaluation of the Effects of Favipiravir Combined with Vitamin C on Alveolar Bone in Rats

A. Lektemur Alpan^{a,*}, G. Torumtay Cin^a, Ö. Özmen^b, M. F. Doğan^c, and Y. Şahin^c

^aDepartment of Periodontology, Faculty of Dentistry, Pamukkale University, Denizli, Turkey

^bDepartment of Pathology, Faculty of Veterinary Medicine, Burdur Mehmet Akif Ersoy University, Burdur, Turkey

^cDepartment of Pharmacology, Faculty of Medicine, University of Pamukkale, Denizli, Turkey

*e-mail: ysnlpn@gmail.com

Received June 1, 2022

Revised June 1, 2022

Accepted June 1, 2022

DOI: 10.1134/S0022093022030280

The method of administration of the drugs has been corrected by the authors as follows:

In section *Abstract*:

Favipiravir (20 mg/kg and 100 mg/kg, orally) and Vit C (150 mg/kg/day, i.m.) were administered to the rats for 14 days.

In section *Methods*:

2. Favi 20 group: 0.1 mL (i.m.) of saline and 20 mg/kg orally of Favipiravir

3. Favi 100 group: 0.1 mL (i.m.) of saline and

100 mg/kg orally of Favipiravir

4. Favi 20+Vit C group: 20 mg/kg orally of Favipiravir and i.m. 150 mg/kg of Vit C

5. Favi 100+Vit C group: 100 mg/kg orally of Favipiravir and i.m. 150 mg/kg of Vit C

Favipiravir (20 or 100 mg/kg; orally, T705, Merck) and Vit C (150 mg/kg; i.m., L Ascorbic acid, Merck) [12].

The original article can be found online at <https://doi.org/10.1134/S0022093022020119>