

# PRPS1 Gene Mutation Causes Complex X-Linked Adult-Onset Cerebellar Ataxia in Women

Flávio M. Rezende Filho, MD, PhD, Mariana M. Palma, MD, José Luiz Pedroso, MD, PhD, Orlando G. Barsottini, MD, PhD, and Juliana M. Sallum, MD, PhD

## Correspondence

Dr. Pedroso  
jlpedroso.neuro@gmail.com

*Neurol Genet* 2021;7:e563. doi:10.1212/NXG.000000000000563

Inherited retinal dystrophies (IRD) comprise a heterogeneous group of disorders that affect visual function. IRD occur in isolated forms or in association with systemic abnormalities.<sup>1</sup> Over 300 disease-causing genes have been identified in IRD.

We report an unusual form of syndromic IRD with a complex neurologic phenotype caused by a mutation in *PRPS1* gene. Pathogenic variants in *PRPS1* lead to Arts syndrome, X-linked deafness 1, *PRPS*-related gout, or Charcot-Marie-Tooth 5 (*CMTX5*) in boys,<sup>2</sup> but it has been recently shown that women may also be symptomatic and display hearing loss, ophthalmologic abnormalities, and neurologic manifestations.<sup>3</sup>

## Case Report

A 30-year-old woman, born from nonconsanguineous parents, presented with a 5-year history of progressive ataxia. She also had congenital strabismus, infantile-onset hearing loss, and a retinal dystrophy with progressive visual loss for the past 10 years. The visual acuity was 20/200 on the right eye and 20/40 on the left eye. Fundus examination revealed pale optic discs, attenuated vessels, and atrophic areas containing whitish fine dots, reticular pigmentation, and bone spicules bilaterally. Examination showed cerebellar ataxia, dysarthria, and gaze-evoked nystagmus. Audiometry confirmed bilateral sensorineural hearing loss. Brain MRI showed atrophy in the pons, middle cerebellar peduncles, vermis, and cerebellar hemispheres (figure). The genetic test for Friedreich ataxia was negative. Alpha-fetoprotein, albumin, vitamin E, creatine kinase, and lactate were normal. Whole-exome sequencing disclosed a heterozygous missense variant c.359G>T(p.Gly120Val) in one of her *PRPS1* alleles.

## Discussion

*PRPS1* is located in X chromosome and codes the enzyme phosphoribosyl pyrophosphate (PRPP) synthetase 1. This enzyme is involved in the synthesis of PRPP from ATP and ribose-5-phosphate, which is essential in the production of purines (adenine and guanine) and pyrimidines (cytosine, thymine, and uracil) nucleotides, the building blocks of life. Pathogenic variants in *PRPS1* cause 4 distinct conditions in males: *CMTX5*, X-linked deafness 1 (*DFNX1*), Arts syndrome, and an inherited form of gout (*PRPS*-related gout). Sensorineural hearing loss is the only common manifestation of these disorders and the isolated finding of *DFNX1*. Patients with *CMTX5* and Arts syndrome additionally have peripheral neuropathy, ataxia, and optic atrophy.<sup>2,4</sup> Severe cognitive decline, recurrent infections, and early death complicate the clinical course in Arts syndrome, the most severe phenotype.<sup>5</sup> Individuals with *PRPS*-related

---


From the Department of Neurology (F.M.R.F., J.L.P., O.G.B.), and Department of Ophthalmology (M.M.P., J.M.S.), Universidade Federal de São Paulo (UNIFESP), Brazil.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP).

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## Figure Ophthalmologic and Radiologic Findings



Color fundus photographs of both eyes depicting pale optic discs, attenuated retinal vessels, and atrophic areas containing whitish fine dots with reticular pigmentation (A). The pattern of retinal dystrophy is remarkably asymmetric and more severe on the right eye, in which the macula is affected. Axial T2-weighted sequence of brain MRI shows atrophy in the cerebellar peduncles and pons (B) and in cerebellar hemispheres (C) (white arrows).

gout, the most common form, have hyperuricemia, but some exhibit mental retardation, ataxia, and hypotonia.<sup>2</sup>

The variant found in our patient, c.359G>T(p.Gly120Val), is novel and was confirmed by Sanger sequencing. Also, it was absent in the parents characterizing a *de novo* mutation. Moreover, this variant is absent in 178,000 X chromosomes. It affects a highly conserved residue, and *in silico* prediction indicates a deleterious effect. The combination of the molecular mechanism, location, and the correlation with clinical features suggests that this variant is most likely pathogenic.

Female patients with *PRPS1* mutations are usually asymptomatic and rarely present with neurologic symptoms.<sup>6</sup> The female disease spectrum ranges from isolated sensorineural hearing loss to the combination of retinal dystrophy, optic atrophy, peripheral neuropathy, ataxia with cerebellar atrophy in MRI, and developmental delay.<sup>3,4,7</sup> Our patient presented early-onset hearing loss, retinal dystrophy, and cerebellar ataxia. Her phenotype is consistent with the one described in 2 women from a Spanish family.<sup>3</sup> Fiorentino et al.<sup>7</sup> reported similar phenotypic features in a Japanese mother and her daughter, who had retinal dystrophy, optic atrophy, sensorineural hearing loss, and gait ataxia. A pathophysiologic explanation for a late-onset phenotype in a woman related to

*PRPS1* is the X inactivation. *De novo* variants and skewed inactivation of X chromosome have been described.

The work by Fiorentino et al.<sup>7</sup> demonstrated 7 of 9 patients with *PRPS1*-related retinal dystrophy had asymmetric retinal abnormalities. A high proportion of women with X-linked IRD have fundus changes, which are frequently asymmetric. This may be a consequence of skewed inactivation of X chromosomes in females, causing a higher proportion of retinal cells to express the pathogenic variant of *PRPS1* gene, rather than the wild-type allele. The degree of skewed X-chromosome inactivation and PRPP synthetase residual activity are the likely determinants of phenotype severity in women with pathogenic variants of *PRPS1*.<sup>4</sup>

It is noteworthy that *PRPS1*-related disorders might be treatable. Dietary S-adenosylmethionine supplementation (SAM) exploits an alternative route of ATP biosynthesis, which may alleviate the shortage of adenosine-derived nucleotides. Of interest, in 2 individuals with Arts syndrome, SAM supplementation resulted in clinical improvement.<sup>2</sup>

In conclusion, this report illustrates the remarkable phenotypic variability of *PRPS1* mutations, which may cause a rare form of X-linked adult-onset cerebellar ataxia. Asymmetric retinal dystrophy might represent a clue for the diagnosis of *PRPS1* spectrum disease in women with hearing loss and cerebellar signs.

## Acknowledgment

The authors thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for have funded this study.

## Study Funding

This study was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (grant number: 2019/15298-1).

## Disclosure

F.M. Rezende Filho and M.M. Palma report no disclosures relevant to the manuscript. J.L. Pedroso received grant from FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo). O.G.P. Barsottini and J.M.F. Sallum report no disclosures relevant to the manuscript. Go to [Neurology.org/NG](http://Neurology.org/NG) for full disclosures.

## Publication History

Received by *Neurology: Genetics* September 7, 2020. Accepted in final form December 8, 2020.

## Appendix Authors

| Name                             | Location                                  | Contribution  |
|----------------------------------|---|---|
| Flávio M. Rezende Filho, MD, PhD | Universidade Federal de São Paulo, Brazil | Conception, organization, and execution; writing of the first draft |
| Mariana M. da Paula, MD          | Universidade Federal de São Paulo, Brazil | Conception, organization, and execution; writing of the first draft |

## Appendix *(continued)*

| Name                                  | Location                                  | Contribution   |
|---------------------------------------|---|--|
| <b>José Luiz Pedroso, MD, PhD</b>     | Universidade Federal de São Paulo, Brazil | Conception, organization, and execution; review and critique |
| <b>Orlando G. Barsottini, MD, PhD</b> | Universidade Federal de São Paulo, Brazil | Conception, organization, and execution; review and critique |
| <b>Juliana M.F. Sallum, MD, PhD</b>   | Universidade Federal de São Paulo, Brazil | Conception, organization, and execution; review and critique |

## References

1. Ellingford JM, Barton S, Bhaskar S, et al. Molecular findings from 537 individuals with inherited retinal disease. *J Med Genet* 2016;53:761–767.
2. de Brouwer APM, van Bokhoven H, Nabuurs SB, Arts WF, Christodoulou J, Duley J. PRPS1 mutations: four distinct syndromes and potential treatment. *Am J Hum Genet* 2010;86:506–518. doi: 10.1016/j.ajhg.2010.02.024.
3. Almoquera B, He S, Corton M, et al. Expanding the phenotype of PRPS1 syndromes in females: neuropathy, hearing loss and retinopathy. *Orphanet J Rare Dis* 2014;9:190.
4. Synofzik M, Müller J, Haack TB, et al. X-linked Charcot-Marie-Tooth disease, Arts syndrome, and prelingual non-syndromic deafness form a disease continuum: evidence from a family with a novel PRPS1 mutation. *Orphanet J Rare Dis* 2014;9:1–8.
5. De Brouwer APM, Williams KL, Duley JA, et al. Arts syndrome is caused by loss-of-function mutations in PRPS1. *Am J Hum Genet* 2007;81:507–518.
6. Kim HJ, Sohn KM, Shy ME, et al. Mutations in PRPS1, which encodes the phosphoribosyl pyrophosphate synthetase enzyme critical for nucleotide biosynthesis, cause hereditary peripheral neuropathy with hearing loss and optic neuropathy (CMTX5). *Am J Hum Genet* 2007;81:552–558.
7. Fiorentino A, Fujinami K, Arno G, et al. Missense variants in the X-linked gene PRPS1 cause retinal degeneration in females. *Hum Mutat* 2018;39:80–91.