

# Defending the genome from the enemy within: mechanisms of retrotransposon suppression in the mouse germline

James H. Crichton · Donncha S. Dunican ·  
Marie MacLennan · Richard R. Meehan · Ian R. Adams

Received: 31 May 2013 / Revised: 27 August 2013 / Accepted: 29 August 2013 / Published online: 18 September 2013  
© The Author(s) 2013. This article is published with open access at Springerlink.com

**Abstract** The viability of any species requires that the genome is kept stable as it is transmitted from generation to generation by the germ cells. One of the challenges to transgenerational genome stability is the potential mutagenic activity of transposable genetic elements, particularly retrotransposons. There are many different types of retrotransposon in mammalian genomes, and these target different points in germline development to amplify and integrate into new genomic locations. Germ cells, and their pluripotent developmental precursors, have evolved a variety of genome defence mechanisms that suppress retrotransposon activity and maintain genome stability across the generations. Here, we review recent advances in understanding how retrotransposon activity is suppressed in the mammalian germline, how genes involved in germline genome defence mechanisms are regulated, and the consequences of mutating these genome defence genes for the developing germline.

**Keywords** Retrotransposon · Germ cell · Genome defence · Epigenetics · DNA methylation · Mouse · Meiosis

## Abbreviations

5hmC	5-Hydroxymethyl cytosine
5mC	5-Methyl cytosine
CGI	CpG island
ERV	Endogenous retrovirus
ES cells	Embryonic stem cells
H3K27Ac	Histone 3 lysine 27 acetylation
H3K27Me3	Histone 3 lysine 27 trimethylation
H3K4Me	Histone 3 lysine 4 methylation
H3K9Me3	Histone 3 lysine 9 trimethylation
H4K20Me3	Histone 4 lysine 20 trimethylation
IAP	Intracisternal A particle
LINE	Long interspersed nuclear element
LTR	Long terminal repeat
piRNA	PIWI-interacting RNA
SINE	Short interspersed nuclear element
siRNA	Small interfering RNA

J. H. Crichton, D. S. Dunican, and M. MacLennan contributed equally.

J. H. Crichton · D. S. Dunican · M. MacLennan ·  
R. R. Meehan (✉) · I. R. Adams (✉)  
MRC Human Genetics Unit, MRC Institute of Genetics  
and Molecular Medicine, University of Edinburgh, Western  
General Hospital, Crewe Road, Edinburgh EH4 2XU, UK  
e-mail: Richard.Meehan@igmm.ed.ac.uk

I. R. Adams  
e-mail: Ian.Adams@igmm.ed.ac.uk

J. H. Crichton  
e-mail: James.Crichton@igmm.ed.ac.uk

D. S. Dunican  
e-mail: Donncha.Dunican@igmm.ed.ac.uk

M. MacLennan  
e-mail: Marie.MacLennan@igmm.ed.ac.uk

## Introduction

Maintaining genetic stability through the generations is key for the survival of all species. If the germline mutation rate is too low, there will not be sufficient variation within the species to adapt and survive over evolutionary time; yet, if the germline mutation rate is too high, the accumulation of deleterious mutations will impact on viability. One of the major drivers of genetic change in the genome is retrotransposons. Retrotransposons are highly abundant mobile genetic elements that account for around 40 % of the sequenced mammalian genome [1, 2]. Retrotransposons contribute to genome instability by acting as sites for

recombination-mediated chromosomal deletions and rearrangements by influencing expression of nearby genes, and by causing mutations when new retrotransposition events disrupt pre-existing genetic information in the host genome [3].

For retrotransposons to be successful, retrotransposition must occur in the germ cells, or in the pluripotent cells from which germ cells arise. Germ cells and pluripotent cells have evolved multiple genome defence mechanisms to limit the mutagenic activity of these mobile genetic elements [4–6]. A number of genes involved in suppressing retrotransposon activity in the mammalian germline have been identified, and the ways that these genome defence mechanisms are regulated and interact to provide an effective defence against retrotransposons are starting to be understood. In this review, we will describe the different types of retrotransposons in mammalian genomes, the mechanisms that germ cells and pluripotent cells use to suppress the activity of these elements, and the phenotypic defects that arise in the developing germ cells when germline genome defence genes are mutated.

## Mammalian retrotransposons


The mammalian genome contains three major classes of retrotransposon: long interspersed nuclear element (LINE), short interspersed nuclear element (SINE), and long terminal repeat (LTR) retrotransposons (Fig. 1) [1, 2]. Each of

these retrotransposon classes has different characteristics and properties.

## LINE retrotransposons

LINE-1, which accounts for ~17 % of the sequenced genome, is the major family of LINE retrotransposons in mammals. Most copies of LINE-1 in mammalian genomes are 5'-truncated, probably due to incomplete reverse-transcription of this element during retrotransposition [1, 2]. Full-length mouse and human LINE-1 elements are around 6–7 kb in length and contain a variable internal promoter driving expression of these elements (Fig. 1). The internal promoter differs significantly in sequence between mouse and human LINE-1 elements, and between different LINE-1 subfamilies within species [7–9]. LINE-1 promoters may have been co-opted from host sequences, and differences between LINE-1 promoter sequences could lead to variations in the transcription factor requirements and regulation of these elements. The transcribed LINE-1 mRNA encodes two proteins, ORF1 and ORF2, that are each required for LINE-1 retrotransposition [10, 11]. ORF1 encodes a ~40-kDa nucleic acid-binding protein containing a central RNA recognition motif that has nucleic acid chaperone activity, while ORF2 encodes a ~150-kDa endonuclease and reverse-transcriptase [12–15]. Both ORF1 and ORF2 are thought to preferentially interact with the same mRNA molecule from which they are translated, which generates a strong *cis*-preference for that encoding mRNA

**Fig. 1** Structure of the major mammalian retrotransposons. Mouse and human examples of the three different classes of mammalian retrotransposon (*LINE*, *SINE*, *LTR*) are shown. Transcription regulatory regions are indicated with *filled rectangles*, and the main protein coding regions with *open rectangles*. Transcriptional start sites are shown with an *arrow*. Some LTR retrotransposons, e.g. IAP, have lost the *env* gene present in their infectious progenitors [241]. *LTR* long terminal repeat



to be retrotransposed [16]. LINE-1 elements are still retrotranspositionally active in mammals, and spontaneous de novo integrations of LINE-1 elements have been reported to cause phenotypic changes and disease in both mice and humans [17–19].

### SINE retrotransposons

Short interspersed nuclear element elements are non-autonomous retrotransposons that make up around 10 % of the sequenced mammalian genome [1, 2]. SINE elements are derived from a broad range of small cellular RNA polymerase III transcripts including 7SL RNA, 5S rRNA, and tRNAs, and rely on LINE-1-encoded proteins, particularly LINE-1 ORF2, to catalyse their retrotransposition [20–22]. SINE elements typically contain an internal RNA polymerase III promoter and are therefore regulated quite differently from the LINE-1-encoded proteins that allow them to retrotranspose. The most prominent families of SINE elements in human and mouse genomes are the Alu and B1 elements, respectively. Human Alu (~300 bp) and mouse B1 (~140 bp) elements are derived from the signal recognition particle 7SL cellular RNA, but subsequent re-arrangements and duplications has led to these elements having quite different structures (Fig. 1) [23]. SINE elements can also be incorporated into cellular transcripts that run through their integration sites, which can provide a route for new variants of these elements to arise [24]. De novo insertions of SINE elements have been identified as potentially causative mutations in both human and mouse genetic disease [17, 25]. Even though SINE elements require LINE-1-encoded proteins for their mobilisation, the germline SINE retrotransposition rate in humans is estimated to be around five times higher than that of LINE-1 [17].

### LTR retrotransposons

Long terminal repeat retrotransposons, also known as endogenous retroviruses (ERVs), have a typical retroviral structure with protein-coding gag, pol, pro and sometimes env genes flanked by long terminal repeats that act as promoters (Fig. 1) [26]. There are around 150 different types of ERV in a typical mammalian genome, classified into ERV1, ERVK, ERVL and MaLR families depending on their phylogenetic relationship. Like LINE-1 elements, LTR retrotransposons require the activity of the proteins that they encode in order to retrotranspose [27]. However, some LTR retrotransposons are non-autonomous and rely on proteins encoded by a different element to catalyse their retrotransposition [28]. LTR retrotransposons make up around 9 % of the human and mouse genomes; however, most of these LTR retrotransposons are lineage-specific elements that have arisen after the mouse and human

lineages diverged from a common ancestor 65–75 million years ago [2]. Different types of LTR retrotransposon have been successful in colonising mouse and human genomes; ERVK elements are ten times more abundant in mouse than in human genomes, whereas human ERV1 elements are four times more abundant than their mouse counterparts [2]. With the possible exception of HERVK (HML2) elements, the LTR retrotransposons that currently exist in the human genome do not appear to be retrotranspositionally active [26], and de novo insertions of LTR retrotransposons have not yet been identified as mutant alleles in human genetic disease [18]. In contrast, LTR retrotransposons are retrotranspositionally active in the mouse genome, and de novo insertions of ERV1, ERVK, ERVL and MaLR LTR retrotransposons have all been associated with spontaneous mutations in mice [18].


### Retrotransposon expression in germ cells and pluripotent cells

For retrotransposons to be successful, they must be expressed and functional in developing germ cells, or in pluripotent cells, during the germline cycle (Fig. 2). De novo retrotransposition events in the germline cycle have been proposed to shape the genomic landscape through mediating chromosomal re-arrangements and by acting as a reservoir for the emergence of new genes, in addition to providing a source of genetic variation within and between individuals [29, 30]. As the chromatin environment, transcription factor availability and post-transcriptional regulation of gene expression all vary significantly during the germline cycle, different retrotransposons have evolved to target different stages of germ cell development. The resulting variety of complex and dynamic retrotransposon expression patterns means that exaptation of retrotransposon sequences can play a role in the evolution of gene expression networks at multiple points within the germline cycle [29, 31–33].

### Retrotransposon expression in pre-implantation embryos

During pre-implantation development, the zygote undergoes a series of mitotic divisions to generate a ball of pluripotent cells known as a morula (Fig. 2). The morula compacts and cavitates to generate a blastocyst possessing trophectoderm, primitive endoderm and epiblast layers. The trophectoderm and primitive endoderm layers will give rise only to extra-embryonic structures, whereas the epiblast cells retain pluripotency and will give rise to all the tissues in the embryo, including the germ cells, after implantation (Fig. 2) [34]. Pre-implantation development only takes around 4 days in mice, but shows dynamic

**Fig. 2** Overview of the mouse germline cycle. Schematic overview of germ cell development in mice. Pluripotent cells are indicated in *italicised* text and germ cells in *bold* text. DNA is passed between germ cells and pluripotent cells through the generations in a germline cycle (*green arrows*). Differentiation into somatic tissues is indicated by *grey arrows*. The level of DNA methylation at different stages of development is indicated by the level of *shading*



changes in retrotransposon expression. For example, RNA transcripts belonging to one of the most abundant LTR retrotransposons in the mouse genome, IAP elements, are present at high levels in fully-grown oocytes, decrease in 1-cell embryos, then increase again during development to the blastocyst stage [32, 35, 36]. These IAP element transcripts are competent to generate A-type retroviral-like particles, whose abundance follows similar dynamics during these stages of development [36]. In contrast, epsilon-type retroviral-like particles, which are encoded by MuERV1 ERVL LTR retrotransposons, are not present in fully-grown oocytes, transiently increase in abundance in 2-cell embryos, then disappear as pre-implantation development proceeds [36]. Again, the changes in epsilon-type retroviral-like particle abundance are mirrored by changes in abundance of MuERV1 transcripts that encode these elements [32, 37]. The MT MaLR LTR retrotransposon transcripts exhibits yet another distinct expression pattern during pre-implantation development and are highly abundant in mouse oocytes but sharply decrease in abundance as pre-implantation development proceeds [32]. Thus, different types of retrotransposon have evolved to take advantage of the dynamic chromatin modifications and transcription factor profiles present at these stages of

development. Interestingly, chimaeric transcripts originating from retrotransposon promoters but spliced onto host genes are present in oocytes and pre-implantation embryos [32], suggesting that mammalian hosts are co-opting retrotransposons to drive gene expression during these stages of development.

#### IAP LTR retrotransposon expression during germ cell development

Germ cell development starts after implantation when the pluripotent epiblast differentiates into primordial germ cells in addition to the endoderm, mesoderm and ectoderm somatic tissues during gastrulation at E6.5–E7.5 (Fig. 2). The primordial germ cells proliferate and migrate to the genital ridges, colonising these structures around E10.5, and differentiate into meiotic oocytes or quiescent prospermatogonia by E13.5–E14.5 (Fig. 2). In males, the quiescent prospermatogonia can differentiate into spermatogonial stem cells a few days after birth, which will give rise to cells progressing through spermatogenesis (proliferating spermatogonia → meiotic spermatocytes → post-meiotic spermatids → sperm) throughout the adult life of the animal (Fig. 2). In females, the oocytes that initiate meiosis in

the foetus undergo meiotic arrest a few days after birth, and groups of these arrested oocytes are selected to grow and mature during each oestrus cycle. Oocyte meiosis is not completed until the ovulated egg is fertilised to generate a zygote (Fig. 2) [4]. The advent of next generation sequencing technologies, and their application to analyse the transcriptome of small numbers of cells, is likely to generate a wealth of data about genome-wide retrotransposon transcript levels at different stages of the germline cycle [38, 39]. However, many stages of germ cell development have not yet been extensively analysed, and much of our understanding of retrotransposon expression during gametogenesis comes from studies on specific elements.

One of the best-studied LTR retrotransposons in mice is the IAP element. The IAP LTR drives transcription of a *LacZ* reporter preferentially in germ cells rather than somatic cells [40], and contains binding sites for a number of widely-expressed transcription factors, including YY1, SP1, CREB1 and glucocorticoid receptors [41]. Thus, the germline-specific expression of IAP elements does not appear to be caused by germ cell-specific transcription factors. Rather, the preferential expression of IAP elements in the germline appears to be a consequence of IAP element DNA being hypomethylated in germ cells, but methylated and silenced in somatic cells [42–45]. Within the germline, IAP LTR-driven expression of a *LacZ* reporter is restricted to a fairly small window of germ cell development, with activity detectable in quiescent male foetal germ cells from E16, and in undifferentiated spermatogonia in adult testes [40]. IAP DNA is hypomethylated at these stages [39, 40, 46–48]. Although the IAP LTR is sufficient to drive germline expression of a *LacZ* reporter, it does not recapitulate endogenous IAP transcription in adult liver [49] nor in pre-implantation embryos [32, 35, 36]. The liver IAP transcripts mainly originate from a single IAP locus suggesting that the local chromatin environment or *cis*-acting mutations allow this copy of IAP to be expressed in the liver [50]. In contrast, the endogenous IAP transcripts present in pre-implantation embryos arise from at least two different IAP subtypes that have distinct expression patterns at these stages [35]. Sequences outside the LTR region might therefore be important for activating IAP expression in pre-implantation embryos.

#### LINE-1 retrotransposon expression during germ cell development


The non-LTR LINE-1 retrotransposon has also been shown to be expressed at specific points in germ cell development. LINE-1 RNA, protein, ribonucleoprotein particles and de novo retrotransposition events are all detectable in pluripotent stem cell lines, and LINE-1 expression is

downregulated when these cells differentiate into somatic cells [51–53]. A number of widely-expressed transcription factors including RUNX3 and YY1 have been implicated in activating human and/or mouse LINE-1 transcription [8, 54, 55]; however, more restricted transcription factors such as SOX2 could potentially confer some tissue and stage specificity [56]. LINE-1 RNA is also present in E11.5–E13.5 primordial germ cells that have colonised the genital ridge [39, 57]. However, LINE-1 ORF1 protein is not detectable in the germ cells until E15.5, a stage of development when male germ cells are quiescent and female germ cells are in leptotene/zygotene stages of meiotic prophase [58]. Interestingly, LINE-1 ORF1 protein levels decrease in germ cells after birth, but full-length LINE-1 transcripts and ORF1 protein peak again as male spermatocytes pass through the leptotene/zygotene stages of meiotic prophase [59]. Thus, early meiotic prophase may be a common point in male and female germline development for susceptibility to LINE-1 expression. LINE-1 ORF1 protein does not appear to be as abundant during later meiotic stages in either male or female germ cells [58, 59]. Interestingly, despite the high levels of LINE-1 expression during specific stages of male and female germline development, LINE-1 retrotransposition has been proposed to occur mainly in early pre-implantation embryos after fertilisation [60]. The poor correlation between LINE-1 RNA expression, LINE-1 protein expression and LINE-1 retrotransposition at some stages of the germline cycle could represent expression of non-functional full-length copies of LINE-1, or read-through transcription from adjacent host promoters into non-functional 5'-truncated copies of LINE-1. In addition, genome defence mechanisms could be acting to inhibit different stages of the LINE-1 life cycle (Fig. 3) and dampen down the retrotransposition potential of these elements.

#### Retrotransposon activity in somatic tissues

Any de novo retrotransposition occurring in pluripotent cells can potentially generate genetic variation in the somatic and germline components of an individual (Fig. 2). Somatic variation has been observed for de novo LINE-1 integration events in neurons, and this genetic heterogeneity has been proposed to be important for brain function [56, 61–63]. Although de novo LINE-1 integration events in neurons could arise from LINE-1 activity in pluripotent cells [60], LINE-1 expression within the neuronal lineage [56] is also likely to contribute to this phenomenon. Although successful retrotransposons need to be active in pluripotent cells or the developing germline, there is also some retrotransposon expression in somatic tissues. Analysis of LTR retrotransposon expression across multiple tissues using retrotransposon-specific microarrays

**Fig. 3** Retrotransposon suppression mechanisms. A schematic overview of a generalised retrotransposon life cycle encompassing transcription, RNA export, translation, assembly, nuclear import, and de novo integration is shown. Specific elements can show some variation in this process, e.g. RNA from non-autonomous SINE retrotransposons is not translated, and uses LINE-1-encoded proteins to catalyse their integration. Also, the mechanism of integration differs between LINE-1 and LTR retrotransposons. Genes involved in suppressing retrotransposons at specific stages of their life cycle in germ cells and pluripotent cells are indicated



suggests that different LTR retrotransposons have different somatic expression profiles [64, 65]. However, some of this somatic tissue-specific expression could represent the behaviour of a small number of anomalous copies of any particular retrotransposon, rather than an inherent tissue-specificity of the LTR [50, 66]. One somatic tissue that has long been associated with retrotransposon expression is the placenta [67]. The trophoblast-derived cells in the placenta have low levels of DNA methylation (Fig. 2), and their specialised epigenetic state is thought to make these cells permissive for expression of some retrotransposons [68, 69]. Specific subsets of LTR retrotransposons, including ERVWE1 elements in human and IAP elements in mouse [64, 69], are highly expressed in the placenta, and retrotransposon-encoded proteins and LTRs have been co-opted to perform host functions during placenta development [33, 70, 71]. Although retrotransposon expression or activity can be beneficial to the somatic tissues of their mammalian hosts, these elements have also been proposed to drive genetic instability and cancer [3, 19, 72]. Establishing mitotically heritable suppression of retrotransposons in the pluripotent cells early in development can therefore help to maintain genomic stability in the somatic tissues.

### Epigenetic silencing of retrotransposons by DNA methylation

One of the major mechanisms that mammalian cells use to defend their genomes against retrotransposons is transcriptional silencing (Fig. 3; Table 1). The main transcriptional silencing mechanisms known to be operating in mammalian cells typically involve adding and removing covalent modifications to DNA and histones [73]. These modifications can alter chromatin structure, and change the physical associations between proteins and the underlying DNA sequence. One type of modification that is truly epigenetic, in that it can mediate heritable changes in gene activity without changing DNA sequence, is DNA methylation [74].

#### DNA methylation in mammalian genomes

In mammals, DNA methylation occurs almost exclusively on cytosine residues in the context of CpG dinucleotides [74]. Most of the mammalian genome, including repetitive DNA sequences, is heavily methylated [75, 76], but this bulk methylation is interrupted by short stretches of DNA containing high densities of CpGs known as CpG islands

**Table 1** Repression mechanisms affecting different types of retrotransposon in mouse ES cells and germ cells

Retrotransposon family	Retrotransposon class	Enriched chromatin modifications (ES cells)	Repression mechanisms (ES cells)	Repression mechanisms (germ cells)
LINE-1	LINE	5mC	TRIM28, KDM1A	DNA methyltransferases, PIWI-piRNA pathway, TDRD7
ERV1	LTR	5mC, H3K9Me3, H4K20Me3	TRIM28, SETDB1, ZFP809, DNA methyltransferases, polycomb repressive complexes	Unknown
ERVK	LTR	5mC, H3K9Me3, H4K20Me3	TRIM28, SETDB1, HDAC1, DNA methyltransferases, polycomb repressive complexes	DNA methyltransferases, PIWI-piRNA pathway, LSH, TEX19.1, endogenous siRNA
ERVL	LTR	5mC, H3K27Me3	TRIM28, KDM1A, ZFP42, RYBP	Unknown
MaLR	LTR	5mC, H3K27Me3	TRIM28	Endogenous siRNA

Summary of some of the main mechanisms that have been shown to repress retrotransposons in mouse ES cells and germ cells. Different types of retrotransposon are repressed by different mechanisms, and are shown grouped by retrotransposon family

(CGIs) that escape DNA methylation [77]. The density of CpGs in the CGI affects the methylation status of these sequences, which are often found coincident with gene promoters. High density CpG CGI promoters are almost always unmethylated and transcriptionally active except in rare tissue-specific genes and some tumour suppressor genes in cancer cells [75, 76, 78]. Intermediate density CpG CGI promoters are rarer than high density CpG CGI promoters but are more likely to be silenced and methylated in a tissue-specific manner [76]. In contrast, low CpG density CGIs promoters tend to be methylated irrespective of their transcriptional status [76]. The correlation between DNA methylation and transcriptional repression is therefore strongly influenced by CpG density and genomic context.

#### Adding DNA methylation to the genome

Mammals possess three DNA methyltransferases that can catalyse methylation of CpG dinucleotides: DNMT1, DNMT3A and DNMT3B. DNMT1 perpetuates existing DNA methylation marks by methylating hemi-methylated DNA templates [79–81]. DNMT1 is thought to be recruited to sites of DNA replication by its protein partner PCNA [82], where it uses the parental strand as a methylation template to modify the newly synthesised daughter strand, thereby maintaining DNA methylation patterns during proliferation. A second protein, Np95 (also known as UHRF1), helps recruit DNMT1 to hemi-methylated CpGs generated at the replication fork during DNA replication [83, 84]. Np95 binding to heterochromatin-associated histone modifications during DNA replication is essential for the recruitment of this maintenance methyltransferase complex; this ensures that DNMT1 activity is stabilised

when in a complex with other heterochromatin-associated proteins during the late phase of DNA replication to perpetuate a global methylation profile [84, 85]. During early development and gametogenesis, tissue-specific DNA methylation patterns are established in part by the two de novo DNA methyltransferases DNMT3A and DNMT3B [86]. *Dnmt3a*<sup>-/-</sup> *Dnmt3b*<sup>-/-</sup> double-knockout embryos have hypomethylated DNA, start to exhibit developmental retardation by E8.5, and die by E11.5 [86]. Interestingly, the de novo DNA methyltransferases have non-overlapping roles at some genomic sequences: DNMT3B, but not DNMT3A, has a key role in regulating methylation at centromeric minor satellite repeats, and both interspersed C-type endogenous retroviruses (which includes MuLV ERV1 LTR retrotransposons) and IAP elements are slightly hypomethylated in *Dnmt3b*<sup>-/-</sup> embryos but retain methylation in *Dnmt3a*<sup>-/-</sup> embryos [86]. IAP hypomethylation is more pronounced in embryos lacking both DNMT3A and DNMT3B, suggesting a degree of redundancy between these two enzymes at these loci [86]. Detailed analysis of DNA methylation differences between these knockout embryos using current genome-wide approaches might provide some insight into the differences in specificity between the de novo DNA methyltransferases.

Like the *Dnmt3a*<sup>-/-</sup> *Dnmt3b*<sup>-/-</sup> double mutants, embryos that carry hypomorphic mutant alleles of *Dnmt1* become developmentally retarded soon after gastrulation, and die by E12.5 [87]. Hypomethylation at IAP elements and C-type endogenous retroviruses, and increased expression of IAP retrotransposon transcripts, have all been shown to occur in these hypomorphic *Dnmt1* mouse embryos [45, 87]. DNMT1 can act as a transcriptional repressor if it is recruited to unmethylated DNA by protein-interacting partners, and DNA methylation-independent

transcriptional repression by DNMT1 has been demonstrated in frogs and mammals [88, 89]. However, developmental retardation, embryonic lethality and IAP and LINE-1 retrotransposon hypomethylation all occur in embryos that carry catalytically inactive mutant alleles of *Dnmt1* [90], suggesting that these aspects of the *Dnmt1* mutant phenotype are a consequence of a failure to maintain DNA methylation.

#### Removing DNA methylation from the genome

The molecular pathways governing the removal of DNA methylation from mammalian genomes are also starting to be more fully realised, and the TET family of dioxygenases are central to this process [91–94]. The TET enzymes can iteratively oxidise 5mC to generate 5-hydroxymethyl cytosine (5hmC), 5-formyl cytosine (5fC), and 5-carboxyl cytosine (5caC) [95]. 5hmC, 5fC and 5caC are not maintained by DNMT1 during DNA replication, and can therefore lead to demethylation as they are diluted during cell proliferation [96, 97]. Alternatively, the DNA repair machinery can excise 5fC and 5caC, or deaminated derivatives of modified cytosine residues, to bring about demethylation [98, 99]. 5hmC is less abundant than 5mC in the genome and is typically enriched in the bodies of actively transcribed genes, and at active enhancers and sequences surrounding transcriptional start sites [100–102]. The distribution of 5hmC across the genome is distinct between tissues, and 5hmC patterns can be used as an identifier of cell type or disease state [103, 104]. 5fC and 5caC are even less abundant than 5hmC, and can also be detected at some enhancers and gene regulatory elements, particularly when components of the DNA repair machinery are mutated to allow these transient intermediates to accumulate [105, 106].

Mammals possess three TET enzymes. *Tet1*<sup>-/-</sup> and *Tet2*<sup>-/-</sup> mice each exhibit some phenotypic abnormalities, but these mice are viable and fertile [107–110]. In contrast, *Tet3* is essential for viability and *Tet3*<sup>-/-</sup> mice die perinatally [111]. There is redundancy between TET1 and TET2 as *Tet1*<sup>-/-</sup> *Tet2*<sup>-/-</sup> double-knockout mice show some lethality at both embryonic and perinatal stages [112]. However, some *Tet1*<sup>-/-</sup> *Tet2*<sup>-/-</sup> double-knockout mice are viable and, aside from small ovaries, have no overt phenotypic abnormalities [112]. *Tet1*<sup>-/-</sup> *Tet2*<sup>-/-</sup> double-knockout mice have reduced levels of 5hmC and increased levels of 5mC in somatic tissues, consistent with these genes having a role in converting 5mC to 5hmC [112]. TET enzymes play a role in reprogramming methylation patterns at some sequences during development, and are contributing to the general dynamic reprogramming of DNA methylation in the genome, which can possibly include retrotransposon sequences [105, 106, 110–113].

#### DNA methylation and transcriptional repression

DNA methylation is strongly enriched at retrotransposon sequences in mammalian genomes, so much so that the primary role of DNA methylation has been proposed to be transcriptional repression of these elements [114]. DNA methylation could exert its repressive effects on transcription by sterically interfering with transcription factors binding to their cognate binding sites in promoter proximal regulatory regions [115]. Alternatively, a number of methyl-CpG-binding proteins have been identified in mammals that could influence transcription factor accessibility or chromatin structure at genomic sites containing this epigenetic mark [116–119]. For example, the methyl-CpG-binding protein MECP2 is thought to mediate transcriptional repression of methylated DNA at least partly through its interaction with the SIN3A co-repressor complex [120]. It is not yet clear if methyl-CpG-binding proteins are necessary for global silencing of methylated target genes or retrotransposons in mammalian cells [121]. However, the strong correlation between DNA methylation and transcriptional repression in the genome [76] primarily represents DNA methylation acting to ‘lock-down’ transcription after a repressed state has already been established by other mechanisms, and only a small number of mammalian genes appear to rely on DNA methylation as a primary mechanism to silence their expression [43].

Experimentally removing DNA methylation in mouse somatic tissues strongly induces expression of IAP retrotransposons, suggesting that DNA methylation has a primary role in repressing these elements in mouse embryonic fibroblasts [42–44]. It is not clear how many other retrotransposons rely predominantly on DNA methylation for their repression in somatic tissues. MMVL30 and MuLV elements, which are both members of the ERV1 family of LTR retrotransposons, are transcriptionally upregulated in response to DNA hypomethylation in mouse embryonic fibroblasts, suggesting that DNA methylation might be a primary transcriptional silencing mechanism for these elements [122]. However, recent data, showing that re-distribution of polycomb-associated histone modifications is responsible for around a third of the transcriptional changes that occur in hypomethylated fibroblasts [123], illustrate the complex interactions that exist between different chromatin modifying mechanisms and the difficulties associated with interpreting these data mechanistically. Notably, both MMVL30 and MuLV elements are upregulated in ES cells carrying mutations in components of polycomb repressive complexes [124, 125].

Like the LTR retrotransposons, non-LTR LINE and SINE retrotransposons are also highly methylated in somatic tissues [126], but transcriptional silencing of these elements does not strongly depend on DNA methylation



in fibroblasts [122]. DNA methylation and MECP2 have been implicated in repressing mouse and human LINE-1 retrotransposons in neuronal cells [63]. However, MECP2 is a global regulator of neuronal chromatin structure, and the modest increase in retrotransposon expression seen in *Mecp2*<sup>-/-</sup> brains could reflect an increase in transcriptional noise [127].

### Transcriptional silencing of retrotransposon expression in embryonic stem cells

Although a number of studies have reported that loss of DNA methylation causes strong upregulation of IAP elements in somatic tissues, ES cells carrying mutations in all three DNA methyltransferases exhibit a more modest upregulation of these elements, even though IAP DNA methylation is severely reduced [42–45, 125, 128–130]. Therefore, DNA methylation does not play as dominant a role in IAP suppression in ES cells as it does in somatic cells. Microarray and deep-sequencing analysis suggests that a relatively small number of LTR retrotransposons are upregulated in the hypomethylated *Dnmt1*<sup>-/-</sup> *Dnmt3*<sup>-/-</sup> *Dnmt3b*<sup>-/-</sup> triple-knockout (*Dnmt*<sup>TKO</sup>) ES cells [125, 129]. Notably, LINE-1 elements and MuLV elements, which have been reported to be regulated by DNA methylation in neuronal and fibroblast somatic cell types, respectively [63, 122], are not strongly de-repressed in *Dnmt*<sup>TKO</sup> ES cells despite being hypomethylated [129, 130]. In contrast, LTR retrotransposons belonging to ERV1 (MMERGLN, RLTR1B) and ERVK (IAP, RLTR45, MMERVK10C, RMER16) families are all upregulated in *Dnmt*<sup>TKO</sup> ES cells [129]. The level of upregulation for each of these elements (~2.3- to 13-fold) is relatively modest, suggesting that additional retrotransposon silencing mechanisms might be operating in ES cells.

An alternative repressive modification that is associated with retrotransposons in ES cells is histone 3 lysine 9 trimethylation (H3K9Me3) [131]. H3K9Me3 is highly enriched on ERV1 and ERVK families of LTR retrotransposons in ES cells, but not on the ERVL and MaLR families [131]. ES cells that carry mutations in the SETDB1 (also known as ESET) histone methyltransferase have reduced levels of H3K9Me3 at LTR retrotransposon sequences, and a significant de-repression of many different types of LTR retrotransposon [125, 129, 132]. In contrast to the *Dnmt*<sup>TKO</sup> ES cells, 69 different LTR retrotransposons are de-repressed in *Setdb1*<sup>-/-</sup> ES cells [129]. The upregulated LTR retrotransposons primarily belong to the ERV1 and ERVK families, whereas ERVL and LINE-1 retrotransposons are not strongly upregulated in *Setdb1*<sup>-/-</sup> ES cells [129]. Interestingly, each LTR retrotransposon that is upregulated in *Dnmt*<sup>TKO</sup> ES cells is more strongly

upregulated in *Setdb1*<sup>-/-</sup> ES cells. Furthermore, the LTR retrotransposon upregulation in *Dnmt*<sup>TKO</sup> ES cells is not associated with decreased levels of H3K9Me3 at these elements, and DNA methylation at these elements is not strongly affected in *Setdb1*<sup>-/-</sup> ES cells, suggesting that these silencing mechanisms are recruited independently, and function in parallel, to repress transcription of these elements [129]. Although H3K9Me3 is enriched on ERV1 and ERVK LTR retrotransposons in fibroblasts, as it is in ES cells [131], a clear difference exists between these cell types in their requirement for SETDB1 in this process. Whereas SETDB1 is bound to ERV1 and ERVK retrotransposons and mediates H3K9Me3 and repression at these loci in ES cells, SETDB1 is neither bound at these sequences nor required for their H3K9Me3 in fibroblasts [129, 132]. Thus, SETDB1-dependent H3K9Me3 appears to play an important role in repressing ERV1 and ERVK retrotransposons in ES cells, but not in somatic fibroblasts.

Transcriptional silencing of the ERVL family of LTR retrotransposons in ES cells does not appear to depend on SETDB1 [129], but relies instead on the histone demethylase KDM1A [133]. *Kdm1a*<sup>-/-</sup> ES cells de-repress MERVL elements, LINE-1 elements and a number of chimeric transcripts containing endogenous genes driven from ERVL and MaLR promoters similar to those found in zygotes [133]. ERVL and MaLR retrotransposon DNA is highly methylated in ES cells and somatic tissues [39, 126], but DNA methylation at these elements is not perturbed in *Kdm1a*<sup>-/-</sup> ES cells [133]. Rather, loss of KDM1A results in a number of changes to the histone modifications present at ERVL and MaLR LTRs in ES cells, with histone 3 lysine 4 methylation (H3K4Me) and histone 3 lysine 27 acetylation (H3K27Ac) levels both increasing and H3K9Me3 levels decreasing [133]. KDM1A physically interacts with the HDAC family of histone deacetylases, and the increase in H3K4Me and H3K27Ac levels at target retrotransposons in *Kdm1a*<sup>-/-</sup> ES cells might reflect reduced demethylase (KDM1A) and deacetylase (HDAC) activities, respectively, at these sequences [133]. ERVL and MaLR retrotransposons are enriched for the polycomb-associated histone 3 lysine 27 trimethylation (H3K27Me3) modification in wild-type ES cells [131], but H3K27Ac will prevent acquisition of the repressive trimethyl modification at the same residue. ES cells carrying mutations in components of the polycomb repressive complexes de-repress various different retrotransposons, including ERV1 (MuLV, MMVL30) and ERVK (IAP, RLTR45) LTR retrotransposons [124, 125]. However, ERVL and MaLR LTR retrotransposons are not strongly de-repressed in these mutant ES cells, despite being strongly enriched for the H3K27Me3 mark [124, 125, 131]. Thus, the de-repression of ERVL elements and MaLR sequences in *Kdm1a*<sup>-/-</sup> probably reflects multiple changes in histone modification and chromatin structure

at these sequences. The *Kdm1a*-dependent repression of ERVL and MaLR retrotransposons seems to be associated with the zinc finger protein ZFP42 (also known as REX1). ZFP42 physically interacts with KDM1A and is bound to ERVL and MaLR retrotransposon DNA in ES cells [134]. Like *Kdm1a*<sup>-/-</sup> ES cells, *Zfp42*<sup>-/-</sup> ES cells also de-repress MERVL retrotransposons [134]. Thus, ZFP42 could potentially provide sequence specificity to this silencing mechanism and recruit KDM1A to MERVL and MaLR retrotransposon DNA in order to transcriptionally repress these elements. RYBP, a protein that interacts with ZFP42, also plays a role in repressing MERVL retrotransposons in ES cells [135].

Although there are differences in the mechanisms repressing the different LTR retrotransposon families (summarised in Table 1), one factor that has strong effects across the entire class of LTR retrotransposons is TRIM28 (also known as KAP1). TRIM28 plays a role in silencing multiple LTR retrotransposons in ES cells including MuLV, IAP, MERVL and MT elements belonging to each of the ERV1, ERVK, ERVL and MaLR families [136, 137]. *Trim28*<sup>-/-</sup> ES cells also modestly upregulate LINE-1 retrotransposons [136]. TRIM28 can be recruited to specific DNA sequences via its interactions with Krüppel-associated box zinc finger proteins, and has been shown to interact with repressive chromatin-modifying enzymes such as SETDB1 and KDM1A [133, 138]. In ES cells, TRIM28 is recruited to a specific sequence in MuLV retrotransposons by the zinc finger protein ZFP809 [137, 139]. The sequences responsible for TRIM28-dependent repression of MuLV are distinct from the sequences implicated in TRIM28-dependent repression of IAP elements, suggesting that different DNA-binding proteins are probably involved in recruiting TRIM28 to different retrotransposons [136]. *Trim28*<sup>-/-</sup> ES cells have reduced levels of H3K9Me3 at IAP elements, which presumably reflects impaired recruitment of SETDB1 to these loci, and upregulate IAP transcription around 20-fold [129, 132, 136, 138]. Notably, DNA methylation at IAP elements is not strongly affected in *Trim28*<sup>-/-</sup> ES cells [136, 140], suggesting that TRIM28 is silencing retrotransposons independently of DNA methylation in this cell type. Interestingly, the primary transcriptional silencing mechanism for IAP retrotransposons appears to switch from being TRIM28/SETDB1-dependent histone modification in ES cells to DNA methylation in somatic cells [42, 43, 45, 128, 129, 132, 136]. Recent findings that differentiation of pluripotent *Trim28*<sup>-/-</sup> cells into somatic cells results in some reduction in the amount of DNA methylation at IAP retrotransposons suggests that there is a temporal link between TRIM28 and DNA methylation during differentiation and development [140]. These results are consistent with the concept that DNA methylation is generally recruited to chromatin after a repressed

state has been established to stably repress transcription during somatic differentiation.

### Transcriptional silencing of retrotransposons in the germline

Although a number of recent advances have been made in identifying transcriptional silencing mechanisms operating on retrotransposons in pluripotent cells, much less is known about the transcriptional regulation of these elements in developing germ cells (Table 1). DNA methylation and many histone modifications have been shown to undergo transient global decreases at multiple points in the germline cycle [141–144]. Genome-wide analysis of the sequences associated with specific histone modifications during germ cell development is starting to become technically possible [145], and it will be of interest to elucidate whether some of the transcriptional silencing mechanisms operating on retrotransposons in ES cells are also important for retrotransposon silencing in germ cells. However, at present, the best-studied transcriptional silencing mechanism operating at retrotransposon sequences in germ cells is DNA methylation.

#### DNA methylation in the developing germline

DNA methylation patterns undergo dynamic changes during germ cell development (Fig. 2). The global level of DNA methylation in primordial germ cells decreases progressively between E8.5 and E12.5, although different sequences exhibit different dynamics during this process [39, 46, 146]. Genome-wide bisulfite sequencing shows that LINE retrotransposons, SINE retrotransposons and ERV1, ERVK, ERVL and MaLR LTR retrotransposons are all hypomethylated in E13.5 germ cells, and typically only have around 20 % of their CpG's methylated compared to around 80 % in foetal somatic tissues [126]. Some LTR retrotransposons, such as IAP elements, are more resistant to this hypomethylation event than others and still retain an intermediate amount of DNA methylation (~60 %) in E13.5 germ cells [39, 147, 148]. Multiple intersecting pathways are likely to be involved in mediating this global demethylation, and conversion of 5mC to 5hmC by TET1 and TET2 enzymes [113], deamination of 5mC by AID (also known as AICDA) [126], passive loss of cytosine modifications during DNA replication [39, 113, 149], active removal of nucleotides by the DNA repair machinery [150] and down-regulation of DNA methyltransferases and their accessory factors [39, 46, 151] are all implicated in this process.

Once the germ cells have undergone global hypomethylation, DNA methylation patterns are re-established on retrotransposon sequences in a sex-specific manner (Fig. 2).

Male germ cells undergo de novo methylation during late foetal and early postnatal development [152], long before male germ cells initiate meiosis. Genome-wide bisulfite sequencing shows that de novo methylation in the male germline generates sperm with similar, or even slightly higher, genome-wide and retrotransposon methylation levels to foetal somatic tissues [39]. In contrast, de novo methylation in female germ cells occurs in postnatal growing oocytes, after the oocytes have completed the early stages of meiotic prophase and are held in a dictyate stage meiotic arrest [153, 154]. De novo methylation increases the amount of DNA methylation in postnatal oocytes, but mature eggs remain hypomethylated relative to foetal somatic tissues and sperm [154, 155]. Although LINE, SINE, and LTR retrotransposons are all de novo methylated during oocyte growth, some copies of these retrotransposons remain partially unmethylated in mature eggs [154, 155]. Acquisition of de novo methylation at some imprinted genes and intragenic CGIs in oocytes is associated with transcription through these regions [154, 156], but it is not clear whether this association extends to retrotransposons.

After fertilisation, DNA methylation declines during pre-implantation development (Fig. 2), with LINE, SINE and LTR retrotransposon sequences all losing DNA methylation [155, 157, 158]. Some retrotransposons, including IAP elements, are somewhat resistant to this hypomethylation event [148, 158]. At the blastocyst stage of pre-implantation development, DNA methylation levels increase in the pluripotent epiblast cells [157], and genome-wide bisulfite sequencing of ES cells derived from this tissue shows that global and retrotransposon DNA methylation levels are similar between ES cells, post-implantation epiblast, and foetal somatic tissues [39, 126]. In contrast, DNA methylation levels remain low in the trophectoderm layer of the blastocyst [157], and the placenta has moderate levels of methylation (~40–50 %) at LINE, SINE and LTR retrotransposon sequences [126], representing a mixed population of hypomethylated trophoctoderm-derived and hypermethylated epiblast-derived tissues.

#### De novo DNA methylation of retrotransposon sequences during gametogenesis


Mutations that interfere with de novo DNA methylation might be expected to cause defects in epigenetic reprogramming and retrotransposon silencing in the developing germline. The lethality of *Dnmt3a*<sup>-/-</sup> and *Dnmt3b*<sup>-/-</sup> embryos has necessitated the generation of conditional mutants to investigate the function of these genes in the germline [159]. *Dnmt3a*<sup>-/-</sup> and *Dnmt3b*<sup>-/-</sup> prospermatogonia have normal DNA methylation levels at IAP and LINE-1 elements, possibly indicating functional redundancy between DNMT3A and DNMT3B at these loci [152,

159, 160]. In contrast, de novo methylation of B1 SINE retrotransposons in spermatogonia requires DNMT3A but not DNMT3B [152]. In oocytes, there is redundancy between DNMT3A and DNMT3B for de novo DNA methylation at IAP elements, but only DNMT3A is involved in de novo DNA methylation of LINE-1 elements [161]. Thus, the de novo methyltransferases have different specificities for retrotransposon sequences.

The de novo methyltransferase activity of both DNMT3A and DNMT3B is stimulated by the catalytically inactive DNA methyltransferase-like protein DNMT3L [162]. In contrast to *Dnmt3a* and *Dnmt3b*, *Dnmt3l* is not required for viability or retrotransposon DNA methylation in somatic tissues. However, *Dnmt3l*<sup>-/-</sup> mice are infertile and have defects in de novo methylation during spermatogenesis and oogenesis [152, 163–166]. *Dnmt3l*<sup>-/-</sup> prospermatogonia and oocytes have reduced levels of DNA methylation at LINE-1, IAP and B1 SINE retrotransposons [152, 154, 155, 161, 163, 166]. The hypomethylation at IAP and LINE-1 retrotransposons in *Dnmt3l*<sup>-/-</sup> prospermatogonia is associated with transcriptional activation of these elements in foetal and perinatal testes [163]. De novo methylation of IAP and LINE-1 retrotransposons in foetal male germ cells therefore seems to be required for transcriptional silencing of these elements in the prospermatogonia.

Another factor that has been linked with de novo DNA methylation of retrotransposons sequences and genome defence is LSH (also known as HELLS). LSH is not restricted to the germline and is required for normal embryonic DNA methylation of repetitive elements and some single copy genes in somatic tissues [167, 168]. LSH has been proposed to recruit the de novo DNA methyltransferase DNMT3B to specific target sites in the genome [169]. Notably, loss of LSH function is sufficient to de-repress IAP transcription in mouse embryonic fibroblasts, alter pericentromeric heterochromatin, and induce gene expression changes at single copy genes [170, 171]. The DNA hypomethylation at IAP elements that occurs in *Lsh*<sup>-/-</sup> somatic tissue is also evident in *Lsh*<sup>-/-</sup> oocytes, and IAP expression is greatly increased in *Lsh*<sup>-/-</sup> foetal ovaries [172]. As IAP is normally hypomethylated in meiotic oocytes until the postnatal oocyte growth phase [153], either the IAP hypomethylation in *Lsh*<sup>-/-</sup> oocytes is affecting a subset of CpGs critical for transcriptional repression of these elements, or the transcriptional upregulation of IAP in *Lsh*<sup>-/-</sup> ovaries is a feature of the somatic cells in the ovary rather than oocytes. The role of *Lsh* in de novo DNA methylation in foetal prospermatogonia and growing oocytes has not yet been studied, and it will be of interest to determine whether de novo methylation events occurring at different stages of the germline cycle (Fig. 2) have different requirements to direct the DNA methylation machinery to target loci.

**Fig. 4** Expression patterns and mutant phenotypes of germline genome defence genes. The stages of spermatogenesis are indicated along the top of the diagram, and expression patterns of the indicated germline genome defence genes are indicated by blue bars. The stages at which mutant mice are reported to have defects in progression through spermatogenesis are indicated with crosses



#### PIWI proteins and retrotransposon DNA methylation

A number of additional genes are also required for de novo methylation and transcriptional silencing of retrotransposons in male germ cells. PIWI proteins belong to a subclade of the Argonaute family of small RNA binding proteins and contain RNA binding and RNaseH-like endonuclease domains [173]. Mice possess three PIWI proteins: MIWI (also known as PIWIL1), MILI (also known as PIWIL2), and MIWI2 (also known as PIWIL4), that each physically interact with populations of germline-restricted small, single-stranded RNAs known as PIWI-interacting RNAs (piRNAs) [174–178]. Expression of the three PIWI proteins is largely restricted to the germline, but each PIWI protein has a different expression pattern during germ cell development (Fig. 4). *Mili* and *Miwi2* are expressed in late foetal male germ cells when de novo methylation of retrotransposon DNA occurs [176, 178]. *Mili*<sup>-/-</sup> and *Miwi2*<sup>-/-</sup> prospermatogonia fail to de novo methylate IAP and LINE-1 elements, and de-repress transcription of these elements [47, 175–177]. It is not clear whether the methylation state of other retrotransposons is affected in these mutant germ cells, and genome-wide analysis of their DNA methylation patterns would be of interest in this regard.

Although *Mili* and *Miwi2* are required genetically for the de novo methylation of retrotransposons [47, 175–177], the mechanistic role that PIWI proteins play in this process is not fully understood. MILI and MIWI2 physically interact with piRNAs that originate from genomic piRNA clusters,

retrotransposons and other genomic regions [47, 175, 176]. Ping-pong amplification of piRNA sequences occurs when an antisense piRNAs guides the cleavage of a complementary mRNA to generate sense piRNAs, which in turn can guide cleavage of complementary sequences in piRNA cluster transcripts to generate more antisense piRNAs to feed back into the system. Molecular chaperones are thought to be important to dissociate PIWI-piRNA complexes, allowing ping-pong amplification to occur [179]. MILI and MIWI2 physically associate with different subpopulations of piRNA in foetal prospermatogonia, and have different roles in de novo methylation of retrotransposon sequences [47, 176]. The putative endonuclease activity of MILI appears to be required for ping-pong amplification of retrotransposon piRNAs in foetal prospermatogonia, and de novo methylation and transcriptional silencing of LINE-1, but not IAP elements [180]. In contrast, the putative endonuclease activity of MIWI2 is not required for these events [180], and MIWI2 has been proposed to function by translocating into the nucleus when loaded with piRNA, where it uses the piRNA sequence as a guide to recruit the de novo methylation machinery to actively transcribing retrotransposons [176]. However, physical interactions between MIWI2 and either DNMT3A or DNMT3B have not been detected in foetal prospermatogonia [176]. It is possible that the MIWI2/piRNA complexes recruit the de novo DNA methylation machinery indirectly, perhaps through inducing repressive histone modifications at retrotransposon loci. However, although there is strong genetic evidence that

*Mili* and *Miwi2* are required for de novo DNA methylation of LINE-1 and IAP retrotransposons in male germ cells, the requirement for piRNA in this phenomenon is yet to be formally demonstrated. In this respect, it is notable that the third mouse PIWI protein, MIWI, has recently been shown to have some piRNA-independent functions in regulating gene expression [181].

De novo methylation of retrotransposons in foetal spermatogonia also requires a number of additional factors that act in the PIWI–piRNA pathway. Tudor domain-containing proteins physically interact with methylated arginine residues in PIWI proteins [182–184], and co-localise with PIWI proteins and piRNAs in electron-dense cytoplasmic structures termed nuage [185–188]. The formation of these structures and interactions between them are thought to be crucial for the biogenesis of piRNAs and functional piRNA–PIWI complexes, and TDRD1, TDRD5 and TDRD9 are all required for normal DNA methylation of LINE-1 retrotransposons [186, 187, 189]. *Tdrd1*<sup>-/-</sup>, *Tdrd5*<sup>-/-</sup> and *Tdrd9*<sup>-/-</sup> mutations all have stronger effects on transcriptional silencing of LINE-1 than IAP retrotransposons [186, 187, 189], again suggesting that there are differences in the requirements for de novo methylation of these elements.

Additional nuage-associated factors, including MITOPLD (also known as PLD6), MOV10L1, MVH (also known as DDX4), MAEL, and ASZ1 (also known as GASZ) are all required for DNA methylation and transcriptional silencing of LINE-1 retrotransposons through their involvement in the piRNA–PIWI pathway [190–194]. MITOPLD and MOV10L1 appear to be required for piRNA biogenesis [192, 193], and MVH is reported to be involved in piRNA processing [194]. MAEL does not appear to be required for piRNA biogenesis, but is speculated to play a role in assembly of PIWI–piRNA complexes or shuttling PIWI–piRNA complexes to or from the nuage [191], whereas ASZ1 is required to stabilise MILI [190].

Although each of the three mouse PIWI proteins is expressed in male germ cells and required for spermatogenesis, only MILI is expressed in the female germline [176–178, 195, 196]. MILI is not required for oogenesis but does appear to play a role in retrotransposon suppression in oocytes [197]. The levels of IAP transcript are modestly upregulated 3.5-fold in *Mili*<sup>-/-</sup> oocytes, but, in contrast to *Mili*<sup>-/-</sup> male germ cells, LINE-1 transcript levels are not affected. The level of IAP upregulation in *Mili*<sup>-/-</sup> oocytes is around 10-fold lower than the level of upregulation in *Mili*<sup>-/-</sup> testes [175], and it remains to be determined if the increase in IAP expression in *Mili*<sup>-/-</sup> mutant oocytes is due to defects in de novo methylation at these elements, or is a result of MILI's proposed additional roles in post-transcriptional regulation [47, 198].

### Post-transcriptional suppression of retrotransposon RNAs in the germline

There is some evidence that retrotransposon RNAs can be suppressed at a post-transcriptional level in germ cells (Fig. 3). The putative endonuclease activity of PIWI proteins could potentially allow PIWI–piRNA complexes to destabilise retrotransposon RNAs via endonucleolytic cleavage. This ‘slicer’ activity towards retrotransposon mRNAs has recently been demonstrated for MIWI [199]. MIWI is expressed postnatally from the zygotene stage of meiosis until the haploid round spermatid stage [195], although the piRNA populations in post-pachytene male germ cells show little evidence of ping-pong amplification [181, 200]. MIWI-associated piRNAs are able to guide MIWI to cleave target RNAs with extensive complementarity [199]. LINE-1 RNA is a target for MIWI slicer activity and LINE-1 transcript levels are increased ~10-fold in round spermatids carrying a catalytically inactive allele of *Miwi* that lacks endonuclease activity [199]. LINE-1 DNA methylation and repeat-derived piRNA abundance are not altered in the *Miwi* endonuclease mutant round spermatids; however, the likely post-transcriptional increase in LINE-1 transcripts in these mutants could represent a combination of piRNA-dependent and piRNA-independent functions of MIWI [199, 201]. It is not clear if MIWI's slicer activity is also important for post-transcriptional suppression of other types of retrotransposon, but this activity is likely to act as a back-up genome defence mechanism to target LINE-1 retrotransposons that have escaped suppression by transcriptional silencing mechanisms.

Post-transcriptional suppression of retrotransposons also occurs in female germ cells. Endogenous double-stranded small interfering RNAs (siRNAs) are present in oocytes, where they play a role in post-transcriptional silencing of retrotransposons [197, 202, 203]. Production of these endogenous siRNAs requires DICER1, an endoribonuclease that is expressed at high levels throughout oocyte growth [197]. The endogenous siRNAs in oocytes physically associate with AGO2 [197], an Argonaute protein that forms the catalytic component of the RNA-induced silencing complex [204]. Members of the ERVK and MaLR families of LTR retrotransposons (RLTR10 and MTA elements, respectively) are upregulated ~3- to 5-fold in *Dicer1*<sup>-/-</sup> and *Ago2*<sup>-/-</sup> growing oocytes, but LINE-1 and IAP transcript abundance does not change [197, 203]. Interestingly, whereas *Dicer1/Ago2* suppress MTA and RLTR10 but not IAP retrotransposons in growing oocytes, *Mili* suppresses IAP but not MTA or RLTR10 retrotransposons in these cells [197]. Thus, the *Mili*/piRNA and *Dicer1/Ago2*/endogenous siRNA pathways are associated with silencing different retrotransposons in growing oocytes.

Interestingly, LINE-1 expression in growing oocytes is not affected by any of the *Mili*<sup>-/-</sup>, *Dicer1*<sup>-/-</sup> or *Ago2*<sup>-/-</sup> mutations [197]. However, post-transcriptional silencing of LINE-1 retrotransposons in the fully-grown oocytes depends on the putative RNA-binding protein MARF1 [205]. *Marf1*<sup>-/-</sup> oocytes exhibit numerous changes in mRNA abundance, including an upregulation in the levels of LINE-1 and IAP transcripts, despite there being no significant changes in IAP or LINE-1 DNA methylation [205]. It will be of interest to determine whether MARF1 directly interacts with LINE-1, IAP and any other retrotransposon mRNA in oocytes, and how any sequence specificity is determined.

### Other mechanisms of retrotransposon suppression

Although germ cells possess multiple mechanisms to prevent the accumulation of retrotransposon transcripts (Fig. 3), retrotransposon RNAs are readily detectable and even abundant in germ cells and pluripotent cells in wild-type mice. Furthermore, discrepancies between full-length retrotransposon transcript abundance, protein abundance, and retrotransposition activity at different stages of the germline cycle suggest that additional mechanisms that suppress retrotransposons at later stages of their life cycles are operating in the germline [32, 35, 38, 40, 52, 59, 60]. The Tudor domain-containing protein TDRD7 plays a role in translational regulation of LINE-1 retrotransposons in male germ cells [206], and two of the PIWI proteins, MILI and MIWI, physically associate with actively translating mRNAs in polysomes and are implicated in translational regulation [198, 207]. Interestingly, some piRNAs are also physically associated with polysomes in mouse testes [207]. It is not known whether retrotransposon mRNAs are targets for translational regulation by MILI or MIWI, but there is evidence that piRNAs can suppress translation of LINE-1 mRNAs. Deletion of a genomic cluster of piRNAs (*Nct1/2*) that includes an antisense LINE-1 sequence results in a modest ~1.5-fold increase in LINE-1 mRNA, but a striking ~15-fold increase in levels of LINE-1-encoded ORF1 protein [208]. piRNA-containing protein complexes could suppress LINE-1 translation in a similar way that microRNAs suppress translation of their target mRNAs [204].

Genome defence mechanisms operating at even later stages of the retroviral life cycle have been demonstrated in somatic tissues. MOV10, a putative RNA helicase, physically associates with human LINE-1 ribonucleoprotein particles and mouse IAP retroviral-like particles, and inhibits retrotransposition of both these elements in cellular assays, and can affect multiple stages of the retroviral life cycle [209–211]. Interestingly, germ cells express a paralog of the somatic *Mov10* gene, *Mov10l1*. MOV10L1

physically interacts with MILI, MIWI2 and MIWI proteins, and *Mov10l1*<sup>-/-</sup> testes have reduced amounts of piRNA, hypomethylation of LINE-1 DNA and de-repression of LINE-1 and IAP elements [193, 212]. Conditional deletion of *Mov10l1* from the pachytene stage of spermatogenesis onwards reveals that MOV10L1 has additional functions in late spermatogenesis [201]. *Mov10l1*<sup>-/-</sup> round spermatids display elevated DNA damage, which could indicate increased retrotransposition in these cells. Although LINE-1 and IAP RNA levels are not elevated in *Mov10l1*<sup>-/-</sup> round spermatids [201], MOV10L1 could, like its somatic paralog, be suppressing LINE-1 and IAP activity at later stages of the retrotransposon life cycle.

Another protein that is operating to repress retrotransposons in the developing germline is TEX19.1, a predominantly cytoplasmic protein in germ cells and pluripotent cells that represses the ERVK LTR retrotransposon MMERVK10C in postnatal testes [213]. Unlike a number of genes implicated in germline genome defence, *Tex19.1* does not appear to have a strong effect on endogenous gene expression, and the largest changes in gene expression in *Tex19.1*<sup>-/-</sup> testes are retrotransposon transcripts [125, 213]. LINE-1, B1 SINE and IAP retrotransposons are not upregulated in *Tex19.1*<sup>-/-</sup> knockout testes, and microarray analysis suggests that retrotransposon upregulation in *Tex19.1*<sup>-/-</sup> testes is primarily restricted to MMERVK10C [125, 213]. The exact molecular mechanisms of how TEX19.1 suppresses accumulation of MMERVK10C transcripts is not yet known, but studies on the role of *Tex19.1* in somatic placenta tissue illustrate one of the complexities in determining retrotransposon targets for genome defence mechanisms: in contrast to *Tex19.1*<sup>-/-</sup> testes, multiple retrotransposons including MMVL30 LTR retrotransposons and LINE-1 elements are upregulated in *Tex19.1*<sup>-/-</sup> placenta [69]. The presence of multiple complementary genome defence mechanisms in the developing germline is likely to be masking some of the effects of mutating individual retrotransposon suppression mechanisms in these cells.

### Regulation of germline genome defence genes by DNA methylation

A number of genes involved in suppressing retrotransposons in the germline have been shown to be regulated by DNA methylation. *Dazl*, *Zfp42*, *Mvh* and *Mael* have all been identified as belonging to groups of genes that require *Dnmt3b*-dependent promoter DNA methylation to silence their expression [214, 215]. Furthermore, analysis of gene expression changes in multiple somatic cell models for hypomethylation identified a core subset of 26 methylation-sensitive genes, which is highly enriched for genes involved in suppressing retrotransposons in the

germline [43]. This group of methylation-sensitive germline genome defence genes, which includes *Tex19.1*, *Mili*, *Dazl*, *Asz1* and *Mov10l1*, is strongly and heritably upregulated in response to DNA hypomethylation [43]. Interestingly, methylation-sensitive germline genome defence gene promoters are not enriched for commonly studied repressive histone modifications, such as H3K27Me3, H3K9Me3 and histone 4 lysine 20 trimethylation (H4K20Me3), when they are transcriptionally repressed in somatic cells [43]. The lack of enrichment for these repressive histone modifications could explain why silencing of these genes is so dependent on DNA methylation.

During germline development, expression of the methylation-sensitive germline genome defence genes is initiated during the global wave of demethylation that occurs progressively in primordial germ cells between E8.5 and E12.5 (Fig. 4) [43, 46, 146]. Hypomethylation and expression of *Tex19.1* and *Mili* initiates early during this reprogramming event at ~E8.5–E9.5 while the primordial germ cells are migrating [43]. Hypomethylation and initiation of *Dazl*, *Mov10l1* and *Asz1* expression occurs towards the end of this epigenetic reprogramming event at ~E10.5–E11.5, after the germ cells have colonised the gonad [43, 216]. It is not known whether the differences in the timing of hypomethylation at different germline genome defence gene promoters reflect recruitment of different demethylation machineries to these genes [98]. Generally, expression of the germline genome defence genes is maintained throughout foetal germ cell development (Fig. 4) and in oocytes in adult females. In males, postnatal expression of the germline genome defence genes starts to decline in pachytene spermatocytes (*Tex19.1*, *Mili* and *Dazl*) and in round spermatids (*Asz1* and *Mov10l1*) (Fig. 4) [43, 176, 178, 190, 193, 217–219]. It is not clear whether DNA methylation transcriptionally represses these genes in these later stages of spermatogenesis.

Global DNA hypomethylation in the primordial germ cells extends to retrotransposon sequences [39, 126, 147, 148], and removing this extra layer of repression could potentially lead to variant copies of these elements being transcribed. The activation of IAP LTR transgenes in male foetal germ cells from ~E16 [40] is consistent with the hypothesis that at least some copies of these methylation-sensitive retrotransposons are becoming transcriptionally de-repressed in hypomethylated germ cells. Indeed, it is possible that some transcriptional de-repression of retrotransposons is a pre-requisite for the generation of piRNA, and PIWI–piRNA-mediated targeting of de novo methylation to retrotransposon sequences [176]. Notably, germline genome defence genes that act at post-transcriptional stages of the retrotransposon life cycle (Fig. 3) could play an important role in limiting retrotransposon activity during these periods of global hypomethylation. Coupling

expression of post-transcriptional genome defence mechanisms to the initiation of this epigenetic reprogramming process provides an effective way to ensure that appropriate genome defence mechanisms are active during the developmental window when the potential for retrotransposons to become active is high.

Hypomethylated cells in the mammalian placenta (Fig. 2) are another potential site for transcriptional de-repression of retrotransposons and activation of methylation-sensitive germline genome defence genes. Retrotransposons are hypomethylated in the placenta [68, 126, 220], but *Tex19.1* is the only one of the germline genome defence genes to be strongly hypomethylated and expressed at similar levels in the placenta and the testis [69]. LINE-1 retrotransposons are upregulated in the hypomethylated trophoblast-derived component of *Tex19.1*<sup>-/-</sup> placentas, suggesting that this germline genome defence gene is functionally repressing retrotransposons in this hypomethylated somatic context [69]. Although the germline genome defence genes are normally methylated and silenced in embryo-derived somatic tissues, expression of this group of genes can be induced in somatic contexts in response to environmental or toxicological insults that perturb DNA methylation. Widespread changes in DNA methylation have been reported to occur in numerous cancers [221, 222], and genetic instability caused by retrotransposon activity can drive tumourigenesis [19]. The potential functional roles of the germline genome defence genes in hypomethylated somatic contexts is clearly an exciting area that requires further investigation.

### Consequences of failures in genome defence

Defects in germline genome defence might be expected to result in increased rates of retrotransposition, and new retrotransposon integrations being passed on to the next generation. However, mutations in most of the germline genome defence genes result in male infertility, making it difficult to assess de novo retrotransposition rates in their offspring. Curiously, many of the germline genome defence mutants have strong phenotypic similarities and common arrest points in spermatogenesis (Fig. 4). De-repression of retrotransposons might therefore have additional and more immediate consequences for germ cell development than generating new retrotransposition events in the next generation.

Consequences of mutations in the DNA methylation machinery

At a cellular level, *Dnmt3l*<sup>-/-</sup> mice display defective spermatogonial proliferation, delayed entry into meiosis and

arrested meiotic progression at pachytene in young adults. Older mutant adults become completely azoospermic due to progressive loss of spermatogonia, suggesting defects in the maintenance of spermatogonial stem cells [163, 223]. Although *Dnmt3l*<sup>-/-</sup> mutant prospermatogonia have defects in DNA methylation of multiple genomic features [152, 163], it is not clear how DNA hypomethylation causes the spermatogonial stem cell or meiotic phenotypes in these mice.

Impaired maintenance of spermatogonial stem cells and delayed entry into meiosis also occur in *Dnmt3a*<sup>-/-</sup> germ cells, whereas *Dnmt3b*<sup>-/-</sup> germ cells have no severe disruptions in spermatogenesis and are able to form functional sperm [159, 160]. Thus, methylation at DNMT3A-specific targets such as B1 SINE retrotransposons or some imprinted genes might be important for spermatogonial stem cell maintenance and timing of meiotic entry, but hypomethylation at DNMT3B-specific targets such as satellite sequences [152] does not disrupt spermatogenesis. Notably, the meiotic arrest seen in *Dnmt3l*<sup>-/-</sup> testes is not recapitulated in either the *Dnmt3a*<sup>-/-</sup> or *Dnmt3b*<sup>-/-</sup> single-knockout male germ cells [159, 160]. Thus, hypomethylation at redundant genomic targets of DNMT3A and DNMT3B, such as LINE-1 retrotransposons, IAP retrotransposons or some imprinted genes [152], are likely to be responsible for the meiotic defects in *Dnmt3l*<sup>-/-</sup> testes.

The meiotic arrest evident in *Dnmt3l*<sup>-/-</sup> spermatocytes is characterised by defective synaptonemal complex formation, with widespread chromosome asynapsis and non-homologous synapsis [163]. Recombination proteins RAD51 and RPA1 localise to chromosome axis-associated foci at a similar frequency in wild-type and *Dnmt3l*<sup>-/-</sup> spermatocytes [224], indicating that the chromosome synapsis defect is not a downstream consequence of a failure to initiate meiotic recombination. The presence of asynapsed chromosomes in *Dnmt3l*<sup>-/-</sup> pachytene spermatocytes sequesters the transcriptional silencing machinery, impairing meiotic sex chromosome inactivation and leading to spermatocyte apoptosis [224]. The meiotic arrest in *Dnmt3l*<sup>-/-</sup> testes is perhaps surprising given that expression of *Dnmt3l* is so low during meiosis [223], and presumably reflects defects in de novo methylation that arise earlier in the foetal prospermatogonia.

Although *Dnmt3l*<sup>-/-</sup> male mice have numerous cellular defects in spermatogenesis, meiosis and oogenesis proceed normally in females, notwithstanding the failure in maternal imprinting [164, 165]. The differential requirement for DNMT3L in meiotic chromosome synapsis between males and females likely reflects de novo methylation occurring before the initiation of meiosis in males, but after meiotic chromosome synapsis is complete in females [152, 153]. Oocytes therefore have low levels of genomic DNA methylation while they progress through the early stages of

meiotic prophase [39]. The low level of DNA methylation that is present in meiotic oocytes appears to be required for normal chromosome synapsis and progression through meiosis. *Lsh*<sup>-/-</sup> oocytes have reduced levels of DNA methylation, and, although these oocytes initiate meiotic recombination normally, they fail to synapse their homologous chromosomes properly [172]. The meiotic defects in hypomethylated *Lsh*<sup>-/-</sup> oocytes bear some similarity to those in hypomethylated *Dnmt3l*<sup>-/-</sup> spermatocytes [163, 172, 224]. Thus, even though there are sex-specific differences in the mechanisms that generate the correct DNA methylation patterns for meiosis, reducing DNA methylation has similar consequences for meiotic chromosome synapsis in both spermatocytes and oocytes.

### Consequences of mutations in the PIWI–piRNA system

The PIWI–piRNA pathway acts genetically upstream of *Dnmt3l*-dependent de novo DNA methylation, and therefore many of the genome defence genes that are important for PIWI–piRNA function in foetal prospermatogonia share cellular spermatogenic defects with *Dnmt3l*<sup>-/-</sup> mice. Like *Dnmt3l*<sup>-/-</sup> testes, self-renewal/maintenance of spermatogonial stem populations are reduced in *Mili*<sup>-/-</sup> testes and in *Miwi2*<sup>-/-</sup> testes [177, 198]. The male germ cells that progress to meiosis in *Miwi2*<sup>-/-</sup> testes are able to initiate meiotic recombination, but chromosome synapsis is severely defective in these cells [177]. Interestingly, these meiotic defects arise some time after *Miwi2* is expressed during spermatogenesis (Fig. 4), suggesting that defects in *Miwi2*<sup>-/-</sup> prospermatogonia are being heritably transmitted through multiple rounds of mitosis. *Mili*<sup>-/-</sup> spermatocytes also fail to progress through pachytene, although the precise nature of the meiotic defects in these cells has not been determined [196]. At least some *Mili*<sup>-/-</sup> single-knockout and *Mili*<sup>-/-</sup> *Miwi*<sup>-/-</sup> double-knockout spermatocytes appear to successfully synapse their chromosomes, and progress further into pachytene than *Miwi2*<sup>-/-</sup> spermatocytes [177, 196, 225]. As MILI is required for localisation of MIWI2 to cytoplasmic granules and accumulation of the MIWI2-interacting piRNA population [176], it is not clear whether the apparent differences between the *Mili*<sup>-/-</sup> and *Miwi2*<sup>-/-</sup> meiotic phenotypes reflect complex interactions between these PIWI proteins during meiosis, or differences in the phenotypic analyses. Taken together, these data suggest that the PIWI–piRNA pathway and de novo DNA methylation in foetal prospermatogonia facilitate chromosome synapsis and progression through pachytene in post-natal meiotic spermatocytes.

Male mice with mutations in other genes required for piRNA biogenesis or PIWI–piRNA function, including *Mov10l1*, *Asz1*, *Mvh* and *Mitopl1d*, all have defects in progression through the zygotene/pachytene stages of meiotic



prophase where chromosome synapsis occurs [190, 192, 193, 212, 226]. Where studied, chromosome synapsis is defective in these mutant spermatocytes, and the meiotic phenotype resembles *Miwi2*<sup>-/-</sup> mice [192, 193]. Female mice are fertile for each of these mutants, suggesting that chromosome synapsis is able to occur in meiotic oocytes in the absence of PIWI-piRNA function. *Mael* is also implicated in PIWI-piRNA pathways for germline genome defence, and is one of the few germline genome defence mutants where there is evidence of novel retrotransposition events taking place in the germline [191]. In the absence of MAEL, male mice are infertile, have defects in meiotic chromosome synapsis, and arrest at pachytene. A large increase in the levels of DNA damage is present in *Mael*<sup>-/-</sup> spermatocytes, and elegant genetic experiments have shown that the generation of DNA damage in *Mael*<sup>-/-</sup> spermatocytes is independent of the SPO11 endonuclease that generates the meiotic DNA double-strand breaks required for recombination [191]. The presence of nuclear LINE-1 ORF1 protein correlates with high levels of the DNA damage marker  $\gamma$ H2AX in *Mael*<sup>-/-</sup> spermatocytes, implying that novel DNA damage is being generated by retrotransposons integrating into the genome.

Maintaining suppression of retrotransposons during spermatogenesis appears to be important even after meiotic chromosome synapsis. Spermatogenesis in *Miwi*<sup>-/-</sup> testes arrests at the round spermatid stage with increased abundance of LINE-1 transcripts, and elevated DNA damage [195, 199]. Impairing post-pachytene piRNA populations by conditionally deleting *Mov10l1* in pachytene spermatocytes also causes elevated DNA damage in round spermatids and spermatogenic arrest at this stage [201]. Although increased DNA damage and round spermatid arrest appear to be a consequence of a loss of piRNA-dependent MIWI slicer activity [199], further work is needed to determine whether these cellular phenotypes are caused by increased retrotransposon activity, or by other changes in these mutants. In contrast to spermatogenesis, mutations in the PIWI-piRNA system do not have severe defects on post-pachytene progression through oogenesis, despite *Mili*<sup>-/-</sup> oocytes exhibiting a modest increase in retrotransposon expression [197].

#### Consequences of mutations in other germline genome defence mechanisms

DAZL is a germ cell-specific RNA binding protein that is essential for fertility in both male and female mice [217, 227–229]. Spermatogonial differentiation is greatly impaired in *Dazl*<sup>-/-</sup> testis, and, although a small proportion of germ cells progress to meiosis, they arrest at pachytene [230]. Various genes have been identified that are translationally regulated by DAZL, including the germline

genome defence genes *Mvh* and *Tex19.1*, and genes known to be important for meiosis such as *Sycp3* [227, 228]. As DAZL translationally regulates *Mvh* and *Tex19.1*, it is likely that impaired translation of these genome defence genes accounts for some aspects of the *Dazl*<sup>-/-</sup> phenotype.

Unlike many of the other germline genome defence mutants, loss of *Tex19.1* causes defects during both oogenesis and spermatogenesis [213]. Approximately half the *Tex19.1*<sup>-/-</sup> pachytene spermatocytes have chromosome synapsis defects and exhibit apoptosis at this stage, while two-thirds of the nuclei progressing through to metaphase I contain at least one set of unpaired univalent chromosomes. Although the retrotransposon de-repression, chromosome asynapsis and pachytene arrest in *Tex19.1*<sup>-/-</sup> males are all shared with the other male germline genome defence mutants, the spectrum of retrotransposons de-repressed in *Tex19.1*<sup>-/-</sup> testes is distinct from PIWI-piRNA pathway mutants [213]. TEX19.1 is stabilised by the E3 ubiquitin ligase UBR2 [231], but no PIWI-piRNA pathway components have been connected to TEX19.1 or UBR2. Interestingly, the de-repression of MMERVK10C retrotransposons in *Tex19.1*<sup>-/-</sup> testes occurs in pachytene spermatocytes [213]; however, it is currently unclear what causes the chromosome synapsis defects in *Tex19.1*<sup>-/-</sup> spermatocytes.

#### Mechanisms linking retrotransposon de-repression to meiotic chromosome asynapsis

Mutations in germline genome defence genes tend to cause male-specific defects in chromosome synapsis and arrest meiosis during pachytene (Fig. 4), although most female mutants do not have a severe phenotype at this stage of meiosis. The cause of the chromosome asynapsis in these male mutants is not well understood. Where reported, DNA double-strand breaks are generated at the start of meiosis in the genome defence mutants, indicating successful initiation of meiotic recombination [177, 191, 213, 224]. These meiotic DNA double-strand breaks repair as chromosomes synapse, but remain unrepaired on asynapsed regions [191, 213, 224], as would be expected from analysis of other meiotic mutants [232]. One informative feature of the chromosome asynapsis in the germline genome defence mutants is that the asynapsed chromosomes in these mutants are not paired, suggesting that the asynapsis is caused by a defect in the homology search rather than synaptonemal complex assembly.

Interestingly, *Mael*<sup>-/-</sup> mutants have increased amounts of SPO11-independent DNA damage, possibly caused by increased retrotransposition of LINE-1 elements, and these SPO11-independent DNA breaks recruit the homologous recombination protein RAD51 [191]. Thus, the increased DNA damage could be sequestering RAD51 and other meiotic recombination proteins away from meiotic

double-strand DNA breaks generated by SPO11, thereby impairing the homology search. While moderate levels of additional DNA double-strand breaks generated by ionising radiation are well tolerated in spermatocytes [233], there appears to be a level at which DNA double-strand breaks become disruptive to meiotic progression. *Atm*<sup>-/-</sup> spermatocytes exhibit a tenfold increase in DNA double-strand breaks, which is accompanied by chromosome asynapsis [234]. Reducing the number of DNA double-strand breaks in *Atm*<sup>-/-</sup> spermatocytes by reducing the dosage of *Spo11* rescues the asynapsis [234]. Therefore, increased retrotransposition creating elevated levels DNA double-strand breaks could be contributing to meiotic chromosome asynapsis in genome defence mutants.

Although *Mael*<sup>-/-</sup> spermatocytes have elevated levels of DNA double-strand breaks, not all genome defence mutants exhibit this phenotype. Unlike *Mael*<sup>-/-</sup> spermatocytes,  $\gamma$ H2AX and RAD51 staining in *Dnmt3l*<sup>-/-</sup> spermatocytes is relatively normal [224]. An alternative explanation for the defects in the meiotic homology search in genome defence mutants is that global changes in chromosome structure or organisation caused by widespread DNA hypomethylation could be generating an inappropriate chromosomal environment for the homology search. Alternatively, it is possible that raised levels of retrotransposon proteins present in genome defence mutants might be sufficient to disrupt the cellular environment of the developing male germ cells. Proteins encoded by human LTR retrotransposons physically interact with endogenous transcription factors present in germ cells and can disrupt spermatogenesis when ectopically expressed in transgenic mice [235–237]. Physical interactions between retrotransposon proteins and germ cell proteins involved in the meiotic homology search could explain the meiotic chromosome asynapsis in germline genome defence mutants. Further work is clearly needed to dissect out the mechanism causing the spermatocyte asynapsis and male infertility in the genome defence mutants.

## Concluding remarks

Germline genome defence mechanisms would be expected to play a role in maintaining genome stability over evolutionary time scales. However, it is becoming apparent that mutations in germline genome defence genes cause infertility and defects in progression through meiosis that are precluding any analysis of genome stability in their offspring. The mechanistic explanation for this unexpected requirement for the germline genome defence genes is not understood, and it is not clear if the meiotic defects in these mutants are related to retrotransposon de-repression, or to uncharacterised meiotic functions of the germline genome

defence genes. Imbalances between genome defence and retrotransposon activities could be causing infertility in mouse models in a similar way that incompatibilities between retrotransposon activity and germline genome defence systems cause sterility and hybrid dysgenesis in fruit flies [238]. It will be of interest to determine whether failures in germline genome defence systems, or their epigenetic regulation [239], could be contributing to infertility in humans.

The regulatory coupling of genome defence mechanisms to the potential for retrotransposon activity is an interesting paradigm that raises a number of questions. The epigenetic disruption and recovery screen used to identify these genes was performed in fibroblasts and used stringent thresholds for activation of gene expression [43]. Additional genome defence genes, such as *Mael* and *Mvh*, could be regulated by DNA methylation in germ cells but not identified in this screen due to the absence of tissue-specific transcription factors, or the presence of additional layers of regulation in fibroblasts. Therefore, more work is needed to better understand tissue-specific responses to epigenetic disruption. Furthermore, there are likely to be some differences between the genome defence mechanisms operating in different species. Indeed, human retrotransposons are de-repressed in transchromosomal mice, suggesting that mouse genome defence mechanisms cannot effectively suppress human retrotransposon sequences [240]. Therefore, although the links between DNA methylation, retrotransposon suppression and genome defence genes are starting to be uncovered in mice, further work will be required to investigate whether these associations exist in other species, and whether similar associations exist in general between different retrotransposon transcriptional silencing mechanisms and different groups of genome defence genes.

**Acknowledgments** We thank Abby Wilson (MRC HGU) for the chromosome image used in Fig. 3, and Duncan Sproul and Ruchi Shukla (both MRC HGU) for comments on the manuscript. Research in our laboratories is funded by the MRC (I.R.A., R.R.M.) and the BBSRC (R.R.M.).

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

1. Lander ES, Linton LM, Birren B et al (2001) Initial sequencing and analysis of the human genome. *Nature* 409:860–921. doi:10.1038/35057062
2. Waterston RH, Lindblad-Toh K, Birney E et al (2002) Initial sequencing and comparative analysis of the mouse genome. *Nature* 420:520–562. doi:10.1038/nature01262

3. Romanish MT, Cohen CJ, Mager DL (2010) Potential mechanisms of endogenous retroviral-mediated genomic instability in human cancer. *Semin Cancer Biol* 20:246–253. doi:[10.1016/j.semcancer.2010.05.005](https://doi.org/10.1016/j.semcancer.2010.05.005)
4. Öllinger R, Reichmann J, Adams IR (2010) Meiosis and retrotransposon silencing during germ cell development in mice. *Differentiation* 79:147–158. doi:[10.1016/j.diff.2009.10.004](https://doi.org/10.1016/j.diff.2009.10.004)
5. Zamudio N, Bourc'his D (2010) Transposable elements in the mammalian germline: a comfortable niche or a deadly trap? *Heredity* 105:92–104. doi:[10.1038/hdy.2010.53](https://doi.org/10.1038/hdy.2010.53)
6. Rowe HM, Trono D (2011) Dynamic control of endogenous retroviruses during development. *Virology* 411:273–287. doi:[10.1016/j.virol.2010.12.007](https://doi.org/10.1016/j.virol.2010.12.007)
7. Khan H, Smit A, Boissinot S (2006) Molecular evolution and tempo of amplification of human LINE-1 retrotransposons since the origin of primates. *Genome Res* 16:78–87. doi:[10.1101/gr.4001406](https://doi.org/10.1101/gr.4001406)
8. Lee S-H, Cho S-Y, Shannon MF et al (2010) The impact of CpG island on defining transcriptional activation of the mouse L1 retrotransposable elements. *PLoS ONE* 5:e11353. doi:[10.1371/journal.pone.0011353](https://doi.org/10.1371/journal.pone.0011353)
9. Sookdeo A, Hepp CM, McClure MA, Boissinot S (2013) Revisiting the evolution of mouse LINE-1 in the genomic era. *Mob DNA* 4:3. doi:[10.1186/1759-8753-4-3](https://doi.org/10.1186/1759-8753-4-3)
10. DeBerardinis RJ, Goodier JL, Ostertag EM, Kazazian HH Jr (1998) Rapid amplification of a retrotransposon subfamily is evolving the mouse genome. *Nat Genet* 20:288–290. doi:[10.1038/3104](https://doi.org/10.1038/3104)
11. Moran JV, Holmes SE, Naas TP et al (1996) High frequency retrotransposition in cultured mammalian cells. *Cell* 87:917–927. doi:[10.1016/S0092-8674\(00\)81998-4](https://doi.org/10.1016/S0092-8674(00)81998-4)
12. Feng Q, Moran JV, Kazazian HH Jr, Boeke JD (1996) Human L1 retrotransposon encodes a conserved endonuclease required for retrotransposition. *Cell* 87:905–916
13. Khazina E, Weichenrieder O (2009) Non-LTR retrotransposons encode noncanonical RRM domains in their first open reading frame. *Proc Natl Acad Sci USA* 106:731–736. doi:[10.1073/pnas.0809964106](https://doi.org/10.1073/pnas.0809964106)
14. Martin SL, Bushman FD (2001) Nucleic acid chaperone activity of the ORF1 protein from the mouse LINE-1 retrotransposon. *Mol Cell Biol* 21:467–475. doi:[10.1128/MCB.21.2.467-475.2001](https://doi.org/10.1128/MCB.21.2.467-475.2001)
15. Mathias SL, Scott AF, Kazazian HH Jr et al (1991) Reverse transcriptase encoded by a human transposable element. *Science* 254:1808–1810
16. Wei W, Gilbert N, Ooi SL et al (2001) Human L1 retrotransposition: cis preference versus trans complementation. *Mol Cell Biol* 21:1429–1439. doi:[10.1128/MCB.21.4.1429-1439.2001](https://doi.org/10.1128/MCB.21.4.1429-1439.2001)
17. Hancks DC, Kazazian HH Jr (2012) Active human retrotransposons: variation and disease. *Curr Opin Genet Dev* 22:191–203. doi:[10.1016/j.gde.2012.02.006](https://doi.org/10.1016/j.gde.2012.02.006)
18. Maksakova IA, Romanish MT, Gagnier L et al (2006) Retroviral elements and their hosts: insertional mutagenesis in the mouse germ line. *PLoS Genet* 2:e2. doi:[10.1371/journal.pgen.0020002](https://doi.org/10.1371/journal.pgen.0020002)
19. Shukla R, Upton KR, Muñoz-Lopez M et al (2013) Endogenous retrotransposition activates oncogenic pathways in hepatocellular carcinoma. *Cell* 153:101–111. doi:[10.1016/j.cell.2013.02.032](https://doi.org/10.1016/j.cell.2013.02.032)
20. Dewannieux M, Heidmann T (2005) L1-mediated retrotransposition of murine B1 and B2 SINEs recapitulated in cultured cells. *J Mol Biol* 349:241–247. doi:[10.1016/j.jmb.2005.03.068](https://doi.org/10.1016/j.jmb.2005.03.068)
21. Dewannieux M, Esnault C, Heidmann T (2003) LINE-mediated retrotransposition of marked Alu sequences. *Nat Genet* 35:41–48. doi:[10.1038/ng1223](https://doi.org/10.1038/ng1223)
22. Kramerov DA, Vassetzky NS (2011) SINEs. *Wiley Interdiscip Rev RNA* 2:772–786. doi:[10.1002/wrna.91](https://doi.org/10.1002/wrna.91)
23. Kriegs JO, Churakov G, Jurka J et al (2007) Evolutionary history of 7SL RNA-derived SINEs in Supraprimates. *Trends Genet* 23:158–161. doi:[10.1016/j.tig.2007.02.002](https://doi.org/10.1016/j.tig.2007.02.002)
24. Zabolotneva AA, Bantysh O, Suntsova MV et al (2012) Transcriptional regulation of human-specific SVAF<sub>1</sub> retrotransposons by cis-regulatory MAST2 sequences. *Gene* 505:128–136. doi:[10.1016/j.gene.2012.05.016](https://doi.org/10.1016/j.gene.2012.05.016)
25. Gilbert N, Bomar JM, Burmeister M, Moran JV (2004) Characterization of a mutagenic B1 retrotransposon insertion in the jittery mouse. *Hum Mutat* 24:9–13. doi:[10.1002/humu.20060](https://doi.org/10.1002/humu.20060)
26. Bannert N, Kurth R (2006) The evolutionary dynamics of human endogenous retroviral families. *Annu Rev Genomics Hum Genet* 7:149–173. doi:[10.1146/annurev.genom.7.080505.115700](https://doi.org/10.1146/annurev.genom.7.080505.115700)
27. Dewannieux M, Dupressoir A, Harper F et al (2004) Identification of autonomous IAP LTR retrotransposons mobile in mammalian cells. *Nat Genet* 36:534–539. doi:[10.1038/ng1353](https://doi.org/10.1038/ng1353)
28. Ribet D, Dewannieux M, Heidmann T (2004) An active murine transposon family pair: retrotransposition of “master” MusD copies and ETn trans-mobilization. *Genome Res* 14:2261–2267. doi:[10.1101/gr.2924904](https://doi.org/10.1101/gr.2924904)
29. Muotri AR, Marchetto MCN, Coufal NG, Gage FH (2007) The necessary junk: new functions for transposable elements. *Hum Mol Genet* 16 Spec No. 2:R159–R167. doi:[10.1093/hmg/ddm196](https://doi.org/10.1093/hmg/ddm196)
30. Singer T, McConnell MJ, Marchetto MCN et al (2010) LINE-1 retrotransposons: mediators of somatic variation in neuronal genomes? *Trends Neurosci* 33:345–354. doi:[10.1016/j.tins.2010.04.001](https://doi.org/10.1016/j.tins.2010.04.001)
31. Rebollo R, Romanish MT, Mager DL (2012) Transposable elements: an abundant and natural source of regulatory sequences for host genes. *Annu Rev Genet* 46:21–42. doi:[10.1146/annurev-genet-110711-155621](https://doi.org/10.1146/annurev-genet-110711-155621)
32. Peaston AE, Evsikov AV, Graber JH et al (2004) Retrotransposons regulate host genes in mouse oocytes and preimplantation embryos. *Dev Cell* 7:597–606. doi:[10.1016/j.devcel.2004.09.004](https://doi.org/10.1016/j.devcel.2004.09.004)
33. Chuong EB, Rumi MAK, Soares MJ, Baker JC (2013) Endogenous retroviruses function as species-specific enhancer elements in the placenta. *Nat Genet* 45:325–329. doi:[10.1038/ng.2553](https://doi.org/10.1038/ng.2553)
34. Rossant J, Tam PPL (2009) Blastocyst lineage formation, early embryonic asymmetries and axis patterning in the mouse. *Development* 136:701–713. doi:[10.1242/dev.017178](https://doi.org/10.1242/dev.017178)
35. Piko L, Hammons MD, Taylor KD (1984) Amounts, synthesis, and some properties of intracisternal A particle-related RNA in early mouse embryos. *Proc Natl Acad Sci USA* 81:488–492
36. Yotsuyanagi Y, Szöllösi D (1981) Early mouse embryo intracisternal particle: fourth type of retrovirus-like particle associated with the mouse. *J Natl Cancer Inst* 67:677–685
37. Ribet D, Louvet-Vallée S, Harper F et al (2008) Murine endogenous retrovirus MuERV-L is the progenitor of the “orphan” epsilon viruslike particles of the early mouse embryo. *J Virol* 82:1622–1625. doi:[10.1128/JVI.02097-07](https://doi.org/10.1128/JVI.02097-07)
38. Fadloun A, Le Gras S, Jost B et al (2013) Chromatin signatures and retrotransposon profiling in mouse embryos reveal regulation of LINE-1 by RNA. *Nat Struct Mol Biol* 20:332–338. doi:[10.1038/nsmb.2495](https://doi.org/10.1038/nsmb.2495)
39. Seisenberger S, Andrews S, Krueger F et al (2012) The dynamics of genome-wide DNA methylation reprogramming in mouse primordial germ cells. *Mol Cell* 48:849–862. doi:[10.1016/j.molcel.2012.11.001](https://doi.org/10.1016/j.molcel.2012.11.001)
40. Dupressoir A, Heidmann T (1996) Germ line-specific expression of intracisternal A-particle retrotransposons in transgenic mice. *Mol Cell Biol* 16:4495–4503
41. Kuff EL, Lueders KK (1988) The intracisternal A-particle gene family: structure and functional aspects. *Adv Cancer Res* 51:183–276

42. Davis CM, Constantinides PG, van der Riet F et al (1989) Activation and demethylation of the intracisternal A particle genes by 5-azacytidine. *Cell Differ Dev* 27:83–93
43. Hackett JA, Reddington JP, Nestor CE et al (2012) Promoter DNA methylation couples genome-defence mechanisms to epigenetic reprogramming in the mouse germline. *Development* 139:3623–3632. doi:10.1242/dev.081661
44. Jackson-Grusby L, Beard C, Possemato R et al (2001) Loss of genomic methylation causes p53-dependent apoptosis and epigenetic deregulation. *Nat Genet* 27:31–39. doi:10.1038/83730
45. Walsh CP, Chaillet JR, Bestor TH (1998) Transcription of IAP endogenous retroviruses is constrained by cytosine methylation. *Nat Genet* 20:116–117. doi:10.1038/2413
46. Hajkova P, Erhardt S, Lane N et al (2002) Epigenetic reprogramming in mouse primordial germ cells. *Mech Dev* 117:15–23. doi:10.1016/S0925-4773(02)00181-8
47. Kuramochi-Miyagawa S, Watanabe T, Gotoh K et al (2008) DNA methylation of retrotransposon genes is regulated by Piwi family members MIL1 and MIWI2 in murine fetal testes. *Genes Dev* 22:908–917. doi:10.1101/gad.1640708
48. Lees-Murdock DJ, De Felici M, Walsh CP (2003) Methylation dynamics of repetitive DNA elements in the mouse germ cell lineage. *Genomics* 82:230–237
49. Dupressoir A, Puech A, Heidmann T (1995) IAP retrotransposons in the mouse liver as reporters of ageing. *Biochim Biophys Acta* 1264:397–402
50. Puech A, Dupressoir A, Loireau MP et al (1997) Characterization of two age-induced intracisternal A-particle-related transcripts in the mouse liver. Transcriptional read-through into an open reading frame with similarities to the yeast ccr4 transcription factor. *J Biol Chem* 272:5995–6003
51. Martin SL (1991) Ribonucleoprotein particles with LINE-1 RNA in mouse embryonal carcinoma cells. *Mol Cell Biol* 11:4804–4807
52. Martin SL, Branciforte D (1993) Synchronous expression of LINE-1 RNA and protein in mouse embryonal carcinoma cells. *Mol Cell Biol* 13:5383–5392
53. Garcia-Perez JL, Morell M, Scheys JO et al (2010) Epigenetic silencing of engineered L1 retrotransposition events in human embryonic carcinoma cells. *Nature* 466:769–773. doi:10.1038/nature09209
54. Athanikar JN, Badge RM, Moran JV (2004) A YY1-binding site is required for accurate human LINE-1 transcription initiation. *Nucleic Acids Res* 32:3846–3855. doi:10.1093/nar/gkh698
55. Yang N, Zhang L, Zhang Y, Kazazian HH Jr (2003) An important role for RUNX3 in human L1 transcription and retrotransposition. *Nucleic Acids Res* 31:4929–4940
56. Muotri AR, Chu VT, Marchetto MCN et al (2005) Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. *Nature* 435:903–910. doi:10.1038/nature03663
57. Hayashi K, Chuva de Sousa Lopes SM, Kaneda M et al (2008) MicroRNA biogenesis is required for mouse primordial germ cell development and spermatogenesis. *PLoS ONE* 3:e1738. doi:10.1371/journal.pone.0001738
58. Trelogan SA, Martin SL (1995) Tightly regulated, developmentally specific expression of the first open reading frame from LINE-1 during mouse embryogenesis. *Proc Natl Acad Sci USA* 92:1520–1524
59. Branciforte D, Martin SL (1994) Developmental and cell type specificity of LINE-1 expression in mouse testis: implications for transposition. *Mol Cell Biol* 14:2584–2592
60. Kano H, Godoy I, Courtney C et al (2009) L1 retrotransposition occurs mainly in embryogenesis and creates somatic mosaicism. *Genes Dev* 23:1303–1312. doi:10.1101/gad.1803909
61. Baillie JK, Barnett MW, Upton KR et al (2011) Somatic retrotransposition alters the genetic landscape of the human brain. *Nature* 479:534–537. doi:10.1038/nature10531
62. Evrony GD, Cai X, Lee E et al (2012) Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell* 151:483–496. doi:10.1016/j.cell.2012.09.035
63. Muotri AR, Marchetto MCN, Coufal NG et al (2010) L1 retrotransposition in neurons is modulated by MeCP2. *Nature* 468:443–446. doi:10.1038/nature09544
64. Gimenez J, Montgiraud C, Pichon J-P et al (2010) Custom human endogenous retroviruses dedicated microarray identifies self-induced HERV-W family elements reactivated in testicular cancer upon methylation control. *Nucleic Acids Res* 38:2229–2246. doi:10.1093/nar/gkp1214
65. Seifarth W, Frank O, Zeilfelder U et al (2005) Comprehensive analysis of human endogenous retrovirus transcriptional activity in human tissues with a retrovirus-specific microarray. *J Virol* 79:341–352. doi:10.1128/JVI.79.1.341-352.2005
66. Reiss D, Zhang Y, Mager DL (2007) Widely variable endogenous retroviral methylation levels in human placenta. *Nucleic Acids Res* 35:4743–4754. doi:10.1093/nar/gkm455
67. Haig D (2012) Retroviruses and the placenta. *Curr Biol* 22:R609–R613. doi:10.1016/j.cub.2012.06.002
68. Chapman V, Forrester L, Sanford J et al (1984) Cell lineage-specific undermethylation of mouse repetitive DNA. *Nature* 307:284–286
69. Reichmann J, Reddington JP, Best D et al (2013) The genome-defence gene *Tex19.1* suppresses LINE-1 retrotransposons in the placenta and prevents intra-uterine growth retardation in mice. *Hum Mol Genet* 22:1791–1806. doi:10.1093/hmg/ddt029
70. Dupressoir A, Vernochet C, Harper F et al (2011) A pair of co-opted retroviral envelope syncytin genes is required for formation of the two-layered murine placental syncytiotrophoblast. *Proc Natl Acad Sci USA* 108:E1164–E1173. doi:10.1073/pnas.1112304108
71. Mi S, Lee X, Li X et al (2000) Syncytin is a captive retroviral envelope protein involved in human placental morphogenesis. *Nature* 403:785–789. doi:10.1038/35001608
72. Solyom S, Ewing AD, Rahrman EP et al (2012) Extensive somatic L1 retrotransposition in colorectal tumors. *Genome Res*. doi:10.1101/gr.145235.112
73. Cedar H, Bergman Y (2009) Linking DNA methylation and histone modification: patterns and paradigms. *Nat Rev Genet* 10:295–304. doi:10.1038/nrg2540
74. Reddington JP, Pennings S, Meehan RR (2013) Non-canonical functions of the DNA methylome in gene regulation. *Biochem J* 451:13–23. doi:10.1042/BJ20121585
75. Lister R, Pelizzola M, Dowen RH et al (2009) Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* 462:315–322. doi:10.1038/nature08514
76. Meissner A, Mikkelsen TS, Gu H et al (2008) Genome-scale DNA methylation maps of pluripotent and differentiated cells. *Nature* 454:766–770. doi:10.1038/nature07107
77. Thomson JP, Skene PJ, Selfridge J et al (2010) CpG islands influence chromatin structure via the CpG-binding protein Cfp1. *Nature* 464:1082–1086. doi:10.1038/nature08924
78. Illingworth R, Kerr A, Desousa D et al (2008) A novel CpG island set identifies tissue-specific methylation at developmental gene loci. *PLoS Biol* 6:e22. doi:10.1371/journal.pbio.0060022
79. Cedar H, Bergman Y (2012) Programming of DNA methylation patterns. *Annu Rev Biochem* 81:97–117. doi:10.1146/annurev-biochem-052610-091920
80. Freitag M, Selker EU (2005) Controlling DNA methylation: many roads to one modification. *Curr Opin Genet Dev* 15:191–199. doi:10.1016/j.gde.2005.02.003

81. Weber M, Schübeler D (2007) Genomic patterns of DNA methylation: targets and function of an epigenetic mark. *Curr Opin Cell Biol* 19:273–280. doi:[10.1016/j.ceb.2007.04.011](https://doi.org/10.1016/j.ceb.2007.04.011)
82. Araujo FD, Croteau S, Slack AD et al (2001) The DNMT1 target recognition domain resides in the N terminus. *J Biol Chem* 276:6930–6936. doi:[10.1074/jbc.M009037200](https://doi.org/10.1074/jbc.M009037200)
83. Sharif J, Muto M, Takebayashi S et al (2007) The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature* 450:908–912. doi:[10.1038/nature06397](https://doi.org/10.1038/nature06397)
84. Smith ZD, Meissner A (2013) DNA methylation: roles in mammalian development. *Nat Rev Genet* 14:204–220. doi:[10.1038/nrg3354](https://doi.org/10.1038/nrg3354)
85. Rothbart SB, Krajewski K, Nady N et al (2012) Association of UHRF1 with methylated H3K9 directs the maintenance of DNA methylation. *Nat Struct Mol Biol* 19:1155–1160. doi:[10.1038/nsmb.2391](https://doi.org/10.1038/nsmb.2391)
86. Okano M, Bell DW, Haber DA, Li E (1999) DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* 99:247–257
87. Li E, Bestor TH, Jaenisch R (1992) Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* 69:915–926
88. Dunican DS, Ruzov A, Hackett JA, Meehan RR (2008) xDnmt1 regulates transcriptional silencing in pre-MBT *Xenopus* embryos independently of its catalytic function. *Development* 135:1295–1302. doi:[10.1242/dev.016402](https://doi.org/10.1242/dev.016402)
89. Milutinovic S, Brown SE, Zhuang Q, Szyf M (2004) DNA methyltransferase 1 knock down induces gene expression by a mechanism independent of DNA methylation and histone deacetylation. *J Biol Chem* 279:27915–27927. doi:[10.1074/jbc.M312823200](https://doi.org/10.1074/jbc.M312823200)
90. Takebayashi S, Tamura T, Matsuoka C, Okano M (2007) Major and essential role for the DNA methylation mark in mouse embryogenesis and stable association of DNMT1 with newly replicated regions. *Mol Cell Biol* 27:8243–8258. doi:[10.1128/MCB.00899-07](https://doi.org/10.1128/MCB.00899-07)
91. Ito S, D'Alessio AC, Taranova OV et al (2010) Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature* 466:1129–1133. doi:[10.1038/nature09303](https://doi.org/10.1038/nature09303)
92. Ko M, Huang Y, Jankowska AM et al (2010) Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature* 468:839–843. doi:[10.1038/nature09586](https://doi.org/10.1038/nature09586)
93. Koh KP, Yabuuchi A, Rao S et al (2011) Tet1 and Tet2 regulate 5-hydroxymethylcytosine production and cell lineage specification in mouse embryonic stem cells. *Cell Stem Cell* 8:200–213. doi:[10.1016/j.stem.2011.01.008](https://doi.org/10.1016/j.stem.2011.01.008)
94. Tahiliani M, Koh KP, Shen Y et al (2009) Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* 324:930–935. doi:[10.1126/science.1170116](https://doi.org/10.1126/science.1170116)
95. Ito S, Shen L, Dai Q et al (2011) Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* 333:1300–1303. doi:[10.1126/science.1210597](https://doi.org/10.1126/science.1210597)
96. Inoue A, Zhang Y (2011) Replication-dependent loss of 5-hydroxymethylcytosine in mouse preimplantation embryos. *Science* 334:194. doi:[10.1126/science.1212483](https://doi.org/10.1126/science.1212483)
97. Inoue A, Shen L, Dai Q et al (2011) Generation and replication-dependent dilution of 5fC and 5caC during mouse preimplantation development. *Cell Res* 21:1670–1676. doi:[10.1038/cr.2011.189](https://doi.org/10.1038/cr.2011.189)
98. Hackett JA, Surani MA (2013) DNA methylation dynamics during the mammalian life cycle. *Philos Trans R Soc Lond B* 368:20110328. doi:[10.1098/rstb.2011.0328](https://doi.org/10.1098/rstb.2011.0328)
99. Pastor WA, Aravind L, Rao A (2013) TETonic shift: biological roles of TET proteins in DNA demethylation and transcription. *Nat Rev Mol Cell Biol* 14:341–356. doi:[10.1038/nrm3589](https://doi.org/10.1038/nrm3589)
100. Song C-X, Szulwach KE, Fu Y et al (2011) Selective chemical labeling reveals the genome-wide distribution of 5-hydroxymethylcytosine. *Nat Biotechnol* 29:68–72. doi:[10.1038/nbt.1732](https://doi.org/10.1038/nbt.1732)
101. Stroud H, Feng S, Morey Kinney S et al (2011) 5-Hydroxymethylcytosine is associated with enhancers and gene bodies in human embryonic stem cells. *Genome Biol* 12:R54. doi:[10.1186/gb-2011-12-6-r54](https://doi.org/10.1186/gb-2011-12-6-r54)
102. Thomson JP, Lempiäinen H, Hackett JA et al (2012) Non-genotoxic carcinogen exposure induces defined changes in the 5-hydroxymethylome. *Genome Biol* 13:R93. doi:[10.1186/gb-2012-13-10-r93](https://doi.org/10.1186/gb-2012-13-10-r93)
103. Lian CG, Xu Y, Ceol C et al (2012) Loss of 5-hydroxymethylcytosine is an epigenetic hallmark of melanoma. *Cell* 150:1135–1146. doi:[10.1016/j.cell.2012.07.033](https://doi.org/10.1016/j.cell.2012.07.033)
104. Nestor CE, Ottaviano R, Reddington J et al (2012) Tissue type is a major modifier of the 5-hydroxymethylcytosine content of human genes. *Genome Res* 22:467–477. doi:[10.1101/gr.126417.111](https://doi.org/10.1101/gr.126417.111)
105. Song C-X, Szulwach KE, Dai Q et al (2013) Genome-wide profiling of 5-formylcytosine reveals its roles in epigenetic priming. *Cell* 153:678–691. doi:[10.1016/j.cell.2013.04.001](https://doi.org/10.1016/j.cell.2013.04.001)
106. Shen L, Wu H, Diep D et al (2013) Genome-wide analysis reveals TET- and TDG-dependent 5-methylcytosine oxidation dynamics. *Cell* 153:692–706. doi:[10.1016/j.cell.2013.04.002](https://doi.org/10.1016/j.cell.2013.04.002)
107. Dawlaty MM, Ganz K, Powell BE et al (2011) Tet1 is dispensable for maintaining pluripotency and its loss is compatible with embryonic and postnatal development. *Cell Stem Cell* 9:166–175. doi:[10.1016/j.stem.2011.07.010](https://doi.org/10.1016/j.stem.2011.07.010)
108. Ko M, Bandukwala HS, An J et al (2011) Ten-eleven-translocation 2 (TET2) negatively regulates homeostasis and differentiation of hematopoietic stem cells in mice. *Proc Natl Acad Sci USA* 108:14566–14571. doi:[10.1073/pnas.1112317108](https://doi.org/10.1073/pnas.1112317108)
109. Li Z, Cai X, Cai C-L et al (2011) Deletion of Tet2 in mice leads to dysregulated hematopoietic stem cells and subsequent development of myeloid malignancies. *Blood* 118:4509–4518. doi:[10.1182/blood-2010-12-325241](https://doi.org/10.1182/blood-2010-12-325241)
110. Yamaguchi S, Hong K, Liu R et al (2012) Tet1 controls meiosis by regulating meiotic gene expression. *Nature* 492:443–447. doi:[10.1038/nature11709](https://doi.org/10.1038/nature11709)
111. Gu T-P, Guo F, Yang H et al (2011) The role of Tet3 DNA dioxygenase in epigenetic reprogramming by oocytes. *Nature* 477:606–610. doi:[10.1038/nature10443](https://doi.org/10.1038/nature10443)
112. Dawlaty MM, Breiling A, Le T et al (2013) Combined deficiency of Tet1 and Tet2 causes epigenetic abnormalities but is compatible with postnatal development. *Dev Cell* 24:310–323. doi:[10.1016/j.devcel.2012.12.015](https://doi.org/10.1016/j.devcel.2012.12.015)
113. Hackett JA, Sengupta R, Zyllic JJ et al (2013) Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. *Science* 339:448–452. doi:[10.1126/science.1229277](https://doi.org/10.1126/science.1229277)
114. Yoder JA, Walsh CP, Bestor TH (1997) Cytosine methylation and the ecology of intragenomic parasites. *Trends Genet* 13:335–340. doi:[10.1016/S0168-9525\(97\)01181-5](https://doi.org/10.1016/S0168-9525(97)01181-5)
115. Wiench M, John S, Baek S et al (2011) DNA methylation status predicts cell type-specific enhancer activity. *EMBO J* 30:3028–3039. doi:[10.1038/emboj.2011.210](https://doi.org/10.1038/emboj.2011.210)
116. Hendrich B, Bird A (1998) Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Mol Cell Biol* 18:6538–6547
117. Lewis JD, Meehan RR, Henzel WJ et al (1992) Purification, sequence, and cellular localization of a novel chromosomal protein that binds to methylated DNA. *Cell* 69:905–914

118. Meehan RR, Lewis JD, Bird AP (1992) Characterization of MeCP2, a vertebrate DNA binding protein with affinity for methylated DNA. *Nucleic Acids Res* 20:5085–5092
119. Prokhorchouk A, Hendrich B, Jørgensen H et al (2001) The p120 catenin partner Kaiso is a DNA methylation-dependent transcriptional repressor. *Genes Dev* 15:1613–1618. doi:10.1101/gad.198501
120. Nan X, Ng HH, Johnson CA et al (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393:386–389. doi:10.1038/30764
121. Martín Caballero I, Hansen J, Leaford D et al (2009) The methyl-CpG binding proteins Mecp2, Mbd2 and Kaiso are dispensable for mouse embryogenesis, but play a redundant function in neural differentiation. *PLoS ONE* 4:e4315. doi:10.1371/journal.pone.0004315
122. Brunmeir R, Lagger S, Simboeck E et al (2010) Epigenetic regulation of a murine retrotransposon by a dual histone modification mark. *PLoS Genet* 6:e1000927. doi:10.1371/journal.pgen.1000927
123. Reddington JP, Perricone SM, Nestor CE et al (2013) Redistribution of H3K27me3 upon DNA hypomethylation results in de-repression of Polycomb-target genes. *Genome Biol* 14:R25. doi:10.1186/gb-2013-14-3-r25
124. Leeb M, Pasini D, Novatchkova M et al (2010) Polycomb complexes act redundantly to repress genomic repeats and genes. *Genes Dev* 24:265–276. doi:10.1101/gad.544410
125. Reichmann J, Crichton JH, Madej MJ et al (2012) Microarray analysis of LTR retrotransposon silencing identifies Hdac1 as a regulator of retrotransposon expression in mouse embryonic stem cells. *PLoS Comput Biol* 8:e1002486. doi:10.1371/journal.pcbi.1002486
126. Popp C, Dean W, Feng S et al (2010) Genome-wide erasure of DNA methylation in mouse primordial germ cells is affected by AID deficiency. *Nature* 463:1101–1105. doi:10.1038/nature08829
127. Skene PJ, Illingworth RS, Webb S et al (2010) Neuronal MeCP2 is expressed at near histone-octamer levels and globally alters the chromatin state. *Mol Cell* 37:457–468. doi:10.1016/j.molcel.2010.01.030
128. Hutnick LK, Huang X, Loo T-C et al (2010) Repression of retrotransposal elements in mouse embryonic stem cells is primarily mediated by a DNA methylation-independent mechanism. *J Biol Chem* 285:21082–21091. doi:10.1074/jbc.M110.125674
129. Karimi MM, Goyal P, Maksakova IA et al (2011) DNA methylation and SETDB1/H3K9me3 regulate predominantly distinct sets of genes, retroelements, and chimeric transcripts in mESCs. *Cell Stem Cell* 8:676–687. doi:10.1016/j.stem.2011.04.004
130. Tsumura A, Hayakawa T, Kumaki Y et al (2006) Maintenance of self-renewal ability of mouse embryonic stem cells in the absence of DNA methyltransferases Dnmt1, Dnmt3a and Dnmt3b. *Genes Cells* 11:805–814. doi:10.1111/j.1365-2443.2006.00984.x
131. Day DS, Luquette LJ, Park PJ, Kharchenko PV (2010) Estimating enrichment of repetitive elements from high-throughput sequence data. *Genome Biol* 11:R69. doi:10.1186/gb-2010-11-6-r69
132. Matsui T, Leung D, Miyashita H et al (2010) Proviral silencing in embryonic stem cells requires the histone methyltransferase ESET. *Nature* 464:927–931. doi:10.1038/nature08858
133. Macfarlan TS, Gifford WD, Agarwal S et al (2011) Endogenous retroviruses and neighboring genes are coordinately repressed by LSD1/KDM1A. *Genes Dev* 25:594–607. doi:10.1101/gad.2008511
134. Guallar D, Pérez-Palacios R, Climent M et al (2012) Expression of endogenous retroviruses is negatively regulated by the pluripotency marker Rex1/Zfp42. *Nucleic Acids Res* 40:8993–9007. doi:10.1093/nar/gks686
135. Hisada K, Sánchez C, Endo TA et al (2012) RYBP represses endogenous retroviruses and preimplantation- and germ line-specific genes in mouse embryonic stem cells. *Mol Cell Biol* 32:1139–1149. doi:10.1128/MCB.06441-11
136. Rowe HM, Jakobsson J, Mesnard D et al (2010) KAP1 controls endogenous retroviruses in embryonic stem cells. *Nature* 463:237–240. doi:10.1038/nature08674
137. Wolf D, Goff SP (2007) TRIM28 mediates primer binding site-targeted silencing of murine leukemia virus in embryonic cells. *Cell* 131:46–57. doi:10.1016/j.cell.2007.07.026
138. Sripathy SP, Stevens J, Schultz DC (2006) The KAP1 corepressor functions to coordinate the assembly of de novo HP1-demarcated microenvironments of heterochromatin required for KRAB zinc finger protein-mediated transcriptional repression. *Mol Cell Biol* 26:8623–8638. doi:10.1128/MCB.00487-06
139. Wolf D, Goff SP (2009) Embryonic stem cells use ZFP809 to silence retroviral DNAs. *Nature* 458:1201–1204. doi:10.1038/nature07844
140. Rowe HM, Friedli M, Offner S et al (2013) De novo DNA methylation of endogenous retroviruses is shaped by KRAB-ZFPs/KAP1 and ESET. *Development* 140:519–529. doi:10.1242/dev.087585
141. Cantone I, Fisher AG (2013) Epigenetic programming and reprogramming during development. *Nat Struct Mol Biol* 20:282–289. doi:10.1038/nsmb.2489
142. Saitou M, Kagiwada S, Kurimoto K (2012) Epigenetic reprogramming in mouse pre-implantation development and primordial germ cells. *Development* 139:15–31. doi:10.1242/dev.050849
143. Abe M, Tsai SY, Jin S-G et al (2011) Sex-specific dynamics of global chromatin changes in fetal mouse germ cells. *PLoS ONE*. doi:10.1371/journal.pone.0023848
144. Tachibana M, Nozaki M, Takeda N, Shinkai Y (2007) Functional dynamics of H3K9 methylation during meiotic prophase progression. *EMBO J* 26:3346–3359. doi:10.1038/sj.emboj.7601767
145. Ng J-H, Kumar V, Muratani M et al (2013) In vivo epigenomic profiling of germ cells reveals germ cell molecular signatures. *Dev Cell* 24:324–333. doi:10.1016/j.devcel.2012.12.011
146. Seki Y, Hayashi K, Itoh K et al (2005) Extensive and orderly reprogramming of genome-wide chromatin modifications associated with specification and early development of germ cells in mice. *Dev Biol* 278:440–458. doi:10.1016/j.ydbio.2004.11.025
147. Guibert S, Forné T, Weber M (2012) Global profiling of DNA methylation erasure in mouse primordial germ cells. *Genome Res* 22:633–641. doi:10.1101/gr.130997.111
148. Lane N, Dean W, Erhardt S et al (2003) Resistance of IAPs to methylation reprogramming may provide a mechanism for epigenetic inheritance in the mouse. *Genesis* 35:88–93. doi:10.1002/gene.10168
149. Kagiwada S, Kurimoto K, Hirota T et al (2013) Replication-coupled passive DNA demethylation for the erasure of genome imprints in mice. *EMBO J* 32:340–353. doi:10.1038/emboj.2012.331
150. Hajkova P, Jeffries SJ, Lee C et al (2010) Genome-wide reprogramming in the mouse germ line entails the base excision repair pathway. *Science* 329:78–82. doi:10.1126/science.1187945
151. Kurimoto K, Yabuta Y, Ohinata Y et al (2008) Complex genome-wide transcription dynamics orchestrated by Blimp1 for the specification of the germ cell lineage in mice. *Genes Dev* 22:1617–1635. doi:10.1101/gad.1649908
152. Kato Y, Kaneda M, Hata K et al (2007) Role of the Dnmt3 family in de novo methylation of imprinted and repetitive sequences during male germ cell development in the mouse. *Hum Mol Genet* 16:2272–2280. doi:10.1093/hmg/ddm179
153. Lucifero D, Mann MRW, Bartolomei MS, Trasler JM (2004) Gene-specific timing and epigenetic memory in oocyte

- imprinting. *Hum Mol Genet* 13:839–849. doi:[10.1093/hmg/ddh104](https://doi.org/10.1093/hmg/ddh104)
154. Smallwood SA, Tomizawa S-I, Krueger F et al (2011) Dynamic CpG island methylation landscape in oocytes and preimplantation embryos. *Nat Genet* 43:811–814. doi:[10.1038/ng.864](https://doi.org/10.1038/ng.864)
  155. Kobayashi H, Sakurai T, Imai M et al (2012) Contribution of intragenic DNA methylation in mouse gametic DNA methylomes to establish oocyte-specific heritable marks. *PLoS Genet* 8:e1002440. doi:[10.1371/journal.pgen.1002440](https://doi.org/10.1371/journal.pgen.1002440)
  156. Chotalia M, Smallwood SA, Ruf N et al (2009) Transcription is required for establishment of germline methylation marks at imprinted genes. *Genes Dev* 23:105–117. doi:[10.1101/gad.495809](https://doi.org/10.1101/gad.495809)
  157. Santos F, Hendrich B, Reik W, Dean W (2002) Dynamic reprogramming of DNA methylation in the early mouse embryo. *Dev Biol* 241:172–182. doi:[10.1006/dbio.2001.0501](https://doi.org/10.1006/dbio.2001.0501)
  158. Smith ZD, Chan MM, Mikkelsen TS et al (2012) A unique regulatory phase of DNA methylation in the early mammalian embryo. *Nature* 484:339–344. doi:[10.1038/nature10960](https://doi.org/10.1038/nature10960)
  159. Kaneda M, Okano M, Hata K et al (2004) Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* 429:900–903. doi:[10.1038/nature02633](https://doi.org/10.1038/nature02633)
  160. Yaman R, Grandjean V (2006) Timing of entry of meiosis depends on a mark generated by DNA methyltransferase 3a in testis. *Mol Reprod Dev* 73:390–397. doi:[10.1002/mrd.20430](https://doi.org/10.1002/mrd.20430)
  161. Kaneda M, Hirasawa R, Chiba H et al (2010) Genetic evidence for Dnmt3a-dependent imprinting during oocyte growth obtained by conditional knockout with Zp3-Cre and complete exclusion of Dnmt3b by chimera formation. *Genes Cells*. doi:[10.1111/j.1365-2443.2009.01374.x](https://doi.org/10.1111/j.1365-2443.2009.01374.x)
  162. Suetake I, Shinozaki F, Miyagawa J et al (2004) DNMT3L stimulates the DNA methylation activity of Dnmt3a and Dnmt3b through a direct interaction. *J Biol Chem* 279:27816–27823. doi:[10.1074/jbc.M400181200](https://doi.org/10.1074/jbc.M400181200)
  163. Bourc'his D, Bestor TH (2004) Meiotic catastrophe and retrotransposon reactivation in male germ cells lacking Dnmt3L. *Nature* 431:96–99. doi:[10.1038/nature02886](https://doi.org/10.1038/nature02886)
  164. Bourc'his D, Xu GL, Lin CS et al (2001) Dnmt3L and the establishment of maternal genomic imprints. *Science* 294:2536–2539. doi:[10.1126/science.1065848](https://doi.org/10.1126/science.1065848)
  165. Hata K, Okano M, Lei H, Li E (2002) Dnmt3L cooperates with the Dnmt3 family of de novo DNA methyltransferases to establish maternal imprints in mice. *Development* 129:1983–1993
  166. Lucifero D, La Salle S, Bourc'his D et al (2007) Coordinate regulation of DNA methyltransferase expression during oogenesis. *BMC Dev Biol* 7:36. doi:[10.1186/1471-213X-7-36](https://doi.org/10.1186/1471-213X-7-36)
  167. Dennis K, Fan T, Geiman T et al (2001) Lsh, a member of the SNF2 family, is required for genome-wide methylation. *Genes Dev* 15:2940–2944. doi:[10.1101/gad.929101](https://doi.org/10.1101/gad.929101)
  168. Sun L-Q, Lee DW, Zhang Q et al (2004) Growth retardation and premature aging phenotypes in mice with disruption of the SNF2-like gene, PASG. *Genes Dev* 18:1035–1046. doi:[10.1101/gad.1176104](https://doi.org/10.1101/gad.1176104)
  169. Myant K, Stancheva I (2008) LSH cooperates with DNA methyltransferases to repress transcription. *Mol Cell Biol* 28:215–226. doi:[10.1128/MCB.01073-07](https://doi.org/10.1128/MCB.01073-07)
  170. Tao Y, Xi S, Shan J et al (2011) Lsh, chromatin remodeling family member, modulates genome-wide cytosine methylation patterns at nonrepeat sequences. *Proc Natl Acad Sci USA* 108:5626–5631. doi:[10.1073/pnas.1017000108](https://doi.org/10.1073/pnas.1017000108)
  171. Yan Q, Huang J, Fan T et al (2003) Lsh, a modulator of CpG methylation, is crucial for normal histone methylation. *EMBO J* 22:5154–5162. doi:[10.1093/emboj/cdg493](https://doi.org/10.1093/emboj/cdg493)
  172. De La Fuente R, Baumann C, Fan T et al (2006) Lsh is required for meiotic chromosome synapsis and retrotransposon silencing in female germ cells. *Nat Cell Biol* 8:1448–1454. doi:[10.1038/ncb1513](https://doi.org/10.1038/ncb1513)
  173. Hutvagner G, Simard MJ (2008) Argonaute proteins: key players in RNA silencing. *Nat Rev Mol Cell Biol* 9:22–32. doi:[10.1038/nrm2321](https://doi.org/10.1038/nrm2321)
  174. Aravin A, Gaidatzis D, Pfeffer S et al (2006) A novel class of small RNAs bind to MILI protein in mouse testes. *Nature* 442:203–207. doi:[10.1038/nature04916](https://doi.org/10.1038/nature04916)
  175. Aravin AA, Sachidanandam R, Girard A et al (2007) Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* 316:744–747. doi:[10.1126/science.1142612](https://doi.org/10.1126/science.1142612)
  176. Aravin AA, Sachidanandam R, Bourc'his D et al (2008) A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell* 31:785–799. doi:[10.1016/j.molcel.2008.09.003](https://doi.org/10.1016/j.molcel.2008.09.003)
  177. Carmell MA, Girard A, van de Kant HJG et al (2007) MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* 12:503–514. doi:[10.1016/j.devcel.2007.03.001](https://doi.org/10.1016/j.devcel.2007.03.001)
  178. Kuramochi-Miyagawa S, Kimura T, Yomogida K et al (2001) Two mouse piwi-related genes: miwi and mili. *Mech Dev* 108:121–133
  179. Xiol J, Cora E, Kogelgruber R et al (2012) A role for Fkbp6 and the chaperone machinery in piRNA amplification and transposon silencing. *Mol Cell* 47:970–979. doi:[10.1016/j.molcel.2012.07.019](https://doi.org/10.1016/j.molcel.2012.07.019)
  180. De Fazio S, Bartonicek N, Di Giacomo M et al (2011) The endonuclease activity of Mili fuels piRNA amplification that silences LINE1 elements. *Nature* 480:259–263. doi:[10.1038/nature10547](https://doi.org/10.1038/nature10547)
  181. Vourekas A, Zheng Q, Alexiou P et al (2012) Mili and Miwi target RNA repertoire reveals piRNA biogenesis and function of Miwi in spermiogenesis. *Nat Struct Mol Biol* 19:773–781. doi:[10.1038/nsmb.2347](https://doi.org/10.1038/nsmb.2347)
  182. Chen C, Jin J, James DA et al (2009) Mouse Piwi interactome identifies binding mechanism of Tdrkh Tudor domain to arginine methylated Miwi. *Proc Natl Acad Sci USA* 106:20336–20341. doi:[10.1073/pnas.0911640106](https://doi.org/10.1073/pnas.0911640106)
  183. Mathioudakis N, Palencia A, Kadlec J et al (2012) The multiple Tudor domain-containing protein TDRD1 is a molecular scaffold for mouse Piwi proteins and piRNA biogenesis factors. *RNA* 18:2056–2072. doi:[10.1261/ma.034181.112](https://doi.org/10.1261/ma.034181.112)
  184. Vagin VV, Wohlschlegel J, Qu J et al (2009) Proteomic analysis of murine Piwi proteins reveals a role for arginine methylation in specifying interaction with Tudor family members. *Genes Dev* 23:1749–1762. doi:[10.1101/gad.1814809](https://doi.org/10.1101/gad.1814809)
  185. Chuma S, Hiyoshi M, Yamamoto A et al (2003) Mouse tudor repeat-1 (MTR-1) is a novel component of chromatoid bodies/nuages in male germ cells and forms a complex with snRNPs. *Mech Dev* 120:979–990
  186. Shoji M, Tanaka T, Hosokawa M et al (2009) The TDRD9-MIWI2 complex is essential for piRNA-mediated retrotransposon silencing in the mouse male germline. *Dev Cell* 17:775–787. doi:[10.1016/j.devcel.2009.10.012](https://doi.org/10.1016/j.devcel.2009.10.012)
  187. Yabuta Y, Ohta H, Abe T et al (2011) TDRD5 is required for retrotransposon silencing, chromatoid body assembly, and spermiogenesis in mice. *J Cell Biol* 192:781–795. doi:[10.1083/jcb.201009043](https://doi.org/10.1083/jcb.201009043)
  188. Aravin AA, van der Heijden GW, Castañeda J et al (2009) Cytoplasmic compartmentalization of the fetal piRNA pathway in mice. *PLoS Genet* 5:e1000764. doi:[10.1371/journal.pgen.1000764](https://doi.org/10.1371/journal.pgen.1000764)
  189. Reuter M, Chuma S, Tanaka T et al (2009) Loss of the Mili-interacting Tudor domain-containing protein-1 activates transposons and alters the Mili-associated small RNA profile. *Nat Struct Mol Biol* 16:639–646. doi:[10.1038/nsmb.1615](https://doi.org/10.1038/nsmb.1615)
  190. Ma L, Buchold GM, Greenbaum MP et al (2009) GASZ is essential for male meiosis and suppression of retrotransposon

- expression in the male germline. *PLoS Genet* 5:e1000635. doi:10.1371/journal.pgen.1000635
191. Soper SFC, van der Heijden GW, Hardiman TC et al (2008) Mouse maelstrom, a component of nuage, is essential for spermatogenesis and transposon repression in meiosis. *Dev Cell* 15:285–297. doi:10.1016/j.devcel.2008.05.015
  192. Watanabe T, Chuma S, Yamamoto Y et al (2011) MITOPLD is a mitochondrial protein essential for nuage formation and piRNA biogenesis in the mouse germline. *Dev Cell* 20:364–375. doi:10.1016/j.devcel.2011.01.005
  193. Zheng K, Xioli J, Reuter M et al (2010) Mouse MOV10L1 associates with Piwi proteins and is an essential component of the Piwi-interacting RNA (piRNA) pathway. *Proc Natl Acad Sci USA* 107:11841–11846. doi:10.1073/pnas.1003953107
  194. Kuramochi-Miyagawa S, Watanabe T, Gotoh K et al (2010) MVH in piRNA processing and gene silencing of retrotransposons. *Genes Dev* 24:887–892. doi:10.1101/gad.1902110
  195. Deng W, Lin H (2002) miwi, a murine homolog of piwi, encodes a cytoplasmic protein essential for spermatogenesis. *Dev Cell* 2:819–830
  196. Kuramochi-Miyagawa S, Kimura T, Ijiri TW et al (2004) Mili, a mammalian member of piwi family gene, is essential for spermatogenesis. *Development* 131:839–849. doi:10.1242/dev.00973
  197. Watanabe T, Totoki Y, Toyoda A et al (2008) Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 453:539–543. doi:10.1038/nature06908
  198. Unhavaithaya Y, Hao Y, Beyret E et al (2009) MILI, a PIWI-interacting RNA-binding protein, is required for germ line stem cell self-renewal and appears to positively regulate translation. *J Biol Chem* 284:6507–6519. doi:10.1074/jbc.M809104200
  199. Reuter M, Berminger P, Chuma S et al (2011) Miwi catalysis is required for piRNA amplification-independent LINE1 transposon silencing. *Nature* 480:264–267. doi:10.1038/nature10672
  200. Beyret E, Liu N, Lin H (2012) piRNA biogenesis during adult spermatogenesis in mice is independent of the ping-pong mechanism. *Cell Res* 22:1429–1439. doi:10.1038/cr.2012.120
  201. Zheng K, Wang PJ (2012) Blockade of pachytene piRNA biogenesis reveals a novel requirement for maintaining post-meiotic germline genome integrity. *PLoS Genet* 8:e1003038. doi:10.1371/journal.pgen.1003038
  202. Tam OH, Aravin AA, Stein P et al (2008) Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 453:534–538. doi:10.1038/nature06904
  203. Watanabe T, Takeda A, Tsukiyama T et al (2006) Identification and characterization of two novel classes of small RNAs in the mouse germline: retrotransposon-derived siRNAs in oocytes and germline small RNAs in testes. *Genes Dev* 20:1732–1743. doi:10.1101/gad.1425706
  204. Filipowicz W, Bhattacharyya SN, Sonenberg N (2008) Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 9:102–114. doi:10.1038/nrg2290
  205. Su Y-Q, Sugiura K, Sun F et al (2012) MARF1 regulates essential oogenic processes in mice. *Science* 335:1496–1499. doi:10.1126/science.1214680
  206. Tanaka T, Hosokawa M, Vagin VV et al (2011) Tudor domain containing 7 (Tdrd7) is essential for dynamic ribonucleoprotein (RNP) remodeling of chromatoid bodies during spermatogenesis. *Proc Natl Acad Sci USA*. doi:10.1073/pnas.1015447108
  207. Grivna ST, Pyhtila B, Lin H (2006) MIWI associates with translational machinery and PIWI-interacting RNAs (piRNAs) in regulating spermatogenesis. *Proc Natl Acad Sci USA* 103:13415–13420. doi:10.1073/pnas.0605506103
  208. Xu M, You Y, Hunsicker P et al (2008) Mice deficient for a small cluster of piwi-Interacting RNAs implicate piwi-interacting RNAs in transposon control. *Biol Reprod* 79:51–57. doi:10.1095/biolreprod.108.068072
  209. Goodier JL, Cheung LE, Kazazian HH Jr (2012) MOV10 RNA helicase is a potent inhibitor of retrotransposition in cells. *PLoS Genet* 8:e1002941. doi:10.1371/journal.pgen.1002941
  210. Lu C, Luo Z, Jäger S et al (2012) Moloney leukemia virus type 10 inhibits reverse transcription and retrotransposition of intracisternal particles. *J Virol* 86:10517–10523. doi:10.1128/JVI.00868-12
  211. Burdick R, Smith JL, Chaipan C et al (2010) P body-associated protein Mov10 inhibits HIV-1 replication at multiple stages. *J Virol* 84:10241–10253. doi:10.1128/JVI.00585-10
  212. Frost RJA, Hamra FK, Richardson JA et al (2010) MOV10L1 is necessary for protection of spermatocytes against retrotransposons by Piwi-interacting RNAs. *Proc Natl Acad Sci USA* 107:11847–11852. doi:10.1073/pnas.1007158107
  213. Öllinger R, Childs AJ, Burgess HM et al (2008) Deletion of the pluripotency-associated Tex19.1 gene causes activation of endogenous retroviruses and defective spermatogenesis in mice. *PLoS Genet* 4:e1000199. doi:10.1371/journal.pgen.1000199
  214. Borgel J, Guibert S, Li Y et al (2010) Targets and dynamics of promoter DNA methylation during early mouse development. *Nat Genet* 42:1093–1100. doi:10.1038/ng.708
  215. Velasco G, Hubé F, Rollin J et al (2010) Dnmt3b recruitment through E2F6 transcriptional repressor mediates germ-line gene silencing in murine somatic tissues. *Proc Natl Acad Sci USA* 107:9281–9286. doi:10.1073/pnas.1000473107
  216. Maatouk DM, Kellam LD, Mann MRW et al (2006) DNA methylation is a primary mechanism for silencing postmigratory primordial germ cell genes in both germ cell and somatic cell lineages. *Development* 133:3411–3418. doi:10.1242/dev.02500
  217. Cooke HJ, Lee M, Kerr S, Ruggiu M (1996) A murine homologue of the human DAZ gene is autosomal and expressed only in male and female gonads. *Hum Mol Genet* 5:513–516
  218. Kuntz S, Kieffer E, Bianchetti L et al (2008) Tex19, a mammalian-specific protein with a restricted expression in pluripotent stem cells and germ line. *Stem Cells* 26:734–744. doi:10.1634/stemcells.2007-0772
  219. Yan W, Rajkovic A, Viveiros MM et al (2002) Identification of Gasz, an evolutionarily conserved gene expressed exclusively in germ cells and encoding a protein with four ankyrin repeats, a sterile-alpha motif, and a basic leucine zipper. *Mol Endocrinol* 16:1168–1184
  220. Schütt S, Florl AR, Shi W et al (2003) DNA methylation in placentas of interspecies mouse hybrids. *Genetics* 165:223–228
  221. Sproul D, Nestor C, Culley J et al (2011) Transcriptionally repressed genes become aberrantly methylated and distinguish tumors of different lineages in breast cancer. *Proc Natl Acad Sci USA* 108:4364–4369. doi:10.1073/pnas.1013224108
  222. Ting DT, Lipson D, Paul S et al (2011) Aberrant overexpression of satellite repeats in pancreatic and other epithelial cancers. *Science*. doi:10.1126/science.1200801
  223. La Salle S, Oakes CC, Neaga OR et al (2007) Loss of spermatogonia and wide-spread DNA methylation defects in newborn male mice deficient in DNMT3L. *BMC Dev Biol* 7:104. doi:10.1186/1471-213X-7-104
  224. Mahadevaiah SK, Bourc'his D, de Rooij DG et al (2008) Extensive meiotic asynapsis in mice antagonises meiotic silencing of unsynapsed chromatin and consequently disrupts meiotic sex chromosome inactivation. *J Cell Biol* 182:263–276. doi:10.1083/jcb.200710195
  225. Beyret E, Lin H (2011) Pinpointing the expression of piRNAs and function of the PIWI protein subfamily during spermatogenesis in the mouse. *Dev Biol* 355:215–226. doi:10.1016/j.ydbio.2011.04.021



226. Tanaka SS, Toyooka Y, Akasu R et al (2000) The mouse homolog of *Drosophila* Vasa is required for the development of male germ cells. *Genes Dev* 14:841–853
227. Chen J, Melton C, Suh N et al (2011) Genome-wide analysis of translation reveals a critical role for deleted in azoospermia-like (Dazl) at the oocyte-to-zygote transition. *Genes Dev* 25:755–766. doi:[10.1101/gad.2028911](https://doi.org/10.1101/gad.2028911)
228. Reynolds N, Collier B, Maratou K et al (2005) Dazl binds in vivo to specific transcripts and can regulate the pre-meiotic translation of Mvh in germ cells. *Hum Mol Genet* 14:3899–3909. doi:[10.1093/hmg/ddi414](https://doi.org/10.1093/hmg/ddi414)
229. Ruggiu M, Speed R, Taggart M et al (1997) The mouse Dazla gene encodes a cytoplasmic protein essential for gametogenesis. *Nature* 389:73–77. doi:[10.1038/37987](https://doi.org/10.1038/37987)
230. Schrans-Stassen BH, Saunders PT, Cooke HJ, de Rooij DG (2001) Nature of the spermatogenic arrest in *Dazl*<sup>-/-</sup> mice. *Biol Reprod* 65:771–776
231. Yang F, Cheng Y, An JY et al (2010) The ubiquitin ligase Ubr2, a recognition E3 component of the N-end rule pathway, stabilizes Tex19.1 during spermatogenesis. *PLoS ONE* 5:e14017. doi:[10.1371/journal.pone.0014017](https://doi.org/10.1371/journal.pone.0014017)
232. Inagaki A, Schoenmakers S, Baarends WM (2010) DNA double strand break repair, chromosome synapsis and transcriptional silencing in meiosis. *Epigenetics* 5:255–266
233. Cai X, Li J, Yang Q, Shi Q (2011) Gamma-irradiation increased meiotic crossovers in mouse spermatocytes. *Mutagenesis* 26:721–727. doi:[10.1093/mutage/ger038](https://doi.org/10.1093/mutage/ger038)
234. Lange J, Pan J, Cole F et al (2011) ATM controls meiotic double-strand-break formation. *Nature* 479:237–240. doi:[10.1038/nature10508](https://doi.org/10.1038/nature10508)
235. Denne M, Sauter M, Armbruster V et al (2007) Physical and functional interactions of human endogenous retrovirus proteins Np9 and rec with the promyelocytic leukemia zinc finger protein. *J Virol* 81:5607–5616. doi:[10.1128/JVI.02771-06](https://doi.org/10.1128/JVI.02771-06)
236. Galli UM, Sauter M, Lecher B et al (2005) Human endogenous retrovirus rec interferes with germ cell development in mice and may cause carcinoma in situ, the predecessor lesion of germ cell tumors. *Oncogene* 24:3223–3228. doi:[10.1038/sj.onc.1208543](https://doi.org/10.1038/sj.onc.1208543)
237. Kaufmann S, Sauter M, Schmitt M et al (2010) Human endogenous retrovirus protein Rec interacts with the testicular zinc-finger protein and androgen receptor. *J Gen Virol* 91:1494–1502. doi:[10.1099/vir.0.014241-0](https://doi.org/10.1099/vir.0.014241-0)
238. Brennecke J, Malone CD, Aravin AA et al (2008) An epigenetic role for maternally inherited piRNAs in transposon silencing. *Science* 322:1387–1392. doi:[10.1126/science.1165171](https://doi.org/10.1126/science.1165171)
239. Heyn H, Ferreira HJ, Bassas L et al (2012) Epigenetic disruption of the PIWI pathway in human spermatogenic disorders. *PLoS ONE* 7:e47892. doi:[10.1371/journal.pone.0047892](https://doi.org/10.1371/journal.pone.0047892)
240. Ward MC, Wilson MD, Barbosa-Morais NL et al (2013) Latent regulatory potential of human-specific repetitive elements. *Mol Cell* 49:262–272. doi:[10.1016/j.molcel.2012.11.013](https://doi.org/10.1016/j.molcel.2012.11.013)
241. Ribet D, Harper F, Dupressoir A et al (2008) An infectious progenitor for the murine IAP retrotransposon: emergence of an intracellular genetic parasite from an ancient retrovirus. *Genome Res* 18:597–609. doi:[10.1101/gr.073486.107](https://doi.org/10.1101/gr.073486.107)