



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Implementation of an ED-based bundled mechanical ventilation protocol improves adherence to lung-protective ventilation☆

Tyler M. Foley^{a,*}, Brittany A. Philpot, MD^a, Alysa S. Davis, DO^a, Morgan B. Swanson^a,
Karisa K. Harland, MPH, PhD^a, Justin D. Kuhn, RRT-NPS/ACCS^b,
Brian M. Fuller, MD, MSCI^c, Nicholas M. Mohr, MD, MS^{a,d}

^a Department of Emergency Medicine, University of Iowa Carver College of Medicine, Iowa City, IA, United States of America

^b Department of Respiratory Care, University of Iowa Carver College of Medicine, Iowa City, IA, United States of America

^c Division of Emergency Medicine, Department of Anesthesiology, Division of Critical Care, Washington University School of Medicine in St. Louis, St. Louis, MO, United States of America

^d Division of Critical Care, Department of Anesthesia, University of Iowa Carver College of Medicine, Iowa City, IA, United States of America

ARTICLE INFO

Article history:

Received 2 December 2019

Received in revised form 11 February 2020

Accepted 25 February 2020

Keywords:

Ventilator-associated lung injury

Acute respiratory distress syndrome

Lung-protective ventilation

Tidal volume

1. Introduction

Mechanical ventilation is a common life-saving protocol, but it has also been linked to ventilator-associated lung injury (VALI), a progressive injury caused by positive pressure ventilation [1]. Despite decades of research, lung-protective ventilation (LPV) with low tidal volume (V_T) remains one of the few effective protocols for patients with acute respiratory distress syndrome (ARDS) [2–5]. Growing evidence suggests that low V_T may also benefit critically ill patients without pre-existing lung injury [6–11]. In addition to low V_T , adequate positive-end expiratory pressure (PEEP) and early titration of oxygen concentration (FiO_2) are important components of LPV that minimize atelectasis [12,13] and oxygen toxicity [14–16], respectively.

The emergency department (ED) is a crucial setting for the prevention and treatment of VALI. ED overcrowding, boarding, and increased ICU admissions have tripled the amount of critical care and prolonged mechanical ventilation delivered in US EDs [17,18]. Of the estimated

240,000 patients mechanically ventilated in US EDs each year, one-quarter are ventilated for more than 5 h [19,20]. These initial hours of care are influential in the outcome of critically ill patients, as VALI has been shown to occur within minutes to hours of initiating mechanical ventilation [21–23] and progression to ARDS occurs early during ventilation of at-risk patients [7]. For these reasons, initial ventilator settings provided in the immediate post-intubation period can be critically important in determining patient outcomes. Observational data show large V_T is commonly used in the ED, ED V_T influences ventilation strategy in the ICU, and injurious early ventilation strategies are associated with worse clinical outcomes, implicating the ED as a vital link in the provision of LPV [24,25].

Standardization of mechanical ventilation through order sets and protocols provides an evidence-based and cost-effective opportunity to reduce variability in the care of ventilated ED patients [26]. Respiratory care protocols administered by non-physician staff have been shown to improve arterial blood gas sampling [27,28], early ventilator weaning, [29,30] and adoption of PEEP- FiO_2 combination guidelines [31]. Successful adherence to a V_T -focused ventilator order set further supports the feasibility of protocol-driven ventilation in the ED [32]. Implementation of a similar mechanical ventilation protocol in an academic ED increased ventilator-free days and hospital-free days while being associated with significantly decreased hospital mortality [33]. Based on the association between lung protective ventilation and improved clinical outcomes demonstrated in the LOV-ED study and the previously demonstrated prevalence of non-protective ventilation in multiple EDs, an RT-driven LPV protocol was implemented in the ED of a large, academic medical center to standardize care of mechanically ventilated ED patients. We hypothesized that implementation of this protocol would improve adherence to LPV ventilation parameters and improve clinical outcomes of patients intubated and mechanically ventilated in the ED.

2. Methods

2.1. Study design

This retrospective before-after observational cohort study was conducted at an academic 60,000-visit ED between March 2016

☆ This research was presented at the Society of Academic Emergency Medicine Great Plains Regional Conference, September 21, 2018, in St. Louis, Missouri.


* Corresponding author at: University of Iowa Carver College of Medicine, Department of Emergency Medicine, 1008 RCP, 200 Hawkins Dr, Iowa City, IA 52242 United States of America.
E-mail address: tyler-foley@uiowa.edu (T.M. Foley).

and July 2018, 15 months before through 13 months after the implementation of a lung-protective mechanical ventilation protocol. Inclusion criteria were adult patients (age ≥ 18 years) receiving mechanical ventilation following intubation while in the ED. Exclusion criteria included death or extubation while in the ED, missing height or ED V_T in the electronic medical record (EMR), non-invasive ventilation only, and transfer from the ED to another facility. This study is reported in accordance with the Standards for Quality Improvement Reporting Excellence (SQUIRE) Statement: Guidelines for Reporting Observational Studies [34] and was approved by the local Institutional Review Board under a waiver of informed consent.

2.2. Protocol

In June 2017, a new mechanical ventilation protocol was implemented in the ED to mimic the Lung Protective Ventilation Initiated in the Emergency Department (LOV-ED) study protocol [33]. The protocol was divided into five components, each intended to prevent a potential mechanism of VALI (Fig. 1). Mode selection of choice for this protocol was assist control-volume control (AC-VC).

Education for respiratory therapists included hands-on sessions with the mechanical ventilator, computer-based tutorial, and quiz. Tools were provided to respiratory therapists including laminated protocol sheets detailing ideal predicted body weight (PBW)-V_T charts



PEEP-FiO₂ Table

FiO₂	30	40	40	50	50	60	70	70	70	80	90	90	90	100
PEEP	5	5	8	8	10	10	10	12	14	14	14	16	18	20-24

Fig. 1. Study protocol diagram.

and tape measures to measure patient height. These tools were attached to the ED ventilators as part of standard protocol procedures. After implementation, weekly audits of all ventilated ED patients were performed to review adherence to the protocol, immediate feedback was provided, and adherence was reported at regular staff meetings.

2.3. Procedures

Baseline patient characteristics were abstracted from the EMR. Sequential organ failure assessment (SOFA) scores were calculated at the time of ED admission and 24 h after admission [35]. The first ventilator settings recorded in the ED and ICU were abstracted from the medical record. Names of the treating respiratory therapist, intubating physician (typically a resident), and attending physician were also recorded. Data were abstracted by trained data abstractors with regular meetings and monitoring of data collection. All variables were collected in a standardized format using a standard case report form.

2.4. Outcomes

The primary outcome of this study was mean difference in V_T (mL/kg) administered in the ED between groups. Difference in V_T was selected as the primary outcome because it is the most widely studied and supported mediator of VALI and would reflect a change in clinical practice induced by the protocol. Secondary outcomes included 24-h change in SOFA score, ventilator-free days, ICU V_T , protocol adherence, mortality, ARDS, hospital-free days, and ventilator-associated pneumonia (VAP). We also tested adherence with individual protocol elements.

2.5. Definitions

Patient height and weight from the hospital encounter were used to calculate BMI and PBW as shown previously [36]. LPV was defined as the use of $V_T \leq 8$ mL/kg PBW.

SOFA scores were calculated as described previously [35]. Saturation by pulse oximetry (SpO_2)/ FiO_2 ratios were used to calculate the SOFA score because of ED arterial blood gas infrequency. SpO_2 / FiO_2 ratios approximated partial pressure of oxygen (PaO_2)/ FiO_2 ratios using methods validated in a prior report [37]. Due to the shape of the oxyhemoglobin dissociation curve, any SpO_2 values above 97% was assigned a zero for the respiratory component of the SOFA score because large PaO_2 changes can occur with small changes in the SpO_2 near 100% [38].

Adherence to each facet of the ventilator protocol was determined according to Fig. 1. If the patient received a tidal volume ≤ 8 mL/kg, they were considered adherent to the “Volutrauma Prevention” facet of the protocol. PEEP was to be ≥ 5 cm H₂O to satisfy “Atelectrauma Prevention.” For “Hyperoxia Prevention,” FiO_2 must have been set to between 0.3 and 0.4 immediately within 15 min of intubation. If a higher FiO_2 was initially selected, the corresponding PEEP value must have been in accordance with the provided PEEP- FiO_2 table (Fig. 1) to be considered adherent. Respiratory rate was to be set between 16 and 30 breaths per minute. Head-of-bed elevation was not included in adherence determinations due to lack of notation in the medical record (98.2% missing). For a patient to be considered adherent to the full protocol, each facet must have been satisfied. If any data value was not adherent, that patient was considered not adherent to the protocol.

ARDS was defined according to the Berlin definition through hospital day five [10,24,39,40]. A panel of three research team clinicians reviewed blinded chest radiographs independently and classified them as “consistent”, “inconsistent”, or “equivocal” for ARDS, and the diagnosis was made by consensus. Each member of the panel reviewed a set of training radiographs [41] prior to study participation and was blinded to protocol group and all other clinical data during the chest radiograph review. Patients were assumed to have clinical evidence of left atrial hypertension if respiratory failure was attributed to congestive

heart failure or dialysis-dependent end-stage renal disease in the EMR, and thus were categorized as “not ARDS.”

VAP was defined as clinical suspicion for pneumonia by a board-certified intensivist with initiation of antibiotic treatment for a lower respiratory tract infection >48 h after initiation of mechanical ventilation, among patients not already being treated for pneumonia [42].

2.6. Sample size

A mean detectable difference for ED V_T of 0.33 mL/kg (SD 1.28) for pre- versus post-protocol (power 0.8, alpha 0.05) required 476 patients (238 pre-protocol patients and 238 post-protocol patients). A mean difference of 0.33 mL/kg was chosen due to the low mean V_T observed in a sample of pre-protocol patients performed prior to final data analysis. Given the already low pre-protocol tidal volume, the likelihood of demonstrating a smaller difference in tidal volume after protocol would require a greater number of patients than would be feasible for the study, but one could not expect a different that would bring the mean tidal volume to <6 mL/kg. We determined that a 0.33 mL/kg change was a compromise value that would detect a meaningful change in clinical practice.

2.7. Analysis

Chi-square tests, Wilcoxon Mann-Whitney tests, and Fisher's exact tests compared distributions of baseline characteristics between the two groups. A scatterplot was used to visualize associations between ED and ICU V_T .

Univariable and multivariable regression assessed the impact of the protocol on clinical and ventilator outcomes. For the primary outcome, a log-transformation of outcome was used, since the error terms on the continuous outcomes followed a right-skewed distribution. Linear, logistic, and negative binomial regression were used to estimate secondary outcomes as appropriate.

Covariate selection was guided by clinical knowledge and previous literature [33], and, stepwise backwards selection guided by Akaike Information Criterion (AIC) was used to select the final model. Respiratory therapist and intubator were also included as fixed effects to account for variance, observed and unobserved, associated with each fixed effect. A post hoc test for effect modification of protocol effect by subject height was conducted by testing an interaction term and then stratifying the analysis by protocol time (pre- vs. post-protocol).

Differences in the time to ARDS and VAP (in days) between the protocol groups were assessed using log-rank tests and univariable Cox proportional hazards regression models. Finally, variation in ED LPV rates ($V_T \leq 8$ mL/kg) by intubator and respiratory therapist were estimated with risk-adjustment (adjusting for: sex, BMI, and initial SOFA score) and reliability-adjustment [43]. Predictive capability of attending physician, intubating physician, and respiratory therapist for ED LPV were separately estimated using univariable logistic regression models to estimate the area under the curve (AUC), using bootstrapping of 1000 independent data samples to estimate precision of the AUC confidence interval. Data analysis was completed in R (version 3.6; RStudio Inc., Vienna, Austria) and SAS (version 9.4; SAS Institute, Inc., Cary, NC), and figures were created in Prism (version 8.2, GraphPad, Inc., San Diego, CA).

2.8. Sensitivity analysis

A post hoc sensitivity analysis was conducted restricting the study population to resemble the eligibility criteria of a previous before-after study of LPV protocols in the ED [33]: excluding subjects that had ARDS at hospital admission or were extubated or died within 24 h.

Table 1
Subject demographics and ventilator settings pre/post.

	Total		Pre-intervention		Post-intervention		p-Value
	N		n		n		
Total	500		272	54.4%	228	45.6%	
Age (yrs.)							
Median (IQR)	58.0	(43.0–71.0)	57.5	(41.5–70.0)	45.0	(45.0–71.0)	0.378
Sex							0.769
Male	293	58.6%	161	59.2%	132	57.9%	
Female	207	41.4%	111	40.8%	96	42.1%	
Height (cm)	172.7	(165.1–180.3)	172.7	(165.1–180.3)	172.7	(165.1–180.2)	0.664
Weight (kg)	82.3	(68.0–97.1)	84.9	(69.9–97.9)	80.6	(66.4–95.2)	0.271
BMI	27.4	(23.7–32.6)	28.0	(23.7–33.1)	27.1	(23.7–32.2)	0.242
Reason for intubation							0.295
Asthma	4	0.8%	3	1.1%	1	0.4%	
Chronic obstructive pulmonary disease	5	1.0%	5	1.8%	0	0.0%	
Chronic heart failure/pulmonary edema	20	4.0%	9	3.3%	11	4.8%	
Sepsis	71	14.2%	42	15.4%	29	12.7%	
Trauma	86	17.2%	51	18.8%	35	15.4%	
Cardiac arrest	15	3.0%	9	3.3%	6	2.6%	
Alcohol/drug overdose	76	15.2%	41	15.1%	35	15.4%	
Neurological	119	23.8%	49	18.0%	55	24.1%	
Other	104	20.8%	63	23.2%	56	24.6%	
SOFA score, median (IQR)							
Initial SOFA	2	(0–4)	0	(0–4)	2	(0–4)	0.419
24-hour SOFA	4	(2–7)	4	(2–6)	4	(2–7)	0.185
ED length of ventilation (min), median (IQR)	106.0	66.0–161.5	101.5	66.5–159.5	107.5	65.5–163.0	0.317
ED ventilator variables							
Tidal volume (mL), median (IQR)	425	(380–450)	440	(400–462.5)	405	(370–450)	<0.001
Tidal volume (ml/kg IBW), median (IQR)	6.4	(5.8–7.1)	6.5	(5.8–7.4)	6.2	(5.8–6.7)	<0.001
PEEP, cmH ₂ O, median (IQR)	5	(5–5)	5	(5–5)	5	(5–5)	0.068
Respiratory rate, median (IQR)	18	(16–20)	18	(15.5–20)	18	(16–20)	0.006
FiO ₂ , median (IQR)	50	(40–75)	60	(40–80)	50	(40–70)	0.029
Lung-protective ventilation** (%)	448	90.7	235	87.7	213	94.3	0.012
Ventilator Mode							0.068
VC-AC	454	90.8	238	87.5	216	94.7	
PC-AC	19	3.8	13	4.8	6	2.6	
VC-SIMV	14	2.8	10	3.7	4	1.8	
PC-SIMV	12	2.4	10	3.7	2	0.9	
PS	1	0.3	1	0.4	0	0.0	
ED protocol adherence	100	20.0	37	13.6	63	27.6	<0.001
ICU ventilator variables							
Tidal volume (mL)	430	380–475	439.5	381–480	420	380–460	0.161
Tidal volume (ml/kg IBW)	6.4	5.9–7.3	6.6	5.9–7.4	6.3	5.9–7.2	0.254
PEEP, cmH ₂ O	5	(5–5)	5	(5–5)	5	(5–5)	0.322
Respiratory rate	18	15–22	18	15–22	18	16–21	0.340
FiO ₂ (%)	40	40–60	40	40–60	40	40–55	0.128
Lung-protective ventilation** (%)	407	82.2	219	82.0	188	84.7	0.433
Ventilator mode							0.002
VC-AC	238	48.2	135	50.0	103	46.0	
PC-AC	37	7.5	23	8.5	14	6.3	
VC-SIMV	4	0.8	2	0.7	2	0.9	
PRVC-AC	97	19.6	38	14.1	59	26.3	
PS	76	15.4	40	14.8	36	16.1	
ICU protocol adherence	132	33.8	55	20.2	77	33.8	0.001

** Lung protective ventilation defined as tidal volume \leq 8 mL/kg.

3. Results

3.1. Characteristics of study subjects

A total of 500 patients were enrolled in this study Supplemental Fig. 1. Baseline characteristics of the study population are shown in Table 1. All measured baseline characteristics, including initial SOFA scores and ED length of ventilation, were similar between the two groups.

3.2. Main results

Of the study cohort, 272 subjects (54.4%) were in the pre-protocol group and 228 (45.6%) were in the post-protocol group (Table 1). Without adjustment, ED V_Ts were smaller in the post-protocol group (6.2 vs. 6.5 mL/kg IBW, $p < 0.001$), and more patients received LPV in the post-protocol group (94.3% vs. 87.7%, $p = 0.012$) (Table 1) than the pre-

protocol group. Time to ARDS and time to VAP did not differ between the pre- and post-protocol groups (ARDS: HR 0.64 [0.34–1.20] and VAP: HR 1.12 [0.58–2.18]). Overall, ICU and ED V_Ts were moderately correlated ($r^2 = 0.484$, $p < 0.001$) (Fig. 2).

After adjustment for sex, BMI, and initial SOFA score, ED V_T remained lower in the post-protocol group when compared to the pre-protocol group (-0.76 mL/kg [95%CI -1.03 to -0.48]) (Table 2). There was no difference in proportion of patients who developed ARDS, VAP, or died between the protocol groups. There were differences in the change in SOFA scores after adjustment with the post-protocol group having greater increases in SOFA score from the ED to 24-hour score (1.40 [95%CI: 0.64 to 2.17]) (Table 2). When GCS was removed from the SOFA calculation, there was no longer a significant difference in the change in SOFA scores.

Subject height modified the relationship between the protocol and use of ED LPV ($p = 0.044$). Before the protocol, a 10% increase in subject height was associated with a 3.39 (95% CI 2.14–4.64) times higher odds




Fig. 2. (A) Upper left: differences in ED tidal volume by pre- and post-intervention; (B) upper right: differences in ICU tidal volume by pre- and post-intervention; (C) lower: plot of differences in individual subjects' ED (black dot) to ICU tidal volume in pre- and post-intervention groups. Differences between ED and ICU tidal volume are represented by the grey line. Subjects are rank-ordered by ED tidal volume.

of ED LPV. Whereas, after the protocol, the association between height and the protocol was no longer significant (95%CI 0.24–2.46).

3.3. Protocol adherence

Protocol adherence in the ED was greater in the post-protocol group (27.6% vs. 13.6%, $p < 0.001$) (Table 1) than the pre-protocol group. Adherence to all components in the protocol was modest in both the pre- and post-group (Table 3). Using the same methods, ICU protocol adherence increased from 20.2% before protocol to 33.8% after protocol ($p = 0.001$). When missing protocol components were assumed adherent for sensitivity analysis, neither relationship changed significantly.

3.4. Variation by intubator and respiratory therapist

Adherence to use of LPV ($V_T \leq 8$ mL/kg IBW) varied by intubating physician and respiratory therapist (Fig. 3). More variation was observed by respiratory therapists than by intubator in adjusted proportion of ED LPV. Across intubators, the median adjusted rate of ED LPV was 90.7% (IQR 84.3–90.9). Whereas among respiratory therapists, the median adjusted proportion was similar, but more variation was observed (median 90.9% [IQR 77.8–91.7]). The impact of the care team in

predicting adherence with the protocol was estimated with univariable models with respiratory therapist or intubator as the predictor and ED LPV as the outcome. Overall, the individual respiratory therapist affected ED LPV more than the intubating physician (Fig. 4), and respiratory therapists become even stronger predictors after the intervention (AUC 0.919 [95% CI 0.917–0.920]). There was no association between attending physician and use of ED LPV.

3.5. Sensitivity analysis

Of this study cohort, 5.2% had ARDS on arrival to the ED and 41.2% were extubated on the date of admission; no subjects died on the date of admission. In the sensitivity analysis including only patients who remained ventilated beyond 24 h, the adjusted associations between the protocol and ED V_T (adj. β -0.16 [95% CI -0.21 to -0.12]) and between the protocol and mortality (aOR 0.28 [95% CI 0.02–3.41]) were similar to the primary analysis.

4. Discussion

This before-after observational cohort study was conducted to determine the impact of a bundled mechanical ventilator protocol on use

Table 2
Regression models.

	n/mean	%/SE	Unadjusted OR/between group difference (95% CI)	Adjusted* OR/between group difference (95% CI)
Primary outcome				
ED tidal volume (mL/kg)	6.6	0.1	-0.37 (-0.58 to -0.15) p = 0.001	-0.71 (-0.94 to -0.48) p < 0.001
Log-transformed ED TV	-	-	-0.05 (-0.08 to -0.02) p = 0.001	-0.10 (-0.13 to -0.06) p < 0.001
Secondary outcomes				
ARDS, n (%)	42	8.4%	0.64 (0.33–1.23) p = 0.179	0.57 (0.08–4.11) p = 0.577
VAP, n (%)	35	7.0%	1.14 (0.58–2.27) p = 0.703	0.76 (0.12–4.61) p = 0.762
28-day hospital free days, mean (SE)	14.7	0.5	-0.03 (-0.26 to 0.20) p = 0.798	-0.30 (-0.65 to 0.04) p = 0.085
Change in SOFA, mean (SE)	-1.0	0.2	0.26 (-0.34 to 0.86) p = 0.392	1.42 (0.67 to 2.18) p < 0.001
Change in SOFA (excluding GCS), mean (SE)	-0.79	0.1	-0.01 (-0.47 to 0.46) p = 0.983	0.50 (-0.02 to 1.02) p = 0.061
Change in SOFA components				
PaO ₂ /FiO ₂ ratio	-1.0	0.1	-0.19 (-0.51 to 0.14) P = 0.260	0.39 (0.12 to 0.67) p = 0.006
Platelets	0.2	0.0	0.06 (-0.05 to 0.16) p = 0.279	0.08 (-0.07 to 0.23) p = 0.304
Glasgow Coma Score	-0.1	0.1	0.27 (-0.06 to 0.59) p = 0.107	0.90 (0.46 to 1.35) p < 0.001
Creatinine	-0.1	0.0	-0.04 (-0.16 to 0.07) p = 0.460	-0.14 (-0.30 to 0.01) p = 0.075
Bilirubin	0.0	0.1	0.17 (-0.09 to 0.43) p = 0.208	0.28 (-7.06 to 7.63) p = 0.940
Mean arterial pressure	0.2	0.1	0.24 (-0.00 to 0.49) p = 0.055	0.30 (-0.03 to 0.63) p = 0.073
Mortality, n (%)	120	24.0%	1.03 (0.68–1.57) p = 0.873	1.22 (0.50–3.00) p = 0.666
Total length of ventilation, (min)	3831	272	0.06 (-0.13 to 0.25) p = 0.527	-0.12 (-0.40 to 0.17) p = 0.424
ED volume settings				
TV cut-point of 8, n (%)	448	90.6%	2.29 (1.17–4.48) p = 0.015	3.92 (0.35–43.99) p = 0.268
ICU volume settings				
Continuous TV, mean (SE)	6.8	0.1	-0.05 (-0.34 to 0.24) p = 0.739	-0.15 (-0.52 to 0.22) p = 0.416
Log-transformed ICU TV	-	-	-0.01 (-0.05 to 0.03) p = 0.566	-0.03 (-0.08 to 0.01) p = 0.180
TV cut-point of 8, n (%)	407	83.2%	1.29 (0.75–1.95) p = 0.435	1.97 (0.59–6.50) p = 0.268

Bold indicates p-value <0.05.

* Adjusted for: sex, BMI (logarithmic transform), height (logarithmic transform), initial SOFA score, respiratory therapist, and intubator.

of LPV in an academic ED. We found that such a protocol can be implemented and influence ventilation practices in the ED. After adjusting for multiple confounders, protocol administration was associated with a 0.76 mL/kg reduction in ED V_T. Though LPV was widely used in the ED pre-protocol group, use of V_T <8 mL/kg was significantly increased after implementation of the protocol. Adherence to PEEP, FiO₂, and respiratory rate protocol parameters were improved in the ED post-

protocol group, though full ED protocol adherence increased from 13.6% to only 27.6%. Despite the observed changes in ventilation practices, protocol implementation did not influence any of the measured clinical outcomes, including ARDS, VAP, hospital-free days, or mortality.

Our study is the third to analyze the effect of a bundled, lung-protective mechanical ventilation protocol administered in the ED. A previous before-after study, the LOV-ED trial, demonstrated a reduction

Table 3
Protocol adherence

	Total		Pre		Post	
	Adherent n (%)	Missing n (%)	Adherent n (%)	Missing n (%)	Adherent n (%)	Missing n (%)
ED protocol components						
PEEP FiO ₂ or FiO ₂ (hyperoxia)	174 (32.8)	1 (0.2)	75 (27.6)	1 (0.4)	99 (43.4)	0 (0.0)
PEEP (atelectrauma)	351 (70.2)	15 (3.0)	181 (66.5)	7 (2.6)	170 (74.6)	8 (3.5)
Respiratory rate (ventilate)	397 (79.4)	2 (0.4)	203 (74.6)	0 (0.0)	194 (85.1)	2 (0.9)
Tidal volume (volutrauma)	448 (89.6)	6 (1.2)	235 (86.4)	4 (1.5)	213 (93.4)	2 (0.9)
ICU protocol components						
PEEP FiO ₂ or FiO ₂ (hyperoxia)	307 (61.4)	6 (1.2)	151 (55.5)	2 (0.7)	156 (68.4)	4 (0.8)
PEEP (atelectrauma)	368 (73.6)	7 (1.4)	195 (71.7)	2 (0.7)	173 (75.9)	5 (2.2)
Respiratory rate (ventilate)	342 (68.4)	6 (1.2)	182 (66.9)	2 (0.7)	160 (70.2)	4 (1.8)
Tidal volume (volutrauma)	407 (81.4)	11 (2.2)	219 (80.5)	5 (1.8)	188 (82.5)	6 (2.6)

in pulmonary complications and mortality following protocol implementation [33]. Our study did not find a significant difference in clinical outcomes between groups despite improvement in overall protocol adherence and reduction in ED V_T. There are multiple potential explanations for a finding of a statistically significant decrease in ED V_T without a change in clinical outcomes. First, our study was primarily quality improvement (QI) in nature, and thus was not adequately powered to detect a change in clinical outcomes. Second, our study included patients who underwent extubation or death at <24 h following ED admission and those meeting ARDS criteria while in the ED. This creates a broader range of illness severity across our patient population and makes direct comparison in clinical outcomes between the studies difficult. However, a sensitivity analysis using the same exclusion criteria as

the LOV-ED study did not significantly change our primary outcome (change in V_T) or mortality. Third, we detected a 0.76 mL/kg adjusted difference in V_T between the pre and post protocol groups, which differs significantly from the 2 mL/kg difference detected in the LOV-ED trial [33]. Though reductions of 1 mL/kg have been shown to decrease mortality in patients with ARDS [44], 0.76 mL/kg may not be sufficient to elicit a change in secondary outcomes in a population that was receiving high rates of LPV even before protocol. The injurious effects of very high tidal volumes may be much more than modest tidal volumes, so the relatively protective settings in the pre-intervention group may not have led to high rates of clinical deterioration. Alternatively, the lack of clinical change could support the results of multiple studies challenging the clinical benefit of low V_T ventilation in populations including patients without ARDS [20,45].

Consistent adherence to all parameters of a complex, bundled protocol can be difficult to attain. Despite improvement in adherence to each facet of the ventilation protocol between the pre- and post-protocol groups, full adherence was seen in only 27.6% of ED patients in the post-protocol group. This finding can be at least partially attributed to protocol dissemination and implementation. Similar before-after respiratory protocol studies were prospective in nature and involved run-in periods of up to 6 months [31–33]. Our study, on the other hand, was retrospective and did not utilize a run-in period. While compliance to the protocol was monitored in the post-protocol group, these differences may have contributed to the observed sub-optimal adherence. The protocol parameter with the lowest adherence involved the setting of FiO₂ between 0.30 and 0.40 or in accordance with ARDSNet PEEP-FiO₂ table for hyperoxia prevention (37.7% in the post-protocol group). This finding is consistent with previous studies demonstrating frequent use of FiO₂ levels >90% with little oxygen titration in mechanically ventilated ED patients [24,25]. While use of low V_Ts had become standard practice in the ED, our results suggest that immediate titration of FiO₂ (Fig. 1) remains poor. Future studies of bundled care protocols should involve active identification of barriers to implementation to ensure successful translation into clinical practice.

ED and ICU patients in both protocol groups received LPV at a much higher proportion than previous reports. We found that 87.7% of ED patients (median V_T of 6.5 mL/kg) and 82.0% of ICU patients (median V_T of 6.6 mL/kg) intubated at our site were ventilated with LPV in the ED prior to protocol implementation. This differs significantly from previous studies demonstrating use of LPV in 23–55.7% of ED patients and 20–46% of ICU patients [24,32,33]. Our data more closely resembles the recent Low Tidal Volume Universal Support (LOTUS) trial feasibility study, which showed an average V_T of 7.1 mL/kg and LPV percentage of 78.2% in the ICUs of 49 hospitals [46]. In this context, our findings could reflect a shift in general practice to low V_T ventilation in all patients undergoing respiratory failure, regardless of ARDS status.

Our study found that variation in V_T administration was greater among RTs than intubating physicians, suggesting that the RTs were




Fig. 3. Variation in ED Lung Protective Ventilation Rates by Intubator and Respiratory Therapist.




Fig. 4. Variation in ED Lung Protective Ventilation Rates by Intubator and Respiratory Therapist.

most directly influencing ventilator settings. Previous survey studies have identified discomfort of emergency medicine residents and physicians in the care of mechanically ventilated patients, causing them to frequently defer care to respiratory therapists [47,48]. Our findings indicate that respiratory therapists are influential in determining ventilator settings, which highlights the importance of tailoring protocols for mechanically ventilated patients to respiratory therapy staff.

This study has several important limitations. As a before-after study, the results may be reflective of temporal changes in care. Since there were no other institutional efforts aimed at prophylactic LPV during the period of this study, however, we think that it remains likely that the ED protocol influenced care. The retrospective design of this study limits data available, but the research team took steps to validate data when possible and selected measures are likely to be recorded accurately. This study was conducted at a single academic teaching facility, which limits generalizability, but since prior reports were also single-center studies, we feel that local factors may lead to heterogeneity in treatment effects. ED providers' and respiratory therapists' awareness of this study may have introduced a Hawthorne effect, in which providers performed differently knowing they were being monitored. However, as quality improvement monitoring is part of the implementation of this intervention, we considered it an important parameter to capture and part of the treatment effect. Data were collected through chart review and providers were never directly observed. Finally, we did not include plateau pressure in our definition of LPV [49–51], because it could not be measured with our standard ED transport ventilators.

5. Conclusion

Implementation of a bundled, ED-based mechanical ventilation protocol is associated with a significant decrease in ED V_T and increase in use of LPV. The protocol was not associated with change in any measured clinical outcomes, including ventilator-free days, ARDS, VAP, or mortality. The results of our study support the use of protocols to standardize care of ventilated ED patients and emphasize the importance of tailoring these protocols to respiratory therapists, who were shown to have a more significant impact on LPV use than intubating physician. Further study to maximize adherence to protocol parameters, particularly the immediate titration of FiO_2 , and clarify the influence of ED V_T on patient-centered outcomes is warranted.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ajem.2020.02.053>.

Financial support

This research was supported by the University of Iowa Carver College of Medicine and Department of Emergency Medicine.

CRedit authorship contribution statement

Tyler M. Foley: Conceptualization, Methodology, Investigation, Data curation, Writing - original draft, Writing - review & editing. **Brittany A. Philpot:** Conceptualization, Methodology, Investigation, Writing - original draft, Writing - review & editing. **Alysa S. Davis:** Conceptualization, Methodology, Investigation, Writing - original draft, Writing - review & editing. **Morgan B. Swanson:** Formal analysis, Data curation, Writing - original draft, Writing - review & editing. **Karisa K. Harland:** Formal analysis, Data curation, Writing - original draft, Writing - review & editing. **Justin D. Kuhn:** Writing - original draft, Writing - review & editing. **Brian M. Fuller:** Conceptualization, Methodology, Writing - original draft, Writing - review & editing. **Nicholas M. Mohr:** Conceptualization, Methodology, Data curation, Writing - original draft, Writing - review & editing.

Declaration of competing interest

None.

References

- [1] Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med* 2013;369(22):2126–36.
- [2] Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342(18):1301–8.
- [3] Meade MO, Cook DJ, Guyatt GH, Slutsky AS, Arabi YM, Cooper DJ, et al. Ventilation strategy using low tidal volumes, recruitment maneuvers, and high positive end-expiratory pressure for acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 2008;299(6):637–45.
- [4] Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998;338(6):347–54.
- [5] Villar J, Kacmarek RM, Pérez-Méndez L, Aguirre-Jaime A. A high positive end-expiratory pressure, low tidal volume ventilatory strategy improves outcome in persistent acute respiratory distress syndrome: a randomized, controlled trial. *Crit Care Med* 2006;34(5):1311–8.
- [6] A. Serpa Neto, S.O. Cardoso, J.A. Manetta, V.G. Pereira, D.C. Espósito, M.e.O. Pasqualucci, M.C. Damasceno, M.J. Schultz, Association between use of lung-protective ventilation with lower tidal volumes and clinical outcomes among patients without acute respiratory distress syndrome: a meta-analysis, *JAMA* 308 (16) (2012) 1651–9.

- [7] Fuller BM, Mohr NM, Drewry AM, Carpenter CR. Lower tidal volume at initiation of mechanical ventilation may reduce progression to acute respiratory distress syndrome: a systematic review. *Crit Care* 2013;17(1):R11.
- [8] Determann RM, Royakkers A, Wolthuis EK, Vlaar AP, Choi G, Paulus F, et al. Ventilation with lower tidal volumes as compared with conventional tidal volumes for patients without acute lung injury: a preventive randomized controlled trial. *Crit Care* 2010;14(1):R1.
- [9] Futier E, Constantin JM, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013;369(5):428–37.
- [10] Gajic O, Dara SI, Mendez JL, Adesanya AO, Festic E, Caples SM, et al. Ventilator-associated lung injury in patients without acute lung injury at the onset of mechanical ventilation. *Crit Care Med* 2004;32(9):1817–24.
- [11] Wolthuis EK, Choi G, Delsing MC, Bresser P, Lutter R, Dzoljic M, et al. Mechanical ventilation with lower tidal volumes and positive end-expiratory pressure prevents pulmonary inflammation in patients without preexisting lung injury. *Anesthesiology* 2008;108(1):46–54.
- [12] Halter JM, Steinberg JM, Gatto LA, DiRocco JD, Pavone LA, Schiller HJ, et al. Effect of positive end-expiratory pressure and tidal volume on lung injury induced by alveolar instability. *Crit Care* 2007;11(1):R20.
- [13] McCann UG, Schiller HJ, Carney DE, Gatto LA, Steinberg JM, Nieman GF. Visual validation of the mechanical stabilizing effects of positive end-expiratory pressure at the alveolar level. *J Surg Res* 2001;99(2):335–42.
- [14] Damiani E, Adrario E, Girardis M, Romano R, Pellaia P, Singer M, et al. Arterial hyperoxia and mortality in critically ill patients: a systematic review and meta-analysis. *Crit Care* 2014;18(6):711.
- [15] Helmerhorst HJ, Roos-Blom MJ, van Westerloo DJ, de Jonge E. Association between arterial hyperoxia and outcome in subsets of critical illness: a systematic review, meta-analysis, and meta-regression of cohort studies. *Crit Care Med* 2015;43(7):1508–19.
- [16] Page D, Ablordeppey E, Wessman BT, Mohr NM, Trzeciak S, Kollef MH, et al. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. *Crit Care* 2018;22(1):9.
- [17] Herring AA, Ginde AA, Fahimi J, Alter HJ, Maselli JH, Espinola JA, et al. Increasing critical care admissions from U.S. emergency departments, 2001–2009. *Crit Care Med* 2013;41(5):1197–204.
- [18] Mullins PM, Goyal M, Pines JM. National growth in intensive care unit admissions from emergency departments in the United States from 2002 to 2009. *Acad Emerg Med* 2013;20(5):479–86.
- [19] Easter BD, Fischer C, Fisher J. The use of mechanical ventilation in the ED. *Am J Emerg Med* 2012;30(7):1183–8.
- [20] Wilcox SR, Richards JB, Fisher DF, Sankoff J, Seigel TA. Initial mechanical ventilator settings and lung protective ventilation in the ED. *Am J Emerg Med* 2016;34(8):1446–51.
- [21] Dreyfuss D, Soler P, Basset G, Saumon G. High inflation pressure pulmonary edema. Respective effects of high airway pressure, high tidal volume, and positive end-expiratory pressure. *Am Rev Respir Dis* 1988;137(5):1159–64.
- [22] Hoegl S, Boost KA, Flondor M, Scheiermann P, Muhl H, Pfeilschifter J, et al. Short-term exposure to high-pressure ventilation leads to pulmonary biotrauma and systemic inflammation in the rat. *Int J Mol Med* 2008;21(4):513–9.
- [23] Ramnath VR, Hess DR, Thompson BT. Conventional mechanical ventilation in acute lung injury and acute respiratory distress syndrome. *Clin Chest Med* 2006;27(4):601–13 [abstract viii].
- [24] Fuller BM, Mohr NM, Miller CN, Deitchman AR, Levine BJ, Castagno N, et al. Mechanical ventilation and ARDS in the ED: a multicenter, observational, prospective, cross-sectional study. *Chest* 2015;148(2):365–74.
- [25] Fuller BM, Mohr NM, Dettmer M, Kennedy S, Cullison K, Bavolek R, et al. Mechanical ventilation and acute lung injury in emergency department patients with severe sepsis and septic shock: an observational study. *Acad Emerg Med* 2013;20(7):659–69.
- [26] Modrykamien AM, Stoller JK. The scientific basis for protocol-directed respiratory care. *Respir Care* 2013;58(10):1662–8.
- [27] Beasley KE, Darin JM, Durbin CG. The effect of respiratory care department management of a blood gas analyzer on the appropriateness of arterial blood gas utilization. *Respir Care* 1992;37(4):343–7.
- [28] Pilon CS, Leathley M, London R, McLean S, Phang PT, Priestley R, et al. Practice guideline for arterial blood gas measurement in the intensive care unit decreases numbers and increases appropriateness of tests. *Crit Care Med* 1997;25(8):1308–13.
- [29] Dries DJ, McGonigal MD, Malian MS, Bor BJ, Sullivan C. Protocol-driven ventilator weaning reduces use of mechanical ventilation, rate of early reintubation, and ventilator-associated pneumonia. *J Trauma* 2004;56(5):943–51 [discussion 951–2].
- [30] Haas CF, Loik PS. Ventilator discontinuation protocols. *Respir Care* 2012;57(10):1649–62.
- [31] Radosevich MA, Wanta BT, Meyer TJ, Weber VW, Brown DR, Smischney NJ, et al. Implementation of a goal-directed mechanical ventilation order set driven by respiratory therapists improves compliance with best practices for mechanical ventilation. *J Intensive Care Med* 2019;34(7):550–6 885066617746089.
- [32] Prekker ME, Donelan C, Ambur S, Driver BE, O'Brien-Lambert A, Hottinger DG, et al. Adoption of low tidal volume ventilation in the emergency department: a quality improvement intervention. *Am J Emerg Med* 2019. <https://doi.org/10.1016/j.ajem.2019.06.026> In Press.
- [33] Fuller BM, Ferguson IT, Mohr NM, Drewry AM, Palmer C, Wessman BT, et al. Lung-protective ventilation initiated in the emergency department (LOV-ED): a quasi-experimental, before-after trial. *Ann Emerg Med* 2017;70(3):406–418.e4.
- [34] Davidoff F, Batalden P, Stevens D, Ogrinc G, Mooney SE, S.D. Group. Publication guidelines for quality improvement studies in health care: evolution of the SQUIRE project. *BMJ* 2009;338:a3152.
- [35] Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996;22(7):707–10.
- [36] Pai MP, Paloucek FP. The origin of the “ideal” body weight equations. *Ann Pharmacother* 2000;34(9):1066–9.
- [37] Rice TW, Wheeler AP, Bernard GR, Hayden DL, Schoenfeld DA, Ware LB, et al. Comparison of the SpO₂/FIO₂ ratio and the PaO₂/FIO₂ ratio in patients with acute lung injury or ARDS. *Chest* 2007;132(2):410–7.
- [38] Severinghaus JW. Simple, accurate equations for human blood O₂ dissociation computations. *J Appl Physiol Respir Environ Exerc Physiol* 1979;46(3):599–602.
- [39] Gajic O, Frutos-Vivar F, Esteban A, Hubmayr RD, Anzueto A. Ventilator settings as a risk factor for acute respiratory distress syndrome in mechanically ventilated patients. *Intensive Care Med* 2005;31(7):922–6.
- [40] Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307(23):2526–33.
- [41] Ferguson ND, Fan E, Camporota L, Antonelli M, Anzueto A, Beale R, et al. The Berlin definition of ARDS: an expanded rationale, justification, and supplementary material. *Intensive Care Med* 2012;38(10):1573–82.
- [42] Wong T, Schlichting AB, Stoltz AJ, Fuller BM, Peacock A, Harland KK, et al. No decrease in early ventilator-associated pneumonia after early use of chlorhexidine. *Am J Crit Care* 2016;25(2):173–7.
- [43] Dimick JB, Staiger DO, Birkmeyer JD. Ranking hospitals on surgical mortality: the importance of reliability adjustment. *Health Serv Res* 2010;45(6):1614–29 Pt 1.
- [44] Needham DM, Yang T, Dinglas VD, Mendez-Tellez PA, Shaholtz C, Sevransky JE, et al. Timing of low tidal volume ventilation and intensive care unit mortality in acute respiratory distress syndrome. A prospective cohort study. *Am J Respir Crit Care Med* 2015;191(2):177–85.
- [45] Simonis FD, Serpa Neto A, Binnekade JM, Braber A, Bruin KCM, Determann RM, et al. Effect of a low vs intermediate tidal volume strategy on ventilator-free days in intensive care unit patients without ARDS: a randomized clinical trial. *JAMA* 2018;320(18):1872–80.
- [46] Lanspa MJ, Gong MN, Schoenfeld DA, Lee KT, Grissom CK, Hou PC, et al. Prospective assessment of the feasibility of a trial of low-tidal volume ventilation for patients with acute respiratory failure. *Ann Am Thorac Soc* 2019;16(3):356–62.
- [47] Wilcox SR, Seigel TA, Strout TD, Schneider JI, Mitchell PM, Marcolini EG, et al. Emergency medicine residents' knowledge of mechanical ventilation. *J Emerg Med* 2015;48(4):481–91.
- [48] Wilcox SR, Strout TD, Schneider JI, Mitchell PM, Smith J, Lutfy-Clayton L, et al. Academic emergency medicine physicians' knowledge of mechanical ventilation. *West J Emerg Med* 2016;17(3):271–9.
- [49] Amato MB, Meade MO, Slutsky AS, Brochard L, Costa EL, Schoenfeld DA, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med* 2015;372(8):747–55.
- [50] Fuller BM, Page D, Stephens RJ, Roberts BW, Drewry AM, Ablordeppey E, et al. Pulmonary mechanics and mortality in mechanically ventilated patients without acute respiratory distress syndrome: a cohort study. *Shock* 2018;49(3):311–6.
- [51] C. Guérin, L. Papazian, J. Reignier, L. Ayzac, A. Loundou, J.M. Forel, I.o.t.A.a.P. trials, Effect of driving pressure on mortality in ARDS patients during lung protective mechanical ventilation in two randomized controlled trials, *Crit Care* 20(1) (2016) 384.