# Review Article **Cytotoxicity of Air Pollutant 9,10-Phenanthrenequinone: Role of Reactive Oxygen Species and Redox Signaling**

## Manli Yang,<sup>1</sup> Hassan Ahmed,<sup>2</sup> Weidong Wu<sup>1</sup>,<sup>1</sup> Bijie Jiang,<sup>1</sup> and Zhenquan Jia<sup>2</sup>

<sup>1</sup>Department of Environmental and Occupational Health, School of Public Health, Xinxiang Medical University, Xinxiang, Henan Province 453003, China <sup>2</sup>Department of Biology, University of North Carolina at Greensboro, Greensboro, NC, USA

Correspondence should be addressed to Weidong Wu; wdwu2013@126.com and Zhenquan Jia; z\_jia@uncg.edu

Received 3 March 2018; Accepted 30 April 2018; Published 10 June 2018

Academic Editor: Lidong Zhai

Copyright © 2018 Manli Yang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Atmospheric pollution has been a principal topic recently in the scientific and political community due to its role and impact on human and ecological health. 9,10-phenanthrenequinone (9,10-PQ) is a quinone molecule found in air pollution abundantly in the diesel exhaust particles (DEP). This compound has studied extensively and has been shown to develop cytotoxic effects both in vitro and in vivo. 9, 10-PQ has been proposed to play a critical role in the development of cytotoxicity via generation of reactive oxygen species (ROS) through redox cycling. This compound also reduces expression of glutathione (GSH), which is critical in Phase II detoxification reactions. Understanding the underlying cellular mechanisms involved in cytotoxicity can allow for the development of therapeutics designed to target specific molecules significantly involved in the 9,10-PQ-induced ROS toxicity. This review highlights the developments in the understanding of the cytotoxic effects of 9, 10-PQ with special emphasis on the possible mechanisms involved.

## 1. Introduction

Atmospheric pollution has been an important topic in the scientific and political community due to its impact on human and ecological health [1]. With industry development, the increase in city heating, and the number of automobiles, there was a severe increase in the concentration of air pollutants[2]. After the Great Smog event in London in 1952[3], the scientific community and politic governments are paying more attention to air pollution on human health [1]. Over the Great Smog event days, mortality in the general population was more than threefold times than expected, leading to thousands of people death[2]. Since this historical air pollution episode, a large number of studies have demonstrated the association between air pollution and human disease including respiratory and lung diseases such as chronic obstructive pulmonary disease, asthma attacks, pulmonary cancer, leukemia, birth defects and immune system defects, cardiovascular problems, heart disease and stroke, neurobehavioral disorders, and liver and other types of cancer [4-18]. Diesel combustion to generate power has existed for well over a century and is used in heavy machinery such as large trucks and oil tankers [19–22]. The mechanism by which diesel exhaust impacts human health is being studied extensively [23–31]. One of the most significant and detrimental impacts diesel combustion is atmospheric pollution.

Diesel particles can either aggregate or remain as individual particles when released into the atmosphere. Upon inhalation, these fine particles can penetrate through the lungs causing many diseases and disorders to develop over time. DEP have been shown to play an important role in the generation of pulmonary-related diseases including bronchitis, asthma, carcinogenesis in the lungs, and other allergic disorders [23–32]. DEPs are composed of complex mixtures of particulates, including elemental carbon, polycyclic aromatic hydrocarbons (PAHs), nitro-PAHs, quinones, heterocyclics, aldehydes, and other hydrocarbons[33–36].

One major particle found in diesel exhaust is 9-10phenanthrenequinone (9,10-PQ, see its structure in Figure 1), a type of quinone. Quinones are electrophilic compounds derived from aromatic organic molecules through oxidation.



FIGURE 1: 9-10-phenanthrenequinone structure.

These molecules have two ketone groups and, thus, are biologically active. Understanding the toxic role of quinones is of great interest as this allows for the development of novel therapeutics designed to specifically target downstream mediators. Exposure to 9,10-PQ containing particulate matter (PM) has been associated with the development of many disease processes including lung cancer, asthma, and allergic inflammation [37–52].

### 2. 9,10-Phenanthrenequinone Is One of the Major Components of Particulate Matter Present in the Environment

As one of the most abundant quinones found in DEPs [43], when inhaled, 9,10-PQ can be generated from phenanthrene by P-450s as below pathway [37]. Phenanthrene is one of major PAHs present in the air pollution [37–52].

In addition to phenanthrene photooxidation, 9,10-PQ could be also directly emitted to the environment from auto exhaust sources such as diesel exhaust particles. The levels of airborne DEP have increased dramatically over the last few decades due to the increase in the use of diesel-based engines. Compared to gasoline-based engines, diesel-based engines can provide higher fuel efficiency and lower carbon dioxide emissions; they also emit about 30–100 times more particulate matter into the atmosphere than gasoline-based engines [53]. It has been estimated that DEP constitute as much as 40% of the respirable particulate matter in a city where the daily human intake has been estimated to be as much as 300  $\mu$ g m<sup>-3</sup> of particulate matter [54]

A few studies determined the 9,10-PQ concentration in the airborne particulates [55, 56]. Various PAH-quinones including 1-acenaphthene, 9-fluorenone, 11H-benzo[a]fluoren-11-one, 7H-benzo[c]fluoren-7-one, 11H-benzo[b]fluoren-11one, benzanthrone, and 6H-benzo[cd]pyrene-6-one, 1,4naphthoquinone, phenanthrenequinone, 5,12-naphthacenequinone, and benzo[a]pyrene-6,12-dione have been detected at less than 1 ppm in the atmospheric particles in Boston, MA [55, 56]. The four quinones, 1,2-naphthoquinone (1,2-NQ), 1,4-naphthoquinone (1,4-NQ), 9,10-phenanthraquinone (9,10-PQ), and 9,10-anthraquinone (9,10-AQ), have been detected in Standard Reference Material (SRM) 1649a and in ambient air samples of PM<sub>2.5</sub> from several rural and urban areas in Central Los Angeles [55, 56]. The mean concentrations of these four quinones are 9–40.4  $\mu$ g/g in the DEP and 5–730 pg/m<sup>3</sup> in the PM<sub>2.5</sub> samples [55, 56]. Among these four detected quinones, 9,10-PQ has been found to have higher concentration [55, 56]. Kishikawa et al. employed a high-performance liquid chromatography with fluorescence detection method for detecting 9,10-PQ in the sample of PM collected at an urban site in Nagasaki city, Japan, weekly over a year from July 1997 to June 1998 [57]. This method is highly sensitive and allows detecting 9,10-PQ at very low concentrations of airborne particulates [57]. The average concentration of 9,10-PQ in airborne particulates was reported to be 0.287+/-0.128 ppm. Interesting, 9,10-PQ concentrations were higher in winter than in summer. Also, 9,10-PQ concentrations were higher during weekdays than at weekends [57].

9,10-PQ concentration in the PM has been investigated in a few areas during the different season. The phenanthrenequinone mean concentrations in Fresno, CA in winter 2004-2005, and spring 2005 are  $1.1\pm0.77$ ng·m<sup>-3</sup> and  $0.31\pm0.13$ ng·m<sup>-3</sup>, respectively [58]. However, phenanthrene average concentrations in urban ambient Southern California are 50.34 ng·m<sup>-3</sup> measured over the period of September 8-9, 1993[59]. 9,10-PQ in Central Los Angeles was found to be 24.19µg/g in DEP[60]. So, the atmospheric PM2.5 and DEP consist of significant levels of 9,10-PQ [61]. High levels of 9,10-PQ were found in samples collected in San Dimas and in Riverside compared to rural Atascadero due to lower automotive exhaust emissions in rural Atascadero than that in San Dimas and in Riverside [60].

#### 3. Redox Cycling of 9,10-PQ in the Generation of Reactive Oxygen Species (ROS)

As a redox-active quinone in diesel exhausts, 9,10-PQ has been shown to generate ROS (Figure 2) via its redox cycling resulting in its toxicity [37-52, 57, 62, 63, 67, 71-88] in vitro and in vivo. 9,10-PQ can accept an electron to produce the semiquinone form of 9,10-PQ [70]. Incomplete reduction of molecular oxygen can cause the redox cycling of 9,10-PQ as the semiquinone form is oxidized back to the original quinone form [70]. During this process, the molecular oxygen becomes incompletely reduced and becomes a superoxide anion radical  $(O_2^{\bullet-})$  [89]. Superoxide radicals, considered as primary ROS, then undergo further metabolism and interact with other molecules to generate secondary ROS such as hydrogen peroxide  $(H_2O_2)$  and highly reactive hydroxyl radicals (\*OH) [90]. ROS are often characterized by their capability to bring an incredible amount of both discriminant and indiscriminate damage to biomolecules including protein, lipids, and nucleic acids. In this context, there are two reduced 9,10-PQ species of biological importance, its semiquinone radical (9,10-phenanthraquinone; PQU-) and its hydroquinone PQH2. The semiquinone form of 9,10-PQ can trigger redox cycling by transferring electrons from a source such as nicotinamide adenine dinucleotide phosphate (NADPH) to oxygen to generate ROS. The semiquinone form of 9,10-PQ was found to be generated via the enzymatic reaction or nonenzymatic reaction. The enzymatic reaction



FIGURE 2: Redox cycling of 9,10-phenanthrenequinone (9,10-PQ) for production of reactive oxygen species (ROS). First, 9,10-PQ quinone undergoes 1 electron reduction to produce semiquinone radical. Semiquinone is unstable and is very reactive to oxygen. It reoxidizes back to quinone and releases superoxide radical. Superoxide molecule is then reduced to hydrogen peroxide by the superoxide dismutase. Hydroxyl radicals are formed when hydrogen peroxide reacts with metals such as ferrous ions via Fenton reaction. Second, semiquinone can also undergo further 1 electron reduction to form 9,10-phenanthrene hydroquinone which can further redox cycles back to semiquinone by losing an electron and yielding superoxide radical. Lastly, 2-electron reduction of 9,10-PQ quinone by NADPH-quinone oxidoreductase-1 (NQO1) will directly form 9,10-phenanthrene hydroquinone. Thus, redox cycle of 9,10-PQ will generate a large amount of ROS.

was catalyzed by flavoenzymes such as NADPH-cytochrome P450 reductase or neuronal nitric oxide synthase and a nonenzymatic reaction was targeted with proximal protein thiols.

 $H_2O_2$  levels within cells can be used to develop an understanding of how oxidative stress and the generation of ROS is involved in cell-induced apoptosis [91–96]. Treatment of human T lymphoblast: acute lymphoblastic leukemia (MOLT-4) cells with DCFH-DA as a fluorogenic probe can detect the levels of hydrogen peroxide within the cell. DCFH-DA is converted to DCFH which additionally is converted to 2',7' –dichlorofluorescein (DCF) by  $H_2O_2$ . If  $H_2O_2$  is present, this conversion will occur, and DCF levels can be measured by fluorescence to determine the levels of hydrogen peroxide within the cell. Cells treated with 9,10-PQ were shown to have a significant increase in  $H_2O_2$  levels than control cells [50]. An interesting approach and technique that can be used to further understand and thereby strengthen the idea of the presence of ROS within the cells are to treat cells with 9,10-PQ as well as ROS scavengers such as polyethylene glycol catalase (PEG-cat) [50]. Upon treatment with PEG-cat and 9,10-PQ, the levels of hydrogen peroxide returned to basal control levels, indicating that 9,10-PQ indeed is involved in the generation of ROS. In addition, pretreatment of cells with PEG-cat and subsequent treatment with 9,10-PQ resulted in increased cellular viability. This further explains the role of 9,10-PQ in the generation of ROS and how ROS scavengers are involved in reducing the detrimental effects posed to the cells upon exposure to 9,10-PQ.

# 4. Role of L-Xylulose Reductase (XR) in 9,10-PQ-Induced Superoxide Production

XR is involved in the reduction of 9,10-PQ at a higher rate than its substrates diacetyl and L-xylulose [50, 97]. This enzyme was thought to be involved in cellular detoxification by reducing  $\alpha$ -dicarbonyl compounds, thus reducing reactive carbonyl compounds. In contrast to what was expected, upon treatment of bovine aortic endothelial cells (BAEC) containing a sixfold overexpression of the XR enzyme with 9,10-PQ, cell viability drastically decreased and significantly enhanced cytotoxicity when compared to wild-type cells treated with only 9,10-PQ [50]. Selective inhibition of XR by 4-methyl-[1,2,3]-thiadiazole-5-carboxylic acid benzyloxy-amide (MTB) eliminated this enhanced cytotoxicity indicating that XR plays a crucial role in decreasing cellular viability when exposed to 9,10-PQ [50]. Furthermore, when XR transformed cells were pretreated with PEG-cat, GSH ester, and N-acetyl-L-cysteine (NAC), the XR induced enhanced cytotoxicity was significantly reduced [50].

Cytochrome C is an enzyme that can be used to measure the production of superoxide anion, a very powerful radical generated in the ROS pathway. As superoxide anion levels increase the reduction rate of cytochrome C also increases. In a reaction mixture containing cytochrome C, XR, and 9,10-PQ, decrease in levels of cytochrome C significantly increased compared to the control mixture lacking 9,10-PQ indicating the generation of the superoxide anion. Also, when the reaction mixture was treated with CuZn-SOD, a compound involved in the conversion of superoxide anion into oxygen and  $H_2O_2$ , cytochrome C reduction levels decreased, further explaining the role of XR in enhancing 9,10-PQ induced cytotoxicity through the generation of superoxide anion [98].

After establishing the relationship between XR and cytotoxicity, it is crucial to understand how the induction levels of XR change upon exposure to 9,10-PQ. When MOLT-4 cells were treated with 1 uM 9,10-PQ, XR mRNA levels were increased 2 hours after treatment, indicating that XR may be an oxidant-inducible enzyme. Cells treated with low concentrations of 9,10-PQ resulted in a greater expression of XR mRNA, as opposed to cells treated with higher concentrations of 9,10-PO. A probable explanation for this phenomenon can be obtained from the DNA fragmentation results. As the cells were treated with higher concentrations of 9,10-PQ increase in DNA fragmentation was observed. This increase in DNA fragmentation can be a cause of reduced expression of XR mRNA levels at higher concentrations of 9,10-PQ. Furthermore, to strengthen the idea of 9,10-PQ induced upregulation of XR mRNA expression, cells exposed to 9,10-PQ were pretreated with radical scavengers PEG-cat and NAC to understand how XR mRNA levels may change [98]. XR mRNA levels were indeed reduced upon pretreatment of cells with radical scavengers; thus, it can be concluded that enhanced expressions of XR mRNA in the presence of 9,10-PQ can be mitigated upon pretreatment with PEG-cat and NAC. Upregulation of XR mRNA was not only observed in cells treated with 9,10-PQ, cells treated with glucose oxidase (GO) and glucose, a hydrogen peroxide generating system, also increased the expression of XR mRNA levels indicating the strong relationship between ROS generation and upregulation of XR. Knowing this relationship between ROS generation and enhanced XR levels, it can be predicted that treatment of cells with XR inhibitory factors can reduce ROS generation levels as well as increasing cell viability levels when compared to cells treated with only 9,10-PQ without XR inhibitors [98]. Indeed, upon treatment of cells with MTB, an inhibitor of XR, cytochrome C reduction significantly

decreased and cellular viability significantly increased. These results further strengthen the important role that XR plays in the molecular pathway of 9,10-PQ induced cellular toxicity.

#### 5. 9,10-PQ and Cellular Toxicity

Many studies demonstrated that PQ can cause toxicity in vivo and in vitro biological system via the redox cycling for the generation of ROS [42, 46, 47, 50, 72, 78, 82]. Excessive ROS is known to cause DNA damage resulting in gene mutation [99-103]. 9,10-PQ has been shown to exert toxic effects in human skin cell lines with low NQO1 activity [45]. Nitric oxide (NO) is widely known to suppress cytotoxic effects of oxidative damage. Pretreatment of endothelial cells resulted in decreased toxicity of 9,10-PQ and conversely, depletion of NO resulted in a greater level of toxicity, signifying the role of NO in cellular detoxification [48]. Interestingly, antitumor studies have been performed using analogs of 9,10-PQ due to their role in apoptosis. In HCT-116 colon tumor cells and HL-60 promyelocytic leukemia cells, 9,10-PQ analogs were successful in inducing apoptosis [43]. In the trophoblast cell line, JEG-3, 9,10-PQ exerted greater amounts of cytotoxicity and increased levels of redox cycling were measured when coupled with copper [67]. 9,10-PQ has been shown to cause iron-mediated cellular damage in human pulmonary epithelial cells. 9,10-PQ induced apoptosis and downregulated the effects of CuZn-SOD, which converts superoxide anion into oxygen or H<sub>2</sub>O<sub>2</sub> [52]. PQH<sub>2</sub>, a hydroquinone of 9,10-PQ, has shown to play a significant role in generating oxidative stress, protein damage, and cellular toxicity [70]. Exposure to 9,10-PQ can be examined in urine specimen by testing for the presence of 9,10-PQHG, a major metabolite generated during the breakdown of 9,10-PQ. This metabolite of 9,10-PQ was detected in rats injected with 9,10-PQ as well as humans exposed to diesel exhaust particles [63].

One of the most crucial experiments highlighting the molecular mechanisms of 9,10-PQ induced cellular toxicity includes a study performed by Matsunaga et al. in 2007. This study evaluated the effect of 9,10-PQ on various cellular mechanisms such as apoptosis, generation of ROS, and the role of the cytoprotective enzyme XR in the apoptotic signaling and ROS generating pathway in BAECs [50].

#### 6. The Role of 9,10-PQ in Cellular Viability and DNA Damage

In a study carried out by Matsunaga et al., it has been shown that 9,10-PQ induces apoptosis in MOLT-4 cells similar to other quinones such as 1,2-naphthoquinone and 1,4-naphthoquinone. MOLT-4 cells are human T lymphoma cells, obtained from cancerous T cell leukemia tissue [50]. The use of cancerous tissue in the study of apoptosis can be very useful in the development of not only anticytotoxic treatments but also antitumor treatments. One of the most important aspects of understanding toxicity is measuring the effects of a compound on cellular toxicity. 9,10-PQ was shown to decrease the viability of MOLT-4 cells as early as 8 hours after treatment. Cells were also treated with phenanthrene to measure cellular toxicity. The purpose of measuring toxicity upon treatment with phenanthrene was to explain the significance the quinone group has on biological activity, as the structure of phenanthrene is similar to 9,10- PQ except it lacks the quinone group. Phenanthrene did not have significant effects on the viability of the cells. Dose-dependent response to cellular toxicity was also measured in this experiment using varying concentrations of 9,10-PQ, 1,2-NQ, 1,4-NQ, acenaphthenequinone, and phenanthrene. The results suggest that all the quinones except acenaphthenequinone contributed significantly to cell viability upon treatment with 1 uM of the quinone [50].

DNA fragmentation occurs when the integrity of the nuclear membrane is compromised by exposure to xenobiotic compounds, which in turn causes morphological changes of the nucleus and DNA. These changes are pivotal features and steps in cells undergoing apoptosis or programmed cell death. Upon treatment of MOLT-4 cells with 9,10-PQ, DNA was extracted and analyzed via gel electrophoresis [50]. A clear and transparent increase in fragmentation of the DNA was observed in cells treated with 0.05 to 5 uM of 9,10-PQ. Treatment with 5 uM of phenanthrene did not have any effect on the fragmentation of DNA, further strengthening the idea that the quinone portion of 9,10-PQ is a vital component of the molecule necessary in the exertion of nuclear toxicity in cells. A separate study performed by Rodriguez et al. on Saccharomyces cerevisiae analyzed the role of 9,10-PQ in DNA damage, specifically examining deletions and point mutations in DNA [49]. The results specify that 9,10-PQ plays a role in DNA damage only when treated to cells in the presence of oxygen. No changes in DNA abnormality were observed within cells treated with 9,10-PQ in anaerobic conditions in yeast cells indicating oxygen is required by cells undergoing 9,10-PQ induced DNA damage. Also, an important protein involved in the repair of DNA is Poly (ADP-ribose) polymerase (PARP) [50]. The levels of activated PARP can give an indication as to the amount of stress, specifically DNA damage, a cell is experiencing [104-107]. PARP repairs DNA and increase in DNA damage triggers an increase in PARP activity. 9,10-PQ has been shown to increase the levels of activated PARP inside cells, further strengthening the role of 9,10-PQ in DNA damage. Nucleic morphology can provide imperative clues in understanding the methodology by which cells are undergoing death.

The two main types of cell death are apoptosis and necrosis [108-111]; noteworthy, molecular and morphological differences are observed in each type of cell death. Apoptosis is referred to as programmed cell death involving well-organized sequence of morphological events [108-111]. Nuclear and cytoplasmic membranes of cells undergoing apoptosis shrink and condense which results in the collapse of the cytoskeleton causing the formation of blebs by the cell membrane. These cells subsequently undergo degradation of the genetic and protein material as the nucleus is condensed and fractured. The cellular debris or blebs are eventually phagocytosed by neighboring cells or macrophages [108-111]. On the contrary, necrosis is referred to as accidental or unscheduled cell death. Necrotic cells swell and distend which drastically alters their structure due to an inability to maintain membrane integrity. This in turn ultimately results

in the damage and destruction of the organelles, and the interior composition of the cells leak out. Nucleic morphology of cells treated with 9,10-PQ displayed the formation of condensed and fragmented nuclei. These changes explain the noteworthy role of 9,10-PQ in the induction of apoptosis [50].

#### 7. The Role of 9,10-PQ on Mitochondria Dysfunction and Caspase Activation

The mitochondria play an essential role in the steps leading towards apoptosis; thus, studying this organelle is vital in understanding how xenobiotic factors impair the activity of cells through this organelle [112-114]. Three key proteins located on the inner mitochondrial membrane include the apoptotic Bax and the antiapoptotic Bcl-2 and Bcl-XL play a crucial role in regulating the permeability of the membrane [115–119]. Upregulating the activity of Bax and downregulating the activity of Bcl-2 and Bcl-XL result in the cell progressing towards apoptosis. Other proteins involved in the apoptotic pathway include cytochrome C and cysteineaspartic proteases (caspase) 3, caspase 8, and caspase 9. Caspases are a family of cysteine-dependent aspartate-directed proteases which play a critical role in the transduction of apoptotic signals [120, 121]. Cytochrome C is a protein normally localized in the mitochondria, but upon receiving intrinsic apoptotic signals, cytochrome C translocates to the cytosol where it interacts with apoptosis-activating factor 1 (Apaf-1) and caspase 9 which play a crucial role in the activation caspase 3 [122-124]. Activation of caspase 3 results in the apoptosis of the cell. Treatment of MOLT-4 cells with 9,10-PQ resulted in the decrease of Bcl-2 and Bcl-XL proteins and an increase in the Bax protein as well as an upregulation of cytosolic cytochrome C levels indicating mitochondrial damage and cell progression towards apoptosis [50]. Also, the membrane potential of the mitochondria was notably reduced. Increase in the activity of caspases 3, 8, and 9 was also observed. Time-dependent experiments were performed to understand the precise time caspases 3, 8, and 9 were activated. To trigger apoptosis caspase 3 should be activated after caspases 8 and 9. Based on the analysis caspase 3 activity increased by approximately 8-fold 8 hours after treatment with 9,10-PQ, whereas caspases 8 and 9 activity was increased by approximately 4-fold, the highest level, after 6 hours [50]. From this data, a possible mechanism involved in triggering apoptosis includes activation of caspase 8 upon exposure to 9,10-PQ. Activated caspase 8 eventually downregulates antiapoptotic Bcl-2 and Bcl-XL proteins and upregulation of the proapoptotic Bax protein. Increase in Bax activity causes Bax to translocate into the mitochondria causing a reduction in the mitochondrial membrane potential and increase in the permeability of the mitochondria, which ultimately results in the release of cytochrome C into the cytoplasm of the cell, triggering caspase activated apoptosis [50].

# 8. The Role of 9,10-PQ in Glutathione and Cellular Detoxification

A major component of cellular detoxification involves Phase II reactions which are responsible for increasing the hydrophilicity of the xenobiotic compounds to aid in the excretion process [125]. Phase II reactions are often referred to as conjugation reactions because of these reactions couple hydrophilic groups to xenobiotic compounds. One of the most studied Phase II coupling reactions involves glutathione conjugation. GSH is a tripeptide which acts as a critical component in the detoxification process of cells undergoing oxidative stress [126-128]. This thiol compound containing a reactive sulfhydryl group is a very powerful and important antioxidant. The enzyme responsible for this crucial reaction is referred to as glutathione transferase (GST), which adds glutathione to the parent xenobiotic compound either by direct addition or by replacement of an electrophilic substituent. Although GSH acts as a cofactor for GST, it can independently act as a free radical scavenger [126–128]. To understand the role of 9,10-PQ in GSH levels, it can be predicted that GSH levels will be low upon exposure to 9,10-PQ. Upon treatment of MOLT-4 cells with 9,10-PQ in a time-dependent manner GSH levels were significantly decreased inside the cells to 40 percent of the original concentration after 4 hours and after 24 hours GSH levels were approximately 10 percent of the original concentration [50]. GSH levels drop significantly when cells are undergoing oxidative stress as this mechanism is very vital in the antioxidant detoxification process. To further prove and confirm the depletion of GSH was indeed caused by 9,10-PQ induced oxidative stress and apoptosis, MOLT-4 cells were pretreated with varying concentrations of cell-permeable GSH monoethyl ester (NAC) to mimic the effects of GSH before being exposed to 5 uM concentrations of 9,10-PQ [50]. Pretreatment of cells with 1 and 2 mM NAC significantly increased cell viability, thereby protecting the cells against 9,10-PQ induced cell death. NAC also protected the cells against DNA fragmentation, although GSH was not directly involved in DNA repair. These results suggest the relative significance and importance of antioxidant compounds, namely, GSH, in the cellular detoxification process as well as maintaining the viability of cells.

# 9. The Role of 9,10-PQ in Cardiovascular Disease

Common types of the cardiovascular disease mainly include hypertension, coronary heart disease (CHD), stroke, rheumatic heart disease, congenital heart disease, and congestive heart failure (CHF)[129]. Despite great progress has been made in the management of CVDs, CVD do harm to human health because of the high morbidity and mortality. It has been shown that about 63% of premature deaths in adults (aged 15–69 years) and three-out-of-four of all adult deaths are mainly attributed to CVD [130]. It is shocking that, by 2030, over 23.3 million people will die annually from CVD[131].A large number of researches have shown that long-term exposure to fine particulate matter, known as PM2.5, can impact heart disease [132–144]. Both acute and chronic PM exposures can significantly increase the cardiovascular morbidity and mortality [133–144].

Previous studies demonstrated that 9,10-PQ can elicit adverse cardiovascular effects such as alterations in vascular

reactivity [48, 72, 82, 145-147] and aortic ring relaxation [53]. Exposure to diesel exhaust containing 9,10-PQ was found to diminish humans forearm blood flow through endotheliumdependent and -independent mechanisms [148]. In the animal experiment, it was found that gaseous components of diesel exhaust can enhance vasoconstriction and suppress vasodilation in septal coronary arteries of mice [146]. An animal study examined the effects of phenanthraquinone on eNOS activity, endothelium-dependent relaxation, and blood pressure [53]. Exposure of rats to phenanthraquinone (0.36 mmol/kg) via an intraperitoneal administration was shown to result in the significant elevation of blood pressure [53]. Administration of phenanthraquinone to rats also decreased the level of NO2<sup>-</sup>/NO3 in plasma. NO2<sup>-</sup>/NO3 has been considered to be an excellent indicator of NO production in vivo [53]. NO production was also suppressed by phenanthraquinone in membrane fraction of BAECs with a concentration-dependent manner by measuring citrulline formation from L-arginine [149]. Thus, these in vitro and in vivo studies demonstrated that 9,10-PQ can cause an impairment of endothelium-dependent vasorelaxation and are associated with cardiopulmonary-related diseases and mortality (Table 1) [48, 72, 82, 147].

#### 10. Conclusion

9,10-PQ is a quinone molecule found in air pollution abundantly in the DEP. This compound has been studied extensively and has been shown to develop cytotoxic effects both in vitro and in vivo. Cytotoxicity generated by 9,10-PQ has been proposed to be mediated by its cycling for ROS generation. Exposure to 9,10-PQ can activate apoptotic factors such as caspases. Understanding the molecular mechanisms involved in the ROS pathway can allow for the development of novel and effective treatment of 9,10-PQ-mediated toxicity.

#### Abbreviations

9,10-PQ:	9,10-phenanthrenequinone
DEP:	Diesel exhaust particles
ROS:	Reactive oxygen species
GSH:	Glutathione
PAHs:	Polycyclic aromatic hydrocarbons
PM:	Particulate matter
Cu,Zn-SOD:	Cu,Zn-superoxide dismutase
NQO1:	Quinone oxidoreductase
XR:	L-Xylulose reductase
9,10-PQH2:	9,10-dihydroxyphenanthrene
1,2-NQ:	1,2-naphthoquinone
1,4-NQ:	1,4-naphthoquinone
9,10-AQ:	9,10-anthraquinone
$H_2O_2$ :	Hydrogen peroxide
•OH:	Hydroxyl radicals
NADPH:	Nicotinamide adenine dinucleotide
	phosphate
MOLT-4 cells:	Acute lymphoblastic leukemia cells
DCF:	2', 7'-dichlorofluorescein
PEG-cat:	Polyethylene glycol catalase
BAEC:	Bovine aortic endothelial cells

Source	Concentrations/Doses of 9,10-PQ	Models (in vitro/in vivo)	Results/ mechanisms of action
Muraki et al. (2017) [62]	$0.1-30\ \mu M$	Human embryonic kidney (HEK) cells and alveolar A549 cells	The results showed that 9,10-PQ activated human TRPA1 via critical cysteine residues at 621 and 665 in the N-terminus of the channel.
Asahi et al. (2014) [63]	50 mg/kg for male rat	Urine samples from Male rats and human	In rat urine samples following 9,10-PQ exposure, the monoglucuronide of 9,10-dihydroxyphenanthrene (9,10-PQHG) was found to be a major metabolite. 9,10-PQHG also present in human urine.
Hatae et al. (2013) [43]	10 <i>µ</i> M	HCT-116 colon tumor cells and HL-60 promyelocytic leukemia cells	Effects of orthoquinone moiety in 9,10-phenanthrenequinone on apoptosis were studied in HCT-116 and HL-60 cells. Results showed that the loss of the cis-orthoquinone unit in 9,10-PQ abrogated its ability to induce apoptosis in HCT-116 and HL-60 cells suggesting that cis-orthoquinine is an essential unit for 9,10-PQ to induce tumor cells apoptosis
Koizumi et al. (2013) [64]	$10\mu M$ or $100\mu M$	In Vitro Enzyme reaction	Redox cycling of 9,10-PQ in the presence of dihydrolipoic acid can cause oxidative damage of Cu, ZnSOD, which are associated with decreased enzyme activity.
Toyooka et al. (2012) [45]	0-25µM	A549 and MCF7 cell lines	In low NQOI-expressing cells, N-acetyl-L-cysteine (NAC) significantly enhanced 9,10-PQ-mediated cytotoxicity and the formation of DNA double-strand breaks with phosphorylation of histone H2AX. In contrast, 9,10-PQ-mediated cytotoxicity and genotoxicity were suppressed in presence of NAC in high NQOI-expressing human adenocarcinoma cell lines, A549 and MCF7. The results suggested that dual effects of NAC on 9,10-PQ-mediated cytotoxicity and genotoxicity are dependent on the NQOI activity
Sugimoto et al. (2005) [65]	5-30µM	A549 human pulmonary epithelial cells	9,10-PQ induced apoptosis in A549 cells and the LC50 is $\sim 7\mu$ M. Furthermore, 9,10-PQ induced the formation of hydroxyl radical in the presence of iron, NADPH, and P450 reductase in A549 cells. Chelating iron provided protection against the PQ-induced cytotoxicity. Treatment of A549 cells with PQ caused a decrease in levels of Cu,Zn-superoxide dismutase (Cu,Zn-SOD) and heme oxygenase-1 (HO-1) indicating that 9,10-PQ exerted oxidative stress in human A549 cells via iron-mediated oxidative damase with down-regulation of Cu, Zn-SOD.

TABLE 1: Major 9,10-phenanthraquinone studies and mechanisms of action.

		TABLE 1: Contin	ued.
Source	Concentrations/Doses of 9,10-PQ	Models (in vitro/in vivo)	Results/ mechanisms of action
Rodriguez et al. (2008) [49]	Aerobic conditions: 2.5-20μM; Anaerobic conditions: 20-50μM	yeast Saccharomyces cerevisiae	9,10 PQ can inhibit yeast S. cerevisiae growth under both aerobic and anaerobic conditions. However, 9,10 PQ induced DNA deletions and point mutations only in the presence of oxygen, not anaerobic conditions.
Taguchi et al. (2007) [66]	1-20μM	A549 human pulmonary epithelial cells	The results showed that PQH <sub>2</sub> is the product of an NADPH-dependent two-electron reduction of 9,10-PQ. NADPH-dependent enzymes such as the AKRIC isozyme can mediate the two-electron reduction of 9,10-PQ to PQH <sub>2</sub> resulting in redox cycling with 9,10-PQ, which is associated with oxidative protein damage
Peters et al. (2007) [67]	$0-100\mu M$	JEG-3 trophoblast cell line	9,10-PQ induced a concentration-dependent decline in Alamar Blue (AB), suggesting an impairment to energy metabolism. 9,10-PQ-mediated cytotoxicity was dramatically enhanced by the addition of copper.
Milko et al. (2009) [68]	TSQ Classic mass spectrometer with a QOQ configuration chemical reaction, 9,10-PQ as a ligand	Gas-phase model complexes [(PQ)FeCl(CH(3)O)](+), [(pdnen)FeCl(CH(3)O)](+), and [(PQ)(phen)FeCl(CH(3)O)](+)	The results showed that there is an interaction between iron(III) and phenanthraquinone in the isolated complexes in the gas phase, which is is driven by the reduction of iron(III) to iron(II) and 9,10-phenanthraquinone to the semihydroquinone radical or semiquinolate.
Hiyoshi et al. (2005) [69]	0.01 nmole/animal	Male ICR mice	Administration of 9,10-PQ (2.1 ng/animal) via intratracheal route dramatically increased allergic airway inflammation in mice and allergen-specific production of IgG1 and IgF.
Matsunag et al. (2008) [50]	1-10µM	human acute T-lymphoblastic leukemia MOLT-4 cells	9,10-PQ generated ROS, depleted cellular glutathione and trigged apoptosis signaling including mitochondrial membrane dysfunction and activation of caspases and poly(ADP-ribose) polymerase. 9,10-PQ-mediated ROS generation and cytotoxicity were increased in an XR-transformed cell line. Furthermore 9,10-PQ-induced cell death was partially inhibited when cells were pretreated with XR-specific inhibitors followed by 9,10-PQ exposure. These results suggest that -Xylulose reductase is critically involved in 9,10-phenanthrenequinone-induced apoptosis in human T lymphoma cells

		TABLE 1: Continu	ied.
Source	Concentrations/Doses of 9,10-PQ	Models ( <i>in vitro/in vivo</i> )	Results/ mechanisms of action
KUMAGAI et al. (2001) [53]	In vitro (0.15-5 µM), in vivo (0.36 mmole/kg body weight)	Bovine aortic endothelial cells; Wistar rats (8–10 wk old)	9,10-PQ was found to significantly inhibit nitric oxide (NO) formation in bovine aortic endothelial cells with an IC(50) value of 0.6 $\mu$ M. 9,10-PQ also inhibited endothelium-dependent relaxation by nitroglycerin was not affected. Furthermore, intraperitoneal injection of 9,10-PQ (0.56 mmol/kg) to rats elevated blood pressure, plasma levels of stable NO metabolites, nitrite/nitrate compared to control group. These results suggested that 9,10-PQ is a potent inhibitory action on eNOS activity resulting in inhibition of NO-mediated vasorelaxation and elevation of blood pressure.
Taguchi et al. (2008) [70]	$1-50 \ \mu M$	Mouse primary hepatocytes	Exposure of human pulmonary epithelial A549 cells to 9,10-PQ resulted in a time-dependent formation of 9,10-PQH2 (9,10-PQHG) and. UGT1A6 found to be particularly involved in 9,10-PQHG formation. In cell-free systems, 9,10-PQ while not 9,10-PQHG, showed a quick thiol oxidation and oxygen consumption in the presence of dithiothreitol.
Oginuma et al. (2005) [71]	$10 \ \mu M$	Pig heart cytosol	9,10-PQ was found to be a potent inhibitor for the 4-benzoylpyridine (4-BP) reduction. and pig heart carbonyl reductase plays a critical role in superoxide generation via redox cycling of 9,10-PQ
Prisby et al. (2005)[32]	5 μM for In Vitro principal nutrient artery studies	Female and male Fischer-344 (F-344) rats	9,10-PQ diminished endothelium-dependent vasodilation of PNA in 14- and 24-mo-old rats of both genders. However, there was no change in femoral PNA in 6-mo-old male rats, suggesting that ovarian hormones play important role in the vascular endothelium at this stage of development.

BioMed Research International

MTB:	4-methyl-[1,2,3]-thiadiazole-5-carboxylic
	acid benzyloxy amide
NAC:	N-acetyl-L-cysteine
GO:	Glucose oxidase
NO:	Nitric oxide
PARP:	Poly (ADP-ribose) polymerase
Caspase:	Cysteine-aspartic proteases
GST:	Glutathione transferase
CHD:	Coronary heart disease
CHF:	Congestive heart failure
CVDs:	Cardiovascular diseases.

#### **Conflicts of Interest**

The authors declare that they have no conflicts of interest.

#### Acknowledgments

This work was supported in part by a grant from UNCG.

#### References

- W. Yang and S. T. Omaye, "Air pollutants, oxidative stress and human health," *Mutation Research - Genetic Toxicology and Environmental Mutagenesis*, vol. 674, no. 1-2, pp. 45–54, 2009.
- [2] B. Brunekreef and S. T. Holgate, "Air pollution and health," *The Lancet*, vol. 360, no. 9341, pp. 1233–1242, 2002.
- W. P. D. Logan, "Mortality from Fog in London, January, 1956," British Medical Journal, vol. 1, no. 4969, pp. 722–725, 1956.
- [4] W. Wu, D. B. Peden, R. McConnell, S. Fruin, and D. Diaz-Sanchez, "Glutathione-S-transferase M1 regulation of diesel exhaust particle-induced pro-inflammatory mediator expression in normal human bronchial epithelial cells," *Particle and Fibre Toxicology*, vol. 9, article 31, 2012.
- [5] Z. An, Y. Jin, J. Li, W. Li, and W. Wu, "Impact of Particulate Air Pollution on Cardiovascular Health," *Current Allergy and Asthma Reports*, vol. 18, no. 3, 2018.
- [6] N. Palacios, "Air pollution and Parkinson's disease Evidence and future directions," *Reviews on Environmental Health*, vol. 32, no. 4, pp. 303–313, 2017.
- [7] G. S. Ajmani, H. H. Suh, and J. M. Pinto, "Effects of ambient air pollution exposure on olfaction: A review," *Environmental Health Perspectives*, vol. 124, no. 11, pp. 1683–1693, 2016.
- [8] J. M. Samet and S. Gruskin, "Air pollution, health, and human rights," *The Lancet Respiratory Medicine*, vol. 3, no. 2, pp. 98–100, 2015.
- [9] X. Rao, J. Zhong, A. Maiseyeu et al., "CD36-dependent 7ketocholesterol accumulation in macrophages mediates progression of atherosclerosis in response to chronic air pollution exposure," *Circulation Research*, vol. 115, no. 9, pp. 770–780, 2014.
- [10] M. Kusha, S. Masse, T. Farid et al., "Controlled exposure study of air pollution and T-Wave alternans in volunteers without cardiovascular disease," *Environmental Health Perspectives*, vol. 120, no. 8, pp. 1157–1161, 2012.
- [11] M. E. Davis, F. Laden, J. E. Hart, E. Garshick, and T. J. Smith, "Economic activity and trends in ambient air pollution," *Environmental Health Perspectives*, vol. 118, no. 5, pp. 614–619, 2010.

- [12] M. Kampa and E. Castanas, "Human health effects of air pollution," *Environmental Pollution*, vol. 151, no. 2, pp. 362–367, 2008.
- [13] H. Yamawaki and N. Iwai, "Mechanisms underlying nano-sized air-pollution-mediated progression of atherosclerosis: Carbon black causes cytotoxic injury/inflammation and inhibits cell growth in vascular endothelial cells," *Circulation Journal*, vol. 70, no. 1, pp. 129–140, 2006.
- [14] D. Schwela, "Air Pollution and Health in Urban Areas," *Reviews on Environmental Health*, vol. 15, no. 1-2, pp. 13–42, 2000.
- [15] C. A. Pope III, "Epidemiology of fine particulate air pollution and human health: Biologic mechanisms and who's at risk?" *Environmental Health Perspectives*, vol. 108, no. 4, pp. 713–723, 2000.
- [16] J. A. Bukowski and D. Wartenberg, "An alternative approach for investigating the carcinogenicity of indoor air pollution: Pets as sentinels of environmental cancer risk," *Environmental Health Perspectives*, vol. 105, no. 12, pp. 1312–1319, 1997.
- [17] D. B. Menzel, "The toxicity of air pollution in experimental animals and humans: the role of oxidative stress," *Toxicology Letters*, vol. 72, no. 1–3, pp. 269–277, 1994.
- [18] M. Lippmann and P. J. Lioy, "Critical issues in air pollution epidemiology," *Environmental Health Perspectives*, vol. 62, pp. 243–258, 1985.
- [19] E. J. Esswein, M. Alexander-Scott, J. Snawder, and M. Breitenstein, "Measurement of area and personal breathing zone concentrations of diesel particulate matter (DPM) during oil and gas extraction operations, including hydraulic fracturing," *Journal of Occupational and Environmental Hygiene*, vol. 15, no. 1, pp. 63–70, 2018.
- [20] J. Bünger, J. Krahl, A. Munack et al., "Strong mutagenic effects of diesel engine emissions using vegetable oil as fuel," *Archives* of *Toxicology*, vol. 81, no. 8, pp. 599–603, 2007.
- [21] S. M. Toner, D. A. Sodeman, and K. A. Prather, "Single particle characterization of ultrafine and accumulation mode particles from heavy duty diesel vehicles using aerosol time-of-flight mass spectrometry," *Environmental Science & Technology*, vol. 40, no. 12, pp. 3912–3921, 2006.
- [22] J. Zhu, X.-L. Cao, R. Pigeon, and K. Mitchell, "Comparison of vehicle exhaust emissions from modified diesel fuels," *Journal* of the Air & Waste Management Association, vol. 53, no. 1, pp. 67–76, 2003.
- [23] R. L. Clifford, M. J. Jones, J. L. MacIsaac et al., "Inhalation of diesel exhaust and allergen alters human bronchial epithelium DNA methylation," *The Journal of Allergy and Clinical Immunol*ogy, vol. 139, no. 1, pp. 112–121, 2017.
- [24] C. Carlsten, A. Blomberg, M. Pui et al., "Diesel exhaust augments allergen-induced lower airway inflammation in allergic individuals: A controlled human exposure study," *Thorax*, vol. 71, no. 1, pp. 35–44, 2016.
- [25] A. Mastrofrancesco, M. Alfè, E. Rosato et al., "Proinflammatory effects of diesel exhaust nanoparticles on scleroderma skin cells," *Journal of Immunology Research*, vol. 2014, Article ID 138751, 9 pages, 2014.
- [26] K. E. Cosselman, R. M. Krishnan, A. P. Oron et al., "Blood pressure response to controlled diesel exhaust exposure in human subjects," *Hypertension*, vol. 59, no. 5, pp. 943–948, 2012.
- [27] K. A. Mattingly and C. M. Klinge, "Diesel exhaust particulate extracts inhibit transcription of nuclear respiratory factor-1 and cell viability in human umbilical vein endothelial cells," *Archives* of *Toxicology*, vol. 86, no. 4, pp. 633–642, 2012.

- [28] J. Li, P. Kanju, M. Patterson et al., "TRPV4-mediated calcium influx into human bronchial epithelia upon exposure to diesel exhaust particles," *Environmental Health Perspectives*, vol. 119, no. 6, pp. 784–793, 2011.
- [29] P. R. Reynolds, K. M. Wasley, and C. H. Allison, "Diesel particulate matter induces receptor for advanced glycation endproducts (RAGE) expression in pulmonary epithelial cells, and RAGE signaling influences NF-κB-mediated inflammation," *Environmental Health Perspectives*, vol. 119, no. 3, pp. 332–336, 2011.
- [30] J. Li, A. J. Ghio, S.-H. Cho, C. E. Brinckerhoff, S. A. Simon, and W. Liedtke, "Diesel exhaust particles activate the matrixmetalloproteinase-1 gene in human bronchial epithelial in a β-arrestin-dependent manner via activation of RAS," *Environmental Health Perspectives*, vol. 117, no. 3, pp. 400–409, 2009.
- [31] F. Anselme, S. Loriot, J.-P. Henry et al., "Inhalation of diluted diesel engine emission impacts heart rate variability and arrhythmia occurrence in a rat model of chronic ischemic heart failure," *Archives of Toxicology*, vol. 81, no. 4, pp. 299–307, 2007.
- [32] R. D. Prisby, J. Muller-Delp, M. D. Delp, and T. R. Nurkiewicz, "Age, gender, and hormonal status modulate the vascular toxicity of the diesel exhaust extract phenanthraquinone," *Journal* of Toxicology and Environmental Health, Part A. Current Issues, vol. 71, no. 7, pp. 464–470, 2008.
- [33] S. Malgorzata and P. Malgorzata, "Polycyclic aromatic hydrocarbons in ultrafine particles of diesel exhaust fumes – The use of ultrafast liquid chromatography," *Medycyna Pracy*, vol. 65, no. 5, pp. 601–608, 2014.
- [34] A. I. Totlandsdal, J. Øvrevik, R. E. Cochran et al., "The occurrence of polycyclic aromatic hydrocarbons and their derivatives and the proinflammatory potential of fractionated extracts of diesel exhaust and wood smoke particles," *Journal* of Environmental Science and Health, Part A: Toxic/Hazardous Substances and Environmental Engineering, vol. 49, no. 4, pp. 383–396, 2014.
- [35] A. I. Totlandsdal, J. I. Herseth, A. K. Bølling et al., "Differential effects of the particle core and organic extract of diesel exhaust particles," *Toxicology Letters*, vol. 208, no. 3, pp. 262–268, 2012.
- [36] H.-E. Wichmann, "Diesel exhaust particles," *Inhalation Toxicology*, vol. 19, no. 1, pp. 241–244, 2007.
- [37] R. D. Brown, R. C. Quardokus, N. A. Wasio et al., "Non-intuitive clustering of 9,10-phenanthrenequinone on Au(111)," *Beilstein Journal of Nanotechnology*, vol. 8, no. 1, pp. 1801–1807, 2017.
- [38] M. El-Maghrabey, N. Kishikawa, and N. Kuroda, "9,10-Phenanthrenequinone as a mass-tagging reagent for ultrasensitive liquid chromatography-tandem mass spectrometry assay of aliphatic aldehydes in human serum," *Journal of Chromatography A*, vol. 1462, pp. 80–89, 2016.
- [39] V. R. Kumar, N. Rajkumar, and S. Umapathy, "Solvatochromism of 9,10-phenanthrenequinone: An electronic and resonance Raman spectroscopic study," *The Journal of Chemical Physics*, vol. 142, no. 2, Article ID 024305, 2015.
- [40] S. Mallakpour, M. T. Behnamfar, M. Dinari, and H. Hadadzadeh, "Preparation of new fluorophore lanthanide complexes-Cloisite nanohybrids using the tricationic Pr(III), Gd(III) and Dy(III) complexes with 9,10-phenanthrenequinone," Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 137, pp. 1206–1212, 2015.
- [41] V. R. Kumar, N. Rajkumar, F. Ariese, and S. Umapathy, "Direct Observation of Thermal Equilibrium of Excited Triplet States of 9,10-Phenanthrenequinone. A Time-Resolved Resonance

Raman Study," *The Journal of Physical Chemistry A*, vol. 119, no. 40, pp. 10147–10157, 2015.

- [42] T. Matsunaga, Y. Morikawa, M. Haga et al., "Exposure to 9,10phenanthrenequinone accelerates malignant progression of lung cancer cells through up-regulation of aldo-keto reductase 1B10," *Toxicology and Applied Pharmacology*, vol. 278, no. 2, pp. 180–189, 2014.
- [43] N. Hatae, J. Nakamura, T. Okujima et al., "Effect of the orthoquinone moiety in 9,10-phenanthrenequinone on its ability to induce apoptosis in HCT-116 and HL-60 cells," *Bioorganic & Medicinal Chemistry Letters*, vol. 23, no. 16, pp. 4637–4640, 2013.
- [44] Z. Afrasiabi, P. Stovall, K. Finley et al., "Targeting Triple Negative Breast Cancer cells by N3-substituted 9, 10-Phenanthrenequinone thiosemicarbazones and their metal complexes," *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 114, pp. 114–119, 2013.
- [45] T. Toyooka, T. Shinmen, J. M. M. J. G. Aarts, and Y. Ibuki, "Dual effects of N-acetyl-l-cysteine dependent on NQO1 activity: Suppressive or promotive of 9,10-phenanthrenequinone-induced toxicity," *Toxicology and Applied Pharmacology*, vol. 264, no. 3, pp. 404–412, 2012.
- [46] T. Matsunaga, M. Hosogai, M. Arakaki, S. Endo, O. El-Kabbani, and A. Hara, "9,10-Phenanthrenequinone induces monocytic differentiation of U937 cells through regulating expression of aldo-keto reductase 1C3," *Biological & Pharmaceutical Bulletin*, vol. 35, no. 9, pp. 1598–1602, 2012.
- [47] T. Matsunaga, M. Haga, G. Watanabe et al., "9,10-Phenanthrenequinone promotes secretion of pulmonary aldo-keto reductases with surfactant," *Cell and Tissue Research*, vol. 347, no. 2, pp. 407–417, 2012.
- [48] T. Matsunaga, M. Arakaki, T. Kamiya et al., "Nitric oxide mitigates apoptosis in human endothelial cells induced by 9,10phenanthrenequinone: Role of proteasomal function," *Toxicol*ogy, vol. 268, no. 3, pp. 191–197, 2010.
- [49] C. E. Rodriguez, Z. Sobol, and R. H. Schiestl, "9,10-Phenanthrenequinone induces DNA deletions and forward mutations via oxidative mechanisms in the yeast Saccharomyces cerevisiae," *Toxicology in Vitro*, vol. 22, no. 2, pp. 296–300, 2008.
- [50] T. Matsunaga, T. Kamiya, D. Sumi, Y. Kumagai, B. Kalyanaraman, and A. Hara, "L-Xylulose reductase is involved in 9,10-phenanthrenequinone-induced apoptosis in human T lymphoma cells," *Free Radical Biology & Medicine*, vol. 44, no. 6, pp. 1191–1202, 2008.
- [51] J. Holt, S. Hothem, H. Howerton, R. Larson, and R. Sanford, "9,10-Phenanthrenequinone photoautocatalyzes its formation from phenanthrene, and inhibits biodegradation of naphthalene," *Journal of Environmental Quality*, vol. 34, no. 2, pp. 462– 468, 2005.
- [52] H. Shimada, M. Oginuma, A. Hara, and Y. Imamura, "9,10-Phenanthrenequinone, a component of diesel exhaust particles, inhibits the reduction of 4-benzoylpyridine and all-transretinal and mediates superoxide formation through its redox cycling in pig heart," *Chemical Research in Toxicology*, vol. 17, no. 8, pp. 1145–1150, 2004.
- [53] Y. Kumagai, T. Hayashi, T. Miyauchi et al., "Phenanthraquinone inhibits eNOS activity and suppresses vasorelaxation," *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, vol. 281, no. 1, pp. R25–R30, 2001.
- [54] B. Peterson and A. Saxon, "Global increases in allergic respiratory disease: the possible role of diesel exhaust particles. Annals of allergy," Annals of Allergy, Asthma & Immunology:

Official Publication of the American College of Allergy, Asthma, & Immunology, vol. 77, pp. 263–268; quiz 269-270, 1996.

- [55] J. O. Allen, N. M. Dookeran, K. Taghizadeh, A. L. Lafleur, K. A. Smith, and A. F. Sarofim, "Measurement of oxygenated polycyclic aromatic hydrocarbons associated with a size-segregated urban aerosol," *Environmental Science & Technology*, vol. 31, no. 7, pp. 2064–2070, 1997.
- [56] A. K. Cho, E. Di Stefano, Y. You et al., "Determination of four quinones in diesel exhaust particles, SRM 1649a, and atmospheric PM2.5," *Aerosol Science and Technology*, vol. 38, no. 1, pp. 68–81, 2004.
- [57] N. Kishikawa, M. Nakao, Y. Ohba, K. Nakashima, and N. Kuroda, "Concentration and trend of 9,10-phenanthrenequinone in airborne particulates collected in Nagasaki city, Japan," *Chemosphere*, vol. 64, no. 5, pp. 834–838, 2006.
- [58] S.-H. Kim, K.-M. Lim, J.-Y. Noh et al., "Doxorubicininduced platelet procoagulant activities: An important clue for chemotherapy-associated thrombosis," *Toxicological Sciences*, vol. 124, no. 1, Article ID kfr222, pp. 215–224, 2011.
- [59] E.-J. Kim, K.-M. Lim, K.-Y. Kim et al., "Doxorubicininduced platelet cytotoxicity: A new contributory factor for doxorubicin-mediated thrombocytopenia," *Journal of Thrombosis and Haemostasis*, vol. 7, no. 7, pp. 1172–1183, 2009.
- [60] A. K. Cho, E. Di Stefano, Y. You et al., "Determination of four quinones in diesel exhaust particles, SRM 1649a, and atmospheric PM2.5: Special Issue of Aerosol Science and Technologyon Findings from the Fine Particulate Matter Supersites Program," *Aerosol Science and Technology*, vol. 38, no. 1, pp. 68– 81, 2004.
- [61] D. Schuetzle, "Sampling of vehicle emissions for chemical analysis and biological testing," *Environmental Health Perspectives*, vol. 47, pp. 65–80, 1983.
- [62] K. Muraki, T. Sekine, Y. Ando et al., "An environmental pollutant, 9,10-phenanthrenequinone, activates human TRPA1 via critical cysteines 621 and 665," *Pharmacology Research and Perspectives*, vol. 5, no. 4, Article ID e00342, 2017.
- [63] M. Asahi, M. Kawai, T. Toyama et al., "Identification and quantification of in vivo metabolites of 9,10-phenanthrenequinone in human urine associated with producing reactive oxygen species," *Chemical Research in Toxicology*, vol. 27, no. 1, pp. 76– 85, 2014.
- [64] R. Koizumi, K. Taguchi, M. Hisamori, and Y. Kumagai, "Interaction of 9,10-phenanthraquinone with dithiol causes oxidative modification of Cu,Zn-superoxide dismutase (SOD) through redox cycling," *Journal of Toxicological Sciences*, vol. 38, no. 3, pp. 317–324, 2013.
- [65] R. Sugimoto, Y. Kumagai, Y. Nakai, and T. Ishii, "9,10-Phenanthraquinone in diesel exhaust particles downregulates Cu,Zn-SOD and HO-1 in human pulmonary epithelial cells: Intracellular iron scavenger 1,10-phenanthroline affords protection against apoptosis," *Free Radical Biology & Medicine*, vol. 38, no. 3, pp. 388–395, 2005.
- [66] K. Taguchi, S. Fujii, S. Yamano et al., "An approach to evaluate two-electron reduction of 9,10-phenanthraquinone and redox activity of the hydroquinone associated with oxidative stress," *Free Radical Biology & Medicine*, vol. 43, no. 5, pp. 789–799, 2007.
- [67] Z. J. Peters, J. A. Nykamp, K. Passaperuma et al., "Effect of copper on the cytotoxicity of phenanthrene and 9,10phenanthrenequinone to the human placental cell line, JEG-3," *Reproductive Toxicology*, vol. 23, no. 4, pp. 513–520, 2007.

- [68] P. Milko and J. Roithová, "Redox processes in the iron(III)/9,10phenanthraquinone system," *Inorganic Chemistry*, vol. 48, no. 24, pp. 11734–11742, 2009.
- [69] K. Hiyoshi, H. Takano, K.-I. Inoue et al., "Effects of phenanthraquinone on allergic airway inflammation in mice," *Clinical* & Experimental Allergy, vol. 35, no. 9, pp. 1243–1248, 2005.
- [70] K. Taguchi, M. Shimada, S. Fujii et al., "Redox cycling of 9,10-phenanthraquinone to cause oxidative stress is terminated through its monoglucuronide conjugation in human pulmonary epithelial A549 cells," *Free Radical Biology & Medicine*, vol. 44, no. 8, pp. 1645–1655, 2008.
- [71] M. Oginuma, H. Shimada, and Y. Imamura, "Involvement of carbonyl reductase in superoxide formation through redox cycling of adrenochrome and 9,10-phenanthrenequinone in pig heart," *Chemico-Biological Interactions*, vol. 155, no. 3, pp. 148– 154, 2005.
- [72] T. Matsunaga, K. Kamase, H. Takasawa et al., "Facilitation of 9,10-phenanthrenequinone-elicited neuroblastoma cell apoptosis by NAD(P)H:quinone oxidoreductase 1," *Chemico-Biological Interactions*, vol. 279, pp. 10–20, 2018.
- [73] H. Peng, D. Zhang, B. Pan, and J. Peng, "Contribution of hydrophobic effect to the sorption of phenanthrene, 9phenanthrol and 9, 10-phenanthrenequinone on carbon nanotubes," *Chemosphere*, vol. 168, pp. 739–747, 2017.
- [74] A. Honda, P. H. Chowdhury, S. Ito et al., "Synergic effects of 9,10-phenanthrenequinone and cadmium on pro-inflammatory responses in airway epithelial cells," *Environmental Toxicology* and Pharmacology, vol. 52, pp. 276–279, 2017.
- [75] P. Anitha, N. Chitrapriya, Y. J. Jang, and P. Viswanathamurthi, "Synthesis, characterization, DNA interaction, antioxidant and anticancer activity of new ruthenium(II) complexes of thiosemicarbazone/semicarbazone bearing 9,10-phenanthrenequinone," *Journal of Photochemistry and Photobiology B: Biology*, vol. 129, pp. 17–26, 2013.
- [76] D. Xiao, B. Pan, M. Wu, Y. Liu, D. Zhang, and H. Peng, "Sorption comparison between phenanthrene and its degradation intermediates, 9,10-phenanthrenequinone and 9-phenanthrol in soils/sediments," *Chemosphere*, vol. 86, no. 2, pp. 183–189, 2012.
- [77] M. K. Biswas, S. C. Patra, A. N. Maity, S.-C. Ke, N. D. Adhikary, and P. Ghosh, "Electronic structures of ruthenium and osmium complexes of 9,10-phenanthrenequinone," *Inorganic Chemistry*, vol. 51, no. 12, pp. 6687–6699, 2012.
- [78] T. Matsunaga, Y. Shinoda, Y. Inoue et al., "Aldo-keto reductase 1C15 as a quinone reductase in rat endothelial cell: Its involvement in redox cycling of 9,10-phenanthrenequinone," *Free Radical Research*, vol. 45, no. 7, pp. 848–857, 2011.
- [79] N. Kishikawa, M. Nakao, M. S. Elgawish, K. Ohyama, K. Nakashima, and N. Kuroda, "4-Carbomethoxybenzaldehyde as a highly sensitive pre-column fluorescence derivatization reagent for 9,10-phenanthrenequinone," *Talanta*, vol. 85, no. 1, pp. 809–812, 2011.
- [80] N. Kishikawa, H. Nakashima, K. Ohyama, K. Nakashima, and N. Kuroda, "Determination of 9, 10-phenanthrenequinone in airborne particulates by high-performance liquid chromatography with post-column fluorescence derivatization using 2aminothiophenol," *Talanta*, vol. 81, no. 4-5, pp. 1852–1855, 2010.
- [81] S. L. Pival, M. Klimacek, and B. Nidetzky, "The catalytic mechanism of NADH-dependent reduction of 9,10-phenanthrenequinone by Candida tenuis xylose reductase reveals plasticity in an aldo-keto reductase active site," *Biochemical Journal*, vol. 421, no. 1, pp. 43–49, 2009.

- [82] T. Matsunaga, M. Arakaki, T. Kamiya, S. Endo, O. El-Kabbani, and A. Hara, "Involvement of an aldo-keto reductase (AKRIC3) in redox cycling of 9,10-phenanthrenequinone leading to apoptosis in human endothelial cells," *Chemico-Biological Interactions*, vol. 181, no. 1, pp. 52–60, 2009.
- [83] F. Xie, M. A. Lampi, D. G. Dixon, and B. M. Greenberg, "Assessment of the toxicity of mixtures of nickel or cadmium with 9,10-phenanthrenequinone to Daphnia magna: Impact of a reactive oxygen-mediated mechanism with different redoxactive metals," *Environmental Toxicology and Chemistry*, vol. 26, no. 7, pp. 1425–1432, 2007.
- [84] F. Xie, S. A. Koziar, M. A. Lampi et al., "Assessment of the toxicity of mixtures of copper, 9,10-phenanthrenequinone, and phenanthrene to Daphnia magna: Evidence for a reactive oxygen mechanism," *Environmental Toxicology and Chemistry*, vol. 25, no. 2, pp. 613–622, 2006.
- [85] C. E. Rodriguez, J. M. Fukuto, K. Taguchi, J. Froines, and A. K. Cho, "The interactions of 9,10-phenanthrenequinone with glyceraldehyde-3- phosphate dehydrogenase (GAPDH), a potential site for toxic actions," *Chemico-Biological Interactions*, vol. 155, no. 1-2, pp. 97–110, 2005.
- [86] N. Kishikawa, M. Wada, Y. Ohba, K. Nakashima, and N. Kuroda, "Highly sensitive and selective determination of 9,10-phenanthrenequinone in airborne particulates using high-performance liquid chromatography with pre-column derivatization and fluorescence detection," *Journal of Chromatography* A, vol. 1057, no. 1-2, pp. 83–88, 2004.
- [87] Z. Yang, P. Wang, W. Zhang, and G. Zhu, "Sol-gel-derived carbon ceramic electrode containing 9,10-phenanthrenequinone, and its electrocatalytic activity toward iodate," *Fresenius' Journal* of Analytical Chemistry, vol. 371, no. 3, pp. 337–341, 2001.
- [88] G. Speier, Z. Tyeklár, P. Tóth et al., "Valence tautomerism and metal-mediated catechol oxidation for complexes of copper prepared with 9,10-phenanthrenequinone," *Inorganic Chemistry*, vol. 40, no. 22, pp. 5653–5659, 2001.
- [89] B. D'Autréaux and M. B. Toledano, "ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 10, pp. 813–824, 2007.
- [90] D. M. Miller, G. R. Buettner, and S. D. Aust, "Transition metals as catalysts of autoxidation reactions," *Free Radical Biology & Medicine*, vol. 8, no. 1, pp. 95–108, 1990.
- [91] S. K. Chaube, P. V. Prasad, S. C. Thakur, and T. G. Shrivastav, "Hydrogen peroxide modulates meiotic cell cycle and induces morphological features characteristic of apoptosis in rat oocytes cultured in vitro," *Apoptosis*, vol. 10, no. 4, pp. 863–874, 2005.
- [92] H. Miyazato, J. Taira, and K. Ueda, "Hydrogen peroxide derived from marine peroxy sesquiterpenoids induces apoptosis in HCT116 human colon cancer cells," *Bioorganic & Medicinal Chemistry Letters*, vol. 26, no. 19, pp. 4641–4644, 2016.
- [93] X. Lu, F. Kambe, X. Cao et al., "3β-hydroxysteroid-Δ24 reductase is a hydrogen peroxide scavenger, protecting cells from oxidative stress-induced apoptosis," *Endocrinology*, vol. 149, no. 7, pp. 3267–3273, 2008.
- [94] S. R. Räisänen, P. Lehenkari, M. Tasanen et al., "Carbonic anhydrase III protects cells from hydrogen peroxide-induced apoptosis," *The FASEB Journal*, vol. 13, no. 3, pp. 513–522, 1999.
- [95] V. Borutaite and G. C. Brown, "Nitric oxide induces apoptosis via hydrogen peroxide, but necrosis via energy and thiol depletion," *Free Radical Biology & Medicine*, vol. 35, no. 11, pp. 1457–1468, 2003.

- [96] H. Wei, Z. Li, S. Hu, X. Chen, and X. Cong, "Apoptosis of mesenchymal stem cells induced by hydrogen peroxide concerns both endoplasmic reticulum stress and mitochondrial death pathway through regulation of caspases, p38 and JNK," *Journal of Cellular Biochemistry*, vol. 111, no. 4, pp. 967–978, 2010.
- [97] S. Yang, Y.-H. Jan, V. Mishin, D. E. Heck, D. L. Laskin, and J. D. Laskin, "Diacetyl/1-Xylulose Reductase Mediates Chemical Redox Cycling in Lung Epithelial Cells," *Chemical Research in Toxicology*, vol. 30, no. 7, pp. 1406–1418, 2017.
- [98] J. Y. Abrams, R. J. Weber, M. Klein et al., "Associations between ambient fine particulate oxidative potential and cardiorespiratory emergency department visits," *Environmental Health Perspectives*, vol. 125, no. 10, Article ID 107008, 2017.
- [99] J. Yang, X. Zhao, M. Tang et al., "The role of ROS and subsequent DNA-damage response in PUMAinduced apoptosis of ovarian cancer cells," *Oncotarget*, vol. 8, no. 14, pp. 23492–23506, 2017.
- [100] C.-H. Lee and H.-S. Yu, "Role of mitochondria ROS and DNA damage in arsenic induced carcinogenesis," *Frontiers in Bioscience*, vol. 8, no. 2, pp. 312–320, 2016.
- [101] J. Li, K. Zhou, X. Meng et al., "Increased ROS generation and SOD activity in heteroplasmic tissues of transmitochondrial mice with A3243G mitochondrial DNA mutation," *Genetics and Molecular Research*, vol. 7, no. 4, pp. 1054–1062, 2008.
- [102] J. Zielonka and B. Kalyanaraman, ""ROS-generating mitochondrial DNA mutations can regulate tumor cell metastasis"-a critical commentary," *Free Radical Biology & Medicine*, vol. 45, no. 9, pp. 1217–1219, 2008.
- [103] K. Ishikawa, K. Takenaga, M. Akimoto et al., "ROS-generating mitochondrial DNA mutations can regulate tumor cell metastasis," *Science*, vol. 320, no. 5876, pp. 661–664, 2008.
- [104] Q. Luo, Y. Li, J. Deng, and Z. Zhang, "PARP-1 inhibitor sensitizes arsenic trioxide in hepatocellular carcinoma cells via abrogation of G2/M checkpoint and suppression of DNA damage repair," *Chemico-Biological Interactions*, vol. 226, pp. 12–22, 2015.
- [105] M.-F. Langelier, A. A. Riccio, and J. M. Pascal, "PARP-2 and PARP-3 are selectively activated by 5' phosphorylated DNA breaks through an allosteric regulatory mechanism shared with PARP-1," *Nucleic Acids Research*, vol. 42, no. 12, pp. 7762–7775, 2014.
- [106] L. Booth, N. Cruickshanks, T. Ridder, Y. Dai, S. Grant, and P. Dent, "PARP and CHK inhibitors interact to cause DNA damage and cell death in mammary carcinoma cells," *Cancer Biology & Therapy*, vol. 14, no. 5, pp. 458–465, 2013.
- [107] A. Bürkle, "DNA repair and PARP in aging," *Free Radical Research*, vol. 40, no. 12, pp. 1295–1302, 2006.
- [108] P. Davidovich, C. J. Kearney, and S. J. Martin, "Inflammatory outcomes of apoptosis, necrosis and necroptosis," *Biological Chemistry*, vol. 395, no. 10, pp. 1163–1171, 2014.
- [109] K. Wang, "Molecular mechanisms of liver injury: apoptosis or necrosis," *Experimental and Toxicologic Pathology*, vol. 66, no. 8, pp. 351–356, 2014.
- [110] C. Günther, H. Neumann, M. F. Neurath, and C. Becker, "Apoptosis, necrosis and necroptosis: cell death regulation in the intestinal epithelium," *Gut*, vol. 62, no. 7, pp. 1062–1071, 2013.
- [111] R. S. Fernandes and T. G. Cotter, "Apoptosis or necrosis: Intracellular levels of glutathione influence mode of cell death," *Biochemical Pharmacology*, vol. 48, no. 4, pp. 675–681, 1994.
- [112] M. P. Kashyap, A. K. Singh, V. Kumar et al., "Monocrotophos induced apoptosis in PC12 cells: Role of xenobiotic metabolizing cytochrome P450S," *PLoS ONE*, vol. 6, no. 3, Article ID e17757, 2011.

- [113] D. E. Frigo, K. A. Vigh, A. P. Struckhoff et al., "Xenobioticinduced TNF-α expression and apoptosis through the p38 MAPK signaling pathway," *Toxicology Letters*, vol. 155, no. 2, pp. 227–238, 2005.
- [114] M. Pallardy, A. Biola, H. Lebrec, and J. Bréard, "Assessment of apoptosis in xenobiotic-induced immunotoxicity," *Methods: A Companion to Methods in Enzymology*, vol. 19, no. 1, pp. 36–47, 1999.
- [115] M. E. Soriano and L. Scorrano, "Traveling Bax and forth from mitochondria to control apoptosis," *Cell*, vol. 145, no. 1, pp. 15–17, 2011.
- [116] Y. Wang, B. Zhang, X. Peng, M. Perpetua, and B. G. Harbrecht, "Bcl-xL prevents staurosporine-induced hepatocyte apoptosis by restoring protein kinase B/mitogen-activated protein kinase activity and mitochondria integrity," *Journal of Cellular Physiology*, vol. 215, no. 3, pp. 676–683, 2008.
- [117] Y. Tsujimoto, "Role of Bcl-2 family proteins in apoptosis: apoptosomes or mitochondria?" *Genes to Cells*, vol. 3, no. 11, pp. 697–707, 1998.
- [118] D. M. Finucane, E. Bossy-Wetzel, N. J. Waterhouse, T. G. Cotter, and D. R. Green, "Bax-induced caspase activation and apoptosis via cytochrome c release from mitochondria is inhibitable by Bcl-xL," *The Journal of Biological Chemistry*, vol. 274, no. 4, pp. 2225–2233, 1999.
- [119] K. G. Wolter, Y.-T. Hsu, C. L. Smith, A. Nechushtan, X.-G. Xi, and R. J. Youle, "Movement of Bax from the cytosol to mitochondria during apoptosis," *The Journal of Cell Biology*, vol. 139, no. 5, pp. 1281–1292, 1997.
- [120] Y. Huang, C. Kuo, K. Lu et al., "18α-Glycyrrhetinic Acid Induces Apoptosis of HL-60 Human Leukemia Cells through Caspasesand Mitochondria-Dependent Signaling Pathways," *Molecules*, vol. 21, no. 7, p. 872, 2016.
- [121] J.-M. Hwang, H.-C. Kuo, T.-H. Tseng, J.-Y. Liu, and C.-Y. Chu, "Berberine induces apoptosis through a mitochondria/caspases pathway in human hepatoma cells," *Archives of Toxicology*, vol. 80, no. 2, pp. 62–73, 2006.
- [122] R. A. Kirkland and J. L. Franklin, "Bax and caspases regulate increased production of mitochondria-derived reactive species in neuronal apoptosis: LACK of A role for depletion of cytochrome c from the mitochondrial electron transport chain," *Biochemistry and Biophysics Reports*, vol. 4, pp. 158–168, 2015.
- [123] H. El-Hassan, K. Anwar, P. Macanas-Pirard et al., "Involvement of mitochondria in acetaminophen-induced apoptosis and hepatic injury: Roles of cytochrome c, Bax, Bid, and caspases," *Toxicology and Applied Pharmacology*, vol. 191, no. 2, pp. 118–129, 2003.
- [124] S. Létuvé, A. Druilhe, M. Grandsaigne, M. Aubier, and M. Pretolani, "Involvement of caspases and of mitochondria in Fas ligation-induced eosinophil apoptosis: Modulation by interleukin-5 and interferon-γ," *Journal of Leukocyte Biology*, vol. 70, no. 5, pp. 767–775, 2001.
- [125] V. Roncalli, M. C. Cieslak, Y. Passamaneck, A. E. Christie, and P. H. Lenz, "Glutathione S-transferase (GST) gene diversity in the crustacean Calanus finmarchicus - Contributors to cellular detoxification," *PLoS ONE*, vol. 10, no. 5, Article ID e0123322, 2015.
- [126] F. J. Sánchez-Gómez, B. Díez-Dacal, E. García-Martín, J. A. G. Agúndez, M. A. Pajares, and D. Pérez-Sala, "Detoxifying enzymes at the cross-roads of inflammation, oxidative stress, and drug hypersensitivity: Role of glutathione transferase P1-1 and aldose reductase," *Frontiers in Pharmacology*, vol. 7, article no. 237, 2016.

- [127] K. D. Tew and D. M. Townsend, "Glutathione-S-transferases as determinants of cell survival and death," *Antioxidants & Redox Signaling*, vol. 17, no. 12, pp. 1728–1737, 2012.
- [128] A. M. Schimel, L. Abraham, D. Cox et al., "N-acetylcysteine amide (NACA) prevents retinal degeneration by up-regulating reduced glutathione production and reversing lipid peroxidation," *The American Journal of Pathology*, vol. 178, no. 5, pp. 2032–2043, 2011.
- [129] Z. Jia, P. Nallasamy, D. Liu et al., "Luteolin protects against vascular inflammation in mice and TNF-alpha-induced monocyte adhesion to endothelial cells via suppressing IKBα/NF-κB signaling pathway," *The Journal of Nutritional Biochemistry*, vol. 26, no. 3, pp. 293–302, 2015.
- [130] M. Sardarinia, S. Akbarpour, M. Lotfaliany et al., "Risk factors for incidence of cardiovascular diseases and all-cause mortality in a middle eastern population over a decade follow-up: Tehran lipid and glucose study," *PLoS ONE*, vol. 11, no. 12, Article ID e0167623, 2016.
- [131] C. D. Mathers and D. Loncar, "Projections of global mortality and burden of disease from 2002 to 2030," *PLoS Medicine*, vol. 3, no. 11, article e442, 2006.
- [132] Y. Wang, Z. Jia, and D. Liu, "Advances in insulin-like growth factor binding proteins and skeleton development and bone tissue regeneration," *Zhonghua Kou Qiang Yi Xue Za Zhi*, vol. 49, no. 7, pp. 444–446, 2014.
- [133] M. S. Loop, L. A. McClure, E. B. Levitan, M. Z. Al-Hamdan, W. L. Crosson, and M. M. Safford, "Fine particulate matter and incident coronary heart disease in the REGARDS cohort," *American Heart Journal*, vol. 197, pp. 94–102, 2018.
- [134] A. M. Neophytou, E. M. Noth, S. Liu et al., "Ischemic heart disease incidence in relation to fine versus total particulate matter exposure in a U.S. aluminum industry cohort," *PLoS ONE*, vol. 11, no. 6, Article ID e0156613, 2016.
- [135] S. Costello, A. M. Neophytou, D. M. Brown et al., "Incident Ischemic Heart Disease after Long-Term Occupational Exposure to Fine Particulate Matter: Accounting for 2 Forms of Survivor Bias," *American Journal of Epidemiology*, vol. 183, no. 9, pp. 861–868, 2016.
- [136] M.-S. Lee, K.-D. Eum, E. G. Rodrigues et al., "Effects of Personal Exposure to Ambient Fine Particulate Matter on Acute Change in Nocturnal Heart Rate Variability in Subjects Without Overt Heart Disease," *American Journal of Cardiology*, vol. 117, no. 1, pp. 151–156, 2016.
- [137] T. Siponen, T. Yli-Tuomi, M. Aurela et al., "Source-specific fine particulate air pollution and systemic inflammation in ischaemic heart disease patients," *Occupational and Environmental Medicine*, vol. 72, no. 4, pp. 277–283, 2015.
- [138] C. A. Pope, "Ischaemic heart disease and fine particulate air pollution," *Heart*, vol. 101, no. 4, pp. 248-249, 2015.
- [139] W. Xie, G. Li, D. Zhao et al., "Relationship between fine particulate air pollution and ischaemic heart disease morbidity and mortality," *Heart*, vol. 101, no. 4, pp. 257–263, 2015.
- [140] R. C. Puett, J. E. Hart, J. D. Yanosky et al., "Chronic fine and coarse particulate exposure, mortality, and coronary heart disease in the Nurses' Health Study," *Environmental Health Perspectives*, vol. 117, no. 11, pp. 1697–1701, 2009.
- [141] C. A. Pope III, J. B. Muhlestein, H. T. May, D. G. Renlund, J. L. Anderson, and B. D. Horne, "Ischemic heart disease events triggered by short-term exposure to fine particulate air pollution," *Circulation*, vol. 114, no. 23, pp. 2443–2448, 2006.
- [142] J. H. Sullivan, A. B. Schreuder, C. A. Trenga et al., "Association between short term exposure to fine particulate matter and

heart rate variability in older subjects with and without heart disease," *Thorax*, vol. 60, no. 6, pp. 462–466, 2005.

- [143] J. Sullivan, N. Ishikawa, L. Sheppard, D. Siscovick, H. Checkoway, and J. Kaufman, "Exposure to ambient fine particulate matter and primary cardiac arrest among persons with and without clinically recognized heart disease," *American Journal* of *Epidemiology*, vol. 157, no. 6, pp. 501–509, 2003.
- [144] J. Pekkanen, A. Peters, G. Hoek et al., "Particulate air pollution and risk of ST-segment depression during repeated submaximal exercise tests among subjects with coronary heart disease: the exposure and risk assessment for fine and ultrafine particles in ambient air (ULTRA) study," *Circulation*, vol. 106, no. 8, pp. 933– 938, 2002.
- [145] C. S. Hansen, M. Sheykhzade, P. Møller et al., "Diesel exhaust particles induce endothelial dysfunction in apoE-/- mice," *Toxicology and Applied Pharmacology*, vol. 219, no. 1, pp. 24–32, 2007.
- [146] M. J. Campen, N. S. Babu, G. A. Helms et al., "Nonparticulate components of diesel exhaust promote constriction in coronary arteries from ApoE-/- mice," *Toxicological Sciences*, vol. 88, no. 1, pp. 95–102, 2005.
- [147] H.-M. Lo, T.-L. Hwang, and W.-B. Wu, "A Phenanthrene Derivative, 5,7-Dimethoxy-1,4-Phenanthrenequinone, Inhibits Cell Adhesion Molecule Expression and Migration in Vascular Endothelial and Smooth Muscle Cells," *Pharmacology*, vol. 99, no. 5-6, pp. 291–302, 2017.
- [148] N. L. Mills, H. Törnqvist, S. D. Robinson et al., "Diesel exhaust inhalation causes vascular dysfunction and impaired endogenous fibrinolysis," *Circulation*, vol. 112, no. 25, pp. 3930– 3936, 2005.
- [149] X. P. Zhou, P. Bai, W. J. Zhang et al., "Genetic data of two STR loci D4S2367, D20S601 in Chinese population," *Journal of Forensic Sciences*, vol. 49, no. 6, p. 1380, 2004.