

Notes from the Field

Identification and Evolution of the First Isolated *Brucella abortus* Strain from a Human Brucellosis Case — Tianjin Municipality, China, August 2024

Xin Gao^{1,2}; Wei Zhang^{1,2}; Guangwen Liu^{1,2}; Wen Li^{1,2}; Aiping Yu^{1,2}; Yuan Wang^{1,2}; Xiaoqing Fan^{1,2}; Xiaohui Lu^{1,2}; Xuan Chen^{1,2}; Xiaoyan Li^{1,2,#}

Brucellosis remains a significant global zoonosis with profound negative implications for human health and economic systems worldwide (1). The disease is caused by Gram-negative, facultative intracellular bacteria belonging to the genus *Brucella* (2). Currently, 13 recognized *Brucella* species exist, with *B. melitensis*, *B. abortus*, and *B. suis* representing the predominant human pathogens (3). Human infection with *Brucella* spp. typically occurs through consumption of contaminated animal products or direct contact with infected animals (4).

In August 2024, a 71-year-old male farmer from Baodi District, Tianjin Municipality, China presented with persistent fever, excessive sweating, muscle and joint pain, and fatigue. The patient reported regularly entering a sheep pen without protective equipment approximately every 7 days, though he had no history of travel to known brucellosis-endemic regions. A *Brucella* strain, designated “BD002,” was isolated from the patient, confirming the diagnosis of brucellosis. Epidemiological investigation suggested that the sheep in the patient’s care were the likely source of infection.

For isolation of the *Brucella* strain, 500 µL of blood sample was inoculated onto a blood agar plate and incubated at 37 °C with 5% carbon dioxide for 72 hours, yielding off-white single colonies. The bacterial suspension was subsequently prepared and inactivated for DNA extraction and polymerase chain reaction (PCR) analysis. The brucella cell surface protein 31 polymerase chain reaction (BCSP31-PCR) test for genus identification revealed the characteristic 224-bp DNA band, while the abortus, melitensis, ovis, and suis polymerase chain reaction (AMOS-PCR) species-identification test yielded no detectable DNA bands (5). Notably, AMOS-PCR can identify *B. abortus* biovars 1, 2, and 4, *B. melitensis*, and *B. ovis*, but does not encompass all biovars of *Brucella* spp.

To further characterize strain BD002, we conducted both multi-locus sequence typing (MLST) and next-

generation sequencing (NGS) through GENEWIZ Co. and BGI-Shenzhen, China, respectively. For MLST, 9 *Brucella* housekeeping genes were amplified and subjected to Sanger sequencing, which identified BD002 as sequence type 2 according to the PubMLST database. NGS analysis generated 23 contigs using SPAdes v3.7, with a total genome length of 3,252,744 base pairs (6). The assembled genomic sequences have been deposited in the National Genomics Data Center (Accession No. GWHFJIX000000000.1) (7). We conducted phylogenetic analysis using the BD002 genome alongside 386 *B. abortus* genomic sequences from the NCBI Genome database, employing Snippy v4.6 and RAxML-ng v1.2.2 with strain 544 (Accession No. GCA_000369945.1) as the reference sequence for core-genome single-nucleotide polymorphism analysis (8). The resulting maximum likelihood phylogenetic tree was visualized using R v4.3.1 and *ggtree* v3.10.1 (9). Phylogenetic analysis revealed 4 distinct clades, with BD002 clustering within Clade B (Figure 1A). Notably, BD002 formed a sub-clade with strains from other Chinese provincial-level administrative divisions, including Inner Mongolia, Jilin, Heilongjiang, Beijing, Hebei, and Shanxi, as well as strains from Russia, Georgia, Mongolia, and other countries (Figure 1B).

Human brucellosis in Tianjin affects approximately 200 individuals annually, with *B. melitensis* identified as the predominant causative agent. This study documents the first isolation of *B. abortus* from a human case in Tianjin, indicating the circulation of multiple *Brucella* species within the region.

Compared with traditional AMOS-PCR methodology, MLST and NGS techniques provide superior accuracy in *Brucella* spp. identification and offer substantially higher resolution for phylogenetic analysis. These advanced molecular approaches should be fully integrated into routine etiological surveillance of human brucellosis.

Additionally, particular attention should be directed




FIGURE 1. The ML phylogenetic tree of *B. abortus* genomes based on cg-SNP. (A) The ML phylogenetic tree was constructed using the genomic sequence of strain BD002 integrated with 386 additional *B. abortus* genomic sequences and associated metadata obtained from the NCBI genome database. (B) A detailed view of the subclade within clade B containing strain BD002 is presented, with countries of strain origin indicated by differently colored circles. The BD002 strain is specifically highlighted with a red circle.

Note: For (A), the analysis revealed four distinct worldwide clades of *B. abortus* strains, highlighted in orange. The continental origin, host species, and sequence types of the strains are represented by differently colored squares. Abbreviation: ML=maximum likelihood; cg-SNP=core-genome single-nucleotide polymorphism.

toward livestock handlers who neglect to appropriate personal protective equipment and fail to implement adequate disinfection and hygiene practices following occupational exposure (10).

doi: 10.46234/ccdcw2025.067

Corresponding author: Xiaoyan Li, xiaoyanli1291@163.com.

¹ Tianjin Centers for Disease Control and Prevention, Tianjin, China;

² Tianjin Key Laboratory of Pathogenic Microbiology of Infectious Disease, Tianjin Centers for Disease Control and Prevention, Tianjin, China.

Copyright © 2025 by Chinese Center for Disease Control and Prevention. All content is distributed under a Creative Commons Attribution Non Commercial License 4.0 (CC BY-NC).

Submitted: December 30, 2024

Accepted: March 13, 2025

Issued: March 21, 2025

REFERENCES

- Franc KA, Kreck RC, Häslér BN, Arenas-Gamboa AM. Brucellosis remains a neglected disease in the developing world: a call for interdisciplinary action. *BMC Public Health* 2018;18(1):125. <https://doi.org/10.1186/s12889-017-5016-y>.
- Suárez-Esquivel M, Chaves-Olarte E, Moreno E, Guzmán-Verri C. *Brucella* genomics: macro and micro evolution. *Int J Mol Sci* 2020;21(20):7749. <https://doi.org/10.3390/ijms21207749>.
- An CH, Nie SM, Luo BY, Zhou DJ, Wang WJ, Sun YX, et al. Multiple *Brucella melitensis* lineages are driving the human brucellosis epidemic in Shaanxi Province, China: evidence from whole genome sequencing-based analysis. *Front Cell Infect Microbiol* 2024;14:1452143. <https://doi.org/10.3389/fcimb.2024.1452143>.
- Qureshi KA, Parvez A, Fahmy NA, Abdel Hady BH, Kumar S, Ganguly A, et al. Brucellosis: epidemiology, pathogenesis, diagnosis and treatment-a comprehensive review. *Ann Med* 2023;55(2):2295398. <https://doi.org/10.1080/07853890.2023.2295398>.
- Abd Eltawab A, El-Hofy F, Hamdy M, Moustafa S, Soliman E, Ahmed W, et al. Isolation and molecular identification of *Brucella* spp. in bovine herds kept at householders in the Delta region of Egypt by MALDI-TOF and AMOS-PCR. *Vet Ital* 2020;56(4):297 – 300. <https://doi.org/10.12834/VetIt.1980.10596.3>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012;19(5):455 – 77. <https://doi.org/10.1089/cmb.2012.0021>.
- CNCB-NGDC Members and Partners. Database Resources of the National Genomics Data Center, China National Center for Bioinformatics in 2024. *Nucleic Acids Res* 2024;52(D1):D18 – 32. <https://doi.org/10.1093/nar/gkad1078>.
- Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAXML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 2019;35(21):4453 – 5. <https://doi.org/10.1093/bioinformatics/btz305>.
- Xu SB, Li L, Luo X, Chen MJ, Tang WL, Zhan L, et al. *Ggtree*: a serialized data object for visualization of a phylogenetic tree and annotation data. *iMeta* 2022;1(4):e56. <https://doi.org/10.1002/imt2.56>.
- Lounes N, Yahiaoui D, Taftaf D, Zenia S. A survey on the occupational exposure of veterinarians to brucellosis in Algeria. *Ger J Microbiol* 2022;2(2):28 – 35. <https://doi.org/10.51585/gjm.2022.2.0017>.