

Evaluating the Relationship between Competition and Productivity within a Native Grassland

Jonathan A. Bennett*, James F. Cahill Jr.

Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

Abstract

Ideas about how plant competition varies with productivity are rooted in classic theories that predict either increasing (Grime) or invariant (Tilman) competition with increasing productivity. Both predictions have received experimental support, although a decade-old meta-analysis supports neither. Attempts to reconcile the conflicting predictions and evidence include: expanding the theory to include other conditions (e.g. stress gradient hypothesis), development of indices to differentiate either the 'intensity' or 'importance' of competition, a focus on resource supply and demand, and explicit recognition that both growth and survival may exhibit different relationships with productivity. To determine which of these theories accurately predict how competition varies with productivity within a native grassland site, we estimated competitive intensity and relative competitive importance using 22 species across the range of productivity naturally occurring within that site. Plant performance was measured as survival and size with and without neighbours and the local environment was quantified according to variability in standing crop, gross water supply, and net water supply. On average, neighbours weakly facilitated seedling survival, but strongly reduced seedling growth. For both seedling survival and growth, relative competitive importance and competitive intensity declined with some measure of productivity; neighbour effects on survival declined with standing crop, while effects on growth declined with gross water supply. These results add to the growing evidence that plant-plant interactions vary among life history components with different life history components contingent upon separate environmental factors. Although the range of productivity measured in this study was not large, our results do not support the theories of Grime or Tilman. However, our results are consistent with the meta-analysis and parts of other theories, although no single theory is capable of explaining the entirety of these results. This suggests that, at least in moderately productive grasslands, new theory needs to be developed.

Citation: Bennett JA, Cahill JF Jr (2012) Evaluating the Relationship between Competition and Productivity within a Native Grassland. PLoS ONE 7(8): e43703. doi:10.1371/journal.pone.0043703

Editor: Han Y.H. Chen, Lakehead University, Canada

Received: May 30, 2012; **Accepted:** July 23, 2012; **Published:** August 21, 2012

Copyright: © 2012 Bennett, Cahill Jr. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by a NSERC Discovery Grant to JFC Jr. and a NSERC Post-Graduate Scholarship to JAB. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Co-author James F. Cahill Jr. is a PLoS ONE Editorial board member. This does not alter the authors' adherence to all the PLoS ONE policies on sharing data and materials.

* E-mail: jon.bennett@ualberta.ca

Introduction

Plant competition is regularly seen as a factor influencing the structure of natural communities. However, the intensity of competition is not constant through space nor time [1,2,3,4,5], and understanding the factors that influence its intensity is critical to understanding the assembly of plant communities. One factor that plant ecologists have long focused on is the relationship between plant competition and productivity. This issue has been studied for so long and so intensely that it is a common topic in introductory ecology textbooks [6,7,8,9].

Textbook authors often present competition-productivity relationships in the context of the theories of Grime [10,11,12], Newman [13], and Tilman [3,14]. Many aspects of the disagreements amongst these authors are presented elsewhere [15,16,17]. In brief, Grime [10,11,12] asserted that competition and competitive exclusion is only important in structuring communities at high productivity, with "stress" more important at low productivity. Newman [13] suggested that competition was strong at both low and high productivity, though it switched from root to shoot competition as resources increased. However, Newman (1973) agreed that it was likely that only shoot

competition would reduce species diversity, a prediction consistent with later experimental work [18,19]. Tilman [3,14] also suggested the intensity of competition would be invariant along productivity gradients, but expected competition to be an important structuring force for communities at all levels of productivity [3]. Thus these sets of theories make separate predictions about when competition will be strong, and when it will be important for structuring communities along productivity gradients (Table 1).

Debate and data focused on competition-productivity (C-P) relationships has appeared in the scientific literature for over twenty years [1,16,17,20,21,22,23,24,25,26,27,28,29,30,31], including experimental efforts which have spanned continents [32]. One of the first meta-analyses in ecology was published by Goldberg and colleagues [1] focused on this issue. They found the intensity of competition for plant survival and growth *declined* with productivity; an outcome that supports none of the theories described above. Nonetheless, these unexpected and synthetic empirical results do not appear with the same frequency as the original theories in modern textbooks, even though another meta-analysis found similar results [33]. Despite the relative commonness of competition declining with productivity, these results are

Table 1. C–P relationship predicted for each major theory and the Goldberg meta-analysis.

Response metric	Gradient	Grime	SGH	Newman	Tilman	Goldberg	Davis	This study
Intensity	Standing crop/Gross supply	+	+	0	0	–	0/–	–
	Net supply						–	0
Competitive frequency	Standing crop/Gross supply		+					–/0
Relative importance	Standing crop/Gross supply	+	+			–		–
Demographic importance	Standing crop/Gross supply	+	+	+	0	–		

Note: Predictions are based on Grime [10,11,12], Bertness and Callaway [37], Maestre *et al.* [40], Newman [13], Tilman [3,14], Goldberg *et al.* [1], and Davis *et al.* [5]. Cells containing a + mean that we expect increasing competition along the gradient, cells with a 0 mean that we expect a non-significant relationship, and cells with a – mean we expect a negative relationship. If a cell is left blank, then that particular metric or gradient does not apply to that theory. The column labelled this study refers to our findings and will be explained further in the results and discussion.
doi:10.1371/journal.pone.0043703.t001

often disputed [34,35] and Grime and Tilman typically remain the starting point for new theory on this topic [16,36].

One such expansion of Grime’s theory, the stress-gradient hypothesis (SGH), relates facilitation to stress [37,38,39], where the level of stress within a community is often approximated by measuring productivity [15,40] and facilitation is measured using metrics similar to those used to study competition. Thus, in practice, SGH studies are empirically identical to C–P studies, although the research focus is more often on the outcome of pairwise interactions rather than effects on community structure [40]. SGH is often only applied to low productivity areas, with the primary prediction being that facilitation should be most common [37,40], intense [39] or important [40,41] at high to intermediate stress and thus that competition should be lowest in unproductive areas. Attempts have been made to apply this hypothesis to more productive areas [42], including an additional, yet little explored, aspect of the initial SGH which predicts that associational defences can cause facilitation to increase with productivity when herbivore pressure is intense [37]. Thus the predictions of SGH are multifaceted and can be consistent with the predictions of Grime, Newman, Tilman, or the results presented by Goldberg, depending on the range of productivity encountered.

Alternate hypotheses regarding the relationship between competition and productivity focus not on the biomass of neighbours, but on the supply and demand of resources [5,43]. In these theories, it is either difference between supply and demand [5] or the ratio of supply and demand [43] that determines the outcome of competitive interactions, with competition declining as either net resource supply (supply - demand) or the ratio of supply to demand increases. Thus in both these cases, competition can decrease with productivity. Although these theories are rooted in the foundations of plant ecology [44], they do not receive as much attention as theories relating to the work of Grime and Tilman.

C–P relationships can be expected to vary as a function of many factors. For example, plant survival and growth may respond differentially to the presence of neighbours [40,45,46,47], and may have differential effects on species exclusion [45]. These differences are rarely addressed by theory [40,48], though they have been addressed empirically [1,5,49]. Additionally, theory predicts different relationships for competitive *intensity* and *importance* [15,16,17,20,30,40]. Intensity is typically defined as the absolute magnitude of an effect [17,50], while importance has highly variable and contentious points of view among researchers. For this manuscript, we will measure importance as the magnitude of competition relative to maximum plant performance [15,17,51]. We recognize competitive importance can also be viewed in terms of the ultimate effects of competition on population growth rates

[52], competitive exclusion and community structure [53,54]. When differentiating between these definitions of importance, we will refer to them as relative and demographic importance respectively. However, demographic importance is difficult to measure in a perennial plant community given the long life spans of some plants and as such we cannot test them with the data at hand [but see 53 for a potential method of looking at community consequences of competition].

Finally, though C–P relationships are presented as an aspect of a community, they may be highly variable among species within a community. To date, most experimental studies of competition have used one or a small number of *phytometers*, species intended to be representative of the entire community [5,15,23,26,55], with exceptions including studies such as Wilson and Tilman [29] with eight and Callaway *et al.* [56] with more than sixty. What is lacking are experimental studies with a large number of species measuring multiple responses within a single community - allowing a true test of the overall relationship within that community.

The aim of this paper is to investigate the predictions of each of the aforementioned theories as outlined in Table 1 for the plant community within a single site using a large number of species. We attempt to address some of the potential causes of disagreement among empirical results by differentiating between size and survival using indices of both competitive intensity and competitive importance among plots naturally varying in both neighbour biomass and resource availability.

Methods

Study Site and Species

The study occurred in an unbroken and unseeded 50 ha section of native prairie at the University of Alberta research ranch at Kinsella, Alberta, Canada (53°05’N, 111°33’W). The field site is a savannah type habitat with mixed grass prairie interspersed with stands of aspen (*Populus tremuloides*). Standing crop within the prairie area naturally varies from 100 to 800 g/m², with *Hesperostipa curtiseta* and *Festuca hallii* being the most common species at low to moderate productivity, switching to *Poa pratensis* and *Galium boreale* at moderate to high productivity (J.A. Bennett, unpublished data). The site is co-limited by water and nitrogen [57], although competitive intensity is more closely linked to water availability [53]. Competition is generally intense; often reducing plant growth by approximately 90% during seedling establishment [57,58] and 50% in established plants [53,59]. Neighbour effects on seedling survival are typically near neutral [58,59], except during extended drought, when competition greatly increases seedling mortality [60].

The site receives an average of 418 mm total precipitation annually, which includes 155 mm of snow and rain from first snowfall through spring and 217 mm rain over the summer months (June through August). However, there is never an average year in a continental grassland, and precipitation was approximately 75% of average both leading up to and during the experiment. Daily temperatures over the experimental period were similar to the thirty year average, with mean daily temperatures of 15.2°C, average highs of 22.3°C and lows of 8.0°C. Though the site has historically been grazed by cattle, grazing had been halted four years prior to the onset of the experiment and did not occur for the duration of the study. Insect herbivory has little effect on the plant community at this site [61], while grazing by free-ranging ungulates (e.g. deer) and small mammals is infrequent (J.A. Bennett, personal observation).

We recognize that the range of productivity within our study is smaller than would be found in a transcontinental study [32,56]. However, none of the C-P theories except the SGH [39,40] suggest they operate only under specific ranges of productivity, and even the SGH has been extended into more productive environments [37,42]. We also recognize that non-linearities in relationships [51,55] would be difficult to detect using narrow productivity ranges.

Twenty-two plant species were selected for the experiment, including four annual species and eighteen perennial species (Table 2). Perennial species were chosen to be representative of the species naturally occurring across the range of productivity within this grassland. Annuals are uncommon at the site, yet are present and could respond to competition differently than perennials [62]. Due to limited seed availability for the annual species present at the site, annual species were chosen from the regional pool and according to seed availability. Combined, these 22 species represent approximately 25% of the total vegetative cover and 15% of the total vascular species richness at the site.

Experimental Design

Twenty replicate blocks, each consisting of two 2 m×2 m plots, were established in the summer of 2008. To test for the effect of neighbours on plant growth and survival, one plot of each pair was randomly assigned a neighbours removed treatment and the other a neighbour intact treatment. Neighbour removal was initially accomplished through a combination of mowing and the application of a broad spectrum herbicide (Round-up®, Monsanto), and maintained by applying herbicide to non-target plants using a paint brush and hand weeding as needed. These plots were surrounded by a 0.5 m vegetation free buffer zone at the edge of which roots were severed to a depth of 0.1 m to minimize interactions with vegetation surrounding the plots. This edge was re-cut on an approximately monthly basis. As the removal of vegetation in neighbours removed plots resulted in the removal of much of the litter layer, neighbours intact plots were also raked to remove an equivalent portion of the litter layer. Standing biomass was low at the time of transplanting, minimizing the amount of damage to aboveground plant structures; however, we kept raking intensity low to reduce damage to the soil surface and existing vegetation.

Each plot was divided into sixty-four 0.25 m by 0.25 m cells in an 8×8 grid. In total there were 20 blocks ×2 plots ×64 cells for a total of 2560 planting locations in this experiment. One seedling of a given species was transplanted into each cell such that each species had two to three individuals in each plot. Species' positions within the grid were assigned randomly for each block, so that the identity of the planted neighbour was consistent between plots within the block, but varied among blocks. This design ensured

Table 2. A list of species used within the experiment by growth form and family.

Life history	Family	Species	Frequency
Annual	Brassicaceae	<i>Capsella bursa-pastoris</i>	<1
		<i>Lepidium densiflorum</i>	<1
	Chenopodiaceae	<i>Chenopodium album</i>	6
		<i>Monolepis nuttalliana</i>	<1
Perennial	Apiaceae	<i>Zizia aptera</i>	2
	Asteraceae	<i>Artemisia ludoviciana</i>	57
		<i>Gaillardia aristata</i>	7
		<i>Heterotheca villosa</i>	<1
		<i>Solidago missouriensis</i>	69
	Campanulaceae	<i>Campanula rotundifolia</i>	52
	Fabaceae	<i>Hedysarum alpinum</i>	5
	Lamiaceae	<i>Monarda fistulosa</i>	4
	Linaceae	<i>Linum lewisii</i>	<1
	Poaceae	<i>Bouteloua gracilis</i>	28
		<i>Bromus inermis</i>	26
		<i>Elymus trachycaulus</i>	85
		<i>Nasella viridula</i>	<1
		<i>Poa pratensis</i>	95
Rosaceae	<i>Geum triflorum</i>	23	
	<i>Potentilla arguta</i>	12	
Scrophulariaceae	<i>Penstemon gracilis</i>	14	

Note: Growth forms were based upon observed morphologies under competition and determined following Cornelissen *et al.* (2003). Frequency of occurrence was determined by a 2009 survey of 100 2×2 m plots spread across the field site. Values of <1 denote plants that are known to occur at the field site, but were not observed within the plots.
doi:10.1371/journal.pone.0043703.t002

that, if competition occurred between seedlings that we planted, any neighbour-specific competitive effects [63,64] on target plant performance would remain consistent within blocks, but that species-specific responses would not be confounded by the identity of the planted neighbour when comparing across species and across blocks.

Seedlings were started in the greenhouse and transplanted into the field at the beginning of June 2009 at approximately four weeks of age. All seedlings at the time of transplanting had at least their first two true leaves, although most had at least four leaves. When transplanting, a 2 cm wide and 5 cm deep circular hole was made using a soil step sampler and the seedling was inserted along with the propagation soil. The narrow hole diameter was chosen to minimize damage to surrounding roots, although some trampling of the surrounding vegetation did occur. To increase establishment success, plots were watered with approximately 2 L/m²/day for the first 5 days following seedling transplanting and 1 L/m²/day for the next 5 days, but received no supplemental water after 10 days. All seedlings that died within the first ten days of the initial transplanting were replaced with new individuals from the trays grown in the greenhouse. Seedling mortality was monitored for all plants approximately biweekly following the initial replacement period for transplants, with percent survival of transplants calculated in late August 2009 after 13 weeks of growth.

Plant Growth

To measure biomass and estimate growth of annual plants, all annuals were clipped in early August 2009 to avoid mass loss due to seed dispersal. Plants were then dried at 65°C for at least 72 h and weighed. This study is intended to persist long-term, and thus destructive measures of perennial plants could not be taken. Instead, perennial growth was estimated using species-specific biomass regressions and plant measurements taken in late August 2009, prior to senescence. For these plants, we measured the maximum width (w_1), width perpendicular to maximum (w_2), and height (h) of each plant. We took the same measurements on a second smaller set of plants also grown with and without neighbours, and clipped these plants for the development of our biomass regressions. For 3 of the 18 species, survival of these plants was too low ($N < 5$) to estimate biomass and these species were removed from all analyses concerning plant growth. We used backwards step-wise regression to estimate biomass with \ln (biomass) as the dependent variable and \ln (basal area), \ln (height), and \ln (flowering stems) as independent variables. For *Bouteloua gracilis*, our step-wise regression model selected only \ln (flowering stems), causing plants without flowers to be underestimated; therefore we removed flowering stems from our starting model, which still gave acceptable results ($R^2 = 0.919$). For *Hedysarum alpinum*, we were unable to obtain a suitable regression ($P > 0.05$); therefore we removed *H. alpinum* from analyses related to growth. The full set of equations and the regression results can be found in Table 3. Similar model selection analyses using mixed models and small sample AIC (AIC_c) selected identical parameters and gave identical coefficient estimates as the regression approach. We chose not to harvest roots or monitor root growth due to our desire to avoid destructive sampling and the inherent logistical issues of having more accurate measures of root biomass in the neighbours removed than the neighbours intact treatment [65]. However, our estimates of shoot growth should provide adequate estimates of the total effect of neighbours on plant performance, which presumably is a proxy for fitness [65].

Productivity Estimation

Aboveground net primary productivity was estimated as standing biomass in grams dry weight/m² (g/m²) for each block at peak biomass in late July 2010. We could not harvest biomass from within the plots with neighbours intact as it would disrupt the long term goals of the study. Therefore, vegetation was clipped in four 0.1 m by 1 m quadrats surrounding the block, with individual quadrats placed on the north, south, east and west sides of each block. Samples were then sorted to remove dead material, dried at 65°C for at least 72 h, and weighed. Values for the individual quadrats ranged between 130 and 630 g/m²; however, productivity is naturally spatially heterogeneous at the site. We therefore used the average biomass of the four quadrats as our estimate of productivity for that block. Averaging among the quadrats restricted the range of productivity to 225–460 g/m². This could underestimate the absolute range of productivity between blocks, but should still represent the relative differences among blocks. We recognize that given the range of productivity covered, our test is not a definitive test of the associated theories; however, it is a test of the relationship between competition and productivity for this site.

Soil moisture was measured in both neighbours intact and neighbours removed plots using a ML2x – ThetaProbe Soil Moisture Sensor (Delta-T Devices) in late May 2010. Within each plot soil moisture was measured 5 times, once in each corner and the center. Within neighbours removed plots, care was taken to avoid sampling within the immediate vicinity of a seedling. Soil moisture in neighbours removed plots approximates the moisture retention capacity of the soil (gross water supply), whereas soil moisture where neighbours are intact approximates difference between supply and demand (net water supply) [5]. We did not calculate the ratio of supply to demand as suggested by Taylor [43] because we did not directly measure demand. Similarly, we chose not to estimate demand using the method laid out by Davis [5] due to potentially confounding instances when neighbours increase water availability as seen in studies of facilitation [37]. We

Table 3. Biomass regression coefficient estimates and significance tests.

Species	Regression coefficients				Regression results			
	Intercept	\ln (height)	\ln (flowers)	\ln (basal area)	Adjusted R ²	F	df	P
<i>Zizia aptera</i>	-6.76			1.11	0.964	240.61	1,8	<0.001
<i>Artemisia ludoviciana</i>	-8.03			1.33	0.944	153.23	1,8	<0.001
<i>Gaillardia aristata</i>	-8.92			1.49	0.953	182.42	1,8	<0.001
<i>Solidago missouriensis</i>	-11.01	3.85			0.645	15.54	1,7	0.006
<i>Symphotrichum laeve</i>	-7.65		0.47	1.14	0.977	193.27	2,7	<0.001
<i>Campanula rotundifolia</i>	-11.01		-1.48	2.10	0.892	29.964	2,5	0.002
<i>Monarda fistulosa</i>	-9.14	1.14		0.95	0.991	343.13	2,4	<0.001
<i>Bouteloua gracilis</i>	-6.981			1.00	0.903	56.99	1,5	0.001
<i>Bromus inermis</i>	-7.01	1.1	0.48	0.54	0.993	437.36	3,6	<0.001
<i>Elymus trachycaulus</i>	-11.53	3.32			0.986	572.31	1,7	<0.001
<i>Nasella viridula</i>	-8.95	1.86		0.56	0.97	147.94	2,7	<0.001
<i>Poa pratensis</i>	-8.21	1.44		0.68	0.966	39.21	2,6	<0.001
<i>Geum triflorum</i>	-8.23			1.43	0.979	416.72	1,8	<0.001
<i>Potentilla arguta</i>	-7.22	0.62	2.41	0.76	0.971	67.17	3,3	0.003
<i>Penstemon gracilis</i>	-7.17		2.39	1.08	0.995	527.78	2,3	<0.001

Note: For each species, if a particular regression coefficient was removed from the regression model by backward step-wise regression, then it is left blank in the table below. For *Bouteloua gracilis*, \ln (flowers) was not included in the regression model as it caused underestimation of biomass for plants without flowers.
doi:10.1371/journal.pone.0043703.t003

recognize that the effect of the plant community on soil moisture in late May is less than can be expected in mid-July, but soil moisture measurements in July were not taken. However, among other plots at the field site within the same growing season, net water supply in May was highly correlated with net water supply in July ($r = 0.782, P < 0.001$).

Although neighbour biomass can be expected to vary with water availability, this correlation is not perfect. Across a survey of 100 sites, net water supply was correlated with standing crop ($r = -0.321, P = 0.003$; J.A. Bennett, unpublished data). However, this does not explain the majority of the variation in standing crop. In this particular study, standing crop was uncorrelated with either net ($r = 0.146, P = 0.538$) or gross water supply ($r = 0.178, P = 0.453$), although net and gross water supply were highly correlated ($r = 0.697, P = 0.001$). This suggests that other factors are potentially limiting to plant growth, including, but not limited to, nitrogen [57]. Determining these factors and their role in competition is the subject of future research.

Competition Metrics

Competition can differentially affect survival and growth [1,45,47]. Further, both survival and growth are important components of fitness [1,45,66,67], with species-specific competitive effects on seedling growth being a strong predictor of species' abundances in the field [45]. Thus we estimated the effect of competition for both survival and size separately.

The choice of response metric can influence the form of the C–P relationship [1,30,50,68,69]. We chose the log response ratio (lnRR) to estimate competitive intensity [1,24] as it is an unbiased estimate of the effect of competition which is usually normally distributed and symmetrical around zero [68,69]. The ratio was calculated such that positive response ratios indicate competition and negative ratios indicate facilitation:

$$\ln RR = \ln \left(\frac{NR}{NI} \right)$$

From the response ratios, we classified interactions as positive, neutral or negative to determine interaction frequency. Interactions were classified as positive if neighbours increased plant survival or size by greater than 10% ($\ln RR < -0.0953$) and negative if neighbours reduced survival or size by greater than 10% ($\ln RR > 0.0953$). All other interactions were classified as neutral.

We calculated the relative importance of competition using the importance index (I_{imp}) [70], modified so that competition would be positive and facilitation negative:

$$I_{imp} = \frac{(NR - NI)}{|NR - NI| + |NR - N_{max}|}$$

This index was chosen as it is symmetrical around zero for competitive and facilitative interactions although there have been some concerns made regarding its utility in some situations [69]. Here N_{max} refers to the maximum performance of a given species in either the neighbours removed or neighbours intact treatments. These indices were calculated for each species within each block except under specific conditions. For survival, indices were not calculated for a given species within a specific block if mortality was complete for that species within that block. Similarly, indices were not calculated for size if mortality for a given species reached

100% in either neighbours intact or neighbours removed plots within that block.

Statistical Analyses

To determine if the effects of competition on survival and size differed, we used two mixed models, one for competitive intensity and one for competitive importance. These models included plant response measure (survival vs. size) as a fixed factor and species and block as random factors, with either competitive intensity or importance as the response variable in SPSS (v18.0). All plant response measures for a given species were pooled at the plot level prior to analysis; survival represents the proportion of seedlings that survived the year and size represents the average size of surviving individuals within that plot.

To test whether neighbour standing crop, gross water supply, or net water supply were associated with changes in competitive intensity or importance for either plant survival or growth, we used twelve mixed models specifying different independent and response variables in SPSS (v18.0). For each independent variable (standing crop, gross water supply, net water supply) we ran four models: $\ln RR - survival, I_{imp} - survival, \ln RR - growth,$ and $I_{imp} - growth$. Although gross water supply is not technically a measure of productivity, it represents potential productivity. Thus for ease of comparison, we will refer to it as a measure of productivity. Variation in competition-productivity (C–P) slopes among species was accounted for with a random interaction between species and productivity that allowed the C–P slopes to vary randomly by species. Estimation method (regression or weighed) was also included as a random effect for these analyses.

To test for changes in the frequency of interaction types (competitive, neutral or facilitative) across the range of biomass and resource availability found for both plant size and survival, we used six generalized linear mixed models with PROC GLIMMIX in SAS (v9.2) specifying the multinomial distribution. Interaction type was used as the response variable with either neighbour biomass, gross water supply, or net water supply as continuous fixed effects and species as a random effect.

Results

Across all species and blocks, plants were 17.3 times larger in no competition plots than competition plots; whereas survival was 1.2 times higher with neighbours (55% survival) than without (47% survival). This resulted in variation in the magnitude of competitive intensity ($F_{1,614} = 420.85, P < 0.001$, Fig. 1A) and importance ($F_{1,610} = 49.92, P < 0.001$, Fig. 1B) between survival and size. The magnitude of competitive intensity were comparable to other competition studies on seedlings conducted at the site for both size and survival [57,58], suggesting these results are 'typical' for this location.

Both competitive intensity and importance declined with increasing neighbour biomass when considering seedling survival (Fig. 2A, C, Table 4); however neither gross nor net water supply significantly affected competitive effects on seedling survival (Table 4). For seedling growth, both competitive intensity and importance declined with increasing gross water supply (Fig. 2B, D, Table 4), but neither neighbour biomass nor net water supply had a significant effect (Table 4).

Across all species, facilitative interactions were common for plant survival (45%), but rare for size (14%). The remaining neighbour effects on survival were both competitive (30%) and neutral (25%), whereas, neighbours were largely competitive when measuring growth (85%), and rarely neutral (<1%). Echoing the declines in competitive intensity and importance, the frequency of




Figure 1. Competitive effects on separate seedling life history aspects. Shown are the mean competitive intensity (A) and competitive importance (B) for seedling survival and size. Responses were calculated such that competition is represented by positive values and facilitation by negative values. Means represent the average of all species and error bars indicate one standard error. Note that the y-axes in the two panels use different scales.

doi:10.1371/journal.pone.0043703.g001

competitive effects on survival decreased with standing crop, with a concurrent increase in the frequency of facilitative effects (Table 4). Interaction frequency was not found to change for survival with either gross or net water supply. Given the rarity of non-competitive effects on plant growth, it is unsurprising that the relative frequency of competitive and facilitative interactions on growth did not change regardless of the productivity estimate used (Table 4).

Discussion

Neighbour Interactions and Productivity

Across all models, we found that competitive intensity declined with productivity, but that the type of productivity measurement that was associated with this decline depended on the plant response. However, we found no evidence of increasing competi-

tive effects across the range of productivity used as predicted by Grime [11,12] and parts of the SGH [37,39,40,51]. This was consistent regardless of the competition metric, plant response, or productivity measurement used. As competition was neither intense nor relatively important at high productivity for either plant survival or size, we suggest that these results are also not consistent with an increase in the likelihood of competitive exclusion as predicted by Newman [13], at least not over this range of productivity.

Given that we found a decline in competitive intensity with at least one measure of productivity for both plant growth and survival, we find little support for Tilman [3,14] either. However, we did find that competition was invariant when measuring plant growth and standing crop as well as survival and gross water supply. This could be construed as support for Tilman's




Figure 2. Changes in competitive intensity and importance with productivity. Competitive intensity (A) and relative competitive importance (C) decline as a function of standing crop for survival. Similarly, competitive intensity (B) and importance (D) decline for plant growth with gross water supply. Horizontal solid lines denote zero on the y-axis. Values above this line are competitive and below this line are facilitative. Dashed lines represent best fit lines. Each panel has a different scale for the y axis and that x axes are the same for A and C and for B and D. doi:10.1371/journal.pone.0043703.g002

prediction, but given that C–P relationships were so often negative, the support is marginal at best.

As previously mentioned, we found little support for the most common prediction of the SGH: that facilitation will increase with increasing stress [36,37,39,40,42], nor did we see evidence for a hump-shaped distribution of facilitation along a stress gradient [40,42]. It is possible that a hump-shaped relationship would appear if our range of productivity covered lower productivity areas. However, across the range of productivity measured, facilitation of survival always decreased with stress and we found no evidence of net facilitation in relation to plant growth at any productivity level. Our results are consistent with the earliest version of the SGH [37] which predicted that facilitation could increase with productivity under high herbivore pressure due to associational defences which lead to a reduced risk of herbivory [37,39]. Although this mechanism is likely active at the site to some extent, it is unlikely to be dominant as herbivory remains low when cattle grazing is not active [61]. However, other mechanisms of facilitation including protection from light damage and desiccation [38] could have led to this pattern (see below).

Although the SGH cannot explain the decline in competitive effects on plant growth and survival with productivity found in this system, these results are consistent with separate meta-analyses [1,33]. Both our results and the meta-analyses are at least partially consistent with theories that account for variation in both resource

supply and demand [5,43,71]. The decline in competition with productivity we observed is consistent with his predictions, but we did not find a relationship between competition and net resource supply as predicted by Davis [5]. This suggests that the supply and demand theory is also unable to predict the outcome of competition in this grassland community. However, we cannot rule out this theory because our measurements of net resource supply were not perfectly timed. Although net resource supply is correlated between early and mid-growing season, this correlation is not perfect and it is possible that a relationship exists between net resource supply and competition at peak biomass.

There are other pieces of evidence that suggest that resource supply and demand are important in determining the relative effects of competition. Nutrient uptake is not perfectly efficient [72] and light is not always limiting at high productivity [73,74]. This can result in increasing net nutrient availability at high productivity [72], which does not necessarily coincide with an increase in aboveground competition [73,74], and can cause a decrease in total competition with productivity. Of course an increase in aboveground competition is going to be dependent on the range of productivity explored. However, even if at higher productivity, competition does increase, this would suggest that the relationship between competition and productivity would have to be non-linear for this site.

Table 4. Results of mixed model analysis of competition productivity relationships.

Plant response	Competition metric	Response variable	F	d.f.	P
Survival	Intensity (lnRR)	Gross water supply	0.69	1,306	0.408
		Net water supply	0.65	1,362	0.422
		Standing crop	18.00	1,250	<0.001
	Importance (limp)	Gross water supply	0.61	1,359	0.434
		Net water supply	0.12	1,392	0.728
		Standing crop	12.71	1,384	<0.001
	Interaction frequency	Gross water supply	1.07	1,381	0.301
		Net water supply	0.49	1,381	0.486
		Standing crop	16.92	1,381	<0.001
Biomass	Intensity (lnRR)	Gross water supply	6.55	1,225	0.011
		Net water supply	0.15	1,226	0.700
		Standing crop	0.42	1,221	0.516
	Importance (limp)	Gross water supply	4.35	1,223	0.038
		Net water supply	0.19	1,222	0.666
		Standing crop	0.02	1,225	0.880
	Interaction frequency	Gross water supply	0.32	1,205	0.573
		Net water supply	0.39	1,205	0.536
		Standing crop	0.91	1,205	0.342

Note: Significant results are in bold.
doi:10.1371/journal.pone.0043703.t004

We also found a number of differences between plant survival and growth which are consistent with previous findings; neighbours tend to have neutral to positive effects on plant survival and competitive effects on plant growth [40,45,46,75]. We also found that neighbour effects were associated with resource supply for growth and standing crop for survival. These differences between seedling survival and growth are consistent with the concept that environmental effects on plant-plant interactions can vary depending on the life history stage of the plant [40,47,48,49]; however, this aspect of theory is not fully developed.

Some have hypothesized that neighbour effects can become more competitive as plants grow, in part due to differences in resource requirements [48]. In the current study, the association between resource supply and competitive effects on growth suggests that neighbour effects on growth are largely determined by resources. Resource interactions in mild environments are thought to be mostly negative [40], which explains the large competitive effects on growth. However, seedling survival can be facilitated through a number of mechanisms including reduced probabilities of desiccation, photoinhibition and herbivory [38]. These mechanisms do not necessarily affect available resources, which may explain why seedling survival increased with standing crop and not resource supply. In some cases, this can lead to an increase in facilitation of seedling survival with productivity as seen in this study and others [1], perhaps because low productivity sites

have too little vegetative cover to provide these benefits to seedlings, making facilitation more likely at higher productivities.

Neither our results nor those of the meta-analyses [1,33] fully support any of the major theories, suggesting that new theory regarding the relationship between plant-plant interactions and environmental gradients should be developed. These theories should incorporate both competition and facilitation as both are occurring simultaneously within most sites and should also account for the effects of multiple environmental gradients on different life history components. Some work has been done in this direction [2,40], although these theories must become more mechanistic and explicit in their predictions.

Acknowledgments

We would like to P. Belter, M. Clarke, N. Mahaffi, K. Peck, P. Rosa, B. Shore, J. Sired, and C. Zinyemba for help in the establishment and maintenance of the experiment. We also thank A. Nixon for helpful comments on an earlier version of this manuscript.

Author Contributions

Conceived and designed the experiments: JAB JFC. Performed the experiments: JAB. Analyzed the data: JAB. Contributed reagents/materials/analysis tools: JFC. Wrote the paper: JAB JFC.

References

- Goldberg DE, Rajaniemi T, Gurevitch J, Stewart-Oaten A (1999) Empirical approaches to quantifying interaction intensity: Competition and facilitation along productivity gradients. *Ecology* 80: 1118–1131.
- Craine JM (2009) *Resource Strategies of Wild Plants*. Princeton, NJ: Princeton University Press.
- Tilman D (1988) *Plant strategies and the dynamics and structure of plant communities*. Princeton, NJ: Princeton University Press.
- Grime JP (2001) *Plant Strategies, Vegetation Processes, and Ecosystem Properties*. Chichester, UK: John Wiley and Sons.
- Davis MA, Wragge KJ, Reich PB (1998) Competition between tree seedlings and herbaceous vegetation: support for a theory of resource supply and demand. *J Ecol* 86: 652–661.
- Cain ML, Bowman WD, and Hacker SD (2008) *Ecology*. Sunderland, MA: Sinauer Associates. 621 p.
- Molles MC, Cahill JF Jr. (2011) *Ecology Concepts and Applications*. Toronto, Canada: McGraw-Hill Ryerson. 654 p.
- Stiling PD (2002) *Ecology: Theories and Applications*. Upper Saddle River, NJ: Prentice Hall. 403 p.

9. Ricklefs RE (2008) *The Economy of Nature*. New York, NY: W.H. Freeman and Co. xix, 620 p.
10. Grime JP (1977) Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *The American Naturalist* 111: 1169–1194.
11. Grime JP (1979) *Plant Strategies, Vegetation Processes, and Ecosystem Properties*. Chichester, UK: John Wiley and Sons.
12. Grime JP (1973) Competitive exclusion in herbaceous vegetation. *Nature* 242: 344–347.
13. Newman EI (1973) Competition and diversity in herbaceous vegetation. *Nature* 244: 310–310.
14. Tilman D (1982) *Resource Competition and Community Structure*. Princeton, NJ: Princeton University Press.
15. Brooker R, Kikvidze Z, Pugnaire FI, Callaway RM, Choler P, et al. (2005) The importance of importance. *Oikos* 109: 63–70.
16. Craine JM (2005) Reconciling plant strategy theories of Grime and Tilman. *J Ecol* 93: 1041–1052.
17. Grace JB (1991) A clarification of the debate between Grime and Tilman. *Funct Ecol* 5: 583–587.
18. Lamb EG, Kembel SW, Cahill JF Jr. (2009) Shoot, but not root, competition reduces community diversity in experimental mesocosms. *J Ecol* 97: 155–163.
19. Hautier Y, Niklaus PA, Hector A (2009) Competition for light causes plant biodiversity loss after eutrophication. *Science* 324: 636–638.
20. Grime JP (2007) Plant strategy theories: a comment on Craine (2005). *J Ecol* 95: 227–230.
21. Tilman D (2007) Resource competition and plant traits: a response to Craine, et al. 2005. *J Ecol* 95: 231–234.
22. Craine JM (2007) Plant strategy theories: replies to Grime and Tilman. *J Ecol* 95: 235–240.
23. Twolan-Strutt L, Keddy PA (1996) Above- and belowground competition intensity in two contrasting wetland plant communities. *Ecology* 77: 259–270.
24. Cahill JF Jr. (1999) Fertilization effects on interactions between above- and belowground competition in an old field. *Ecology* 80: 466–480.
25. Peltzer DA, Wilson SD, Gerry AK (1998) Competition intensity along a productivity gradient in a low-diversity grassland. *The American Naturalist* 151: 465–476.
26. Wilson SD, Tilman D (1993) Plant competition and resource availability in response to disturbance and fertilization. *Ecology* 74: 599–611.
27. Campbell BD, Grime JP (1992) An Experimental Test of Plant Strategy Theory. *Ecology* 73: 15–29.
28. Wilson SD, Tilman D (1991) Component of plant competition along an experimental gradient of nitrogen availability. *Ecology* 72: 1050–1065.
29. Wilson SD, Tilman D (1995) Competitive responses of eight old-field plant species in four environments. *Ecology* 76: 1169–1180.
30. Brooker RW, Kikvidze Z (2008) Importance: an overlooked concept in plant interaction research. *J Ecol* 96: 703–708.
31. Wilson MV (2007) Measuring the components of competition along productivity gradients. *J Ecol* 95: 301–308.
32. Reader RJ, Wilson SD, Belcher JW, Wisheu I, Keddy PA, et al. (1994) Plant competition in relation to neighbor biomass - An intercontinental study with *Poa pratensis*. *Ecology* 75: 1753–1760.
33. Maestre FT, Valladares F, Reynolds JF (2005) Is the change of plant–plant interactions with abiotic stress predictable? A meta-analysis of field results in arid environments. *J Ecol* 93: 748–757.
34. Lortie CJ, Callaway RM (2006) Re-analysis of meta-analysis: support for the stress-gradient hypothesis. *J Ecol* 94: 7–16.
35. Maestre FT, Valladares F, Reynolds JF (2006) The stress-gradient hypothesis does not fit all relationships between plant–plant interactions and abiotic stress: further insights from arid environments. *J Ecol* 94: 17–22.
36. Michalet R, Brooker RW, Cavieres LA, Kikvidze Z, Lortie CJ, et al. (2006) Do biotic interactions shape both sides of the humped-back model of species richness in plant communities? *Ecol Lett* 9: 767–773.
37. Bertness MD, Callaway R (1994) Positive interactions in communities. *Trends Ecol Evol* 9: 191–193.
38. Callaway RM (1995) Positive interactions among plants. *Bot Rev* 61: 306–349.
39. Callaway RM, Walker LR (1997) Competition and facilitation: A synthetic approach to interactions in plant communities. *Ecology* 78: 1958–1965.
40. Maestre FT, Callaway RM, Valladares F, Lortie CJ (2009) Refining the stress-gradient hypothesis for competition and facilitation in plant communities. *J Ecol* 97: 199–205.
41. Malkinson D, Tielbörger K (2010) What does the stress-gradient hypothesis predict? Resolving the discrepancies. *Oikos* 119: 1546–1552.
42. Holmgren M, Scheffer M (2010) Strong facilitation in mild environments: the stress gradient hypothesis revisited. *J Ecol* 98: 1269–1275.
43. Taylor DR, Aarssen LW, Loehle C (1990) On the relationship between r/K selection and environmental carrying-capacity - A new habitat templet for plant life-history strategies. *Oikos* 58: 239–250.
44. Weaver JE, Clements FE (1938) *Plant ecology*. New York, London: McGraw-Hill Book Company, inc. xxii, 601 p. incl. front. (map) illus., diags. p.
45. Howard TG, Goldberg DE (2001) Competitive response hierarchies for germination, growth, and survival and their influence on abundance. *Ecology* 82: 979–990.
46. Wilson SD (1993) Competition and resource availability in heath and grassland in the Snowy Mountains of Australia. *J Ecol* 81: 445–451.
47. Goldberg D, Novoplansky A (1997) On the relative importance of competition in unproductive environments. *J Ecol* 85: 409–418.
48. Miriti MN (2006) Ontogenetic shift from facilitation to competition in a desert shrub. *J Ecol* 94: 973–979.
49. Schiffers K, Tielbörger K (2006) Ontogenetic shifts in interactions among annual plants. *J Ecol* 94: 336–341.
50. Grace JB (1995) On the measurement of plant competition intensity. *Ecology* 76: 305–308.
51. Kikvidze Z, Suzuki M, Brooker R (2011) Importance versus intensity of ecological effects: why context matters. *Trends in Ecology & Evolution* 26: 383–388.
52. Freckleton RP, Watkinson AR, Rees M (2009) Measuring the importance of competition in plant communities. *J Ecol* 97: 379–384.
53. Lamb EG, Cahill JF (2008) When competition does not matter: Grassland diversity and community composition. *Am Nat* 171: 777–787.
54. Damgaard C, Fayolle A (2010) Measuring the importance of competition: a new formulation of the problem. *J Ecol* 98: 1–6.
55. Belcher JW, Keddy PA, Twolan-Strutt L (1995) Root and shoot competition intensity along a soil depth gradient. *J Ecol* 83: 673–682.
56. Callaway RM, Brooker RW, Choler P, Kikvidze Z, Lortie CJ, et al. (2002) Positive interactions among alpine plants increase with stress. *Nature* 417: 844–848.
57. Lamb EG, Shore BH, Cahill JF (2007) Water and nitrogen addition differentially impact plant competition in a native rough fescue grassland. *Plant Ecol* 192: 21–33.
58. Haag JJ, Coupe MD, Cahill JF (2004) Antagonistic interactions between competition and insect herbivory on plant growth. *J Ecol* 92: 156–167.
59. Cahill JFJ (2003) Lack of relationship between below-ground competition and allocation to roots in 10 grassland species. *J Ecol* 91: 532–540.
60. Cahill JF (2003) Neighbourhood-scale diversity, composition and root crowding do not alter competition during drought in a native grassland. *Ecol Lett* 6: 599–603.
61. Coupe MD, Stacey JN, Cahill JF (2009) Limited effects of above- and belowground insects on community structure and function in a species-rich grassland. *Journal of Vegetation Science* 20: 121–129.
62. Gomez-Aparicio L (2009) The role of plant interactions in the restoration of degraded ecosystems: a meta-analysis across life-forms and ecosystems. *J Ecol* 97: 1202–1214.
63. Goldberg DE (1990) Components of resource competition in plant communities. In: Grace JB, Tilman D, editors. *Perspectives on Plant Competition*. Caldwell, NJ: The Blackburn Press. 27–49.
64. Goldberg DE (1996) Competitive ability: Definitions, contingency and correlated traits. *Philosophical Transactions: Biological Sciences* 351: 1377–1385.
65. Cahill JF (2002) What evidence is necessary in studies which separate root and shoot competition along productivity gradients? *J Ecol* 90: 201–205.
66. Aarssen LW, Keogh T (2002) Conundrums of competitive ability in plants: what to measure? *Oikos* 96: 531–542.
67. Neytcheva MS, Aarssen LW (2008) More plant biomass results in more offspring production in annuals, or does it? *Oikos* 117: 1298–1307.
68. Hedges LV, Gurevitch J, Curtis PS (1999) The meta-analysis of response ratios in experimental ecology. *Ecology* 80: 1150–1156.
69. Rees M, Childs DZ, Freckleton RP (2012) Assessing the role of competition and stress: a critique of importance indices and the development of a new approach. *J Ecol* 100: 577–585.
70. Seifan M, Seifan T, Ariza C, Tielbörger K (2010) Facilitating an importance index. *J Ecol* 98: 356–361.
71. Aarii K, Turkington R (2001) Assessing competition intensity along productivity gradients using a simple model. *Canadian Journal of Botany* 79: 1486–1491.
72. Yuan Z-Y, Li L-H, Han X-G, Chen S-P, Wang Z-W, et al. (2006) Nitrogen response efficiency increased monotonically with decreasing soil resource availability: a case study from a semiarid grassland in northern China. *Oecologia* 148: 564–572.
73. Abrams PA (1995) Monotonic or unimodal diversity-productivity gradients: What does competition theory predict? *Ecology* 76: 2019–2027.
74. Dickson TL, Foster BL (2011) Fertilization decreases plant biodiversity even when light is not limiting. *Ecol Lett* 14: 380–388.
75. Dyer AR, Goldberg DE, Turkington R, Sayre C (2001) Effects of growing conditions and source habitat on plant traits and functional group definition. *Funct Ecol* 15: 85–95.