

# The seroprevalence of Rubella in pregnant women in Turkey: a meta-analysis research of 90988 Rubella IgM, 84398 Rubella IgG, and 522 avidity results

## *Türkiye’de gebelerdeki Rubella seroprevalansı: 90988 Rubella IgM, 84398 Rubella IgG ve 522 avidite sonucunun meta-analiz araştırması*

© Rıza Aytaç Çetinkaya, © Ercan Yenilmez

University of Health Sciences, Sultan Abdülhamid Han Training and Research Hospital, Clinic of Infectious Diseases and Clinical Microbiology, İstanbul, Turkey

### Abstract

**Objective:** Rubella infection prevalence in pregnant women can vary from country to country, or even across regions in the same country. In this meta-analysis, the seroprevalence Rubella among pregnant women in Turkey in the last decade was evaluated.

**Materials and Methods:** Studies conducted in Turkey between 2007 and 2017 were analyzed, and differences in seroprevalence between provinces were compared by evaluating Rubella immunoglobulin (Ig)-G, IgM, and IgG avidity results in pregnancy in this period. A data search was performed using the search terms Rubella, kızamıkçık, gebe, hamile, pregnancy, Türkiye, Turkey in Google Scholar, PubMed, Web of Science, Türk Medline, and the YÖK thesis database center.

**Results:** A total of 26 articles associated with the seroprevalence of Rubella among pregnant women in Turkey were enrolled in the meta-analysis. As a result of an analysis of 84398 Rubella IgG, and 90988 Rubella IgM serology tests among pregnant women in 26 studies; Rubella IgG and IgM seroprevalence rates in pregnant woman in Turkey were found as 93.47% (95% CI: 91.72 to 95.03) and 0.783% (95% CI: 0.505 to 1.120), respectively. Rubella IgG low, intermediate, and high avidity rates were 4.66% (95% CI: 0.969 to 10.906), 7.51% (95% CI: 5.101 to 10.345), and 93.55% (95% CI: 82.584 to 99.311), respectively.

**Conclusion:** The Rubella IgG seropositivity rate in Turkey among pregnant woman is high, whereas it is low for IgM. These rates may be considered as the result of successful immunization policies and practices. In a few provinces, it is necessary to revise the Rubella immunization procedures and adult vaccination strategies should be developed in order to control Rubella infections in adults, including pregnant women.

**Keywords:** Rubella, pregnant women, meta-analysis, Turkey

### Öz

**Giriş:** Hamilelerde Rubella prevalansı ülkeler arasında, hatta aynı ülke içinde farklı bölgelerde değişiklik göstermektedir. Bu meta-analiz ile son on yılda Türkiye’deki gebelerde Rubella seroprevalansı araştırıldı.

**Gereç ve Yöntemler:** Türkiye’de 2007 ve 2017 yılları arasında gebelerde Rubella immunoglobulin (Ig)-G, IgM ve IgG avidite sonuçları araştırıldı. Bu işlem Google Scholar, PubMed, Web of Science, Türk Medline ve YÖK tez merkezi veri tabanlarında Rubella, kızamıkçık, gebe, hamile, pregnancy, Türkiye, Turkey anahtar kelimeleri kullanılarak gerçekleştirildi.

**Bulgular:** Ülkemizde gebe kadınlar arasında Rubella seroprevalansı ile ilişkili toplam 26 makale meta-analize dâhil edildi. 26 çalışmada gebe kadınlar arasında 84398 Rubella IgG ve 90988 Rubella IgM seroloji sonucuna göre; IgG ve IgM seroprevalans oranları sırasıyla %93,47 (%95 CI: 91,72-95,03) ve %0,783 (%95 CI: 0,505-1,120) saptandı. Rubella IgG düşük, orta ve yüksek avidite oranları sırasıyla %4,66 (%95 CI: 0,969 ila 10,906), %7,51 (%95 CI: 5,101 ila 10,345), %93,549 (%95 CI: 82,584 ila 99,311) saptandı.

**PRECIS:** In this meta-analysis, studies conducted in Turkey between 2007 and 2017 were analyzed, and differences in seroprevalence between provinces were revealed by evaluating Rubella IgG, IgM and IgG avidity results in pregnancy in this period.

Address for Correspondence/Yazışma Adresi: Rıza Aytaç Çetinkaya, MD,

University of Health Sciences, Sultan Abdülhamid Han Research and Training Hospital, Clinic of Infectious Diseases and Clinical Microbiology, İstanbul, Turkey

Phone: +90 505 438 28 14 E-mail: aytacetinkaya@yahoo.com ORCID ID: orcid.org/0000-0002-5676-9527

Received/Geliş Tarihi: 14.11.2018 Accepted/Kabul Tarihi: 12.02.2019

©Copyright 2019 by Turkish Society of Obstetrics and Gynecology

Turkish Journal of Obstetrics and Gynecology published by Galenos Publishing House.

**Sonuç:** Türkiye'deki gebelerde Rubella IgG seropozitiflik oranı yüksek, IgM ise düşüktür. Bu oranlar ülkemizdeki başarılı bağışıklama politikaları ve uygulamalarının bir sonucudur. Fakat birkaç ilde çocukluk çağı ve yetişkin Rubella aşılama prosedürlerinin yeniden gözden geçirilmesinin faydalı olabileceği değerlendirilmektedir.

**Anahtar Kelimeler:** Rubella, gebe, meta-analiz, Türkiye

## Introduction

Rubella is a vaccine-preventable disease and is one of the most infectious viral diseases known in humans. Congenital Rubella syndrome (CRS), consisting of cardiac disorders, cataract, deafness, cleft palate, autism, and fetal death can occur when the Rubella virus vertically infects the fetus during pregnancy<sup>(1)</sup>. Owing to the vaccination practices in childhood and adults, the prevalence of Rubella in pregnant women decreased and Rubella has become a rare infection in many developed and some developing countries. Some countries in the Western hemisphere and Europe have eliminated Rubella and CRS<sup>(2)</sup>.

Immune status can be evaluated by enzyme-linked immunoreactive techniques (enzyme immunoassay, ELISA). From these tests, immunoglobulin (Ig)-G antibody shows a previous infection or immunization by vaccines. The Rubella serum IgM test indicates an acute Rubella infection and must be confirmed by at least one of these tests: Rubella-RNA polymerase chain reaction or Rubella IgG-avidity or western-blot<sup>(2)</sup>.

The prevalence of Rubella infection in pregnant women can vary from country to country, and even across regions in the same country. In this meta-analysis, Rubella antibody tests in the last 11 years in pregnant women in Turkey were examined, and the differences in prevalence between provinces in our country, and between our country and other countries were compared.

## Materials and Methods

In this meta-analysis, studies conducted in Turkey between 2007 and 2017 were analyzed, and differences in seroprevalences between provinces were compared by evaluating Rubella IgG, IgM, and IgG avidity results in pregnancy in this period. Data in these studies were screened and evaluated using the preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow-chart according to the inclusion criteria (Figure 1).

### Source of data

A data search was performed using the search terms Rubella, kızamıkçık, gebe, hamile, pregnancy, Türkiye, Turkey in Google Scholar, PubMed, Web of Science, Türk Medline, and the YÖK thesis database center by two independent researcher in 2018.

### Inclusion and exclusion criteria

Studies with Rubella IgM, IgG, and IgG avidity test results in pregnant women in Turkey between 2007 and 2017 were included in the study. Original articles with antibody test results of at least 150 pregnant women with full text in

Turkish or English were recorded. Studies were excluded if they included the results of the antibodies in non-pregnant women and male or child patients, and if the studies did not include the study period or were published after 2007 but the data were collected before 2007 (Figure 1).

### Data search and collection of data

Considering the criteria, data were screened and evaluated by two different researchers (RAC, EY) in order to prevent publication bias. Study data: author's surname, date of publication, years of tests performed, numbers (n) and rates (%) of Rubella IgG, IgM, and IgG-avidity tests, and province of the study performed were recorded in the Microsoft Office 2016 Professional Plus Excel program. Before the meta-analysis, all data were listed in alphabetical order according to the author's surname in the extended format. Disputes between researchers were resolved by mutual discussion.

### Statistical Analysis


Medcalc© software version 17.9.7 program was used for meta-analysis. Data were transferred from the Excel program. A funnel plot was used to evaluate possible bias and the results were interpreted.

A statistical test for heterogeneity was performed to measure the data heterogeneity. According to this;  $I^2 \leq 25\%$  heterogeneity was assumed to be insignificant and a fixed effect was used.  $I^2 > 25\%$  heterogeneity was assumed to be insignificant; the study data were considered as nonhomogeneous and the random effect value was used.  $P < 0.01$  was considered to be no need to add more studies.

## Results

In this meta-analysis, 884 articles were found in accordance with the research criteria (Figure 1). A total of 681 articles were excluded from the study because of repetition in two or more different databases. After the removal, we had 203 studies, 201 of which we could screen. After evaluation of the study title and summary, another 31 were excluded from the study. After full text evaluations, 144 of the 170 studies were excluded from the study according to the determined criteria. As a result, a total of 26 articles associated with Rubella seroprevalence in pregnant women in Turkey were enrolled in the meta-analysis.

The studies included in the meta-analysis were from Afyon (n=2), Artvin (n=1), Bingöl (n=1), Denizli (n=1), Edirne (n=1), Isparta (n=1), Istanbul (n=4), Kahramanmaraş (n=1), Konya (n=2), Manisa (n=1), Muğla (n=1), Middle Black Sea (n=1), Rize (n=1), Uşak (n=1), Van (n=2), Yozgat (n=1), Zonguldak (n=1).



**Figure 1.** Flow chart for study selection and literature review. Summary of the literature search and study selection on Rubella antibodies in pregnant women

The study of Sargin and Saygan<sup>(3)</sup> from Ankara had the maximum number of cases with 31385 pregnant women, and the study of Bakacak et al.<sup>(4)</sup> from Kahramanmaraş had high number of cases with 11823 pregnant women.

According to the meta-analysis of 84398 serologic tests of pregnant women in the 26 studies, the seroprevalence rate of Rubella IgG in pregnant woman in Turkey was 93.47% (95% CI: 91.72 to 95.03). The Cochran Q test was 2032,5378;

$I^2=98.77\%$  and  $p<0.0001$ , respectively (Table 1). In funnel plot analysis, minimal asymmetry was found in the studies of Varol et al.,<sup>(5)</sup> Nazik et al.,<sup>(6)</sup> Başkesen et al.,<sup>(7)</sup> and Çeltek et al.<sup>(8)</sup> (Figure 2,3). Overall, the asymmetry test showed no bias.

According to a meta-analysis of 90,988 serologic tests of pregnant women in the 26 studies, the seroprevalence rate of Rubella IgM in pregnant woman in Turkey was 0.783%

**Table 1.** Meta-analysis of anti-Rubella IgG among pregnant women in Turkey

Study name, year, reference	Province	Sample size	Proportion (%)	95% CI	Weight (%)	
					Fixed	Random
Akpınar (2017) <sup>(9)</sup>	Isparta	805	97.52	96.19 to 98.48	0.95	3.84
Aşık (2013) <sup>(10)</sup>	Afyon	505	92.08	89.37 to 94.28	0.60	3.75
Aynioğlu (2015) <sup>(22)</sup>	Zonguldak	910	93.85	92.08 to 95.32	1.08	3.86
Bakacak (2014) <sup>(4)</sup>	Kahramanmaraş	11.823	93.20	92.73 to 93.65	14.01	3.99
Başkesen (2010) <sup>(7)</sup>	Manisa	1202	83.69	81.48 to 85.74	1.42	3.89
Çeltek (2014) <sup>(8)</sup>	Middle Black Sea	3162	99.68	99.42 to 99.85	3.75	3.96
Doğan (2014) <sup>(23)</sup>	İstanbul	1641	95.73	94.64 to 96.66	1.94	3.92
Efe (2009) <sup>(12)</sup>	Van	613	99.51	98.58 to 99.90	0.73	3.79
Gündem (2014) <sup>(24)</sup>	Konya	419	92.84	89.94 to 95.12	0.50	3.70
İnci (2014) <sup>(25)</sup>	Artvin	1292	95.20	93.89 to 96.30	1.53	3.90
Karabulut (2011) <sup>(26)</sup>	Denizli	1268	95.11	93.77 to 96.23	1.50	3.90
Karacan (2014) <sup>(27)</sup>	İstanbul	1258	95.55	94.26 to 96.62	1.49	3.90
Kasap (2017) <sup>(28)</sup>	Muğla	189	89.95	84.75 to 93.84	0.23	3.39
Keskin (2013) <sup>(29)</sup>	İstanbul	1926	95.74	94.74 to 96.60	2.28	3.93
Nazik (2017) <sup>(6)</sup>	Bingöl	10.178	84.31	83.59 to 85.01	12.06	3.99
Numan (2015) <sup>(30)</sup>	İstanbul	1101	94.19	92.64 to 95.50	1.31	3.88
Özdemir (2011) <sup>(31)</sup>	Konya	249	95.98	92.74 to 98.06	0.30	3.52
Parlak (2015) <sup>(14)</sup>	Van	416	86.54	82.88 to 89.67	0.49	3.70
Satılmış (2014) <sup>(32)</sup>	Yozgat	804	94.03	92.16 to 95.57	0.95	3.84
Şentürk (2016) <sup>(33)</sup>	Rize	424	93.87	91.14 to 95.96	0.50	3.71
Şevki (2013) <sup>(3)</sup>	Ankara	31.385	93.92	93.65 to 94.19	37.18	4.00
Şimşek (2016) <sup>(11)</sup>	Afyon	1076	94.52	92.98 to 95.80	1.28	3.88
Şirin (2017) <sup>(34)</sup>	İzmir	7189	93.49	92.89 to 94.05	8.52	3.99
Toklu (2013) <sup>(35)</sup>	Uşak	1465	92.15	90.65 to 93.48	1.74	3.91
Varıcı-Balcı (2014) <sup>(13)</sup>	İzmir	1871	93.21	91.98 to 94.31	2.22	3.93
Varol (2011) <sup>(5)</sup>	Edirne	1227	76.61	74.14 to 78.95	1.45	3.90
Total (fixed effects)	TURKEY	84.398	93.06	92.89 to 93.23	100.00	100.00
Total (random effects)			93.48	91.727 to 95.033	100.00	100.00

## Test for heterogeneity

Q 2032.5378

DF 25

Significance level p&lt;0.0001

I<sup>2</sup> (inconsistency) 98.77%95% CI for I<sup>2</sup> 98.57 to 98.94

DF: Dickey-Fuller test

(95% CI: 0.505 to 1.120). The Cochrane Q test was 583,6836; I<sup>2</sup>=95,72% and p<0.0001 (Table 2). A negligible asymmetry was found in the funnel plot analysis, and the asymmetry test showed no bias (Figure 4). The results of Başkesen et al.<sup>(7)</sup> and Akpınar et al.<sup>(9)</sup> were most distant from the average value

of Turkey.

Additionally, Rubella IgG-avidity rates were analyzed in the present meta-analysis (Table 3). Accordingly, the low avidity rate was 4.66% (95% CI: 0.969 to 10.906) and the Cochrane Q test was 10,3230; I<sup>2</sup>=70.94% (p<0.0001) in 427 pregnant in four studies<sup>(10-13)</sup>, intermediate avidity rate was 7.51% (95% CI: 5.101 to 10.345), and the Cochrane Q test was 1,8145; I<sup>2</sup>=0.00 (p=0.404) in 384 pregnant in three studies<sup>(11-13)</sup>, the high avidity rate was 93.55% (95% CI: 82.584 to 99.311), and the Cochrane Q test was 46,4845, I<sup>2</sup>=91.39% (p<0.0001) in 522 pregnant women in five studies<sup>(3,10-13)</sup>.

**Discussion**

Rubella infection is usually subclinical in childhood, but may be more severe at older ages in life. It can also lead to




Figure 2. Funnel plot analysis graph of anti-Rubella immunoglobulin-G in Turkey




Figure 3. Meta-analysis graph of anti-Rubella immunoglobulin-G severe anomalies or death in the fetus in the first trimester of pregnancy<sup>(1,14)</sup>.

The serologic test showing previous Rubella infection or immunization status alone is the Rubella IgG antibody. Due to the increase in the awareness of pregnant women and socioeconomic developments in our country, there has been an increase in prenatal screening tests in recent years. According to the meta-analysis of 26 studies analyzed, the Rubella IgG seropositivity rate in Turkey was 93.4%. This rate was higher than 44% of the studies, and lower than 56% of the studies included in the meta-analysis. The study of Şevki et al.<sup>(3)</sup> in Ankara with 31,385 cases represents the Turkey's average best; the seropositivity rate of 93.9% and 37.1% weight in the meta-analysis<sup>(3)</sup>.

In a general perspective, it is considered that the vaccination rates in the west part of Turkey is higher, and in the east part of Turkey, immunity is gained after acquiring infections against infections that can be prevented by vaccination. The study of Varol et al.<sup>(5)</sup> conducted in the Thrace region




Figure 4. Funnel plot analysis graph of anti-Rubella immunoglobulin-M

of Turkey revealed the lowest IgG ratio (76.6%), and also Baskesen et al.<sup>(7)</sup> from Manisa revealed a lower rate (83.6%) than the average of Turkey. In these two studies, it was not possible to make an inference because the age status of the pregnant women was not given in cross-sectional intervals. However, these rates raise doubts about effective vaccination strategies in the regions where both studies were conducted. More comprehensive randomized controlled prospective research is needed for these two regions.

Rubella IgG seropositivity rate was 84.3% in the study conducted by Nazik et al.<sup>(6)</sup> in the Bingöl province in 10,178 pregnant women. Similarly, the study of Parlak et al.<sup>(14)</sup> in the Van province in 416 pregnant women, the rate was 86.5% for IgG seropositivity. These rates could be considered to be due to the low level of vaccinations of the people living in the region, but it is not possible to form a definite opinion on this issue.

In the study of Çetlek et al.<sup>(8)</sup> with 3162 pregnant women representing the Middle Black Sea region, Rubella IgG seropositivity rate was found as 99%. Çetlek et al.<sup>(8)</sup> interpreted their result as close to the average of Turkey. However, the ratio in their study was found to be higher than the average of Turkey according to our meta-analysis. The high rate in this region indicates that the number of pregnant women who could be infected with Rubella during pregnancy was low. In order to use the results across the country, the characteristics of the cases should be evaluated further.

The study of Özdemir et al.<sup>(15)</sup> was not included in the meta-analysis because the test and technique used in the multicenter study conducted in seven provinces were not explicitly written. In this multicenter study, Rubella IgG positivity was between 76-96.4% and the results varied significantly in different provinces. Our meta-analysis, or similarly, this multicenter study shows that different results can be obtained in different regions. This difference can be



**Table 2.** Meta-analysis of anti-Rubella immunoglobulin-M among pregnant women in Turkey

Study Name, Year, References	Province	Sample size	Proportion (%)	95% CI	Weight (%)	
					Fixed	Random
Akpınar (2017) <sup>(9)</sup>	Isparta	1829	4.97	4.02 to 6.07	2.01	4.04
Aşık (2013) <sup>(10)</sup>	Afyon	552	1.81	0.87 to 3.31	0.61	3.48
Aymoğlu (2015) <sup>(22)</sup>	Zonguldak	933	1.50	0.82 to 2.50	1.03	3.79
Bakacak (2014) <sup>(4)</sup>	Kahramanmaraş	7733	0.19	0.11 to 0.32	8.50	4.27
Başkesen (2010) <sup>(7)</sup>	Manisa	1202	7.65	6.21 to 9.30	1.32	3.90
Çeltek (2014) <sup>(8)</sup>	Middle Black Sea	3162	0.25	0.09 to 0.50	3.48	4.16
Doğan (2014) <sup>(23)</sup>	İstanbul	1714	0.23	0.064 to 0.60	1.88	4.02
Efe (2009) <sup>(12)</sup>	Van	613	0.33	0.039 to 1.17	0.67	3.55
Gündem (2014) <sup>(24)</sup>	Konya	419	0.95	0.26 to 2.43	0.46	3.28
İnci (2014) <sup>(25)</sup>	Artvin	1292	0.31	0.084 to 0.79	1.42	3.93
Karabulut (2011) <sup>(26)</sup>	Denizli	1268	0.00	0.00 to 0.29	1.39	3.92
Karacan (2014) <sup>(27)</sup>	Istanbul	1258	0.48	0.17 to 1.03	1.38	3.92
Kasap (2017) <sup>(28)</sup>	Muğla	189	0.53	0.013 to 2.91	0.21	2.52
Keskin (2013) <sup>(29)</sup>	Istanbul	1926	0.16	0.032 to 0.45	2.12	4.06
Nazik (2017) <sup>(6)</sup>	Bingöl	10.178	0.78	0.61 to 0.97	11.18	4.29
Numan (2015) <sup>(30)</sup>	İstanbul	1101	0.18	0.02 to 0.65	1.21	3.86
Özdemir (2011) <sup>(31)</sup>	Konya	249	0.40	0.010 to 2.22	0.27	2.81
Parlak (2015) <sup>(14)</sup>	Van	9340	0.47	0.34 to 0.63	10.26	4.28
Satılmış (2014) <sup>(32)</sup>	Yozgat	804	0.12	0.003 to 0.69	0.88	3.71
Şentürk (2016) <sup>(33)</sup>	Rize	1037	0.29	0.06 to 0.84	1.14	3.84
Şevki (2013) <sup>(3)</sup>	Ankara	31.385	0.39	0.33 to 0.47	34.48	4.32
Şimşek (2016) <sup>(11)</sup>	Afyon	1112	2.52	1.68 to 3.62	1.22	3.87
Şirin (2017) <sup>(34)</sup>	İzmir	7189	1.33	1.08 to 1.63	7.90	4.26
Toklu (2013) <sup>(35)</sup>	Uşak	1465	0.96	0.52 to 1.60	1.61	3.97
Varıcı-Balcı (2014) <sup>(13)</sup>	İzmir	1784	0.17	0.035 to 0.49	1.96	4.03
Varol (2011) <sup>(5)</sup>	Edirne	1254	0.72	0.33 to 1.36	1.38	3.92
<b>Total (fixed effects)</b>	<b>TURKEY</b>	<b>90.988</b>	<b>0.58</b>	<b>0.53 to 0.63</b>	<b>100.0</b>	<b>100.0</b>
<b>Total (random effects)</b>			<b>0.78</b>	<b>0.50 to 1.12</b>	<b>100.0</b>	<b>100.0</b>

explained by the wide geographic structure of Turkey and by the sociocultural differences that exist between the regions. Therefore, we believe that accurate rates throughout the country can only be obtained by meta-analysis studies.

The age and dose of the Rubella vaccination may vary depending on a country's vaccination policy. In general, an 84.7% seropositivity rate is achieved by single-dose vaccination, and this rate reaches 90% with two doses of vaccine (ages 1 and 5)<sup>(16)</sup>. In our country, Rubella vaccines are administered as two doses in children at 12 months and 7 years. Turkey, with a rate of 93.4% Rubella IgG seropositivity, has a rate just above that in the general literature. Turkey's neighboring Iran has a seropositivity rate of 94%,<sup>(17)</sup> and Greece, which is a member of the European Union, has a seropositivity rate of 97% in women with childbearing age<sup>(18)</sup>.

#### Test for heterogeneity

Q 583.6836

DF 25

Significance level p<0.0001

I<sup>2</sup> (inconsistency) 95.72%

95% CI for I<sup>2</sup> 94.62 to 96.59

DF: Dickey-Fuller test

This ratio is 89.3% in Brazil and 99.3% in the United States of America<sup>(16)</sup>. In India, which is a Far East country, it is 68.3%.

The reason for this low rate is that Rubella vaccine is currently not included in the national immunization program<sup>(19)</sup>. In

**Table 3.** Meta-analysis of Rubella immunoglobulin-G-avidity test results

Study name and year	Rubella IgG-avidity positive								
	Low avidity			Intermediate avidity			High avidity		
	Sample size	Pro-portion (%)	95% CI	Sample size	Proportion (%)	95% CI	Sample size	Proportion (%)	95% CI
ASIK (2013) <sup>(10)</sup>	43	2.33	0.060 to 12.29	N/A	N/A	N/A	43	97.67	87.71 to 99.94
SEVKI (2013) <sup>(3)</sup>	N/A	N/A	N/A	N/A	N/A	N/A	95	100.00	96.19 to 100.00
ŞİMSEK (2016) <sup>(11)</sup>	16	12.50	1.55 to 38.35	16	12.50	1.55 to 38.35	16	75.00	47.62 to 92.73
SIRIN (2017) <sup>(34)</sup>	54	0.00	0.00 to 6.60	54	3.70	0.45 to 12.75	54	98.15	90.11 to 99.95
VARICI-BALCI (2014) <sup>(13)</sup>	314	7.32	4.70 to 10.79	314	7.64	4.96 to 11.16	314	85.03	80.60 to 88.79
Total (fixed effects)	427	5.88	3.85 to 8.54	384	7.51	5.09 to 10.61	522	91.41	88.68 to 93.66
Total (random effects)	427	4.66	0.97 to 10.91	384	7.51	5.10 to 10.34	522	93.55	82.58 to 99.31
Q	10.3230			1.8145					
DF	3			2					
Significance level	p=0.0160			p=0.4036					
I <sup>2</sup> (inconsistency)	70.94%			0.00%					
95% CI for I <sup>2</sup>	16.99 to 89.83			0.00 to 96.30					

NA: Not applicable

China, although this rate varies according to age in pregnant women, this rate is approximately 84.0%<sup>(20)</sup>. The high level of IgG antibody seropositivity in our country reveals that we have a successful vaccination policy compared with other developing countries.

It is reported that if immunity against Rubella is below 90%, the risk of acute infection and CRS might increase in childbearing age<sup>(9)</sup>. Rubella IgM antibody is the first-step test in determining acute infection. In our meta-analysis, the IgM seroprevalence rate in Turkey was about 0.8%. The study of Başkesen et al.,<sup>(7)</sup> which included 1202 pregnant women in the Manisa region, is one of the most important studies on this subject with 7.6% IgM seropositivity rate and 1.3% weight in meta-analysis. In patients in Manisa, clinical evaluations should be made, IgM results, cross-reactions or primary/re-infection should be confirmed, and it should be kept in mind that antibody positivity may persist for a year following vaccination or asymptomatic infection. There is a need for a prospective randomized controlled trial in the Manisa region to identify these patients.

As a part of the “Elimination of Rubella and CRS Prevention Program”, Rubella vaccine has administered in Turkey, including Isparta, since 2006<sup>(21)</sup>. The Rubella IgM seropositivity rate was found as 4.9% in the retrospective

study of Akpınar et al.<sup>(9)</sup> in Isparta, which comprised 1829 pregnant women. There is not enough information about whether pregnant women are vaccinated because of the fact that the study was retrospective; however, it can still be concluded that the routine vaccination program is not adequately successful in the Isparta province.

In pregnant woman with both IgG and IgM positivity, an IgG avidity test should be performed to estimate the time of infection. According to our analysis, Rubella IgG avidity was studied in only 5 studies in Turkey, and the low avidity rate was 4.6% in a total of 427 pregnant women. One of the reasons for this high rate was the study of Şimsek et al.<sup>(11)</sup> conducted in Afyon. In this study, there were 28 patients with IgM positivity and 44 patients with gray-zone IgM positivity, but avidity was studied in only 12 of them. Two patients with low avidity in this group increased the rate of high avidity in the analysis.

It is not possible to make a detailed and direct comparison through avidity results because publications about CRS are generally case studies in our country. There is a need for a multicenter prospective study for our country including CRS patient series with validation tests, advanced diagnostic tests, and large-scale IgG-avidity results.

There are some restrictions for meta-analyses; there may be

studies that have not completed the publication procedure in the analysis period and also there may be publications for which the full text cannot be reached. There may also be changes in the interpretation of the study results due to differences in the kit and devices used in the studies analyzed. In summary, our study cannot show the whole the Rubella seroprevalence in Turkey. The high seroprevalence of IgG antibodies in Turkey may be considered as the result of successful immunization policies and practices. In a few provinces, it is necessary to revise the Rubella immunization procedures and adult vaccination strategies should be developed in order to control Rubella infections in adults including pregnant woman.

**Acknowledgement:** The authors are very thankful to Prof. Dr. Arzu Kanik for her contribution for the statistical analysis.

**Ethics Committee Approval:** Meta-analysis studies are accepted as the original article. However, ethics committee approval document has not been submitted since it is not required to obtain an ethics committee approval in the meta-analysis studies.

**Informed Consent:** Meta-analysis study.

**Peer-review:** External and internal peer-reviewed.

#### Authorship Contributions

Concept: R.A.C., E.Y., Design R.A.C., E.Y., Data Collection or Processing: R.A.C., E.Y., Analysis or Interpretation: R.A.C., E.Y., Literature Search: R.A.C., E.Y., Writing: R.A.C., E.Y.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

#### References

- Orenstein WA, Cairns L, Hinman A, Nkowane B, Olive JM, Reingold AL. Measles and Rubella Global Strategic Plan 2012-2020 midterm review report: Background and summary. *Vaccine* 2018;36(Suppl1):A35-A42.
- Berger A, Doerr HW. Preventing vertical virus infections: the role of serologic screening of pregnant women. *Med Microbiol Immunol* 2018;207:249-53.
- Sargın A., Saygın S. Seroprevalence of *Toxoplasma gondii* and rubella among pregnant women in central Turkey. *African Journal of Microbiology Research* 2013;7:2524-9.
- Bakacak M, Bostancı MS, Köstü B, Ercan Ö, Serin S, Avcı F. Gebelerde *Toxoplasma gondii*, rubella ve sitomegalovirüs seroprevalansı. *Dicle Med J* 2014;41:326-31.
- Varol FG, Sayin NC, Soysuren S. Seroprevalance of toxoplasma gondii antibodies in antenatal population of Trakya region. *J Turk Soc Obstet Gynecol* 2011;8:93-9.
- Nazik S, Duran İ, Nazik H, Duran Ş. Gebelikte Toksoplazma ve Rubella seropozitifliğinin değerlendirilmesi. *Balıkesir Medical Journal* 2017;1:22-5.
- Başkesen T, Ecemiş T, Şanlıdağ T. Gebelerde rubella bağışıklığının değerlendirilmesi. *Kocatepe Tıp Dergisi* 2010;11:19-21.
- Çeltik NY, Tetikçok R, Günal Ö, Demirtürk F, Duygu F, Barut HŞ, et al. Türkiye'nin Orta Karadeniz Bölgesi'nde Gebelerde Rubella, CMV ve Toksoplazmozis Seroprevalansı Gaziosmanpaşa Üniversitesi Tıp Fakültesi Dergisi 2014;6:54-62.
- Akpınar O, Akpınar H. Gebe Kadınlarda Rubella ve Sitomegalovirus Seroprevalansının ELISA yöntemi ile Araştırılması. *Balıkesir Sağlık Bilimleri Dergisi* 2017;6:11-5.
- Aşık G, Ünlü BS, Er H, Yoldaş Ö, Köken G, Çufalı D, et al. Afyon bölgesinde gebelerde Toksoplazma ve Rubella seroprevalansı. *Pamukkale Tıp Dergisi* 2013;6:128-32.
- Şimşek M, Keşli R, Demir C, Çetinkaya Ö, Ariöz DT. Afyon Kocatepe Üniversitesi, Araştırma ve Uygulama Hastanesinde takip edilen gebelerde Toksoplazma, Rubella, Sitomegalovirus ve Herpes Simpleks Virus Tip 2 seroprevalansının incelenmesi. *Ortadoğu Medical Journal/Ortadoğu Tıp Dergisi* 2016;8:1-6.
- Efe Ş, Kurdoğlu Z, Korkmaz G. Van yöresindeki gebelerde Sitomegalovirüs, Rubella ve Toksoplazma antikorlarının seroprevalansı. *Van Tıp Derg* 2009;16:6-9.
- Varıcı BF, Arslan A, Sertöz R, Altuğlu İ. Rubella and cytomegalovirus seroprevalance among pregnant women admitted to Ege University Hospital. *Ege Tıp Dergisi* 2014;53:179-83.
- Parlak M, Çim N, Erdin BN, Güven A, Bayram Y, Yıldızhan R. Seroprevalence of *Toxoplasma*, Rubella, and Cytomegalovirus among pregnant women in Van. *Turkish journal of obstetrics and gynecology* 2015;12:79.
- Özdemir M, Taşbent FE, Terzi HA, Çetinkol Y, Uysal EB, Aşık G, et al. Seroprevalence of major viral pathogens during pregnancy: A multicenter study in Turkey. *Adv ClinMed Microbiol* 2016;1:1-5.
- Mongua-Rodriguez N, Diaz-Ortega JL, Garcia-Garcia L, Pina-Pozas M, Ferreira-Guerrero E, Delgado-Sánchez G, et al. A systematic review of rubella vaccination strategies implemented in the Americas: impact on the incidence and seroprevalence rates of rubella and congenital rubella syndrome. *Vaccine* 2013;31:2145-51.
- Ghaderi R, Ghaderi F. Rubella Immunity among Pregnant Women in Iran. *MOJ Immunol* 2016;4:00118:1-4. 18.
- Gioula G, Exindari M, Melidou A, Minti F, Sidiropoulou E, Dionisopoulou S, et al. Seroprevalence of measles in Northern Greece 2017;62:145-50.
- Jayakrishnan Thayyil VK, Moorkoth AP, Rao B, Selvam P. Prevalence of rubella-specific IgG antibodies in unimmunized young female population. *Journal of family medicine and primary care* 2016;5:658-62.
- Meng Q, Luo J, Li L, Shi W, Yu J, Shen Y, et al. Rubella seroprevalence among pregnant women in Beijing, China. *BMC infectious diseases* 2018;18:130.
- Genişletilmiş Bağışıklama Programı Genelgesi, Sağlık Bakanlığı Temel Sağlık Hizmetleri Genel Müdürlüğü 2006/120 sayılı genelge. Available from: <https://dosyab.saglik.gov.tr/Eklenti/1117,gbp Genelge2008.pdf>. Available date: 15.01.2019.
- Aynioğlu A, Aynioğlu O, Altunok ES. Seroprevalence of *Toxoplasma gondii*, rubella and Cytomegalovirus among pregnant females in north-western Turkey. *Acta Clinica Belgica* 2015;70:321-4.
- Dogan K, Güraslan H, Özel G, Aydan Z, Yasar L. Seroprevalence Rates of *Toxoplasma gondii*, Rubella, Cytomegalovirus, Syphilis, and Hepatitis B, Seroprevalences Rate in The Pregnant Population in Istanbul. *Türkiye Parazitoloji Dergisi* 2014;38:228.
- Gündem NS, Ağır MÇ. Investigation of seroprevalences of Rubella and *Toxoplasma gondii* among pregnant women. *Anatolian Journal of Clinical Investigation* 2014;8:152-4.
- İnci A, Yener C, Güven D. Bir devlet hastanesinde gebe kadınlarda toksoplazma, rubella ve sitomegalovirüs seroprevalansının araştırılması. *Pamukkale Tıp Dergisi* 2014:143-6.



26. Karabulut A, Polat Y, Türk M, Balci YI. Evaluation of rubella, Toxoplasma gondii, and cytomegalovirus seroprevalences among pregnant women in Denizli province. Turkish Journal of Medical Sciences 2011;41:159-64.
27. Karacan M, Batukan M, Çebi Z, Berberoglugil M, Levent S, Kır M, et al. Screening cytomegalovirus, rubella and toxoplasma infections in pregnant women with unknown pre-pregnancy serological status. Archives of gynecology and obstetrics 2014;290:1115-20.
28. Kasap B, Öner G, Küçük M, Öztürk Turhan N, Akın MN, Arıkan S, et al. Muğla'daki gebelerin Toksoplazma, Rubella, Sitomegalovirüs ve Hepatit prevalansının değerlendirilmesi. İzmir Tepecik Eğitim ve Araştırma Hastanesi Dergisi 2017;27:31-6.
29. Deha K. İlk Trimester Gebelerde Toksoplazma, Rubella, CMV, HBV, AntiHBs, HCV, HIV Seroprevelansları. Selçuk Tıp Derg 2013;3:123-6.
30. Numan O, Vural F, Aka N, Alpay M, Ertürk Coşkun AD. İstanbul Anadolu Kuzey Kamu Hastaneleri Birliği Haydarpaşa Numune Eğitim ve Araştırma Hastanesi Gebe Polikliniği'ne Başvuran Hastalarda TORCH Seroprevalansının Araştırılması 2015;2:203-9.
31. Özdemir M, Kalem F, Feyzioğlu B, Baysal B. Investigation of viral pathogens during pregnancy in a city region in Turkey. AJCI 2011;5:78-81.
32. Kiriş Satılmış Ö, Yapça ÖE, Yapça D, Çatma T. Seroprevalance of Rubella, Cytomegalovirus and Toxoplasma Antibodies Among Pregnant Women that Referred to Sorgun State Hospital. Istanbul Kanuni Sultan Süleyman Tıp Dergisi (IKSST) 2014;6:90-6.
33. Şentürk Ş, Kağıtçı M, Balık G, Şahin K, Şahin FK., Doğu Karadeniz Bölgesi'ndeki Gebe Kadınlarda Rubella Virus Seroprevalansı. Van Tıp Derg 2016;23:242-5.
34. Sirin MC, Agus N, Yılmaz N, Bayram A, Derici YK, Samlioglu P, et al. Seroprevalence of Toxoplasma gondii, Rubella virus and Cytomegalovirus among pregnant women and the importance of avidity assays. Saudi Medical Journal 2017;38:727.
35. Toklu GD. Antibodies Frequency Against Toxoplasmosis, Rubella Virus and Cytomegalovirus in Pregnant Women Gebelerde Toksoplazma, Rubella Virus ve Sitomegalovirus'a Karşı Oluşan Antikorların Sıklığı. J Clin Anal Med 2013;4:38-40.