



Comprehensive analysis reveals a four-gene signature in colorectal cancer

Bin Zhao^{1,2}, Zheng Wan^{1,2}, Xiaohong Zhang^{1,2}, Yilin Zhao^{1,2}

¹Department of Oncology and Vascular Interventional Radiology, Zhongshan Hospital, ²School of Medicine, Xiamen University, Xiamen 361005, China

Contributions: (I) Conception and design: B Zhao; (II) Administrative support: Y Zhao; (III) Provision of study materials or patients: B Zhao; (IV) Collection and assembly of data: B Zhao; (V) Data analysis and interpretation: B Zhao, Z Wan, X Zhang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Yilin Zhao. Department of Oncology and Vascular Interventional Radiology, Zhongshan Hospital, Xiamen University, Xiamen 361005, China; School of Medicine, Xiamen University, Xiamen 361005, China. Email: zhaoboxmu@xmu.edu.cn.

Background: Colorectal cancer (CRC) is one of the major malignant diseases of the gastrointestinal system around the world. However, the current therapeutic regimens were not always effective. This study was designed to identify and depict potential molecular biomarkers and correlated signal pathways in CRC.

Methods: The gene expression profiles of GSE21510 were obtained on the Gene Expression Omnibus website, we filtered out 44 samples from the GSE21510 to identify different expression genes (DEGs) between CRC tissues and noncancerous tissues. Subsequently, the function and signal pathways enrichment analyses were implemented, the protein-protein interaction (PPI) networks of DEGs were to be carried out, and the hub genes were screened by MCODE built in Cytoscape software. Lastly, we have validated gene expressions and overall survival analyses of these hub genes in related datasets, such as colon adenocarcinoma (COAD) and rectum adenocarcinoma (READ), built in TCGA/GTEx database.

Results: Results showed that a totally of 166 up-regulated genes and 260 down-regulated genes were identified and met the following criteria: $|\log_2$ fold change ≥ 2 & adjusted P value < 0.01 . Here, we identified *AURKA*, *BUB1*, *DLGAP5* and *HMMR*, which were associated with the regulation of mitotic cycle phase transition and oocyte meiosis pathways.

Conclusions: The findings of these four genes in this study may shed light on the mechanisms of these four genes as drug-sensitive therapeutic targets for the patients of CRC.

Keywords: Differentially expressed genes (DEGs); biomarkers; colorectal cancer (CRC); colon cancer; survival analyses

Submitted Oct 27, 2019. Accepted for publication Jan 06, 2020.

doi: 10.21037/tcr.2020.01.18

View this article at: <http://dx.doi.org/10.21037/tcr.2020.01.18>

Introduction

Colorectal cancer (CRC) is known as one of the most common non-skin cancers diagnosed both in men and women in the world. CRC often begins as a polyp inside the colon or rectum. Adenomas, a form of polyps, are innocent tumors within the tissue of the colon or rectum (1). Though most polyps will stay benign, some of them have the probabilities of converting into cancer as time goes on.

There were reported more than 9.4 million new cases in 2015 and nearly 832,000 deaths in developed countries (2,3).

As a heterogeneous disease, CRC is associated with gene aberration, the microenvironment of tumor initiation, progression and metastasis (4). Over the last decade, many molecular biomarkers and signal pathways associated with the occurrence and progression of CRC have been reported, which has been involved in clinical therapy. To date, owing to the high incidence and mortality in the CRC disease,

uncovering the causes and the molecular characterization of CRC to discover the molecular biomarkers for initial diagnosis, prevention and personalized therapy is quite urgent and demanded.

The recent high-throughput sequence techniques for the analyses of different gene expression and alternative splicing variants, like microarrays or the RNA-seq chip, are increasingly valued as promising techniques in medical oncology with great clinical applications, such as molecular diagnosis, prognosis prediction, drug targets discovery. Many gene expression profiling studies have been focused on CRC in the last decade (5), and hundreds of potential gene biomarkers have been obtained (6), which involved in different functional enrichments [including biological process (BP), cellular components (CC), molecular function (MF)], signal pathways and protein-protein interaction (PPI) networks.

In this study, we have obtained the dataset (GSE21510) from National Center Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) database (NCBI-GEO) (website: <https://www.ncbi.nlm.nih.gov/geo/>), and we have chosen 25 CRC and 19 noncancerous tissue samples for differentially expressed genes (DEGs) analysis, gene ontology (GO) enrichment analysis, Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis, PPI networks complex construction, and hub genes exploration. What's more, we have validated gene expression levels and overall survival (OS) analyses of these hub genes in related datasets [colon adenocarcinoma (COAD) and rectum adenocarcinoma (READ)] built in the TCGA/GTEX database. Moreover, the filtered potential genes associated with CRC be recognized as biomarkers for diagnosis, prognosis and drug targets. *Figure 1A* shows the workflow of this study.

Methods

Data collection

The gene expression microarray data GSE21510 (7) were collected from the NCBI-GEO database (8,9). The GSE21510, taken the Affymetrix GPL570 platform (Affymetrix Human Genome U133 Plus 2.0 Array) as a reference, was submitted by Kaoru Mogushi *et al.*, which featured 104 CRC patients. We choose 19 patients (CRC tissues, cancer group) and 25 patients (noncancerous tissues, normal group) to identify the genes and pathways.

Data preprocessing

After GSE21510 was obtained, probe identification numbers (IDs) were converted into gene symbols or ENTREZID. For multiple probes corresponding to the same one gene, their most significantly expressed value was treated as the gene expression value. Non-mRNA probes were discarded. Then, the gene expression values were normalized by using the Affy package, and RMA signal intensity was performed with log₂ transformation and normalization (10,11).

Identification of DEGs

Linear models for microarray data (*limma*) is an R package applied to analyze gene expression matrix, especially when the linear models are constructed to assess the differentially expressed gene expression under the designed experiment condition (12). *Limma* package (<http://bioconductor.org/packages/2.4/bioc/html/limma.html>) built in R was applied to identify the DEGs between CRC tissues (cancer group) and noncancerous tissues (normal group). Significant DEGs were selected for further analyses with setting the |log₂ fold change (FC)| ≥ 2 & the adjusted P value (adj.P.Val) < 0.01 (13).

Function and signal pathway enrichment analysis

GO provides a controlled vocabulary of terms to elucidate a gene product's characteristics via their annotation. GO terms reflect what is currently known about a gene in terms of BP, CC and MF (14,15). Moreover, KEGG (16) provides data resources of known biological pathways to annotate a gene or a set of genes/proteins with their respective KEGG pathways. In order to illustrate the function and signal pathway analysis of DEGs, GO and KEGG pathway enrichment analyses were carried out with using the *clusterProfiler* package and *ReactomePA* package (17,18) in R and P value < 0.05 was considered significance.

PPI network and gene module analysis

Search Tool for the Retrieval of Interacting Genes (STRING) (19) is an open access database designed to evaluate the PPI messages of DEGs. STRING (version 10.5) covers 9.6 million proteins originated from 2,031 organisms. At first, we uploaded and mapped the list of DEGs to STRING website. Then PPIs of DEGs with a combined score > 0.4 (medium confidence) and genes closely correlated with the other genes were chosen with the degree




Figure 1 The pipeline of screening of differentially expressed genes (DEGs) and some descriptions of DEGs. (A) The workflow of research. (B) Volcano plot of DEGs. Red: up-regulated DEGs; Blue: down-regulated DEGs. (C) Heatmap of the 30 significantly DEGs (15 up-regulated DEGs and 15 down-regulated DEGs, respectively).

≥ 10 (20). After that, PPI networks were constructed by using the Cytoscape software (21). The plug-in Molecular Complex Detection (MCODE) built in Cytoscape was applied to pick out the significant gene modules of the PPI networks. The parameters were set as follows: MCODE scores >3 and the count of nodes >4 . Finally, we selected two significant gene modules (including 46 genes) from the PPI networks for further validation analyses.

Validation of four genes in TCGA/GTEX

To further screen for precise biomarkers, we have validated these 46 genes at gene expression level and OS time on web server GEPIA, which integrated COAD and READ datasets. The gene expression consistency and the survival analyses of these candidate genes were tested and evaluated

in GEPIA. As for gene expression validation involved in 367 tumor samples (COAD: 275, READ: 92) and 667 normal samples (COAD: 349, READ: 318), the threshold with $|\log_2 \text{FC}| \geq 2$ & P value <0.01 was considered statistically significant. For the OS analyses in integrated COAD & READ datasets, the 362 patients with available OS time data were sorted into low- and high-expression groups by the median transcripts per kilobase million (TPM), and significance was decided by the log-rank test with $P < 0.05$.

Results

DEGs identification

Microarray data of 19 CRC tissues (cancer group) and 25 noncancerous tissues (normal group) were analyzed by the

Table 1 The identified differentially expressed genes

DEGs	Gene symbol
Up-regulated	CLDN1, CEMIP, CDH3, ENC1, PPM1H, SLC6A6, ASCL2, VSNL1, FOXQ1, FABP6, PPAT, NUFIP1, NUDCD1, RAD54B, C2, HILPDA, MYC, CRNDE, NFE2L3, CKAP2, AJUBA, IRAK1BP1, CDK4, ATP11A, SLC7A5, EPHX4, LOC101060264, NEBL, PUS7, TRIB3, HOMER1, FAM60A, UTP14A, AXIN2, CSE1L, PPIL1, LYAR, DGAT2, SLC39A10, PAICS, CCDC113, TOP1MT, PROCR, MCM2, ZAK, ZC3HAV1L, HSP90AB1, TMPRSS3, UBE2C, LGR5, NUF2, RFC3, CDC25B, PRMT3, HELLS, DSCC1, PALD1, SLC04A1, PLAU, ATAD2, BICD1, PARBP, E2F7, TGFBI, PMAIP1, RPP40, AZGP1, DPEP1, KRT23, SPC25, MET, SP5, KIF20A, TRIP13, CDKN3, SLC12A2, PABPC1L, MMP1, ANLN, HS2ST1, CDK1, FAM92A1, CXCL3, SLC22A3, RNF43, ZNRF3, DUSP14, PSAT1, TPX2, ACSL4, AURKA, CKS2, MKI67, LDLRAD3, RAD51AP1, PHLDA1, SUPT16H, INHBA, SRPX2, MMP7, CXCL1, RASSF10, KIF4A, CXCL8, MMP3, TOP2A, BMP4, TESC, TTK, CTHRC1, HMMR, CDCA7, BUB1, DACH1, SLC38A5, CYP4X1, C2CD4A, GNG4, ELOVL5, CXCL2, FAM83D, CHI3L1, KLK10, TCFL5, LEF1, TACSTD2, DLGAP5, PRC1, ACSL6, NKD1, CADPS, NEK2, CENPK, MMP12, WT1, SLC35D3, CLDN2, PRR11, GINS1, COL11A1, CXCL10, CCNB1, CEP55, PTPRO, PLEKHB1, RGCC, CXCL11, SLC01B3, GDF15, HS6ST2, RRM2, GAL, APCDD1, CXCL5, KRT6B, CKMT2, COL4A1, COL1A2, EDNRA, SPP1, TCN1, REG3A, AMIGO2, PPBP, WDR72, COL15A1
Down-regulated	UGP2, EPB41L3, GLTP, SEMA6A, SCARA5, PPAP2A, UGCG, IL6R, RELL1, TP53INP2, GCNT2, SPPL2A, ETFDH, ADH1B, SGK1, MXI1, CNTN3, NR3C2, MMP28, ABCA8, ZZEF1, C2orf88, RHOU, TMEM220, SYTL4, KLF4, AGPAT9, SMIM14, KAT2B, ABCG2, METTL7A, TMCC3, FRMD3, SLC2A13, PARM1, SMPD1, ARRDC4, TSPAN7, ABI3BP, RUNDC3B, SLC30A4, SLC4A4, GRAMD3, ADCY9, SRI, HPGD, GUCA2A, MIER3, PLCE1, SLC02A1, LIFR, PCSK5, PRKACB, FAM107B, TMEM100, WDR78, SCIN, FAM46A, MT1M, GUCA2B, SLC51B, PDE9A, PLCL2, SPIB, C1orf115, NR5A2, CDHR5, USP2, CA7, PADI2, CD177, BEST4, CPNE8, ENTPD5, HOXD1, PDE3A, PHLPP2, ANO5, CXCL12, SLC30A10, TEX11, KIAA1211, SEMA6D, SFRP1, TLCD2, TMEM56, RMDN2, MAOA, APPL2, ITM2C, MXD1, RHOF, NPY1R, SCNN1B, LDHD, PDK4, BHLHE41, PTPRH, BMP2, ADTRP, CCDC68, ARHGAP44, CHP2, ARHGAP42, PAG1, MT1E, PKIB, MEIS3P1, MT1HL1, MT1X, HDAC9, SSPN, SMPDL3A, ENDOD1, CLDN23, EDIL3, EPB41L4A, OSBPL1A, MT2A, HRCT1, MT1H, GBA3, CMAHP, SLC17A4, VLDLR, ITM2A, MALL, NAAA, GDDP3, AHCYL2, AQP8, TMEM72, CHST5, THRB, TBC1D9, SLC1A1, HSD17B2, SRPX, FGL2, HAGLR, CLU, CDHR2, KIF16B, VSIG2, CCL28, CA4, BEST2, MT1G, TNFSF10, MT1F, VILL, F2RL1, CES2, LAMA1, SLC41A2, RNF125, EGLN3, DHRS11, RETSAT, EMP1, OGN, CA2, CHRDL1, TMEM171, CWH43, CEACAM7, ST6GALNAC6, B3GALT5, LRRC19, ADH1C, FMO5, LGALS2, PTPRR, PRSS12, ARL14, LRRC66, PTGDR, SPON1, ZBTB7C, ANPEP, PIGR, EDN3, DHRS9, ZG16, HHLA2, PROM2, BCAS1, AKR1B10, CLCA4, LINC01133, BTNL8, MUC12, MALAT1, SLC16A9, TUBAL3, UGT2A3, HEPACAM2, NXPE4, CLIC6, TTC22, FAM134B, MS4A12, CA1, TRPM6, SI, CEACAM1, CYBRD1, IQGAP2, SLC16A14, SLC26A2, LYPD8, FOXP2, MYLK, HMGCS2, CFD, RASSF6, HSD11B2, GHR, AKAP7, DEFB1, DSC2, SULT1B1, PCK1, B3GNT7, NXPE1, MOGAT2, MEIS1, C4orf19, GGT6, IL1R2, MGP, SCNN1G, FHL1, LEPREL1, PLAC8, ENPP3, CLDN8, GCG, TSPAN1, ABCB1, NR1H4, MFAP5, FCGBP, VIP, RARRES1, SLC51A, SPINK5, CAPN13, INSL5, MEP1A, IGJ, GCNT3, ISX, MUC2, SYNPO2, AGR3, SFRP2, CLCA1, ITLN1, HSD3B2

limma package built in R through the linear model and the contrast model to identify DEGs. A total of 426 DEGs were selected by the criteria of adjusted P value <0.01 & $|\log_2 FC| \geq 2$, including 166 up-regulated genes and 260 down-regulated genes (Figure 1B, Table 1). The hierarchical cluster analysis was done to show the most 15 significantly up-regulated genes and 15 down-regulated genes in Figure 1C. The top ten genes with the most significant expression were *CLDN1*, *CEMIP*, *UGP2*, *EPB41L3*, *CDH3*, *ENC1*, *PPM1H*, *GLTP*, *SLC6A6* and *SEMA6A* (Table 1).

Function and signal pathway enrichment analysis

To investigate functions and signal pathway enrichment of identified DEGs, we further analyzed DEGs using the

clusterProfiler package and *ReactomePA* package in R with criteria of $P < 0.05$. Figure 2A show the significant ten BP, CC and MF enrichment terms, respectively. Moreover, Figure 2B shows the top 30 significant GO terms.

As shown in Table 2, in BP term, the up-regulated genes were mainly enriched in the nuclear division, mitotic nuclear division and organelle fission, while the down-regulated genes were focused on the detoxification of copper ion, stress response to copper ion and detoxification of inorganic compound. In CC term, the up-regulated genes were mainly enriched in the spindle pole, spindle, chromosome and centromeric region, while the down-regulated genes were focused on the microvillus membrane, apical part of cell and apical plasma membrane. In MF term, the up-regulated genes were mainly enriched in the




Figure 2 The gene ontology and signal pathway enrichments. (A) The top ten functional enrichment analysis of DEGs in biological process, cellular component and molecular function group, respectively; (B) the top 30 significantly enriched GO terms of DEGs; (C) the significantly enriched signal pathways of DEGs in CRC. DEG, differentially expressed gene; GO, gene ontology; SLC, solute carrier.

CXCR chemokine receptor binding, chemokine activity and chemokine receptor binding, while the down-regulated genes were focused on the oxidoreductase activity, phosphoric diester hydrolase activity and oxidoreductase activity.

Figure 2C shows the significant pathways in which the most signal pathways were enriched in the metallothioneins bind metals, response to metal ions and chemokine receptors bind chemokines (Figure 2C). The up-regulated genes were mainly enriched in the chemokine receptors bind chemokines, collagen degradation and degradation of the extracellular matrix (Table 2), and the down-regulated genes

were enriched in the metallothioneins bind metals, response to metal ions and stimuli-sensing channels (Table 2).

Module screening from the PPI network

Firstly, 426 DEGs were uploaded into the STRING website and analyzed by Cytoscape software. A total of 269 DEGs with score >0.4 (medium confidence) were picked out for the construction of the PPI networks (Figure 3A). Then, two significant gene modules were clustered via MCODE APP built in Cytoscape. Module 1 was made up of 31 up-

Table 2 The top three gene ontology and pathway enrichment terms of up-regulated and down-regulated genes, respectively

Terms	Category	Description	FDR	Count
Up-regulated genes				
GO: 0000280	BP	Nuclear division	4.27E-07	21
GO: 0140014	BP	Mitotic nuclear division	5.16E-07	17
GO: 0048285	BP	Organelle fission	9.75E-07	21
GO: 0000922	CC	Spindle pole	0.000108	10
GO: 0005819	CC	Spindle	0.000108	14
GO: 0000775	CC	Chromosome, centromeric region	0.000574	10
GO: 0045236	MF	CXCR chemokine receptor binding	2.53E-10	8
GO: 0008009	MF	Chemokine activity	1.84E-06	8
GO: 0042379	MF	Chemokine receptor binding	1.28E-05	8
R-HSA-380108	KEGG	Chemokine receptors bind chemokines	7.09E-06	8
R-HSA-1442490	KEGG	Collagen degradation	3.65E-05	8
R-HSA-1474228	KEGG	Degradation of the extracellular matrix	0.001122	9
Down-regulated genes				
GO: 0010273	BP	Detoxification of copper ion	5.97E-09	8
GO: 1990169	BP	Stress response to copper ion	5.97E-09	8
GO: 0061687	BP	Detoxification of inorganic compound	1.10E-08	8
GO: 0031528	CC	Microvillus membrane	5.58E-05	6
GO: 0045177	CC	Apical part of cell	5.58E-05	19
GO: 0016324	CC	Apical plasma membrane	5.58E-05	17
GO: 0016614	MF	Oxidoreductase activity, acting on CH-OH group of donors	0.00075	11
GO: 0008081	MF	Phosphoric diester hydrolase activity	0.00075	9
GO: 0016616	MF	Oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	0.00075	10
R-HSA-5661231	KEGG	Metallothioneins bind metals	1.59E-08	7
R-HSA-5660526	KEGG	Response to metal ions	7.97E-08	7
R-HSA-2672351	KEGG	Stimuli-sensing channels	0.000575	10

GO, gene ontology; FDR, false discovery rate; BP, biological process; CC, cellular components; MF, molecular function; KEGG, Kyoto Encyclopedia of Genes and Genomes.

regulated genes/nodes and 427 edges (*Figure 3B*), while module 2 consisted of 15 genes/nodes (10 up-regulated and 5 down-regulated genes) and 105 edges (*Figure 3C*).

Validation of four genes in TCGA/GTEX

Genes from those two gene modules were chosen for

validation in COAD and READ built in TCGA/GTEX datasets. We further found that only the four gene expressions of *AURKA*, *BUB1*, *DLGAP5* and *HMMR* in COAD and READ datasets have consistency with that in GSE21510 (*Figure 4A*). In addition, their OS is significantly different between high expression and low expression (*Figure 4B*). GO and KEGG pathways show that these four




Figure 3 The protein-protein interaction (PPI) networks construction and significant gene modules analysis. (A) The PPI networks of DEGs (orange and blue represents the most two significant gene modules, respectively); (B) module 1 consists of 31 nodes/genes (orange indicates an up-regulated gene); (C) module 2 consists of 15 nodes/genes (orange indicates an up-regulated gene and light blue indicates a down-regulated gene).




Figure 4 The validation of the final potential four genes and its functional annotation. (A) Validation of the gene expression of *AURKA*, *BUB1*, *DLGAP5* and *HMMR* in COAD & READ datasets. The cutoff: $|\log_2 \text{fold change (FC)}| \geq 2$, and $P < 0.01$ (* indicates $P < 0.01$). (B) Overall survival (OS) analysis of the *AURKA*, *BUB1*, *DLGAP5* and *HMMR* in COAD & READ datasets. (C) Chord plot for functional enrichments of four genes. COAD, colon adenocarcinoma; READ, rectum adenocarcinoma; HR, hazard ratio; TPM, transcripts per kilobase million.

candidate genes are significantly enriched in the regulation of mitotic cycle phase transition and oocyte meiosis (Figure 4C and Table 3).

Discussion

In this study, our purposes were to expound the potential genetic biomarkers and pathways through comparing the array datasets between cancer group (CRC tissues) and Normal group (noncancerous tissues), and 166 up-regulated and 260 down-regulated DEGs were identified. Then the function (GO) and signal pathway (KEGG) annotation analyses have been performed. Moreover, the PPI networks of DEGs were constructed and 269 DEGs/nodes were connected to 1,169 edges, and finally the most two significant modules were chosen from the PPIs, from which 46 central nodes/genes were selected to validate its gene expressions and OS time in TCGA/GTEX.

As we knew that mutations in the Wnt signaling pathway and inflammatory bowel disease are the two major causes of CRC (22). Through bioinformatics analyses, we have identified 46 central nodes/genes, among them, the first

significant gene module (Figure 3B) consists of 31 genes, including *TOP2A*, *CDK1*, *CCNB*, *MYC*, *AURKA*, *BUB1*, etc., and the second significant gene module (Figure 3C) consists of 15 genes, including *CXCL12*, *CCL28*, *CXCL2*, *GNG4*, *CXCL1*, *INSL5*, etc. However, some of genes in these two significant gene modules, associated with CRC, have been researched and identified in the past years. Finally, we screened four genes (*AURKA*, *BUB1*, *DLGAP5* and *HMMR*), which have been consistent with their gene expression level in tumor and normal patients of COAD & READ. Besides, we discarded the other 42 genes which didn't meet the threshold we set.

Aurora A kinase (*AURKA*) is encoded by the *AURKA* gene and a member of the serine/threonine kinases family (23,24). *AURKA* has been shown to interact with *Wnt* and *Ras-MAPK* signaling in CRC (25). What's more, it was reported that *AURKA* has associated with CRC liver metastasis (CRLCM) (26,27). Budding uninhibited by benzimidazoles (*BUB1*) is also a member of the serine/threonine-protein kinase family (28). Over 90 percentages of all human solid tumors have a common feature that mutations occur in the spindle checkpoints (29). Jaffrey *et al.* have suggested that mutations in *BUB1* can lead to

Table 3 Gene ontology and pathways enrichment of identified four genes

Terms	Category	Description	FDR	Gene	Count
GO: 1901990	BP	Regulation of mitotic cell cycle phase transition	0.00025	<i>AURKA, BUB1, DLGAP5, HMMR</i>	4
GO: 0007088	BP	Regulation of mitotic nuclear division	0.00073	<i>AURKA, BUB1, DLGAP5</i>	3
GO: 0000280	BP	Nuclear division	0.001	<i>AURKA, BUB1, DLGAP5</i>	3
GO: 0030071	BP	Regulation of mitotic metaphase/anaphase transition	0.0031	<i>BUB1, DLGAP5</i>	2
GO: 0045840	BP	Positive regulation of mitotic nuclear division	0.0031	<i>AURKA, DLGAP5</i>	2
GO: 0032436	BP	Positive regulation of proteasomal ubiquitin-dependent protein catabolic process	0.0058	<i>AURKA, DLGAP5</i>	2
GO: 1903047	BP	Mitotic cell cycle process	0.0058	<i>AURKA, BUB1, DLGAP5</i>	3
GO: 0051781	BP	Positive regulation of cell division	0.0066	<i>AURKA, DLGAP5</i>	2
GO: 0000819	BP	Sister chromatid segregation	0.0076	<i>BUB1, DLGAP5</i>	2
GO: 0140014	BP	Mitotic nuclear division	0.0082	<i>AURKA, DLGAP5</i>	2
GO: 0007093	BP	Mitotic cell cycle checkpoint	0.0089	<i>AURKA, BUB1</i>	2
GO: 0010389	BP	Regulation of G2/M transition of mitotic cell cycle	0.0089	<i>AURKA, HMMR</i>	2
GO: 0140013	BP	Meiotic nuclear division	0.0089	<i>AURKA, BUB1</i>	2
GO: 1901991	BP	Negative regulation of mitotic cell cycle phase transition	0.0091	<i>AURKA, BUB1</i>	2
GO: 0051276	BP	Chromosome organization	0.0128	<i>AURKA, BUB1, DLGAP5</i>	3
GO: 0051301	BP	Cell division	0.0453	<i>AURKA, BUB1</i>	2
GO: 0031616	CC	Spindle pole centrosome	0.00041	<i>AURKA, DLGAP5</i>	2
GO: 0000780	CC	Condensed nuclear chromosome, centromeric region	0.00062	<i>AURKA, BUB1</i>	2
GO: 0005813	CC	Centrosome	0.0038	<i>AURKA, DLGAP5, HMMR</i>	3
GO: 0043232	CC	Intracellular non-membrane-bounded organelle	0.0395	<i>AURKA, BUB1, DLGAP5, HMMR</i>	4
hsa04114	KEGG	Oocyte meiosis	0.0028	<i>AURKA, BUB1</i>	2
hsa04914	KEGG	Progesterone-mediated oocyte maturation	0.0028	<i>AURKA, BUB1</i>	2

GO, gene ontology; FDR, false discovery rate; BP, biological process; CC, cellular components; KEGG, Kyoto Encyclopedia of Genes and Genomes.

chromosome instability in cancer cell lines (30). Disks large-associated protein 5 (*DLGAP5*) is a kinetochore protein that plays a role in stabilizing microtubules in chromosomes, controlling spindle dynamics, promoting interkinetochore tension and executing efficient kinetochore capture (31). However, Schneider MA's team has predicted that *AURKA* and *DLGAP5* could have a correlation with poor prognosis in non-small cell lung cancer patients (32). By regulating *DLGAP5* gene expression, it could enhance the efficacy of epirubicin for invasive breast cancer (33). Hyaluronan-mediated motility receptor (*HMMR*), known as receptor for hyaluronan mediated motility (*RHAMM*), has a high-

level gene expression and correlation with poor outcome in breast cancer (34). What's more, *HMMR* may be associated with the risk of breast cancer patients who have *BRCA1* mutation (35).

As shown in *Figure 4B*, you will be surprised to find that the low TPM group of these four genes have low percent survival than that in high TPM group, which are contrary to what we expected. But the survival curve of these four genes was statistically significant, and it is essential for us to seek the reasons why it occurs.

The above four genes were chosen via comprehensive bioinformatics analyses and mainly enriched in the

regulation of mitotic cycle phase transition and oocyte meiosis pathway. However, further molecular and cellular experiments are required to verify the function of these four gene biomarkers in CRC.

Acknowledgments

Funding: This project was supported and funded by National Natural Science Foundation of China (No. 81770294).

Footnote

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr.2020.01.18>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. For human datasets mentioned in this study, please refer to the original article (PMID: 21270110). We just re-analyzed the open accessed datasets, and no ethical approval or informed consent was required by the local ethics committees. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Giardiello FM, Hamilton SR, Krush AJ, et al. Treatment of Colonic and Rectal Adenomas with Sulindac in Familial Adenomatous Polyposis. *New Engl J Med* 1993;328:1313-6.
- Siegel RL, Miller KD, Fedewa SA, et al. Colorectal cancer statistics, 2017. *CA Cancer J Clin* 2017;67:177-93.
- Vos T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388:1545-602.
- Aran V, Victorino AP, Thuler LC, et al. Colorectal Cancer: Epidemiology, Disease Mechanisms and Interventions to Reduce Onset and Mortality. *Clin Colorectal Cancer* 2016;15:195-203.
- Isella C, Terrasi A, Bellomo SE, et al. Stromal contribution to the colorectal cancer transcriptome. *Nat Genet* 2015;47:312-9.
- Vogelstein B, Papadopoulos N, Velculescu VE, et al. Cancer genome landscapes. *Science* 2013;339:1546-58.
- Tsukamoto S, Ishikawa T, Iida S, et al. Clinical significance of osteoprotegerin expression in human colorectal cancer. *Clin Cancer Res* 2011;17:2444-50.
- Barrett TT, Dennis B, Wilhite SE, et al. NCBI GEO: archive for high-throughput functional genomic data. *Nucleic Acids Res* 2009;37:D885-90.
- Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* 2002;30:207-10.
- Gautier L, Cope L, Bolstad BM, et al. affy—analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 2004;20:307-15.
- Larriba Y, Rueda C, Fernández MA, et al. Microarray Data Normalization and Robust Detection of Rhythmic Features. In: Bolón-Canedo V, Alonso-Betanzos A. editors. *Microarray Bioinformatics*. New York, NY: Humana, 2019:207-25.
- Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004;3:Article3.
- Reiner A, Yekutieli D, Benjamini Y. Identifying differentially expressed genes using false discovery rate controlling procedures. *Bioinformatics* 2003;19:368-75.
- Hale ML, Thapa I, Ghersi D. FunSet: an open-source software and web server for performing and displaying Gene Ontology enrichment analysis. *BMC Bioinformatics* 2019;20:359.
- Carbon S, Dietze H, Lewis S, et al. Expansion of the Gene Ontology knowledgebase and resources. *Nucleic Acids Res* 2017;45:D331-8.
- Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 2000;28:27-30.
- Yu G, He QY. ReactomePA: an R/Bioconductor package for reactome pathway analysis and visualization. *Mol Biosyst* 2016;12:477-9.
- Yu G, Wang LG, Han Y, et al. clusterProfiler: an R

- package for comparing biological themes among gene clusters. *OMICS* 2012;16:284-7.
19. Franceschini A, Szklarczyk D, Frankild S, et al. STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 2013;41:D808-15.
 20. Zhou Z, Cheng Y, Jiang Y, et al. Ten hub genes associated with progression and prognosis of pancreatic carcinoma identified by co-expression analysis. *Int J Biol Sci* 2018;14:124-36.
 21. Smoot M, Ono K, Ruscheinski J, et al. Cytoscape 2.8. *Bioinformatics* 2011;27:431-2.
 22. Hu T, Li LF, Shen J, et al. Chronic inflammation and colorectal cancer: the role of vascular endothelial growth factor. *Curr Pharm Des* 2015;21:2960-7.
 23. Roskoski R Jr. Cyclin-dependent protein serine/threonine kinase inhibitors as anticancer drugs. *Pharmacol Res* 2019;139:471-88.
 24. Rozpędek W, Pytel D, Nowak-Zduńczyk A, et al. Breaking the DNA Damage Response via Serine/Threonine Kinase Inhibitors to Improve Cancer Treatment. *Curr Med Chem* 2019;26:1425-45.
 25. Jacobsen A, Bosch LJW, Martens-de Kemp SR, et al. Aurora kinase A (AURKA) interaction with Wnt and Ras-MAPK signalling pathways in colorectal cancer. *Sci Rep* 2018;8:7522.
 26. Goos JA, Coupe VM, Diosdado B, et al. Aurora kinase A (AURKA) expression in colorectal cancer liver metastasis is associated with poor prognosis. *Br J Cancer* 2013;109:2445-52.
 27. Goos JA, Coupe VM, van de Wiel MA, et al. A prognostic classifier for patients with colorectal cancer liver metastasis, based on AURKA, PTGS2 and MMP9. *Oncotarget* 2016;7:2123-34.
 28. Matsubara Y, Matsumoto T, Yoshiya K, et al. Budding uninhibited by benzimidazole-1 insufficiency prevents acute renal failure in severe sepsis by maintaining anticoagulant functions of vascular endothelial cells. *Shock* 2019;51:364-71.
 29. Williams BR, Amon A. Aneuploidy: cancer's fatal flaw? *Cancer Res* 2009;69:5289-91.
 30. Jaffrey RG, Pritchard SC, Clark C, et al. Genomic instability at the BUB1 locus in colorectal cancer, but not in non-small cell lung cancer. *Cancer Res* 2000;60:4349-52.
 31. Wilde A. "HURP on" we're off to the kinetochore! *J Cell Biol* 2006;173:829-31.
 32. Schneider MA, Christopoulos P, Muley T, et al. AURKA, DLGAP5, TPX2, KIF11 and CKAP5: Five specific mitosis-associated genes correlate with poor prognosis for non-small cell lung cancer patients. *Int J Oncol* 2017;50:365-72.
 33. Zhang X, Pan Y, Fu H, et al. Nucleolar and Spindle Associated Protein 1 (NUSAP1) Inhibits Cell Proliferation and Enhances Susceptibility to Epirubicin In Invasive Breast Cancer Cells by Regulating Cyclin D Kinase (CDK1) and DLGAP5 Expression. *Med Sci Monit* 2018;24:8553-64.
 34. Rhodes DR, Yu J, Shanker K, et al. Large-scale meta-analysis of cancer microarray data identifies common transcriptional profiles of neoplastic transformation and progression. *Proc Natl Acad Sci U S A* 2004;101:9309-14.
 35. Blanco I, Kuchenbaecker K, Cuadras D, et al. Assessing associations between the AURKA-HMMR-TPX2-TUBG1 functional module and breast cancer risk in BRCA1/2 mutation carriers. *PLoS One* 2015;10:e0120020.

Cite this article as: Zhao B, Wan Z, Zhang X, Zhao Y. Comprehensive analysis reveals a four-gene signature in colorectal cancer. *Transl Cancer Res* 2020;9(3):1395-1405. doi: 10.21037/tcr.2020.01.18