

Comment on 'The impact of vitamin D pathway genetic variation and circulating 25-hydroxyvitamin D on cancer outcome: systematic review and meta-analysis'

Alain Braillon^{*,1}

¹Department of Medicine, University Hospital, 80000 Amiens, France

Sir,

Vaughan-Shaw *et al* (2017) claimed that 25(OH)D concentration is associated with better cancer outcome, and the observed association of functional variants in vitamin D pathway genes with outcome supports a causal link. This deserves a comment.

First, using arbitrary cut-points to derive subgroups for 25(OH)D is not appropriate when there is a continuous distribution of the values with no obvious modal values.

Second, no adjustments for main confounding clinical variables were performed. For smoking it would have been a catch-22: (a) smokers have lower 25(OH)D (Tønnesen *et al*, 2016); (b) 25(OH)D is associated with higher risk of tobacco-related cancers (Afzal *et al*, 2013). Similarly, alcohol consumption, obesity, overweight, insulin resistance, type 2 diabetes have an impact on vitamin D status (Palaniswamy *et al*, 2017).

Long ago, in 1998, a prospective survey (NHANES III) investigated 25(OH)D levels with mortality, accounting for age, sex, ethnicity, diabetes, current smoking, body mass index, physical activity, supplementation, season and so on (Melamed *et al*, 2008). Personalised medicine is first about phenotyping not genotyping!

REFERENCES

- Afzal S, Bojesen SE, Nordestgaard BG (2013) Low plasma 25-hydroxyvitamin D and risk of tobacco-related cancer. *Clin Chem* **59**: 771–780.
- Melamed ML, Michos ED, Post W, Astor B (2008) 25-hydroxyvitamin D levels and the risk of mortality in the general population. *Arch Intern Med* **168**: 1629–1637.
- Palaniswamy S, Hyppönen E, Williams D, Jokelainen J, Lowry E, Keinänen-Kiukaanniemi S, Herzig KH, Järvelin MR, Sebert S (2017) Potential determinants of vitamin D in Finnish adults: a cross-sectional study from the Northern Finland birth cohort 1966. *BMJ Open* **7**: e013161.
- Tønnesen R, Hovind PH, Jensen LT, Schwarz P (2016) Determinants of vitamin D status in young adults: influence of lifestyle, sociodemographic and anthropometric factors. *BMC Public Health* **16**: 385.
- Vaughan-Shaw PG, O'Sullivan F, Farrington SM, Theodoratou E, Campbell H, Dunlop MG, Zgaga L (2017) The impact of vitamin D pathway genetic variation and circulating 25-hydroxyvitamin D on cancer outcome: systematic review and meta-analysis. *Br J Cancer* **116**: 1092–1110.

CONFLICT OF INTEREST

The author declares no conflict of interest.

This work is published under the BJC's standard license to publish agreement. After 12 months the license terms will change to a Creative Commons AttributionNonCommercial-Share Alike 4.0 Unported License.

*Correspondence: Dr A Braillon; E-mail: braillon.alain@gmail.com

Published online 22 June 2017

© 2017 Cancer Research UK. All rights reserved 0007–0920/17