


 Cite this: *RSC Adv.*, 2021, 11, 18040

The graphite-catalyzed *ipso*-functionalization of arylboronic acids in an aqueous medium: metal-free access to phenols, anilines, nitroarenes, and haloarenes†

 Anshu Dandia,^{*a} Ruchi Sharma,^a Pratibha Saini,^a Ranveer Singh Badgoti,^a Kuldeep S. Rathore^b and Vijay Parewa^{id} ^{*a}

An efficient, metal-free, and sustainable strategy has been described for the *ipso*-functionalization of phenylboronic acids using air as an oxidant in an aqueous medium. A range of carbon materials has been tested as carbocatalysts. To our surprise, graphite was found to be the best catalyst in terms of the turnover frequency. A broad range of valuable substituted aromatic compounds, *i.e.*, phenols, anilines, nitroarenes, and haloarenes, has been prepared *via* the functionalization of the C–B bond into C–N, C–O, and many other C–X bonds. The vital role of the aromatic π -conjugation system of graphite in this protocol has been established and was observed *via* numerous analytic techniques. The heterogeneous nature of graphite facilitates the high recyclability of the carbocatalyst. This effective and easy system provides a multipurpose approach for the production of valuable substituted aromatic compounds without using any metals, ligands, bases, or harsh oxidants.

Received 11th March 2021

Accepted 30th April 2021

DOI: 10.1039/d1ra01940f

rsc.li/rsc-advances

1. Introduction

Arylboronic acids and their derivatives have attracted substantial research interest from chemists since they have immense value in chemical synthesis and materials science.¹ They have also been used as synthons for the further construction of therapeutic molecules and value-added chemicals.² Furthermore, arylboronic acids have been extensively used as ubiquitous structural motifs for various functional group transformations *via ipso*-functionalization.³

Phenols are a well-known class of compounds for their biological applications, including antibacterial, antitumor, antiviral, pro-oxidant, cardioprotective, and antimutagenicity activities.⁴ They have also been employed as synthetic intermediates in various chemical transformations.⁵ Due to the wide-ranging applications of phenols, dedicated efforts have been made to advance the preparation of phenols.⁶ Among them, the *ipso*-hydroxylation of arylboronic acid has been the most-studied strategy in the recent past.^{7–9}

This transformation is usually carried out with several equivalents of an oxidant, like hypervalent iodine, *tert*-butyl

hydroperoxide (TBHP), ozone, $\text{PhI}(\text{OAc})_2$, benzoquinone, or hydrogen peroxide (H_2O_2), which can be explosive or have a hazardous environmental impact. These reactions can also entail high temperatures, strongly basic conditions, and the use of metal catalysts (Pd, Au, Cu, Ru, Ag, *etc.*) and microwave irradiation, and they often require the use of expensive specialized ligands. As phenols are susceptible to oxidants, the quantity of oxidant needs to be cautiously determined. Moreover, the removal of trace quantities of metal-containing catalysts from the final compounds can be complicated. Therefore, the development of sustainable protocols for the above transformation presents a difficult task for chemists.

To meet the requirements of green chemistry, carbon-based materials, such as ordered mesoporous carbon, activated carbon, graphene-based materials, graphite, and carbon nanotubes, open up opportunities in the field of catalysis due to their inimitable physical and chemical properties.¹⁰ In recent years, many scientists have established that carbon-based materials can show elevated catalytic efficiencies in organic synthesis compared to traditional metal catalysts.¹¹ The sustainable nature of these carbon-based materials has been well documented in the literature.¹² Among them, the use of graphite as an environmentally benign catalyst in organic synthesis has taken on vast significance.¹³


Continuing our interest in developing sustainable protocols for organic synthesis,¹⁴ herein, we evaluate the catalytic activities of various carbon-based substances for the preparation of a library of phenol derivatives *via* the *ipso*-hydroxylation of

^aCentre of Advanced Studies, Department of Chemistry, University of Rajasthan, Jaipur, India. E-mail: dranshudandia@yahoo.co.in; parewavijay.parewa@gmail.com; parewavijay@uniraj.ac.in

^bDepartment of Physics, Arya College of Engineering and IT, Jaipur, India. E-mail: kuldeep_ssr@yahoo.com

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra01940f





Scheme 1 The synthesis of substituted phenols, anilines, nitroarenes, and haloarenes from arylboronic acids.

arylboronic acid, utilizing water as an environmentally friendly solvent under sustainable conditions. Commercially available graphite was found to be the best metal-free carbocatalyst for the above transformation. Furthermore, we have also extended our strategy to the synthesis of anilines, nitroarenes, and haloarenes *via* transforming the C–B bond of arylboronic acid to C–N and many other C–X bonds (Scheme 1).

2. Results and discussion

Firstly, phenylboronic acid has been selected as a model reactant for *ipso*-hydroxylation in an aqueous medium under air at 80 °C (Table 1). No transformation of phenylboronic acid into phenol was noticed in the absence of catalyst. This outcome

forced us to further optimize the reaction conditions. Recently, we have developed a series of carbon-based catalysts for various organic transformations.¹⁵ Therefore, from a sustainable point of view, the catalytic competences of various carbocatalysts have been studied. To our surprise, graphite was found to be the best catalyst for this transformation in terms of the turnover frequency (TOF^{15b,16}). Moreover, inferior yields were observed upon using other carbon-based materials as catalysts. GO and rGO gave poor product yields. Approximately similar catalytic activities were observed when the reaction was tried with various graphite samples purchased from different vendors.

An improvement in the yield of the desired product to an immense extent (36–92%) was observed when the catalyst loading was increased from 5 to 20 wt% (Table 1). We have tried

Table 1 The catalytic competences of various carbocatalysts for the *ipso*-hydroxylation of phenylboronic acid^a

S. no.	Catalyst	Solvent	Time	Yield ^b (%)	TOF ($\times 10^{-3}$ mol g ⁻¹ min ⁻¹)
1	—	H ₂ O	24 h	—	n.c.
2	Graphite (20 wt%)	H ₂ O	24 h	92	0.021
3	Activated carbon (20 wt%)	H ₂ O	24 h	74	0.017
4	MWCNTs (20 wt%) ^c	H ₂ O	24 h	72	0.016
5	GO (20 wt%)	H ₂ O	24 h	21	0.005
6	rGO (20 wt%)	H ₂ O	24 h	48	0.011
7	Graphite (5 wt%)	H ₂ O	24 h	64	0.069
8	Graphite (10 wt%)	H ₂ O	24 h	79	0.041
9	Graphite (20 wt%) ^d	H ₂ O	24 h	—	—
10	Graphite (20 wt%)	CH ₃ CN	24 h	82	0.019
11	Graphite (20 wt%)	MeOH	24 h	69	0.016
12	Graphite (20 wt%)	THF	24 h	44	0.010
13	Graphite (20 wt%)	DMSO	24 h	38	0.009

^a Reactions were carried out with phenylboronic acid (2 mmol) in H₂O at 80 °C under air. ^b Isolated yield. ^c Multi-walled carbon nanotubes.

^d Conversion conducted under a N₂ atmosphere.

different catalyst loadings (20, 30, 40, and 50 wt%), however further increases in catalyst loading did not affect the yield, and the optimum loading of catalyst was 20 wt%. The desired product was not formed when the model reaction was conducted under a nitrogen atmosphere. These results show that, from a sustainable point of view, air is an effective and environmentally benign oxidant. The desirable transformation was not observed in the absence of graphite, which shows that graphite plays a vital role in the further activation of oxygen molecules from air for the catalytic reaction.









A series of trials was carried out using a variety of solvents. Methanol and acetonitrile were found to give moderate yields among the various organic solvents. Compared to conventional organic solvents, H₂O was found to be a best solvent for the above transformation from an economical perspective. The outcome can be attributed to the excellent dispersion of the catalyst in H₂O. Additionally, the substrates show good dissolving capacities in water. The model reaction was also performed at different temperatures (60 °C, 80 °C, and 100 °C). It was concluded that, to achieve maximum conversion to the product, 80 °C was required.

To recognize the responsible catalytic sites in carbon-based catalysts, various organic compounds were employed as catalysts for the above transformation. Compounds with various oxygen functionalities showed poor catalytic activities. These outcomes confirmed that oxygen-containing functional groups may not be the catalytic sites in the carbon materials for hydroxylation. It was recognized from various catalytic tests that compounds containing aromatic π -conjugation systems demonstrated reasonable catalytic activities (Table 2). We suppose that the aromatic π -conjugation system has intrinsic activity for the above transformation. Thus, carbon materials that have significant oxygen functionality showed poor results compared to graphite.

Carbon materials illustrate metallic features and activate O₂ due to the presence of delocalized π -electrons.¹⁷ The introduction of any surface defects into a carbon material decreases its catalytic activity because this diminishes the mobility of delocalized π -electrons.¹⁸ Activated charcoal showed lower catalytic activity due to its decreased mass transfer efficiency because of its high degree of defects and its microporous structure.¹⁹ It has been revealed in the literature that sp²-hybridized carbon atoms are vital centers for oxygen adsorption and activation.¹⁷ These activated oxygen atoms can move on graphitic planes. As we already showed the necessity of oxygen for the above transformation, we proposed that the activation of oxygen was a decisive step in this transformation. The graphitic planes of the catalyst transform atmospheric O₂ to short-lived O₂^{•-} species that are stored in the carbon pores.¹⁷

There is a possibility that the observed catalytic properties of graphite could be attributed to the presence of trace metal-ion impurities. To establish the presence of trace amounts of metal ions, we have studied graphite catalyst samples (fresh and reused) *via* various analytical techniques. EDAX analysis confirmed the absence of any transition metals in the graphite catalyst, which ruled out catalytic assistance from any metal impurities. Moreover, the carbon : oxygen atomic ratios of fresh

Table 2 The catalytic activities of various organic counterparts^a

S. no.	Catalyst	Time	Yield ^b (%)
1		72 h	8
2		72 h	29
3	HCOOH	72 h	—
4	CH ₃ COOH	72 h	—
5		72 h	4
6		72 h	12
7		72 h	7
8		72 h	5
9		24 h	42
10		24 h	58

^a Reactions were carried out with phenylboronic acid (2 mmol) in H₂O at 80 °C under air. ^b Isolated yield.

and reused graphite were comparable, showing the stable nature of the graphite catalyst during the reaction (see ESI, Fig. S1 and S2†).

The deconvoluted C 1s XPS spectrum of fresh graphite shows a peak at 284.4 eV. Additionally, the O 1s spectrum illustrates a broad peak from 530 to 535 eV. The intensities and energies of the peaks of reused graphite in both XPS spectra were approximately the same (Fig. 1).

In the Fourier transform infrared (FT-IR) spectrum of reused graphite, absorption peak fluctuations from 1443 to 1573 cm⁻¹ are noted because of aromatic C=C stretching (Fig. 2a). We have also carried out solid-state ¹³C-NMR spectroscopy analysis of reused graphite (see ESI, Fig. S5†). The spectrum showed broad resonance associated with sp²-hybridized carbon atoms.²⁰ There were no additional peaks from any oxygen functionalities in the FT-IR and solid-state ¹³C-NMR spectra of reused graphite.




Fig. 1 The XPS spectra of fresh and reused graphite: (a) C 1s spectra; (b) O 1s spectra; and (c) the complete XPS spectra.




Fig. 2 (a) The FT-IR spectrum of reused graphite and (b) Raman spectra of fresh and reused graphite.

The Raman spectra of fresh and reused catalyst graphite samples comprise many overlapping bands (Fig. 2b). The D band, G band, and 2D band arise at around 1341 cm^{-1} , 1570 cm^{-1} , and 2690 cm^{-1} , respectively.

In SEM images, the graphite sheets displayed crumpled and wrinkled surfaces of thin lamellae (Fig. 3a, see ESI, Fig. S3†). The average particle size of the catalyst is about $9.8\text{ }\mu\text{m}$. The morphology and average particle size of the reused catalyst remain unchanged, as shown by SEM images of the reused graphite catalyst after four runs. The XRD pattern of recycled graphite was similarly documented, indicating that the structural integrity remains unchanged after the transformation reaction (Fig. 3b).

These analytical results proved that no oxidation of the catalyst was observed during the reaction, and the aromatic nature of the catalyst remains constant throughout the process. These results demonstrate that the utilized graphite catalyst is a stable and reusable catalyst for the above transformation.

To further confirm the role of the catalyst in the activation of the oxygen, EPR measurements were carried out with DMPO (see ESI, Fig. S5†). When the EPR experiment was performed with catalyst, peaks from the spin adduct of the DMPO superoxide radical were observed,²¹ while in the absence of catalyst, these peaks were absent. This result indicated that the generation of superoxide radicals was indispensable for the reaction to proceed.

These results suggest that a radical process might be involved in the above transformation. We also conducted the reaction with H_2O^{18} as a solvent under optimized conditions. The incorporation of ^{18}O into the product was not viewed, which ruled out the possibility of water acting as an oxygen source (Scheme 2).

In most cases, literature reports reveal the requirement for a base/metal in the reaction when phenylboronic acid reacts with reactive oxygen species.²² The mechanism of the reaction is ambiguous, particularly in light of the fact that the reaction can





Fig. 3 (a) SEM images and (b) XRD patterns of fresh and reused graphite.



Scheme 2 Isotopic experiment results.


occur in the absence of any base/metal under air. *Via* control experiments, we have already confirmed that a radical process might be involved in the transformation. Due to the presence of delocalized π -electrons, graphitic planes convert O_2 to short-lived $O_2^{\cdot-}$ species that assemble in the carbon pores.¹⁷ This short-lived $O_2^{\cdot-}$ was the active sustainable oxidant for the above transformation. Based on the above information and literature reports,²³ we considered that the reaction occurred *via* a radical nucleophilic substitution mechanism²⁴ through an unexplored process (to be the subject of future investigations). Owing to the presence of a delocalized π -electron system, graphite also stabilizes the radical intermediates for better reaction performance.

To elaborate on the possibilities of this approach, numerous electronically diversified phenylboronic acids were used for hydroxylation under the established optimum conditions (Table 3). This approach shows broad functional-group tolerance. Arylboronic acids containing electron-withdrawing and


-donating substituents at the *para* position gave excellent phenol yields. The use of sterically demanding *ortho*-substituted arylboronic acids did not influence the yields drastically, and the final products were achieved in good yields. *meta*-Substituted arylboronic acids gave analogous outcomes. Heteroaryl boronic acids also gave good yields in the optimized reaction environment.

Encouraged by the above results, we would like to examine whether the formation of anilines would happen when aqueous NH_3 solution was inserted into the system due to nucleophilic competition between the nitrogen and oxygen centers. To our delight, the formation of aniline was observed instead of phenol. A probable reason for this is that the superior nucleophilic power of the nitrogen center compared to the oxygen center led to the construction of aniline instead of phenol (Scheme 3). Therefore, the reaction proceeds through the more nucleophilic center (in this case the nitrogen center) instead of the oxygen center. Similarly, when we added $NaNO_2$ to the system instead of aqueous NH_3 , nitroarenes were formed.

Inspired by these outcomes, we used the same sustainable protocol for the iodination of arylboronic acids, using KI as an iodide source. As iodide ions have more nucleophilic power, iodobenzene was formed as the sole product. In the same way, when we used KBr in the system, bromobenzene was formed as the desired product. Unfortunately, KCl failed to give the halogenated product under these conditions. However, NaCl withstands the reaction conditions and gives the chlorinated


Table 3 The graphite-catalyzed *ipso*-hydroxylation of arylboronic acids^a

^a Reactions are performed with various phenylboronic acids in water at 80 °C under air.




Scheme 3 Intermolecular competition between the oxygen and nitrogen centres during the functionalization of phenylboronic acid.

product in good yield. Since the functionalization of phenylboronic acids at the nitrogen center was successful in the presence of an oxygen center, a competitive trial was executed to determine which phenylboronic acid functionalization would happen when three halogen reactants were added at the same time (Scheme 4). Under these conditions, iodobenzene was formed as a major product, with the order I > Br > Cl.



Scheme 4 Intermolecular competition between various halide centres during the functionalization of phenylboronic acid.



Scheme 5 The synthesis of substituted anilines, nitroarenes, and haloarenes from arylboronic acids.

and electron-donating arylboronic acids, but lower and more variable yields were obtained in the case of chlorobenzenes.

The heterogeneous character and excellent recyclability of graphite was confirmed (see ESI†). For recycling experiments, arylboronic acid was selected as the model substrate for all transformations. The used graphite was simply isolated *via* filtration, cleaned with ethanol, and reused for an additional four cycles. As shown in Table 4, graphite was found to be a very stable carbocatalyst for all these transformations.

3. Experimental


3.1. General

See ESI.†

3.2. Preparation of substituted phenols (2)

A round-bottomed flask was loaded with arylboronic acid (2 mmol) and 20 wt% graphite in 20 mL of H₂O, and the mixture was stirred at 80 °C until complete transformation was achieved

Table 4 Recyclability experiments involving graphite for various functional group transformations^a



Functional entity	First cycle [%]	Second cycle [%]	Third cycle [%]	Fourth cycle [%]
Ph-OH	92	88	86	86
Ph-NH ₂	80	78	77	77
Ph-NO ₂	72	70	68	68
Ph-I	76	72	72	71
Ph-Br	73	71	71	68
Ph-Cl	72	67	66	66

^a Isolated product yield.

(checked *via* TLC). The used graphite was removed simply *via* filtration after the transformation was complete. The filtrate was permitted to cool to atmospheric temperature and was then extracted with ethyl acetate (3 × 2 mL). The solution was then concentrated under reduced pressure and purified *via* column chromatography on silica gel (eluent, hexane : EtOAc) to furnish the pure final product.

3.3. Preparation of substituted anilines (3)

A round-bottomed flask was loaded with arylboronic acid (2 mmol), NH₃ solution (1.5 eq.) and 20 wt% graphite in 20 mL of H₂O, and the mixture was stirred at 80 °C until the complete transformation was achieved (checked *via* TLC). The used graphite catalyst was separated simply *via* filtration after the completion of the transformation. The filtrate was permitted to cool to atmospheric temperature and was then extracted with ethyl acetate (3 × 2 mL). The solution was then concentrated under reduced pressure and purified *via* column chromatography on silica gel (eluent, hexane : EtOAc) to deliver the required pure product.

3.4. Preparation of substituted nitroarenes (4)

A round-bottomed flask was loaded with arylboronic acid (2 mmol), NaNO₂ (1.5 eq.) and 20 wt% graphite in 20 mL of H₂O, and the mixture was stirred at 80 °C until the conversion was complete (monitored *via* TLC). The used graphite was separated simply *via* filtration after the completion of the transformation. The filtrate was permitted to cool to atmospheric temperature and was then extracted with ethyl acetate (3 × 2 mL). The solution was then concentrated under reduced pressure and purified *via* column chromatography on silica gel (eluent, hexane : EtOAc) to supply the final pure product.

3.5. Preparation of substituted haloarenes (5–7)

A round-bottomed flask was loaded with arylboronic acid (2 mmol), KCl (1.5 eq.)/KI (1.5 eq.)/KBr (1.5 eq.), and 20 wt% graphite in 20 mL of H₂O, and the mixture was stirred at 80 °C until complete transformation was achieved (checked *via* TLC). Graphite was isolated simply *via* filtration after the transformation was complete. The filtrate was permitted to cool to atmospheric temperature and was then extracted with ethyl acetate (3 × 2 mL). The solution was then concentrated under reduced pressure and purified *via* column chromatography on silica gel (eluent, hexane : EtOAc) to offer the desired pure product.

4. Conclusions

A versatile and simple graphite-catalyzed process has been illustrated for the conversion of numerous functional groups on aromatic rings in an aqueous medium under air *via* the *ipso*-functionalization of phenylboronic acids. Various important aromatic compounds, *e.g.*, phenols, anilines, nitroarenes, and haloarenes, have been synthesized without using any metals, ligands, bases, or oxidants. The vital role of the aromatic π -conjugation system of graphite in assisting the reaction has

been confirmed *via* various analytic techniques, *viz.*, FT-IR, Raman, XPS, SEM, and XRD analysis. *Via* control experiments, we have offered a plausible mechanistic pathway for the above transformation. The main advantage of this protocol is the use of economical and abundant graphite as an efficient carbocatalyst without any prior treatment. Overall, this strategy offers numerous advantages, such as the recyclability of the catalyst, operational simplicity, inexpensiveness, environmental friendliness, a broad substrate scope, and high yields.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

DST-SERB, C.S.I.R., and U.G.C., New Delhi, are acknowledged for financial assistance. The authors are thankful to MNIT, Jaipur, for assistance with the spectral studies.

References

- 1 D. Hall, *Boronic Acids: Preparation and Applications in Organic Synthesis, Medicine and Materials*, ed. D. G. Hall, Wiley-VCH, Weinheim, 2nd edn, 2011.
- 2 (a) C. Kleeberg, L. Dang, Z. Lin and T. B. Marder, *Angew. Chem.*, 2009, **121**, 5454–5458; (b) M. Murata, T. Oyama, S. Watanabe and Y. Masuda, *J. Org. Chem.*, 2000, **65**, 164–168; (c) W. Zhu and D. Ma, *Org. Lett.*, 2006, **8**, 261–263; (d) D. A. Wilson, C. J. Wilson, B. M. Rosen and V. Percec, *Org. Lett.*, 2008, **10**, 4879–4882; (e) F. Mo, Y. Jiang, D. Qiu, Y. Zhang and J. Wang, *Angew. Chem.*, 2010, **49**, 1846–1849; (f) J. Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka and M. R. Smith, *Science*, 2002, **295**, 305–308; (g) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura and N. R. Anastasi, *Science*, 2002, **295**, 305–308; (h) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 390–391; (i) I. A. I. Mkhaliid, J. H. Barnard, T. B. Marder, J. M. Murphy and J. F. Hartwig, *Chem. Rev.*, 2010, **110**, 890–931.
- 3 C. Zhua and R. F. John, *Adv. Synth. Catal.*, 2014, **356**, 2395–2410.
- 4 (a) J. H. P. Tyman, *Synthetic and Natural Phenols*, Elsevier, New York, 1996; (b) Z. Rappoport, *The Chemistry of Phenols*, Wiley-VCH, Weinheim, Germany, 2003.
- 5 (a) R. W. Owen, A. Giacosa, R. Haubner, B. Spiegel-Hader and H. Bartsch, *Eur. J. Cancer.*, 2000, **36**, 1235–1247; (b) L. Pilato, *React. Funct. Polym.*, 2013, **73**, 270–277.
- 6 (a) C. Hoarau and T. R. R. Pettus, *Synlett*, 2003, **1**, 127–137; (b) P. Hanson, J. R. Jones, A. B. Taylor, P. H. Walton and A. W. Timms, *J. Am. Chem. Soc.*, 2002, **2**, 1135–1150; (c) T. George, R. Mabon, G. Sweeney, J. B. Sweeney and A. Tavassoli, *J. Am. Chem. Soc.*, 2000, **1**, 2529–2574; (d) T. Schulz, C. Torborg and B. Schaffner, *Angew. Chem., Int. Ed.*, 2009, **48**, 918–921; (e) K. W. Anderson, T. Ikawa, R. E. Tundel and S. L. Buchwald, *J. Am. Chem. Soc.*, 2006, **128**, 10694–10695.

- 7 (a) B. A. Dar, P. Bhatti, A. P. Singh, A. Lazar, P. R. Sharma, M. Sharma and B. Singh, *Appl. Catal., A*, 2013, **466**, 60–67; (b) D. Yang, B. An, W. Wei, M. Jiang, J. You and H. Wang, *Tetrahedron*, 2014, **70**, 3630–3634; (c) H. Yang, Y. Li, M. Jiang, J. Wang and H. Fu, *Chem.–Eur. J.*, 2011, **17**, 5652–5660; (d) A. Affrose, I. A. Azath, A. Dhakshinamoorthy and K. Pitchumani, *J. Mol. Catal. A: Chem.*, 2014, **395**, 500–505; (e) J. Xu, X. Wang, C. Shao, D. Su, G. Cheng and Y. Hu, *Org. Lett.*, 2010, **12**, 1964–1967; (f) K. Inamoto, K. Nozawa, M. Yonemoto and Y. Kondo, *Chem. Commun.*, 2011, **47**, 11775–11777; (g) A. D. Chowdhury, S. M. Mobin, S. Mukherjee, S. Bhaduri and G. K. Lahiri, *Eur. J. Inorg. Chem.*, 2011, **21**, 3232–3239; (h) J. Zheng, S. Lin, X. Zhu, B. Jiang, Z. Yang and Z. Pan, *Chem. Commun.*, 2012, **48**, 6235–6237; (i) L. Wang, W. Zhang, D. Sheng Su, X. Meng and F.-S. Xiao, *Chem. Commun.*, 2012, **48**, 5476–5478; (j) N. Gogoi, P. K. Gogoi, G. Borah and U. Bora, *Tetrahedron Lett.*, 2016, **57**, 4050–4052; (k) T. Begum, A. Gogoi, P. K. Gogoi and U. Bora, *Tetrahedron Lett.*, 2015, **56**, 95–97; (l) T. Toyao, N. Ueno, K. Miyahara, Y. Matsui, T.-H. Kim, Y. Horiuchi, H. Ikeda and M. Matsuoka, *Chem. Commun.*, 2015, **51**, 16103–16106; (m) M.-J. Zhang, H.-X. Li, H.-Y. Li and J.-P. Lang, *Dalton Trans.*, 2016, **45**, 17759–17769.
- 8 (a) G. K. S. Prakash, S. Chacko, C. Panja, T. E. Thomas, L. Gurung, G. Rasul, T. Mathew and G. A. Olah, *Adv. Synth. Catal.*, 2009, **351**, 1567–1574; (b) L. Wang, D.-Y. Dai, Q. Chen and M.-Y. He, *J. Fluorine Chem.*, 2014, **158**, 44–47; (c) S. Gupta, P. Chaudhary, V. Srivastava and J. Kandasamy, *Tetrahedron Lett.*, 2016, **57**, 2506–2510; (d) S. Guo, L. Lu and H. Cai, *Synlett*, 2013, **24**, 1712–1714; (e) D.-S. Chen and J.-M. Huang, *Synlett*, 2013, **24**, 499–501; (f) K. S. Webb and D. Levy, *Tetrahedron Lett.*, 1995, **36**, 5117–5118; (g) B. R. Travis, B. P. Ciaramitaro and B. Borhan, *Eur. J. Org. Chem.*, 2002, **2002**, 3429–3434; (h) R. E. Maleczka Jr, F. Shi, D. Holmes and M. R. Smith III, *J. Am. Chem. Soc.*, 2003, **125**, 7792–7793; (i) G. A. Molander and L. N. Cavalcanti, *J. Org. Chem.*, 2011, **76**, 623–630; (j) E. Kianmehr, M. Yahyae and K. Tabatabai, *Tetrahedron Lett.*, 2007, **48**, 2713–2715; (k) C. Zhu, R. Wang and J. R. Falck, *Org. Lett.*, 2012, **14**, 3494–3497; (l) Y. K. Bommegowda, N. Mallesha, A. C. Vinayaka and M. P. Sadashiva, *Chem. Lett.*, 2016, **45**, 268–270; (m) K. Murugan and S. Palaniswamy, *ACS Sustainable Chem. Eng.*, 2019, **7**, 9028–9034.
- 9 (a) A. Paul, D. Chatterjee, H. T. Rajkamal, S. Banerjee and S. Yadav, *Tetrahedron Lett.*, 2015, **56**, 2496–2499; (b) N. Chatterjee, H. Chowdhury, K. Sneha and A. Goswami, *Tetrahedron Lett.*, 2015, **56**, 172–174; (c) J. S. Ruso, N. Rajendiran and R. S. Kumaran, *Tetrahedron Lett.*, 2014, **55**, 2345–2347; (d) G. Cheng, X. Zeng and X. Cui, *Synthesis*, 2014, **46**, 295–300; (e) P. Gogoi, P. Bezboruah, J. Gogoi and R. C. Boruah, *Eur. J. Org. Chem.*, 2013, **32**, 7291–7294; (f) K. Hosoi, Y. Kuriyama, S. Inagi and T. Fuchigami, *Chem. Commun.*, 2010, **46**, 1284–1286; (g) H.-L. Qi, D.-S. Chen, J.-S. Ye and J.-M. Huang, *J. Org. Chem.*, 2013, **78**, 7482–7487; (h) H. Jiang, L. Lykke, S. U. Pedersen, W.-J. Xiao and K. A. Jørgensen, *Chem. Commun.*, 2012, **48**, 7203–7205; (i) A. N. Cammidge, V. H. M. Goddard, C. P. J. Schubert, H. Gopee, D. L. Hughes and D. Gonzalez-Lucas, *Org. Lett.*, 2011, **13**, 6034–6037; (j) H. Kotoučová, I. Strnadová, M. Kovandová, J. Chudoba, H. Dvořáková and R. Cibulka, *Org. Biomol. Chem.*, 2014, **12**, 2137–2142; (k) X. Yu and S. M. Cohen, *Chem. Commun.*, 2015, **51**, 9880–9883; (l) J. Luo, X. Zhang and J. Zhang, *ACS Catal.*, 2015, **5**, 2250–2254; (m) S. P. Pitre, C. D. McTiernan, H. Ismaili and J. C. Scaiano, *J. Am. Chem. Soc.*, 2013, **135**, 13286–13289; (n) Y.-Q. Zou, J.-R. Chen, X.-P. Liu, L.-Q. Lu, R. L. Davis, K. A. Jørgensen and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2012, **124**, 802–812; (o) J. A. Johnson, J. Luo, X. Zhang, Y.-S. Chen, M. D. Morton, E. Echeverría, F. E. Torres and J. Zhang, *ACS Catal.*, 2015, **5**, 5283–5291; (p) W. Yina, X. Pana, W. Lenga, J. Chenb and H. Hec, *Green Chem.*, 2019, **21**, 4614–4618.
- 10 (a) L. Xien and L. Dai, *Nat. Rev. Mater.*, 2016, **1**, 1–12; (b) E. Pérez-Mayoral, V. Calvino-Casilda and E. Soriano, *Catal. Sci. Technol.*, 2016, **6**, 1265–1291; (c) E. Lam and H. T. John Luong, *ACS Catal.*, 2014, **4**, 3393–3410.
- 11 (a) M. B. Gawande, P. Fornasiero and R. Zbořil, *ACS Catal.*, 2020, **10**, 2231–2259; (b) Z. Liu, B. Zhang, H. Yu, K. H. Wu and X. Huang, *Front. Chem.*, 2020, **8**, 308.
- 12 (a) K. P. Gopinath, D. V. N. Vo, D. G. Prakash, A. A. Joseph, S. Viswanathan and J. Arun, *Environ. Chem. Lett.*, 2020, 1–26; (b) D. R. Dreyer and C. W. Bielawski, *Chem. Sci.*, 2011, **2**, 1233–1240; (c) S. S. Shang and G. Shuang, *ChemCatChem*, 2019, **11**, 3730–3744; (d) S. Chenliang and K. P. Loh, *Acc. Chem. Res.*, 2013, **46**, 2275–2285.
- 13 (a) M. Nagai, R. Isoe, K. Ishiguro, H. Tominaga and M. Shimizu, *Chem. Eng. J.*, 2012, **207**, 938–942; (b) B. H. Han, D. H. Shin and S. Y. Cho, *Tetrahedron Lett.*, 1985, **26**, 6233–6234; (c) H. K. Kadam, S. Khan, R. A. Kunkalkar and S. G. Tilve, *Tetrahedron Lett.*, 2013, **54**, 1003–1007; (d) M. Kodomari, Y. Suzuki and K. Yoshida, *Chem. Commun.*, 1997, 1567–1568; (e) M. R. Acocella, L. D'Urso, M. Maggio and G. Guerra, *ChemCatChem*, 2016, **8**, 1915–1920; (f) K. N. Nandeesh, G. M. Raghavendra, C. N. Revanna, T. A. J. Vijay, K. S. Rangappa and K. Mantelingu, *Synth. Commun.*, 2014, **44**, 1103–1110.
- 14 (a) A. Dandia, V. Parewa, A. K. Jain and K. S. Rathore, *Green Chem.*, 2011, **13**, 2135–2145; (b) A. Dandia, V. Parewa, S. Kumari, S. Bansal and A. Sharma, *Green Chem.*, 2016, **18**, 2488–2499; (c) A. Dandia, S. L. Gupta, A. Indora, P. Saini, V. Parewa and K. S. Rathore, *Tetrahedron Lett.*, 2017, **58**, 1170–1175; (d) A. Dandia, S. Bansal, R. Sharma, D. K. Mahawar, K. S. Rathore, M. L. Meena and V. Parewa, *J. Photochem. Photobiol., A*, 2020, **389**, 112242; (e) A. Dandia, S. Khan, R. Sharma, S. Parihar and V. Parewa, *ChemistrySelect*, 2017, **2**, 9684–9690; (f) A. Dandia, S. Parihar, R. Sharma, K. S. Rathore and V. Parewa, Nanocatalysis in green organic synthesis, in *Green Sustainable Process for Chemical and Environmental Engineering and Science*, Elsevier, 2020, pp. 71–103; (g) A. Dandia, P. Saini, R. Sharma and V. Parewa, Green organic synthesis by photochemical protocol, in *Green Sustainable Process for Chemical and Environmental Engineering and Science*, Elsevier, 2020, pp. 155–198; (h)

- A. Dandia, R. Sharma, A. Indora and V. Parewa, *ChemistrySelect*, 2018, **3**, 8285–8290.
- 15 (a) A. Dandia, S. Parihar, P. Saini, K. S. Rathore and V. Parewa, *Catal. Lett.*, 2019, **149**, 3169–3175; (b) A. Dandia, D. K. Mahawar, R. Sharma, R. S. Badgoti, K. S. Rathore and V. Parewa, *Appl. Organomet. Chem.*, 2019, **33**, e5232; (c) A. Dandia, S. Bansal, R. Sharma, K. S. Rathore and V. Parewa, *RSC Adv.*, 2018, **8**, 30280–30288; (d) A. Dandia, A. Sharma, A. Indora, K. S. Rathore, A. Jain and V. Parewa, *Mol. Catal.*, 2018, **459**, 97–105.
- 16 A. Dandia, A. Sharma, V. Parewa, B. Kumawat, K. S. Rathore and A. Sharma, *RSC Adv.*, 2015, **5**, 91888–91902.
- 17 F. Atamny, J. Blocker, A. Dubotzky, H. Kurt, O. Timpe and G. Loose, *Mol. Phys.*, 1992, **4**, 851–886.
- 18 H. Yu, F. Peng, J. Tan, X. W. Hu, H. J. Wang and J. Yang, *Angew. Chem.*, 2011, **123**, 4064–4068.
- 19 (a) A. Corma, M. T. Navarro, L. Nemeth and M. Renz, *Chem. Commun.*, 2001, 2190–2191; (b) J. Rivera-Utrilla and M. Sánchez-Polo, *Appl. Catal., B*, 2002, **39**, 319–329.
- 20 (a) A. S. Mazur, M. A. Vovk and P. M. Tolstoy, *Fullerenes, Nanotubes, Carbon Nanostruct.*, 2020, **28**(3), 202–213; (b) M. A. Vieira, G. R. Gonçalves, D. F. Cipriano, M. A. Schettino Jr, E. A. Silva Filho, A. G. Cunha and J. C. Freitas, *Carbon*, 2016, **98**, 496–503.
- 21 H. Li, F. Qin, Z. P. Yang, X. M. Cui, J. F. Wang and L. Z. Zhang, *J. Am. Chem. Soc.*, 2017, **139**, 3513–3521.
- 22 (a) A. Ding, Y. Zhang, Y. Chen, R. Rios, J. Hu and H. Guo, *Tetrahedron Lett.*, 2019, **60**, 660–663; (b) Y. L. Ren, X. Z. Tian, C. Dong, S. Zhao, J. Wang, M. Yan, X. Qi and G. Liu, *Catal. Commun.*, 2013, **32**, 15–17; (c) R. Gujjarappa, N. Vodnala, A. Garg, C. K. Hazra, S. Gupta and C. C. Malakar, *ChemistrySelect*, 2020, **5**, 2419–2423.
- 23 (a) I. B. Seiple, S. Shun, R. A. Rodriguez, R. Gianatassio, Y. Fujiwara, A. L. Sobel and P. S. Baran, *J. Am. Chem. Soc.*, 2010, **132**, 13194–13196; (b) N. R. Patel and R. A. Flowers, *J. Am. Chem. Soc.*, 2013, **135**, 4672–4675; (c) A. S. Demir, O. Reis and M. Emrullahoglu, *J. Org. Chem.*, 2003, **68**, 578–580; (d) G. Yan, M. Yang and X. Wu, *Org. Biomol. Chem.*, 2013, **11**, 7999–8008.
- 24 (a) G. A. Molander and L. N. Cavalcanti, *J. Org. Chem.*, 2011, **76**, 7195–7203; (b) R. W. Alder, *J. Chem. Soc., Chem. Commun.*, 1980, **24**, 1184–1186; (c) S. Hashimoto, I. Kurimoto, Y. Fujii and R. Noyori, *J. Am. Chem. Soc.*, 1985, **107**, 1427–1429.