Original research

Infliximab is associated with attenuated immunogenicity to BNT162b2 and ChAdOx1 nCoV-19 SARS-CoV-2 vaccines in patients with IBD

Nicholas A Kennedy (), ^{1,2} Simeng Lin (), ^{1,2} James R Goodhand (), ^{1,2} Neil Chanchlani (), ^{1,2} Benjamin Hamilton (), ^{1,2} Claire Bewshea (), ² Rachel Nice, ^{2,3} Desmond Chee (), ^{1,2} JR Fraser Cummings (), ⁴ Aileen Fraser (), ⁵ Peter M Irving (), ^{6,7} Nikolaos Kamperidis, ⁸ Klaartje B Kok (), ^{9,10} Christopher Andrew Lamb (), ^{11,12} Jonathan Macdonald (), ^{13,14} Shameer Mehta (), ¹⁵ Richard CG Pollok (), ^{16,17} Tim Raine (), ¹⁸ Philip J Smith (), ¹⁹ Ajay Mark Verma (), ²⁰ Simon Jochum, ²¹ Timothy J McDonald (), ³ Shaji Sebastian (), ^{22,23} Charlie W Lees (), ^{24,25} Nick Powell (), ^{26,27} Tariq Ahmad (), ^{1,2} Contributors to the CLARITY IBD study

ABSTRACT

► Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi.org/10.1136/gutjnl-2021-324789).

For numbered affiliations see end of article.

Correspondence to

Dr Tariq Ahmad, Gastroenterology, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK; tariq.ahmad1@nhs.net

NAK, SL and JRG contributed equally. NP and TA contributed equally.

Received 29 March 2021 Revised 8 April 2021 Accepted 10 April 2021



Check for updates

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Kennedy NA, Lin S, Goodhand JR, *et al. Gut* Epub ahead of print: [*please include* Day Month Year]. doi:10.1136/ gutjnl-2021-324789 **Objective** Delayed second dose SARS-CoV-2 vaccination trades maximal effectiveness for a lower level of immunity across more of the population. We investigated whether patients with inflammatory bowel disease treated with infliximab have attenuated serological responses to a single dose of a SARS-CoV-2 vaccine.

Design Antibody responses and seroconversion rates in infliximab-treated patients (n=865) were compared with a cohort treated with vedolizumab (n=428), a gutselective anti-integrin α 4 β 7 monoclonal antibody. Our primary outcome was anti-SARS-CoV-2 spike (S) antibody concentrations, measured using the Elecsys anti-SARS-CoV-2 spike (S) antibody assay 3–10 weeks after vaccination, in patients without evidence of prior infection. Secondary outcomes were seroconversion rates (defined by a cut-off of 15 U/mL), and antibody responses following past infection or a second dose of the BNT162b2 vaccine.

Results Geometric mean (SD) anti-SARS-CoV-2 antibody concentrations were lower in patients treated with infliximab than vedolizumab, following BNT162b2 (6.0 U/mL (5.9) vs 28.8 U/mL (5.4) p<0.0001) and ChAdOx1 nCoV-19 (4.7 U/ mL (4.9)) vs 13.8 U/mL (5.9) p<0.0001) vaccines. In our multivariable models, antibody concentrations were lower in infliximab-treated compared with vedolizumab-treated patients who received the BNT162b2 (fold change (FC) 0.29 (95% CI 0.21 to 0.40), p<0.0001) and ChAdOx1 nCoV-19 (FC 0.39 (95% CI 0.30 to 0.51), p<0.0001) vaccines. In both models, age ≥ 60 years, immunomodulator use, Crohn's disease and smoking were associated with lower, while nonwhite ethnicity was associated with higher, anti-SARS-CoV-2 antibody concentrations. Seroconversion rates after a single dose of either vaccine were higher in patients with prior SARS-CoV-2 infection and after two doses of BNT162b2 vaccine.

Conclusion Infliximab is associated with attenuated immunogenicity to a single dose of the BNT162b2 and ChAdOx1 nCoV-19 SARS-CoV-2 vaccines. Vaccination after SARS-CoV-2 infection, or a second dose of vaccine,

Significance of this study

What is already known on this subject?

- A growing number of countries, including the UK, have opted to delay second SARS-CoV-2 vaccine doses for all people, trading maximal effectiveness against a lower level of protective immunity across more of the at-risk population. Whether single doses of vaccines are effective in patients treated with antitumour necrosis factor (TNF) therapies is unknown.
- We have previously shown in this cohort that seroprevalence, seroconversion in PCRconfirmed cases and the magnitude of anti-SARS-CoV-2 antibodies following SARS-CoV-2 infection are reduced in infliximab-treated compared with vedolizumab-treated patients.
- Two recent studies have reported that SARS-CoV-2 spike (S) antibody responses are impaired in patients with cancer and transplant recipients treated with chemotherapy and antimetabolite immunosuppressants, respectively. To date, no studies have assessed the effect of anti-TNF therapy on immunogenicity following SARS-CoV-2 vaccination.

led to seroconversion in most patients. Delayed second dosing should be avoided in patients treated with infliximab.

Trial registration number ISRCTN45176516.

INTRODUCTION

Limited SARS-CoV-2 vaccine supplies and pressure on critical care services have forced governments to prioritise primary vaccination to vulnerable groups.

osg

1

Significance of this study

What are the new findings?

- Anti-SARS-CoV-2 spike (S) antibody concentrations and rates of seroconversion were lower following primary vaccination with both the BNT162b2 and ChAdOx1 nCoV-19 vaccines in patients with inflammatory bowel disease treated with infliximab compared with vedolizumab.
- Older age, immunomodulator use, Crohn's disease (vs ulcerative colitis or inflammatory bowel disease unclassified), and current smoking were associated with lower anti-SARS-CoV-2 antibody concentrations, irrespective of vaccine type. Non-white ethnicity was associated with higher anti-SARS-CoV-2 (S) antibody concentrations following primary vaccination with both vaccines.
- Lowest rates of seroconversion were observed in participants treated with infliximab in combination with an immunomodulator with both the BNT162b2 and ChAdOx1 nCoV-19 vaccines, whereas highest rates of seroconversion were seen in patients treated with vedolizumab monotherapy who received either vaccine.
- Antibody concentrations and seroconversion rates were higher in patients with past SARS-CoV-2 infection prior to a single dose of either vaccine, and after two doses of the BNT162b2 vaccine.

How might it impact on clinical practice in the foreseeable future?

- For patients treated with antitumour necrosis factor (TNF) therapy, particularly for those also treated with an immunomodulator, poor antibody responses to a single dose of vaccine exposes them to a potential increased risk of SARS-CoV-2 infection.
- Higher rates of seroconversion in patients with two exposures to SARS-CoV-2 antigen, even in the presence of TNF blockade, suggest that all patients receiving anti-TNF therapy should be prioritised for optimally timed second doses.
- Until patients receive a second vaccine dose, they should consider that they are not protected from SARS-CoV-2 infection and continue to practice enhanced physical distancing and shielding if appropriate.
- Even after two antigen exposures, a small subset of patients failed to mount an antibody response. Antibody testing and adapted vaccine schedules should be considered to protect these at-risk patients.

In the UK, second vaccine doses have also been delayed, trading maximal effectiveness for a lower level of protective immunity across a greater proportion of the most at-risk population.¹ Consequently, more than half of the adult population have received a single dose of either the RNA vaccine, BNT162b2 (Pfizer/BioNTech) or the adenovirus-vector vaccine, ChAdOx1 nCoV-19 (Oxford/AstraZeneca). Faced with further surges of SARS-CoV-2 infection, a growing number of other countries have also opted to delay second vaccine doses.²

The inflammatory bowel diseases (IBD), Crohn's disease and ulcerative colitis (UC) are chronic immune-mediated inflammatory diseases (IMIDs) that affect about 1% of the UK population.^{4 5} Treatment typically requires immunosuppression with immunomodulators (azathioprine, mercaptopurine and methotrexate) and/or biological therapies that target disease relevant cytokines or the immune cells that produce them. Antitumour

necrosis factor (TNF) drugs, such as infliximab and adalimumab, are the most frequently prescribed biopharmaceuticals used in the treatment of IMIDs. These drugs impair immunogenicity following pneumococcal,⁶ influenza⁷ and hepatitis B⁸ vaccinations and increase the risk of serious infection, most notably with respiratory pathogens.⁹ Conversely, vedolizumab, a gut-selective anti-integrin $\alpha 4\beta 7$ monoclonal antibody, is not associated with increased susceptibility to systemic infection or attenuated sero-logical responses to vaccination.¹⁰

We have recently reported that seroprevalence, seroconversion in PCR-confirmed cases, and the magnitude of anti-SARS-CoV-2 antibodies following SARS-CoV-2 infection are reduced in infliximab-treated compared with vedolizumab-treated patients.¹¹ We hypothesised that, following at least a single dose with BNT162b2 or ChAdOx1 nCoV-19 vaccine, serological responses would be similarly impaired in patients treated with infliximab compared with vedolizumab arguing against delaying second doses in these patients.

Objectives

We aimed to define, in patients with IBD who had received a COVID-19 vaccination, whether biological class and concomitant use of an immunomodulator impact:

- i. Anti-SARS-CoV-2 spike (S) antibody levels.
- ii. Rates of seroconversion.
- iii. Antibody responses in patients who had previously been infected with SARS-CoV-2 or who had two doses of vaccine.

METHODS

Patient and settings

ImpaCt of bioLogic therApy on saRs-cov-2 Infection and immuniTY (CLARITY) IBD is a UK-wide, multicentre, prospective observational cohort study investigating the impact of infliximab and vedolizumab and/or concomitant immunomodulators (azathioprine, mercaptopurine and methotrexate) on SARS-CoV-2 acquisition, illness and immunity in patients with IBD.

Study methods have been described in detail previously.¹¹ In brief, consecutive patients were recruited at the time of attendance at infusion units from 92 National Health Service (NHS) hospitals across the UK between 22 September 2020 and 23 December 2020 (online supplemental pp 2–17). The eligibility criteria were age 5 years and over, a diagnosis of IBD, and current treatment with infliximab or vedolizumab for 6 weeks or more, with at least one dose of drug received in the previous 16 weeks. Patients were excluded if they had participated in a SARS-CoV-2 vaccine trial.

Follow-up visits were timed to coincide with biological infusions and occurred approximately 8 weekly. Here, we report vaccine-induced antibody responses at first study visit after primary vaccination, and where possible, after two doses. Participants were eligible for inclusion in our vaccine immunogenicity analysis if they had had a SARS-CoV-2 antibody test within the first 10 weeks after their primary vaccination with any of the available SARS-CoV-2 vaccines.

Outcome measures

Our primary outcome was anti-SARS-CoV-2 anti-spike (S) protein receptor-binding protein antibodies 3–10 weeks after primary vaccination.

Secondary outcomes were:

i. The proportion of participants with seroconversion.

- ii. Antibody concentrations and rate of seroconversion in patients with PCR or serological evidence of past SARS-CoV-2 infection.
- iii. Antibody concentrations and seroconversion after two doses of vaccine.

Variables

Variables recorded by participants were demographics (age, sex, ethnicity, comorbidities, height and weight, smoking status, and postcode), IBD disease activity (PRO2), SARS-CoV-2 symptoms aligned to the COVID-19 symptoms study (symptoms, previous testing and hospital admissions for COVID-19) and vaccine uptake (type and date of primary vaccination). Study sites completed data relating to IBD history (age at diagnosis, disease duration and phenotype according to the Montreal classifications, previous surgeries and duration of current biological and immunomodulator therapy).¹¹ We linked our data by NHS number or Community Health Index to Public Health England, Scotland and Wales who archive dates and results of all SARS-CoV-2 PCR tests undertaken. Data were entered electronically into a purpose-designed REDCap database hosted at the Royal Devon and Exeter NHS Foundation Trust.¹² Participants without access to the internet or electronic device completed their questionnaires on paper case record forms that were subsequently entered by local research teams.

Laboratory methods

Laboratory analyses were performed at the Academic Department of Blood Sciences at the Royal Devon and Exeter NHS Foundation Trust. To determine antibody responses specific to vaccination we used the Roche Elecsys Anti-SARS-CoV-2 spike (S) immunoassay¹³ alongside the nucleocapsid (N) immunoassay.¹⁴ This double sandwich electrochemiluminescence immunoassay uses a recombinant protein of the receptor binding domain on the spike protein as an antigen for the determination of antibodies against SARS-CoV-2. Sample electrochemiluminescence signals are compared with an internal calibration curve and quantitative values are reported as units (U)/mL.

In-house assay validation experiments demonstrated:

- i. The intra-assay and interassay coefficient of variation were 1.3% and 5.6%, respectively.
- ii. Anti-SARS-CoV-2 (S) antibodies were stable in uncentrifuged blood and serum at ambient temperature for up to 7 days permitting postal transport.
- iii. No effect was observed on recovery of anti-SARS-CoV-2 (S) antibodies following four freeze/thaw cycles.
- iv. No analytical interference was observed for the detection of anti-SARS-CoV-2 (S) with infliximab or vedolizumab up to 10 000 mg/L and 60 000 mg/L, respectively, or with antidrug antibodies to infliximab or vedolizumab up to 400 AU/mL and 38 AU/mL, respectively (data not shown).

At entry to CLARITY IBD and at follow-up visits, all patients were tested for previous SARS-CoV-2 infection using the Roche Elecsys anti-SARS-CoV-2 (N) immunoassay. Because antibody responses are impaired following PCR-confirmed natural infection we set a threshold of 0.25 times the Cut-Off Index (COI) at or above which patients were deemed to have had prior infection.¹¹ We defined a second threshold of 0.12 times the COI, below which patients were deemed to have no evidence of prior infection. Patients with a PCR test confirming SARS-CoV-2 infection at any time prior to vaccination were deemed to have evidence of past infection irrespective of any antibody test result.

Our threshold for seroconversion was defined at Roche Diagnostics (Penzberg, Germany). In brief, anti-SARS-CoV-2 (S) antibodies in 534 serum samples from 210 patients (71 hospitalised with severe COVID-19 and 139 patients with milder disease who were not hospitalised) were correlated with results from the cPass SARS-CoV-2 Neutralisation Antibody Detection Kit (Genscript, Netherlands), a competitive ELISA that reports the proportion of anti-SARS-CoV-2 antibodies that are neutralising.¹⁵ While individuals infected with SARS-CoV-2 develop binding antibodies to the virus, not all develop neutralising antibodies which block cellular infiltration and replication of the virus.¹⁶ In both cohorts, Elecsys Anti-SARS-CoV-2 spike (S) concentrations of greater than or equal to 15 U/mL were associated with neutralisation of $\geq 20\%$ with a positive predictive value of 99.10% (95% CI 97.74% to 99.64%) (online supplemental figure 1).

Ethical consideration and role of funders

CLARITY IBD is an investigator-led, UK National Institute for Health Research COVID-19 urgent public health study, funded by the Royal Devon and Exeter NHS Foundation Trust, Hull University Teaching Hospital NHS Trust, and by unrestricted educational grants from F. Hoffmann-La Roche AG (Switzerland), Biogen GmbH (Switzerland), Celltrion Healthcare (South Korea), Takeda (UK) and Galapagos NV (Belgium).

None of our funding bodies had any role in study design, data collection or analysis, writing, or decision to submit for publication. Patients were included after providing informed, written consent. The sponsor was the Royal Devon and Exeter NHS Foundation Trust. The protocol is available online at https:// www.clarityibd.org. The study was registered with the ISRCTN registry.

Statistics

The sample size for CLARITY IBD was based on the number of participants required to demonstrate a difference in the impact of infliximab and vedolizumab on seroprevalence and seroconversion following SARS-CoV-2 infection, with an estimated background seroprevalence of 0.05. We calculated that a sample of 6970 patients would provide 80% power to detect differences in the seroprevalence of anti-SARS-CoV-2 antibodies in infliximab-treated patients compared with vedolizumab-treated patients, while controlling for immunomodulator status at the 0.05 significance level. We stored and then analysed all serum samples as soon as the Roche Elecsys anti-SARS-CoV-2 (S) immunoassay was established in our laboratory.

Statistical analyses were undertaken in R V.4.0.4 (R Foundation for Statistical Computing, Vienna, Austria). All tests were two tailed and values of p < 0.05 were considered significant. We included patients with missing clinical data in analyses for which they had data and have specified the denominator for each variable. Anti-S antibody concentrations are reported as geometric means and SD. Other continuous data are reported as median and IQR, and discrete data as numbers and percentages, unless otherwise stated.

Univariable analyses, using t-tests of log-transformed anti-SARS-CoV-2 (S) antibody concentration and Spearman's rank correlation coefficients, were used to identify demographic, disease, vaccine, and treatment-related factors associated with the concentration of anti-SARS-CoV-2 (S) antibodies. To test our primary outcome, we used multivariable linear regression models to identify factors independently associated with log anti-SARS-CoV-2 (S) levels. A priori, we included age, ethnicity,

Inflammatory bowel disease

biological medication and immunomodulator use. No stepwise regression was performed. Results are presented after exponentiation, so that the coefficients of the model correspond to the fold change (FC) associated with each binary covariate. For age, a cut-off was chosen based on graphical inspection of the relationship between age and anti-SARS-CoV-2 (S) antibody concentrations. We also report the proportions of patients who seroconverted following vaccination. Seroconversion was defined as a threshold of 15 U/mL. We conducted sensitivity analyses to compare antibody responses stratified by participants with serological or PCR evidence of SARS-CoV-2 infection at any time prior to vaccination and in those who had received two doses of vaccine.

RESULTS

Patient characteristics

Between September 22 2020 and December 23 2020, 7226 patients were recruited to the CLARITY study from 92 UK hospitals.¹¹ For the primary immunogenicity analyses we included 865 infliximab-treated and 428 vedolizumab-treated participants without evidence of prior SARS-CoV-2 infection, who had received uninterrupted biological therapy

since recruitment and had an antibody test between 21 and 70 days after primary vaccination. Participant characteristics are shown in table 1.

Anti-SARS-CoV-2 (S) antibody level following primary COVID-19 vaccine

Geometric mean (geometric SD) anti-SARS-CoV-2 (S) antibody concentrations were lower in patients treated with infliximab than vedolizumab, following both the BNT162b2 (6.0 U/mL (5.9) vs 28.8 U/mL (5.4) p<0.0001) and ChAdOx1 nCoV-19 (4.7 U/mL (4.9) vs 13.8 U/mL (5.9) p<0.0001) vaccines (figure 1). Among infliximab-treated patients, the geometric mean (geometric SD) anti-SARS-CoV-2 (S) antibody concentrations were also lower in patients treated with a concomitant immunomodulator. Additional univariable analyses are shown in table 2.

In our multivariable models, anti-SARS-CoV-2 antibody concentrations were lower in infliximab-treated patients compared with vedolizumab-treated patients in participants who received the BNT162b2 (FC 0.29 (95% CI 0.21 to 0.40), p<0.0001) and ChAdOx1 nCoV-19 (FC 0.39 (95% CI 0.30 to 0.51), p<0.0001) vaccines. Age ≥ 60 years, immunomodulator

 Table 1
 Baseline characteristics of participants who had anti-SARS-CoV-2 spike antibodies measured 3–10 weeks following primary vaccination against SARS-CoV-2

Variable		Infliximab	Vedolizumab	Overall	P value
Vaccine	BNT162b2	44.7% (387/865)	47.2% (202/428)	45.6% (589/1293)	0.41
	ChAdOx1 nCoV-19	55.3% (478/865)	52.8% (226/428)	54.4% (704/1293)	
Age (years)		41.4 (31.5–54.8)	49.6 (37.1–63.8)	43.8 (32.8–57.6)	<0.0001
Sex	Female	50.3% (434/863)	47.1% (200/425)	49.2% (634/1288)	0.19
	Male	49.7% (429/863)	52.7% (224/425)	50.7% (653/1288)	
	Intersex	0.0% (0/863)	0.0% (0/425)	0.0% (0/1288)	
	Prefer not to say	0.0% (0/863)	0.2% (1/425)	0.1% (1/1288)	
Ethnicity	White	91.8% (791/862)	89.9% (381/424)	91.1% (1172/1286)	0.62
	Asian	5.3% (46/862)	7.5% (32/424)	6.1% (78/1286)	
	Mixed	1.9% (16/862)	1.9% (8/424)	1.9% (24/1286)	
	Black	0.7% (6/862)	0.5% (2/424)	0.6% (8/1286)	
	Other	0.3% (3/862)	0.2% (1/424)	0.3% (4/1286)	
Diagnosis	Crohn's disease	65.4% (566/865)	40.7% (174/428)	57.2% (740/1293)	0.00050
	Ulcerative colitis or IBD unclassified	34.6% (299/865)	59.3% (254/428)	42.8% (553/1293)	
Duration of IBD (years)		8.0 (4.0–16.0)	10.0 (5.0–17.8)	9.0 (4.0–16.0)	0.0040
Age at IBD diagnosis (years)		28.8 (21.6-41.8)	34.0 (23.3–47.6)	30.3 (21.9–43.7)	< 0.0001
Immunomodulator		61.6% (533/865)	22.0% (94/427)	48.5% (627/1292)	< 0.0001
5-ASA		23.0% (199/865)	31.6% (135/427)	25.9% (334/1292)	0.0012
Steroids		3.0% (26/865)	8.4% (36/427)	4.8% (62/1292)	<0.0001
BMI (kg/m ²)		25.9 (22.8–30.6)	26.1 (23.1–30.1)	26.0 (22.9–30.4)	0.75
Heart disease		3.6% (31/865)	6.5% (28/428)	4.6% (59/1293)	0.023
Diabetes		3.8% (33/865)	7.5% (32/428)	5.0% (65/1293)	0.0065
Lung disease		13.5% (117/865)	18.2% (78/428)	15.1% (195/1293)	0.032
Kidney disease		1.2% (10/865)	2.1% (9/428)	1.5% (19/1293)	0.22
Cancer		0.5% (4/865)	2.1% (9/428)	1.0% (13/1293)	0.013
Smoker	Yes	9.7% (84/862)	5.4% (23/425)	8.3% (107/1287)	0.0010
	Not currently	32.0% (276/862)	41.6% (177/425)	35.2% (453/1287)	
	Never	58.2% (502/862)	52.9% (225/425)	56.5% (727/1287)	
Exposure to documented cases of COVID-19		9.4% (81/862)	8.7% (37/425) 9.2% (118/1287)		0.76
Income deprivation score		0.086 (0.052–0.151)	0.084 (0.054–0.141) 0.086 (0.052–0.14		0.94
Active disease (PRO2)		4.9% (41/831)	11.4% (46/405)	7.0% (87/1236)	<0.0001

Values presented are median (interquartile range) or percentage (numerator/denominator). P values represent the results of a Mann Whitney U, Kruskal Wallis or Fisher's exact test.

5-ASA, 5-aminosalicylic acid; BMI, body mass index; IBD, inflammatory bowel disease; ; PRO2, IBD disease activity.



Figure 1 Anti-SARS-CoV-2 spike antibody concentration stratified by biological therapy (infliximab vs vedolizumab) and type of vaccine. The wider bar represents the geometric mean, while the narrower bars are drawn one geometric SD either side of the geometric mean. The threshold shown of 15 U/mL was used to determine seroconversion.

use and current smoking were also independently associated with lower anti-SARS-CoV-2 antibody concentrations in participants who received either vaccine. Conversely, non-white ethnicity was associated with higher antibody concentrations following both vaccines (figure 2).

To allow us to calculate a 15-day rolling geometric mean of anti-SARS-CoV-2 antibody concentrations, we included 2126 participants who had an antibody test carried out between 1 and 63 days after primary vaccination (1427 treated with infliximab and 699 treated with vedolizumab), as shown in figure 3. Three weeks after vaccination, we observed lower anti-SARS-CoV-2 (S) antibody concentrations in infliximabtreated patients compared with vedolizumab-treated patients following both vaccines. Sustained serological responses were observed in the vedolizumab-treated patients but not infliximab-treated patients.

Seroconversion following primary COVID-19 vaccination

The lowest rates of seroconversion were observed in participants treated with infliximab in combination with an immunomodulator with both the BNT162b2 (27.1%; 65/240) or ChAdOx1 nCoV-19 (20.2%; 60/297) vaccines. Highest rates of seroconversion were seen in patients treated with vedolizumab monotherapy who received the BNT162b2 (74.7%;124/166) or ChAdOx1 nCoV-19 (57.3%; 94/164) vaccines (figure 4).

Antibody responses following prior SARS-CoV-2 infection

Among participants with SARS-CoV-2 infection prior to vaccination, geometric mean (SD) anti-SARS-CoV-2 (S) antibody concentrations were lower in infliximab-treated patients compared with vedolizumab-treated patients in those who received a single dose of BNT162b2 (191 U/mL (12.5) vs 1865 U/mL (8.0) p<0.0001) and ChAdOx1 nCoV-19 (185 U/mL (9.3) vs 752 (12.5) p=0.046) vaccines. In both infliximab-treated patients and vedolizumab-treated patients, antibody concentrations following vaccination were higher than those observed in patients without prior infection (figure 5). Overall, across both vaccines, 81.7% (76/93) patients treated with infliximab and 97.1% (33/34) patients treated with vedolizumab seroconverted (p=0.041).

Antibody responses following two COVID-19 vaccine doses

Antibody responses were assessed in 27 patients following two doses of the BNT162b2 vaccine without serological evidence of prior infection (figure 5). In both infliximab- and vedolizumab-treated patients, antibody levels and seroconversion rates were higher after two doses than after a primary vaccine without prior infection (geometric means infliximab 158 U/mL (7.0) vs 6.0 U/mL (5.9), p<0.0001; vedolizumab 562 U/mL (11.5) vs 28.8 U/mL (5.4), p=0.018). After second-vaccine doses, 85% (17/20) infliximab-treated patients and 86% (6/7) vedolizumab-treated patients seroconverted (p=0.68).

DISCUSSION

We have shown that anti-SARS-CoV-2 spike antibody levels and rates of seroconversion are lower following vaccination with a single dose of either BNT162b2 or ChAdOx1 nCoV-19 vaccines in patients with IBD treated with infliximab than vedolizumab. Combination therapy with an immunomodulator further attenuated immunogenicity to both vaccines in infliximab-treated patients. Reassuringly, however, a second exposure to antigen, either by vaccination after infection, or a second dose of vaccine led to seroconversion in most patients.

Direct comparisons between our data and the antibody responses reported in the vaccine registration trials are limited by differences in the assays used to define immunogenicity and the adoption of different thresholds to define seroconversion. No adequately powered studies have reported the effect of anti-TNF drugs on vaccine responses.¹⁷ Our findings are similar, however, to recent reports of the immunogenicity of the BNT162b2 and mRNA-1273 vaccines in transplant recipients and in patients with malignancy treated with antimetabolite immunosuppression, conventional chemotherapy or immune checkpoint inhibitors.¹⁸¹⁹ The authors showed fewer patients treated with potent immunosuppressants seroconverted than healthy controls. Importantly, as we have also shown here, second vaccine doses led to seroconversion in the cancer cohort. However, even after two antigen exposures, a small subset of patients (18% (20/113) infliximab-treated patients and 5% (2/41) vedolizumab-treated patients) in our study failed to mount an antibody response. To identify this group, and because the sustainability of antibody responses overall is unknown, serial measurement of antibody responses is indicated.

Urgent research is needed to understand the factors linked to non-response and how to potentiate long-term immunogenicity in this group. Strategies to be tested include the manipulation of timing of second vaccinations, booster doses, the use of adjuvants and/or switching between vaccines with different mechanisms of action. Moreover, from the public health standpoint, recent case reports have shown that potent immunosuppression leads to chronic nasopharyngeal carriage and evolution of new SARS-CoV-2 variants.^{20 21} Whether this occurs in patients treated with anti-TNF therapy with impaired antibody response is an important conceptual concern.

Our data have other important findings relating to SARS-CoV-2 vaccine responses. We have demonstrated that antibody responses to SARS-CoV-2 vaccines are reduced in older individuals and current smokers. Smoking has also been associated with lower antibody responses to hepatitis B vaccination and faster decay of antibodies after vaccination with live attenuated and trivalent influenza vaccines.^{22 23} We have also demonstrated higher antibody responses to both the BNT162b2 and ChAdOx1 nCoV-19 vaccines in non-white participants. This might be explained by differences in genetics,²⁴ gut microbiota,²⁵

Table 2 Univariable associations with anti-SARS-CoV-2 spike antibodies, stratified by vaccine type						
		BNT162b2		ChAdOx1 nCoV-19		
Variable		Value	P value	Value	P value	
Biological treatment	Infliximab	6.0 (5.9)	<0.0001	4.7 (4.9)	<0.0001	
	Vedolizumab	28.8 (5.4)		13.8 (5.9)		
Immunomodulator in infliximab-treated participants	No	9.7 (4.7)	< 0.0001	5.7 (5.1)	0.045	
	Yes	4.4 (6.3)		4.2 (4.7)		
Immunomodulator in vedolizumab-treated participants	No	32.4 (5.2)	0.052	15.6 (6.0)	0.082	
	Yes	16.7 (6.3)		10.0 (5.5)		
Age (years)		rho=-0.22	<0.0001	rho=-0.15	<0.0001	
Sex	Female	9.4 (7.0)	0.092	6.6 (5.5)	0.83	
	Male	10.9 (6.3)		6.8 (5.7)		
Ethnicity	White	9.4 (6.6)	0.037	6.2 (5.6)	0.0051	
	Asian	20.9 (7.3)		16.1 (5.2)		
	Mixed	25.7 (6.7)		13.7 (5.3)		
	Black	12.5 (1.6)		19.4 (2.2)		
	Other	22.9 (3.7)		5.7 (3.1)		
Diagnosis	Crohn's disease	7.3 (6.4)	<0.0001	5.6 (5.6)	0.0014	
	Ulcerative colitis or IBD- unclassified	15.6 (6.5)		8.5 (5.5)		
Duration of IBD (years)		rho=-0.16	< 0.0001	rho=-0.12	0.0013	
Age at IBD diagnosis (years)		rho=-0.13	0.0021	rho=-0.04	0.25	
5-ASA	No	9.8 (6.6)	0.40	6.7 (5.5)	0.93	
	Yes	11.5 (7.1)		6.6 (5.9)		
Steroids	No	10.2 (6.7)	0.90	6.8 (5.5)	0.12	
	Yes	10.7 (7.3)		4.1 (6.7)		
BMI (kg/m ²)		rho=-0.08	0.068	rho=-0.01	0.81	
Heart disease	No	10.3 (6.7)	0.65	6.9 (5.6)	0.010	
	Yes	8.7 (7.0)		2.8 (5.2)		
Diabetes	No	10.7 (6.7)	0.0028	6.8 (5.6)	0.066	
	Yes	4.1 (4.6)		4.0 (5.2)		
Lung disease	No	10.1 (6.9)	0.70	6.9 (5.5)	0.31	
	Yes	10.9 (5.7)		5.7 (6.1)		
Kidney disease	No	10.2 (6.6)	0.60	6.7 (5.5)	0.66	
	Yes	15.6 (10.4)		4.7 (12.4)		
Cancer	No	10.4 (6.6)	0.13	6.7 (5.6)	0.069	
	Yes	2.0 (9.2)		2.3 (3.6)		
Smoking	Yes	4.7 (7.1)	0.0077	3.4 (4.8)	0.00077	
	Not currently	9.4 (6.6)		6.1 (5.4)		
	Never	11.8 (6.5)	0.07	8.0 (5.7)	0.50	
Exposure to documented cases of COVID-19	No	10.3 (6.7)	0.87	6.6 (5.5)	0.53	
	Yes	9.8 (6.8)	0.75	7.8 (6.1)		
Income deprivation score		rho=0.01	0.75	rho=0.02	0.65	
Active disease (PRO2)	No	10.1 (6.5)	0.32	6.6 (5.4)	0.51	
	Yes	14.0 (7.6)		8.1 (7.0)		

Values presented are geometric mean antibody concentration (geometric SD) or Spearman's rho. P values represent the results of an unpaired t-test or test of Spearman's rho. 5-ASA, 5-aminosalicylic acid; BMI, body mass index; IBD, inflammatory bowel disease; VAS, Visual Analogue Scale.

nutrition²⁶ and priming of the immune system by prior exposure to SARS-CoV-2 not detected by our prevaccination antibody test. Lower antibody concentrations were also observed in patients with Crohn's disease when compared with patients with UC or IBD unclassified. Despite evidence of defective mucosal immunity, previous vaccine studies involving patients with Crohn's disease or UC have not shown attenuated antibody responses to vaccination in the absence of concomitant immunomodulator or biological therapy.⁶⁷

The cytokine TNF shapes multiple aspects of host immune responses, including T-cell dependent antibody production. Genetic ablation of TNF results in disruption of B-cell follicles in germinal centres with defective induction of antigen-induced antibody production.^{27 28} These biological properties may in part explain why TNF blockade is clinically beneficial in IMIDs, but also explain the increased risk of serious and opportunistic infections and impaired response to other vaccines.

Our findings have important implications for patients treated with anti-TNF drugs particularly those also treated with an immunomodulator. Poor antibody responses to a single dose of vaccine unnecessarily exposes infliximab-treated patients to SARS-CoV-2 infection. However, because we observed higher rates of seroconversion in patients with two exposures to SARS-CoV-2 antigen, even in the presence of TNF blockade,

ln fl	ammat	ory		dicasca
	amma	LUIYI	JUWEI	uisease

BN 11822 Fold change (95% CI) p Infliximab (vs vedolizumab) 386/587 0.29 (0.21, 0.40) <0.0001 Immunomodulator 260/587 0.41 (0.31, 0.55) <0.0001 Crohn's disease (vs UC or IBDU) 327/587 0.41 (0.31, 0.55) <0.0001 Age ≥ 60 113/587 0.75 (0.57, 1.00) 0.046 Mape ≥ 60 113/587 0.34 (0.24, 0.47) <0.0001 Non-white ethnicity 61/587 1.81 (1.16, 2.81) 0.0085 Current smoker 44/587 0.52 (0.31, 0.80) 0.011 Variable N Fold change (95% CI) p Infliximab (vs vedolizumab) 475/697 0.38 (0.50, 0.84) 0.00085 Crohn's disease (vs UC or IBDU) 409/697 0.48 (0.80, 0.65) <0.0011 Non-white ethnicity 55/697 0.48 (0.80, 0.65) 0.0017 Current smoker 63/697 0.48 (0.80, 0.65) 0.0017 Curr							
Variable N Polic Grange 95% CI) p Infliximab (vs vedolizumab) 386/557 0.29 (0.21, 0.40) <0.0001 Crohrs disease (vs UC or IBDU) 327/557 0.75 (0.57, 1.00) 0.046 Age 2 60 113/557 0.75 (0.57, 1.00) 0.046 Non-white ethnicity 61/557 1.81 (1.16, 2.81) 0.00085 Current smoker 44/587 0.39 (0.30, 0.51) <0.0011 ChAdOx1 nCoV-19 0.39 (0.30, 0.51) <0.00085 0.52 (0.31, 0.86) 0.011 Variable N 0.39 (0.30, 0.51) <0.00085 <0.0001 Immunomodulator 355/697 0.39 (0.30, 0.51) <0.0001 Variable N 0.38 (0.50, 0.84) 0.00085 <0.0011 Immunomodulator 355/697 0.49 (0.36, 0.85) <0.0011 Variable N 0.49 (0.36, 0.84) 0.0059 All 0.34 (0.27, 0.41) 0.0017	BN1162D2	N				Fold change (95% CI)	
Immunomodulator 289/367 0.47 (0.21, 0.42) <0.0001	variable	296/697				0.20 (0.21, 0.40)	p <0.0001
Immunomodulator Construction Construct	Innixinab (vs veuolizumab)	300/307				0.29 (0.21, 0.40)	<0.0001
Cronn s disease (vs UC or IBUU) 32/7867 0.75 (0.57, 1.00) 0.048 Age 2 60 13/587 0.75 (0.24, 0.47) <0.0001 Current smoker 44/587 0.52 (0.37, 0.86) 0.011 ChAdOx1 nCoV-19 0.52 (0.31, 0.86) 0.011 Variable N Fold change (95% CI) p Infliximab (vs vedolizumab) 47/5697 0.53 (0.30, 0.51) <0.0001 Immunomodulator 355/697 0.54 (0.36, 0.84) 0.00059 Age 2 60 0.48 (0.36, 0.65) <0.0011 Variable 162/697 0.55 (0.36, 0.84) 0.00059 All 20 0.51 2 0.051 (0.27, 1.00) Variable N 50/697 0.55 (0.36, 0.84) 0.00059 All 20 0.51 2 0.0011 Variable N 50/697 0.55 (0.36, 0.84) 0.0059 All 20 0.52 (0.31, 0.41) <0.0017 Variable N Fold change (95% CI) p 1 Infiximab (vs vedolizumab) 85/1/224 - 0.51 (0.42, 0.62) <0.0011 Variable N Fold change (95% CI) p 0.0017 0.0017 Current smoker 63/697 - 0.51 (0.42, 0.62) <0.0011 <th>Immunomodulator</th> <th>269/587</th> <th>_</th> <th></th> <th></th> <th>0.41 (0.31, 0.55)</th> <th><0.0001</th>	Immunomodulator	269/587	_			0.41 (0.31, 0.55)	<0.0001
Age 2 b0 113/587 0.34 (0.24, 0.47) <0.0001 Non-white ethnicity 61/587 1.81 (1.45, 2.81) 0.00085 Current smoker 44/587 0.52 (0.31, 0.86) 0.011 023 0.5 2 0.52 (0.31, 0.86) 0.011 Variable N Fold change (95% Cl) p Infliximab (vs vedolizumab) 475/697 0.39 (0.30, 0.51) <0.0001 Immunomodulator 355/697 0.65 (0.50, 0.84) 0.00095 Crohn's disease (vs UC or IBDU) 409/697 0.48 (0.36, 0.65) <0.0001 Non-white ethnicity 55/697 0.55 (0.36, 0.84) 0.0001 Variable N 0.45 (0.36, 0.65) <0.0001 Variable N 0.55 (0.36, 0.84) 0.0001 Variable N 0.55 (0.36, 0.84) 0.0001 Variable N 0.55 (0.36, 0.84) 0.0001 Variable N 0.56 (0.36, 0.84) 0.0001 Variable N 0.56 (0.26, 0.44) 0.0001 Immunomodulator 62/1/284	Cronn's disease (vs UC or IBDU)	327/587	-			0.75 (0.57, 1.00)	0.046
Non-white ethnicity 61/587 181 (1.16, 2.81) 0.0085 Current smoker 44/587 2 0.52 (0.31, 0.86) 0.011 ChadOx1 nCoV-19 2 0.52 (0.31, 0.86) 0.011 Uariable N Fold change (95% CI) p Infliximab (vs vedolizumab) 475/697 - 0.33 (0.30, 0.51) < 0.00016 Crohrs disease (vs UC or IBDU) 409/697 - 0.88 (0.66, 1.09) 0.20 Age ≥ 60 0.46 (0.36, 0.65) < 0.0017 - 0.48 (0.36, 0.65) < 0.0017 Current smoker 63/697 - - 0.48 (0.36, 0.65) < 0.0017 Variable N - - 0.48 (0.36, 0.65) < 0.0017 Variable N - - 0.55 (0.36, 0.84) 0.0059 All - - 0.34 (0.27, 0.41) < 0.0017 Variable N - - 0.54 (0.27, 0.21) 0.0017 Immunomodulator 63/1224 - - 0.51 (0.42, 0.62) < 0.0001 <	Age 2 60	113/587			_	0.34 (0.24, 0.47)	<0.0001
Current smoker 44/587 0.52 (0.31, 0.86) 0.011 052 0.5 2 0.5 2 0.52 (0.31, 0.86) 0.011 ChAdOx1 nCoV-19 Variable N Fold change (95% CI) p Imfliximab (vs vedolizumab) 475/697 0.39 (0.30, 0.51) <0.0001 Immunomodulator 355/697 0.65 (0.50, 0.84) 0.00085 Crohn's disease (vs UC or IBDU) 409/897 0.48 (0.36, 0.65) <0.0001 Non-white ethnicity 55/697 0.55 (0.36, 0.84) 0.0001 Variable N 0.55 (0.36, 0.84) 0.0059 d2 0.3 2 0.51 (0.27, 0.14) 0.0017 Current smoker 63/697 0.55 (0.36, 0.84) 0.0059 d1 0.34 (0.27, 0.41) -0.0017 0.55 (0.36, 0.84) 0.0059 d2 0.3 2 0.51 (0.42, 0.62) 0.0011 Immunomodulator 62/1/284 <th>Non-white ethnicity</th> <th>61/587</th> <th></th> <th></th> <th></th> <th>1.81 (1.16, 2.81)</th> <th>0.0085</th>	Non-white ethnicity	61/587				1.81 (1.16, 2.81)	0.0085
Variable N Fold change (95% CI) p Infliximably (vs vedolizumab) 475/697 0.39 (0.30, 0.51) < 0.0001 Inmunomodulator 355/697 0.85 (0.50, 0.44) 0.00005 Crohn's disease (vs UC or IBDU) 409/697 0.48 (0.36, 0.65) < 0.0011 Non-write ethnicity 55/697 0.48 (0.36, 0.65) < 0.00015 Variable 0.48 (0.36, 0.65) < 0.0017 0.0017 Variable 0.55 (0.36, 0.84) 0.0059 Variable N 0.54 (0.27, 0.21) Immunomodulator 65/1/224 0.54 (0.27, 0.21) Variable N 0.54 (0.27, 0.52) Immunomodulator 62/1/224 0.54 (0.27, 0.52) Immunomodulator 62/1/224 0.57 (0.42, 0.62) <0.0001 Crohn's disease	Current smoker	44/587	<u> </u>			0.52 (0.31, 0.86)	0.011
ChAdox1 nCoV-19 Variable N Variable Variable N Variable N Variable N Variable N Variable N Variable N Variable Variable N Variable Variable N Variable N Variable N			0.2	0.5 1	2		
Variable N Fold change (95% CI) p Infliximably (vs vedolizumab) 475/697 - 0.39 (0.30, 0.51) <0.0001 Immunomodulator 355/697 - 0.85 (0.50, 0.84) 0.00085 Crohn's disease (vs UC or IBDU) 49/897 - 0.85 (0.50, 0.84) 0.00085 Crohn's disease (vs UC or IBDU) 49/897 - - 0.48 (0.36, 0.65) <0.0011 Non-white ethnicity 55/697 - - 0.48 (0.36, 0.65) <0.0011 Current smoker 63/697 - - 0.55 (0.36, 0.84) 0.0059 All - - 0.34 (0.27, 0.41) <0.0011 Variable N Fold change (95% CI) p Inmunomodulator 62/1/284 - - 0.54 (0.27, 0.41) <0.0001 Crohn's disease (vs UC or IBDU) 73/1/284 - - 0.79 (0.66, 0.96) 0.015 Age ≥ 60 275/1/284 - - 0.49 (0.32, 0.50) <0.0001 Non-whitie ethnicity	ChAdOx1 nCoV-19						
Infliximab (vs vedolizumab) 475/697 0.39 (0.30, 0.51) <0.0001	Variable	N				Fold change (95% CI)	р
Immunomodulator 355/697 0.65 (0.50, 0.84) 0.00085 Crohn's disease (vs UC or IBDU) 400/697 0.85 (0.66, 1.09) 0.20 Age ≥ 60 162/697 0.85 (0.66, 1.09) 0.20 Non-white ethnicity 55/697 0.48 (0.36, 0.65) <0.0011 Current smoker 63/697 0.55 (0.36, 0.84) 0.0059 All 0.34 (0.27, 0.41) <0.0001 <0.0011 Immunomodulator 62/1/284 0.34 (0.27, 0.41) <0.0011 Crohn's disease (vs UC or IBDU) 736/1284 0.51 (0.42, 0.62) <0.0011 Crohn's disease (vs UC or IBDU) 736/1284 0.40 (0.32, 0.50) <0.0011 Crohn's disease (vs UC or 1BDU) 736/1284 0.40 (0.32, 0.50) <0.0011 Non-white ethnicity 116/1284 0.40 (0.32, 0.50) <0.0001 Crohn's disease (vs UC or 1BDU) 736/1284 0.40 (0.32, 0.50) <0.0001 Non-white ethnicity 116/1284 <t< th=""><th>Infliximab (vs vedolizumab)</th><th>475/697</th><th>-</th><th>H :</th><th></th><th>0.39 (0.30, 0.51)</th><th>< 0.0001</th></t<>	Infliximab (vs vedolizumab)	475/697	-	H :		0.39 (0.30, 0.51)	< 0.0001
Crohr's disease (vs UC or IBDU) 409/697 0.85 (0.66, 1.09) 0.20 Age ≥ 60 162/697 0.48 (0.36, 0.65) <0.0001 Non-white ethnicity 55/697 0.48 (0.36, 0.65) <0.0001 Current smoker 63/697 0.55 (0.36, 0.84) 0.0017 Variable 0.55 (0.36, 0.84) 0.0059 Variable N Fold change (95% CI) p Inmunomodulator 62/1/284 0.54 (0.27, 0.21) <0.0001 Crohr's disease (vs UC or IBDU) 73/1284 0.79 (0.66, 0.96) 0.015 Age ≥ 60 275/1284 0.40 (0.32, 0.57) <0.0001 Non-white ethnicity 116/1284 0.40 (0.32, 0.57) <0.0001	Immunomodulator	355/697		-		0.65 (0.50, 0.84)	0.00085
Age ≥ 60 162/697 0.48 (0.36, 0.65) <0.0001	Crohn's disease (vs UC or IBDU)	409/697				0.85 (0.66, 1.09)	0.20
Non-white ethnicity 55/697 2.04 (1.31, 3.17) 0.0017 Current smoker 63/697 0.55 (0.36, 0.84) 0.0059 All	Age ≥ 60	162/697	-	.		0.48 (0.36, 0.65)	< 0.0001
Current smoker 63/697 - 0.55 (0.36, 0.84) 0.0059 All 0.5 0.5 0.5 0.5 0.5 Variable N Fold change (95% Cl) p Infliximab (vs vedolizumab) 85/11/24 - 0.34 (0.27, 0.41) <0.0001 Immunondulator 624/1284 - - 0.51 (0.42, 0.62) <0.0011 Crohn's disease (vs UC or IBDU) 73/1284 - - 0.40 (0.32, 0.50) <0.0001 Non-white ethnicity 116/1284 - - 0.40 (0.32, 0.50) <0.0001 Current emplore 107/1284 - - 0.40 (0.32, 0.50) <0.0001	Non-white ethnicity	55/697				2.04 (1.31, 3.17)	0.0017
b2 0.5 2 Variable N Fold change (95% Cl) p Infliximab (vs vedolizumab) 661/1284 0.34 (0.27, 0.41) <0.0001 Immunomodulator 624/1284 0.51 (0.42, 0.62) <0.0001 Crohn's disease (vs UC or IBDU) 736/1284 0.79 (0.66, 0.96) 0.015 Age ≥ 60 2257/1284 0.40 (0.32, 0.50) <0.0001 Non-white ethnicity 116/1284 0.51 (4.32, 6.7) <0.0001 Current ethnicity 116/1284 0.53 (0.30, 0.7) <0.0001	Current smoker	63/697	-			0.55 (0.36, 0.84)	0.0059
All Fold change (95% Cl) p Uariable N Fold change (95% Cl) p Infliximab (vs vedolizumab) 861/1284 -■ 0.34 (0.27, 0.41) <0.0001 Immunomodulator 624/1284 -■ 0.51 (0.42, 0.62) <0.0001 Crohn's disease (vs UC or IBDU) 736/1284 -■ 0.79 (0.66, 0.96) 0.015 Age ≥ 60 275/1284 -■ 0.40 (0.32, 0.50) <0.0001 Non-white ethnicity 116/1284 -■ 0.40 (0.32, 0.50) <0.0001 Current emplore 107/1284 -■ 0.51 (0.42, 0.75) <0.0001			0.2	0.5 1	2		
Variable N Fold change (95% CI) p Infliximab (vs vedolizumab) 661/1284 -■ 0.34 (0.27, 0.41) <0.0001 Immunomodulator 624/1284 -■ 0.51 (0.42, 0.62) <0.0001 Crohn's disease (vs UC or IBDU) 736/1284 -■ 0.79 (0.66, 0.96) 0.015 Age ≥ 60 225/1284 -■ 0.40 (0.32, 0.50) <0.0001 Non-white ethnicity 116/1284 -■ 0.40 (0.32, 0.50) <0.0001 Current ethnicity 116/1284 -■ 0.51 (4.32, 2.67) <0.0001	All						
Infliximab (vs vedolizumab) 85/1/284 -■ 0.34 (0.27, 0.41) <0.001 Immunomodulator 65/1/284 -■ 0.51 (0.42, 0.62) <0.001 Crohn's disease (vs UC or IBDU) 736/1284 -■ 0.79 (0.66, 0.96) 0.015 Age ≿ 60 275/1284 -■ 0.49 (0.32, 0.59) <0.001 Unovshite ethnicity 116/1284 -■ 0.49 (0.32, 0.59) <0.001 Unovshite ethnicity 116/1284 -■ 0.49 (0.32, 0.59) <0.001	Variable	N	1			Fold change (95% CI)	n
Immunomodulator 624/1284 - 0.51 (0.42, 0.62) <0.0001	Infliximab (vs vedolizumab)	861/1284				0.34 (0.27 0.41)	<0.0001
Crohn's disease (vs UC or IBDU) 736/1284 - ● 0.79 (0.66, 0.96) 0.015 Age ≿ 60 275/1284 - ● 0.49 (0.82, 0.50) - 0.0001 Non-white ethnicity 116/1284 - ● 0.49 (0.82, 0.50) - 0.0001 Unrest termination 116/1284 - ● - ● 0.59 (1.43, 2.67) - 0.0001	Immunomodulator	624/1284	-	-		0.51 (0.42, 0.62)	<0.0001
Age ≥ 6 257/1224 → 0.47 (0.00, 0.32, 0.5) 0.001 Non-white ethnicity 116/1224 → 0.49 (0.32, 0.5) <0.0011 Non-white ethnicity 116/1224 → 0.57 (0.30, 0.27) <0.0001 Unreader endler 107/1224 → 0.57 (0.30, 0.27) <0.0001	Crohn's disease (vs IIC or IBDII)	736/1284		-		0.79 (0.66, 0.96)	0.015
Non-white ethnicity 116/1284 Image: 100/1284 105 (1.43, 2.67) <0.0001		275/1284				0.40 (0.22, 0.50)	<0.0001
Current employ 1107/1284	Non white ethnicity	116/1204			-	1.05 (1.42, 0.50)	<0.0001
	Our and an alway	10/1204		-		1.55 (1.45, 2.07)	~0.0001
	Current smoker	107/1284				0.53 (0.39, 0.74)	0.00016

Figure 2 Exponentiated coefficients of linear regression models of log(anti-SARS-CoV-2 spike antibody concentration). The resultant values represent the fold change of antibody concentration associated with each variable. Each vaccine was modelled separately, and then a further model was created using all of the available data. IBDU, inflammatory bowel disease unclassified; UC, ulcerative colitis.

these patients should be prioritised for optimally timed second doses. Until patients receive a second vaccine dose they should consider that they are not protected from SARS-CoV-2 infection and continue to practice enhanced physical distancing and shielding if appropriate.

Limitations

While our data are biologically plausible, we acknowledge the following limitations of our study. We have used an electrochemiluminescence immunoassay to measure antibody concentrations rather than using a neutralising assay. Although neutralisation assays are considered more biologically relevant, it is now established that anti receptor-binding domain antibodies, which target the spike protein component that engages host cells through ligation of ACE 2, closely correlate with neutralisation assays.^{29 30} Our validation experiments, comparing anti-SARS-CoV-2 spike (S) concentrations with neutralisation using the cPass test in two cohorts of patients with PCR confirmed SARS-CoV-2 infection,



Figure 3 Rolling geometric mean antibody concentration over time, stratified by biological therapy (infliximab vs vedolizumab) and vaccine. Geometric means are calculated using a rolling 15-day window (ie, 7 days either side of the day indicated). The shaded areas represent the 95% Cls of the geometric means. Overall, data from 2126 participants (1427 on infliximab and 699 on vedolizumab) between 1 and 63 days post vaccination are included in this graph .



Figure 4 Percentages of participants with seroconversion defined by an anti-SARS-CoV-2 spike antibody concentration \geq 15 U/mL, stratified by vaccine, biological and immunomodulator use. Error bars represent the 95% CI of the percentages. IMM, immunomodulator.

confirm this correlation. Second, we only assessed humoral responses to infection, and it is likely that protective immunity additionally requires induction of memory T cell responses. Third, we were unable to investigate whether the timing of biological infusion with respect to vaccination or drug level at the time of vaccination, influences antibody responses. As follow-up blood tests occurred at the time of infusions, which for the vast majority occurred 8 weekly, the time from last infusion to vaccination was negatively correlated with the time from vaccination to the next antibody test, confounding these analyses. Finally, we investigated one anti-TNF drug, infliximab, only. However, we suspect that our key findings will apply to other anti-TNF biologics used to treat IMIDs, including adalimumab, certolizumab, golimumab and etanercept. Further observational data will be required to elucidate the impact of other classes of therapies for IMIDs on SARS-CoV-2 vaccine immunogenicity.

CONCLUSIONS

Infliximab is associated with attenuated immunogenicity to a single dose of the BNT162b2 and ChAdOx1 nCoV-19 SARS-CoV-2 vaccines in patients with IBD. Immunomodulators further blunted immunogenicity rates to both vaccines. Reassuringly, vaccination after infection, or a second dose of vaccine



Figure 5 Anti-SARS-CoV-2 spike antibody concentration, stratified by biological therapy (infliximab vs vedolizumab), prior infection and number of doses and type of vaccine. The wider bar represents the geometric mean, while the narrower bars are drawn one geometric SD either side of the geometric mean. The threshold shown of 15 U/mL is the one used to determine seroconversion.

led to seroconversion in most patients. Delayed second dosing should be avoided in patients treated with infliximab.

Author affiliations

¹Gastroenterology, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK ²Exeter Inflammatory Bowel Disease and Pharmacogenetics Research Group, University of Exeter, Exeter, UK

³Biochemistry, Exeter Clinical Laboratory International, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

⁴Gastroenterology, University Hospital Southampton NHS Foundation Trust, Southampton, UK

 5 Gastroenterology, University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK

⁶Gastroenterology, Guy's and St Thomas' Hospitals NHS Trust, London, UK

⁷School of Immunology & Microbial Sciences, King's College London, London, UK ⁸Gastroenterology, St Marks Hospital and Academic Institute, London, UK, London, UK

⁹Gastroenterology, Barts and The London NHS Trust, London, UK

¹⁰Centre for Immunobiology, Blizard Institute, Barts and The London School of Medicine and Dentistry Blizard Institute, London, UK

¹¹Translational & Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

¹²Gastroenterology, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

¹³Gastroenterology, Queen Elizabeth University Hospital, NHS Greater Glasgow and Clyde, Glasgow, UK

¹⁴School of Medicine, Dentistry and Nursing, University of Glasgow, Glasgow, UK ¹⁵Gastroenterology, University College London Hospitals NHS Foundation Trust, London, UK

 $^{16}\mbox{Gastroenterology},$ St George's University Hospitals NHS Foundation Trust, London, UK

 ¹⁷Institute for Infection & Immunity, University of London St George's, London, UK
 ¹⁸Gastroenterology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

¹⁹Gastroenterology, Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK

²⁰Gastroenterology, Kettering General Hospital NHS Foundation Trust, Kettering, UK
²¹Roche Diagnostics GmbH, Mannheim, Baden-Württemberg, Germany

²²IBD Unit - Gastroenterology, Hull University Teaching Hospitals NHS Trust, Hull, UK
²³Hull York Medical School, University of Hull, Hull, UK

²⁴Gastroenterology, Western General Hospital, Edinburgh, UK

²⁵The University of Edinburgh Centre for Genomic and Experimental Medicine, Edinburgh, UK

²⁶Metabolism, Digestion and Reproduction, Imperial College London, London, UK
²⁷Gastroenterology, Imperial College Healthcare NHS Trust, London, UK

Twitter Nicholas A Kennedy @DrNickKennedy, Simeng Lin @SimengLin, James R Goodhand @JamesGoodhand, Neil Chanchlani @nchanchlani1, Claire Bewshea @ clairebewshea, Rachel Nice @rachelnice3, Aileen Fraser @AileenIBDBRI, Nikolaos Kamperidis @KamperidisNik, Klaartje B Kok @klaartjekok, Christopher Andrew Lamb @DrChrisLamb, Jonathan Macdonald @jOnnymac1, Richard CG Pollok @ RichadPollok, Tim Raine @IBD_MB, Philip J Smith @DrPhilipJSmith, Ajay Mark Verma @UKGastroDr, Timothy J McDonald @T_J_McDonald, Shaji Sebastian @ibdseb, Charlie W Lees @charlie_lees, Nick Powell @NickPowellLab and Tariq Ahmad @ tariqahmadIBD

Acknowledgements CLARITY IBD is a UK National Institute for Health Research (NIHR) Urgent Public Health Study. The NIHR Clinical Research Network supported study setup, site identification and delivery of this study. This was facilitated by Professor Mark Hull, the national specialty lead for gastroenterology. We acknowledge the contribution of our Patient Advisory Group who helped shape the trial design around patient priorities. Our partners, Crohn's and Colitis UK (CCUK), continue to support this group and participate in Study Management Team meetings. We thank Professor Danny Altmann, Professor Rosemary Boyton, Professor Graham Cooke and Dr Katrina Pollock for their helpful discussions and review of the data. Laboratory tests were undertaken by the Exeter Blood Sciences Laboratory at the Royal Devon and Exeter NHS Foundation Trust. The Exeter NIHR Clinical Research Facility coordinated sample storage and management. Tariq Malik and James Thomas from Public Health England, Guy Stevens, Katie Donelon, Elen de Lacy from Public Health Wales and Johanna Bruce from Public Health Scotland supported linkage of central SARS-CoV-2 PCR test results with study data. Roche Diagnostics Limited provided the Elecsys Anti-SARS-CoV-2 immunoassay for the study. SL is supported by a Wellcome GW4-CAT fellowship. NC acknowledges support from CCUK. CAL acknowledges support from the NIHR Newcastle Biomedical Research Centre and the support of the Programmed Investigation Unit at Royal Victoria Infirmary, Newcastle on Tyne. TR acknowledges support with recruitment from the NIHR Cambridge Biomedical Research Centre. CWL is funded by a UKRI Future Leaders

Fellowship. NP is supported by the NIHR Imperial Biomedical Research Centre. We acknowledge the study coordinators of the Exeter Inflammatory Bowel Disease and Pharmacogenetics Research Group: Marian Parkinson and Helen Gardner-Thorpe for their ongoing administrative support to the study. The sponsor of the study was the Royal Devon and Exeter NHS Foundation Trust.

Collaborators Contributors to the CLARITY IBD study: Barts Health NHS Trust Klaartje Kok, Farjhana Bokth, Bessie Cipriano, Caroline Francia, Nosheen Khalid, Hafiza Khatun, Ashley Kingston, Irish Lee, Anouk Lehmann, Kinnari Naik, Elise Pabriaga, Nicolene Plaatijes, Kevin Samuels. Barts Health NHS Trust (paediatric): Bessie Cipriano, Kevin Samuels, Nicolene Plaatjies, Hafiza Khatun, Farjana Bokth, Elise Pabriaga, Caroline Francia. Basingstoke and North Hampshire Hospital: Rebecca Saich, Hayley Cousins, Wendy Fraser, Rachel Thomas, Matthew Brown, Benjamin White. Birmingham Women's and Children's NHS Foundation Trust: Rafeeq Muhammed, Rehana Bi, Catherine Cotter, Jayne Grove, Kate Hong, Ruth Howman, Monica Mitchell, Sugrah Sultan. Bolton NHS Foundation Trust: Salil Singh, Chris Dawe, Robert Hull, Natalie Silva. Borders General Hospital: Manning Jansen, Lauren Jansen. Calderdale and Huddersfield NHS Foundation Trust: Sunil Sonwalkar, Naomi Chambers, Andrew Haigh, Lear Matapure Cambridge University Hospitals NHS Foundation Trust: Tim Raine, Varun George, Christina Kapizioni, Konstantina Strongili. Chelsea and Westminster Hospital NHS Foundation Trust: Tina Thompson, Philip Hendy, Rhian Bull, Patricia Costa, Lisa Davey, Hayley Hannington, Kribashnie Nundlall, Catarina Martins, Laura Avanzi, Jaime Carungcong, Sabrina Barr. Chesterfield Royal Hospital: Kath Phillis, Rachel Gascoyne. Countess Of Chester Hospital NHS Foundation Trust: Ian London, Jenny Grounds, Emmeline Martin, Susie Paiak. Dartford and Gravesham NHS Trust: Ben Warner, Carmel Stuart, Louise Lacev. Darlington Memorial Hospital: Anjan Dhar, Ellen Brown, Amanda Cowton, Kimberley Stamp. The Dudley Group NHS Foundation Trust: Shanika de Silva, Clare Allcock, Philip Harvey. East and North Hertfordshire NHS Trust: Johanne Brooks, Pearl Baker, Hannah Beadle, Carina Cruz, Debbie Potter. East Lancashire Hospitals NHS Trust: Joe Collum, Farzana Masters. East Suffolk and North Essex NHS Foundation Trust: Achuth Shenoy, Alison O'Kelly. Glangwili Hospital: Aashish Kumar, Samantha Coetzee, Mihaela Peiu. Great Ormond Street Hospital: Edward Gaynor, Sibongile Chadokufa, Bonita Huggett, Hamza Meghari, Sara El-Khouly, Fevronia Kiparissi, Waffa Girshab. Great Western Hospitals NHS Foundation Trust: Andrew Claridge, Emily Fowler, Laura McCafferty. Guy's and St Thomas' NHS Foundation Trust: Peter Irving, Karolina Christodoulides, Angela Clifford, Patrick Dawson, Sailish Honap, Samuel Lim, Raphael Luber, Karina Mahiouz, Susanna Meade, Parizade Raymode, Rebecca Reynolds, Anna Stanton, Sherill Tripoli, Naomi Hare. The Hillingdon Hospitals NHS Foundation Trust: Yih Harn Siaw, Lane Manzano, Jonathan Segal, Ibrahim Al-Bakir, Imran Khakoo. Homerton University Hospital Foundation Trust: Nora Thoua, Katherine Davidson, Jagrul Miah, Alex Hall. Hull University Teaching Hospitals NHS Trust: Shaji Sebastian, Melony Hayes, Sally Myers, Alison Talbot, Jack Turnbull, Emma Whitehead, Katie Stamp, Alison Pattinson, Verghese Mathew, Leanne Sherris. Imperial College Healthcare NHS Trust: Lucy Hicks, Tara-Marie Byrne, Leilani Cabreros, Hannah Downing-Wood, Sophie Hunter, Mohammad Aamir Saifuddin, Hemanth Prabhudev, Sharmili Balarajah. James Paget University Hospitals NHS Foundation Trust: Helen Sutherland. Kettering General Hospital: Ajay M Verma, Juliemol Sebastian, Mohammad Farhad Peerally, King's College Hospital NHS Foundation Trust: Alexandra Kent, Lee Meng Choong, Benedetta Pantaloni, Pantelis Ravdas. King's College Hospital NHS Foundation Trust (paediatric): Babu Vadamalayan. King's Mill Hospital: Stephen Foley, Becky Arnold, Cheryl Heeley, Wayne Lovegrove. Liverpool University Hospitals NHS Foundation Trust: Philip J Smith, Giovanna Bretland, Sarah King, Martina Lofthouse, Lindsey Rigby, Sreedhar Subramanian, David Tyrer, Kate Martin, Christopher Probert. London North West University Healthcare NHS Trust: Nik Kamperidis, Temi Adedoyin, Manisha Baden, Jeannette Brown, Feba Chacko, Michela Cicchetti, Mohammad Saifuddin, Priya Yesupatham. Maidstone and Tunbridge Wells NHS Trust: Rohit Gowda, Maureen Williams. Manchester University NHS Foundation Trust: Karen Kemp, Rima Akhand, Glaxy Gray, Anu John, Maya John, Diamond Sathe, Jennifer Soren. The Mid Yorkshire Hospitals NHS Trust: Michael Sprakes, Julie Burton, Patricia Kane, Stephanie Lupton Milton Keynes University Hospital: George MacFaul, Diane Scaletta, Loria Siamia. Newcastle Hospitals NHS Foundation Trust: Chris Lamb, Mary Doona, Ashleigh Hogg, Lesley Jeffrey, Andrew King, R Alexander Speight. Ninewells Hospital & Medical School: Craig Mowat, Debbie Rice, Susan MacFarlane, Anne MacLeod, Samera Mohammed. Norfolk and Norwich University Hospitals NHS Foundation Trust: Mary Anne Morris, Louise Coke, Grace Hindle, Eirini Kolokouri, Catherine Wright. North Bristol NHS Trust: Melanie Lockett, Charlotte Cranfield, Louise Jennings, Ankur Srivastava, Lana Ward, Nouf Jeynes. North Tyneside General Hospital: Praveen Rajasekhar, Lisa Gallagher, Linda Patterson, Jill Ward, Rae Basnett, Judy Murphy, Lauren Parking, Emma Lawson. Nottingham University Hospitals NHS Trust: David Devadason, Gordon Moran, Neelam Khan, Lauren Tarr. The Pennine Acute Hospitals NHS Trust: Jimmy Limdi, Kay Goulden, Asad Javed, Lauren McKenzie. Portsmouth Hospitals NHS Trust: Pradeep Bhandari, Michelle Baker-Moffatt, Joanne Dash. The Queen Elizabeth Hospital Kings Lynn NHS Trust: Alan Wiles, Hannah Bloxham, Jose Dias, Ellie Graham, Oueen Elizabeth University Hospital, Glasgow: Jonathan Macdonald, Shona Finan, Faye McMeeken, Stephanie Shields, John Paul Seenan. Royal Berkshire NHS Foundation Trust: Des DeSilva, Ofori Boateng, Holly Lawrence, Susanna Malkakorpi. The Royal Bournemouth and Christchurch Hospitals

NHS Foundation Trust: Simon Whiteoak, Kelli Edger-Earley. Royal Cornwall Hospitals NHS Trust: Sarah Ingram, Sharon Botfield, Fiona Hammonds, Clare James. Royal Devon and Exeter NHS Foundation Trust: Tarig Ahmad. Gemma Aspinall. Sarah Hawkins, Suzie Marriott, Clare Redstone, Halina Windak. Royal Free London NHS Foundation Trust: Charles Murray, Cynthia Diaba, Fexy Joseph, Glykeria Pakou. Royal Glamorgan Hospital: James Berrill, Natalie Stroud, Carla Pothecary, Lisa Roche, Keri Turner, Lisa Deering, Lynda Israel. Royal Gwent Hospital: Evelyn Baker, Sean Cutler, Rina Mardania Evans, Maxine Nash. Royal Hampshire County Hospital: John Gordon, Emma Levell, Silvia Zagalo. Royal Hospital for Sick Children, Edinburgh: Richard Russell, Paul Henderson, Margaret Millar. Royal Manchester Children's Hospital: Andrew Fagbemi, Felicia Jennings, Imelda Mayor, Jill Wilson. Royal Surrey County Hospital: Christopher Alexakis, Natalia Michalak, Royal United Hospitals Bath: John Saunders, Helen Burton, Vanessa Cambridge, Tonia Clark, Charlotte Ekblad, Sarah Hierons, Joyce Katebe, Emma Saunsbury, Rachel Perry. The Royal Wolverhampton NHS Trust: Matthew Brookes, Kathryn Davies, Marie Green, Ann Plumbe. Salford Royal NHS Foundation Trust: Clare Ormerod, Helen Christensen, Anne Keen, Jonathan Ogor. Salisbury District Hospital: Alpha Anthony, Emily Newitt. Sandwell and West Birmingham NHS Trust: Edward Fogden, Kalisha Russell. Sheffield Teaching Hospitals NHS Foundation Trust: Anne Phillips, Muaad Abdulla. Shrewsbury and Telford Hospital NHS Trust: Jeff Butterworth, Colene Adams, Elizabeth Buckingham, Danielle Childs, Alison Magness, Jo Stickley, Singleton Hospital: Caradog Thomas, Elaine Brinkworth, Lynda Connor, Amanda Cook, Tabitha Rees. Somerset NHS Foundation Trust: Emma Wesley, Alison Moss. South Tees Hospitals NHS Foundation Trust: Arvind Ramadas, Julie Tregonning. Southend University Hospital NHS Foundation Trust: Ioannis Koumoutsos, Viji George, Swapna Kunhunny, Sophie Laverick. St George's University Hospitals NHS Foundation Trust: Kamal Patel, Mariam Ali, Hilda Mhandu, Aleem Rana, Katherine Spears, Joana Teixeira, Richard Pollok, Mark Mencias, Abigail Seaward. St George's University Hospitals NHS Foundation Trust (paediatric): Nicholas Reps, Rebecca Martin. St James's University Hospital: Christian Selinger, Jenelyn Carbonell, Felicia Onovira, Doris Quartey. Stockport NHS Foundation Trust: Zahid Mahmood, Racheal Campbell, Liane Marsh. Surrey and Sussex Healthcare NHS Trust: Monira Rahman, Sarah Davies, Ruth Habibi, Ellen Jessup-Dunton, Teishel Joefield, Reina Layug. Tameside and Glossop Integrated Care NHS Foundation Trust: Vinod Patel, Joanne Vere. Torbay and South Devon NHS Foundation Trust: Gareth Walker, Stacey Atkins, Jasmine Growdon, Charlotte McNeill. University Hospitals Birmingham NHS Foundation Trust: Rachel Cooney, Lillie Bennett, Louise Bowlas, Sharafaath Shariff. University Hospitals Bristol NHS Foundation Trust: Aileen Fraser, Katherine Belfield. University Hospitals of Derby and Burton NHS Foundation Trust: Said Din, Catherine Addleton, Marie Appleby, Johanna Brown, Kathleen Holding. University Hospitals of Leicester NHS Trust: John deCaestecker, Olivia Watchorn. University Hospitals Plymouth NHS Trust: Chris Hayward, Susan Inniss, Lucy Pritchard. University Hospital Southampton NHS Foundation Trust: Fraser Cummings, Clare Harris, Amy Jones, Liga Krauze, Sohail Rahmany, Audrey Torokwa. United Lincolnshire Hospitals NHS Trust: Jervoise Andrevev, Caroline Hayhurst, Carol Lockwood, Lynn Osborne, Amanda Roper, Karen Warner, Julia Hindle. University College London Hospitals NHS Foundation Trust: Shameer Mehta, James Bell, William Blad, Lisa Whitley. University Hospital Llandough: Durai Dhamaraj, Mark Baker. University Hospital of Wales (paediatric): Amar Wahid, Zoe Morrison. West Hertfordshire Hospitals NHS Trust: Rakesh Chaudhary, Melanie Claridge, Chiara Ellis, Cheryl Kemp, Ogwa Tobi. West Middlesex University Hospital: Emma Johnston. Western General Hospital: Metod Oblak, Richard Appleby. West Suffolk NHS Foundation Trust: Marium Asghar, Charlie Lees, Debbie Alexander, Kate Covil, Lauranne Derikx, Sryros Siakavellas, Helen Baxter, Scott Robertson. Withybush General Hospital: Kerrie Johns, Rachel Hughes, Janet Phipps, Abigail Taylor. Yeovil District Hospital NHS Foundation Trust: Katie Smith, Linda Howard, Dianne Wood. York Teaching Hospital NHS Foundation Trust: Ajay Muddu, Laura Barman, Janine Mallinson. Ysbyty Gwynedd: Iona Thomas, Kelly Andrews, Caroline Mulvaney Jones, Julia Roberts.

Contributors NAK, JRG, CB, SS, NP and TA participated in the conception and design of this study. CB was the project manager and coordinated patient recruitment. RN and TM coordinated all biochemical analyses and central laboratory aspects of the project. SJ conducted the experiments at Roche Diagnostics to define the seroconversion threshold used in this manuscript. NAK, SL, JRG, NC, BH, DC, JRFC, AF, PMI, NK, KBK, CAL, JM, SM, RCP, TR, PJS, AMV, TJM, SS, CWL, NP and TA were involved in the acquisition, analysis or interpretation of data. Data analysis was done by NAK. Drafting of the manuscript was done by NAK, SL, JRG, NC, RN, DC, RCP, SS, CWL, NP, TA. SS, NP and TA obtained the funding for the study. All the authors contributed to the critical review and final approval of the manuscript. NAK and TA have verified the underlying data.

Funding This study was funded by F. Hoffmann-La Roche AG (Switzerland), Biogen GmbH (Switzerland), Celltrion Healthcare (South Korea), Takeda (UK), Galapagos NV (Belgium), Hull University Teaching Hospital NHS Trust, and Royal Devon and Exeter NHS Foundation Trust.

Competing interests NAK reports grants from F. Hoffmann-La Roche AG, grants from Biogen Inc, grants from Celltrion Healthcare, grants from Galapagos NV, non-financial support from Immundiagnostik, during the conduct of the study; grants and non-financial support from AbbVie, grants and personal fees from Celltrion,

personal fees and non-financial support from Janssen, personal fees from Takeda, personal fees and non-financial support from Dr Falk, outside the submitted work. SL reports non-financial support from Pfizer, non-financial support from Ferring, outside the submitted work. JRG reports grants from F. Hoffmann-La Roche AG, grants from Biogen, grants from Celltrion Healthcare, grants from Galapagos NV, non-financial support from Immundiagnostik, during the conduct of the study. DC reports non-financial support from Ferring, personal fees and non-financial support from Pfizer, outside the submitted work. JRFC reports grants and personal fees from Samsung, Pfizer & Biogen; personal fees and non-financial support from Janssen & Abbyie: grants, personal fees and non-financial support from Takeda: personal fees from MSD, Sandoz, Celltrion & NAPP, outside the submitted work. AF reports personal fees from Takeda UK Ltd, personal fees from Dr Falk Pharma, personal fees from Tillotts, personal fees from Abbvie Ltd, personal fees from Sheild, personal fees from Ferring, from Pharmacosmos, personal fees from Allergan, personal fees from Janssen, outside the submitted work. PMI reports grants and personal fees from Takeda, grants from MSD, grants and personal fees from Pfizer, personal fees from Galapagos, personal fees from Gilead, personal fees from Abbvie, personal fees from Janssen, personal fees from Boehringer Ingelheim, personal fees from Topivert, personal fees from VH2, personal fees from Celgene, personal fees from Arena, personal fees from Samsung Bioepis, personal fees from Sandoz, personal fees from Procise, personal fees from Prometheus, outside the submitted work. NK reports personal fees from Janssen, outside the submitted work. KBK reports personal fees from Janssen, personal fees from Takeda, personal fees from PredictImmune, personal fees from Amgen, outside the submitted work. CAL reports grants from Genentech, grants and personal fees from Janssen, grants and personal fees from Takeda, grants from AbbVie, personal fees from Ferring, grants from Eli Lilly, grants from Pfizer, grants from Roche, grants from UCB Biopharma, grants from Sanofi Aventis, grants from Biogen IDEC, grants from Orion OYJ, personal fees from Dr Falk Pharma, grants from AstraZeneca, outside the submitted work. JM reports grants and personal fees from Takeda Pharmaceuticals, grants and personal fees from Biogen, personal fees and non-financial support from AbbVie, personal fees from Grifols, personal fees from Sandoz, personal fees from Celltrion, personal fees and non-financial support from Janssen, personal fees from Vifor Pharmaceuticals, personal fees from Predictimmune, personal fees from Bristol Myers Squibb, non-financial support from Ferring Pharmaceuticals, outside the submitted work. RCGP reports acting as consultant, advisory board member, speaker or recipient of educational grant from Dr Falk, Ferring, Janssen, Pharmacosmos and Takeda. TR reports grants and personal fees from Abbvie, personal fees from BMS, personal fees from Celgene, personal fees from Ferring, personal fees from Gilead, personal fees from GSK, personal fees from LabGenius, personal fees from Janssen, personal fees from Mylan, personal fees from MSD, personal fees from Novartis, personal fees from Pfizer, personal fees from Sandoz, personal fees from Takeda, personal fees from Galapagos, personal fees from Arena, outside the submitted work. PJS reports speaker fees and advisory board sponsorship from Janssen, Celltrion and Takeda outside the submitted work. AMV reports personal fees and non-financial support from Takeda, personal fees and non-financial support from Celltrion, personal fees and non-financial support from Merck Sharp & Dohme, outside the submitted work. SJ is an employee of Roche Diagnostics and holds Roche shares. SS reports grants from Takeda, Abbvie, AMGEN, Tillots Pharma, personal fees from Jaansen, Takeda, Galapagos, Celltrion, Falk Pharma, Tillots pharma, Cellgene, Pfizer, Pharmacocosmos, outside the submitted work. CWL reports personal fees from Abbvie, personal fees from Janssen, personal fees from Pfizer, personal fees from Takeda, grants from Gilead, personal fees from Gilead, personal fees from Galapagos, personal fees from Iterative Scopes, personal fees from Trellus Health, personal fees from Celltion, personal fees from Ferring, personal fees from BMS, during the conduct of the study. NP reports personal fees from Takeda, personal fees from Janssen, personal fees from Pfizer, personal fees from Bristol-Myers Squibb, personal fees from Abbvie, personal fees from Roche, personal fees from Lilly, personal fees from Allergan, personal fees from Celgene, outside the submitted work: and NP has served as a speaker/advisory board member for Abbvie, Allergan, Bristol Myers Squibb, Celgene, Falk, Ferring, Janssen, Pfizer, Tillotts, Takeda and Vifor Pharma. TA reports grants and non-financial support from F. Hoffmann-La Roche AG, grants from Biogen Inc, grants from Celltrion Healthcare, grants from Galapagos NV, non-financial support from Immundiagnostik, during the conduct of the study; personal fees from Biogen, grants and personal fees from Celltrion Healthcare, personal fees and non-financial support from Immundiagnostik, personal fees from Takeda, personal fees from ARENA, personal fees from Gilead, personal fees from Adcock Ingram Healthcare, personal fees from Pfizer, personal fees from Genentech, non-financial support from Tillotts, outside the submitted work.

Patient and public involvement statement We conducted an electronic survey to gauge the opinion of patients with IBD on the patient questionnaires to be delivered as part of the CLARITY IBD study. We surveyed 250 patients across 74 hospitals. All our proposed questions for study inclusion were rated as important or very important by at least 83% of participants. The Exeter IBD Patient Panel refined the questions included in the study questionnaire, reviewed the study protocol, supported the writing of the patient information sheets, and participated in testing of the electronic consent form and patient questionnaire. A member of the Exeter

IBD Patient Panel sits on the study management committee, ensuring patient involvement in all aspects of study delivery, data analysis and dissemination of findings.

Patient consent for publication Not required.

Ethics approval The Surrey Borders Research Ethics committee approved the study (REC reference: REC 20/HRA/3114) in September 2020.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The study protocol including the statistical analysis plan is available at www.clarityibd. org. Individual participant deidentified data that underlie the results reported in this article will be available immediately after publication for a period of 5 years. The data will be made available to investigators whose proposed use of the data has been approved by an independent review committee. Analyses will be restricted to the aims in the approved proposal. Proposals should be directed to tariq.ahmad1@ nhs.net. To gain access data requestors will need to sign a data access agreement.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iDs

Nicholas A Kennedy http://orcid.org/0000-0003-4368-1961 Simeng Lin http://orcid.org/0000-0002-4201-4879 James R Goodhand http://orcid.org/0000-0003-3112-376X Neil Chanchlani http://orcid.org/0000-0003-0207-6706 Benjamin Hamilton http://orcid.org/0000-0002-0349-278X Claire Bewshea http://orcid.org/0000-0002-0965-9587 Desmond Chee http://orcid.org/0000-0003-3473-9447 JR Fraser Cummings http://orcid.org/0000-0002-9659-3247 Aileen Fraser http://orcid.org/0000-0001-6462-5091 Peter M Irving http://orcid.org/0000-0003-0972-8148 Klaartje B Kok http://orcid.org/0000-0002-5942-9149 Christopher Andrew Lamb http://orcid.org/0000-0002-7271-4956 Jonathan Macdonald http://orcid.org/0000-0003-0249-107X Shameer Mehta http://orcid.org/0000-0002-7002-293X Richard CG Pollok http://orcid.org/0000-0001-6452-6763 Tim Raine http://orcid.org/0000-0002-5855-9873 Philip J Smith http://orcid.org/0000-0003-1568-3978 Ajay Mark Verma http://orcid.org/0000-0002-6432-3357 Timothy J McDonald http://orcid.org/0000-0003-3559-6660 Shaji Sebastian http://orcid.org/0000-0002-3670-6545 Charlie W Lees http://orcid.org/0000-0002-0732-8215 Nick Powell http://orcid.org/0000-0003-3231-6950 Tariq Ahmad http://orcid.org/0000-0002-6058-5528

REFERENCES

- Joint Committee on Vaccination and Immunisation. Optimising the COVID-19 vaccination programme for maximum short-term impact, 2021. Available: https:// www.gov.uk/government/publications/prioritising-the-first-covid-19-vaccine-dose-jcvistatement/optimising-the-covid-19-vaccination-programme-for-maximum-short-termimpact [Accessed 20 Mar 2021].
- 2 European Centre for Disease Prevention and Control. Overview of the implementation of COVID-19 vaccination strategies and vaccine deployment plans in the EU/EEA, 2021.
- 3 National Advisory Committee on Immunization. COVID-19 vaccine extended dose interval for Canadians, 2021. Available: https://www.canada.ca/en/public-health/ services/immunization/national-advisory-committee-on-immunization-naci/rapidresponse-extended-dose-intervals-covid-19-vaccines-early-rollout-populationprotection.html [Accessed 22 Mar 2021].

- 4 Jones G-R, Lyons M, Plevris N, et al. Ibd prevalence in Lothian, Scotland, derived by capture-recapture methodology. *Gut* 2019;68:1953–60.
- 5 Hamilton B, Green H, Heerasing N, et al. Incidence and prevalence of inflammatory bowel disease in Devon, UK. Frontline Gastroenterol 2020;68:flgastro-2019-101369.
- 6 Melmed GY, Agarwal N, Frenck RW, et al. Immunosuppression impairs response to pneumococcal polysaccharide vaccination in patients with inflammatory bowel disease. Am J Gastroenterol 2010;105:148–54.
- 7 Caldera F, Hillman L, Saha S, *et al.* Immunogenicity of high dose influenza vaccine for patients with inflammatory bowel disease on anti-TNF monotherapy: a randomized clinical trial. *Inflamm Bowel Dis* 2020;26:593–602.
- 8 Pratt PK, David N, Weber HC, et al. Antibody response to hepatitis B virus vaccine is impaired in patients with inflammatory bowel disease on infliximab therapy. Inflamm Bowel Dis 2018;24:380–6.
- 9 Long MD, Martin C, Sandler RS, *et al*. Increased risk of pneumonia among patients with inflammatory bowel disease. *Am J Gastroenterol* 2013;108:240–8.
- 10 Harrington JE, Hamilton RE, Ganley-Leal L, et al. The immunogenicity of the influenza, pneumococcal, and hepatitis B vaccines in patients with inflammatory bowel disease treated with Vedolizumab. Crohns Colitis 360 2020;2:otaa082.
- 11 Kennedy NA, Goodhand JR, Bewshea C, et al. Anti-SARS-CoV-2 antibody responses are attenuated in patients with IBD treated with infliximab. Gut 2021;70:865–75.
- 12 Harris PA, Taylor R, Minor BL, et al. The REDCap Consortium: building an international community of software platform partners. J Biomed Inform 2019;95:103208.
- 13 Roche Diagnostics GmbH. Elecsys® Anti-SARS-CoV-2 S assay method sheet, 2020. Available: https://diagnostics.roche.com/gb/en/products/params/elecsys-anti-sars-cov-2-s.html [Accessed 24 Mar 2021].
- 14 Muench P, Jochum S, Wenderoth V, et al. Development and validation of the elecsys anti-SARS-CoV-2 immunoassay as a highly specific tool for determining past exposure to SARS-CoV-2. J Clin Microbiol 2020;58:1694–714.
- 15 GenScript. SARS-CoV-2 surrogate virus neutralization test (sVNT) kit (RUO), 2021. Available: https://www.genscript.com/covid-19-detection-svnt.html [Accessed 5 Apr 2021].
- 16 Tan CW, Chia WN, Qin X, et al. A SARS-CoV-2 surrogate virus neutralization test based on antibody-mediated blockage of ACE2-spike protein-protein interaction. Nat Biotechnol 2020;38:1073–8.
- 17 Alexander JL, Moran GW, Gaya DR, *et al.* SARS-CoV-2 vaccination for patients with inflammatory bowel disease: a British Society of gastroenterology inflammatory bowel disease section and IBD clinical research Group position statement. *Lancet Gastroenterol Hepatol* 2021;6:218–24.
- 18 Boyarsky BJ, Werbel WA, Avery RK, et al. Immunogenicity of a single dose of SARS-CoV-2 messenger RNA vaccine in solid organ transplant recipients. JAMA 2021:e214385.
- 19 Monin-Aldama L, Laing AG, McKenzie DR. Interim results of the safety and immuneefficacy of 1 versus 2 doses of COVID-19 vaccine BNT162b2 for cancer patients in the context of the UK vaccine priority guidelines. *medRxiv*2021:2021.03.17.21253131.
- 20 Choi B, Choudhary MC, Regan J, et al. Persistence and evolution of SARS-CoV-2 in an immunocompromised host. N Engl J Med 2020;383:2291–3.
- 21 Avanzato VA, Matson MJ, Seifert SN, *et al*. Case study: prolonged infectious SARS-CoV-2 shedding from an asymptomatic immunocompromised individual with cancer. *Cell* 2020;183:1901–12.
- 22 Winter AP, Follett EA, McIntyre J, et al. Influence of smoking on immunological responses to hepatitis B vaccine. Vaccine 1994;12:771–2.
- 23 MacKenzie JS, MacKenzie IH, Holt PG. The effect of cigarette smoking on susceptibility to epidemic influenza and on serological responses to live attenuated and killed subunit influenza vaccines. J Hyg 1976;77:409–17.
- 24 Haralambieva IH, Ovsyannikova IG, Umlauf BJ, et al. Genetic polymorphisms in host antiviral genes: associations with humoral and cellular immunity to measles vaccine. Vaccine 2011;29:8988–97.
- 25 Huda MN, Lewis Z, Kalanetra KM, et al. Stool microbiota and vaccine responses of infants. Pediatrics 2014;134:e362–72.
- 26 Savy M, Edmond K, Fine PEM, et al. Landscape analysis of interactions between nutrition and vaccine responses in children. J Nutr 2009;139:2154S–218.
- 27 Pasparakis M, Alexopoulou L, Episkopou V, et al. Immune and inflammatory responses in TNF alpha-deficient mice: a critical requirement for TNF alpha in the formation of primary B cell follicles, follicular dendritic cell networks and germinal centers, and in the maturation of the humoral immune response. J Exp Med 1996;184:1397–411.
- 28 Ritter U, Meissner A, Ott J, et al. Analysis of the maturation process of dendritic cells deficient for TNF and lymphotoxin-alpha reveals an essential role for TNF. J Leukoc Biol 2003;74:216–22.
- 29 Marot S, Malet I, Leducq V, et al. Rapid decline of neutralizing antibodies against SARS-CoV-2 among infected healthcare workers. Nat Commun 2021;12:1–7.
- 30 Edara VV, Hudson WH, Xie X, et al. Neutralizing antibodies against SARS-CoV-2 variants after infection and vaccination. JAMA 2021. doi:10.1001/jama.2021.4388. [Epub ahead of print: 19 Mar 2021].