

# Investigation of Antioxidant, Hypoglycemic and Anti-Obesity Effects of *Euphorbia Resinifera* L.

Kaoutar Benrahou<sup>1\*</sup>, Otman El Guourami<sup>2</sup>, Hanae Naceiri Mrabti<sup>1</sup>, Yahia Cherrah<sup>1</sup>, My El Abbes Faouzi<sup>1</sup>

<sup>1</sup>Laboratory of Pharmacology and Toxicology, Bio Pharmaceutical and Toxicological Analyzes Research Team, Faculty of Medicine and Pharmacy, Mohammed V University, Rabat, Morocco

<sup>2</sup>Laboratory of Analytical Chemistry and Bromatology, Faculty of Medicine and Pharmacy, Mohammed V University, Rabat, Morocco

**Received** February 25, 2022

**Reviewed** May 22, 2022

**Accepted** June 16, 2022

## \*Corresponding Author

Kaoutar Benrahou  
Laboratory of Pharmacology and  
Toxicology, Bio Pharmaceutical and  
Toxicological Analyzes Research Team,  
Faculty of Medicine and Pharmacy,  
Mohammed V University, BP 6203,  
Rabat 10100, Morocco  
Tel: +212-682-577-700  
E-mail: benrahoukaoutar@gmail.com

**Objectives:** The aim of this work is to evaluate the in vitro antioxidant, hypoglycemic, and antiobesity effects of *Euphorbia resinifera* extracts and investigate the phenolic constituents and the toxicity of these extracts.

**Methods:** Phytochemical screening was performed to detect polyphenols and flavonoids. Antioxidant activity was evaluated by four methods (DPPH, ABTS, H<sub>2</sub>O<sub>2</sub>, and xanthine oxidase inhibition). The hypoglycemic effect was determined by the inhibition of  $\alpha$ -amylase and  $\alpha$ -glucosidase enzymes in vitro and via a starch tolerance study in normal rats. The antiobesity effect was estimated by in vitro inhibition of lipase.

**Results:** Phytochemical screening revealed that the ethanolic extract was rich in polyphenols (99  $\pm$  0.56 mg GEA/g extract) and tannins (55.22  $\pm$  0.17 mg RE/g extract). Moreover, this extract showed higher antioxidant activity in different tests: the DPPH assay (IC<sub>50</sub> = 53.81  $\pm$  1.83  $\mu$ g/mL), ABTS assay (111.4  $\pm$  2.64 mg TE/g extract), H<sub>2</sub>O<sub>2</sub> (IC<sub>50</sub> = 98.15  $\pm$  0.68  $\mu$ g/mL), and xanthine oxidase (IC<sub>50</sub> = 10.26  $\pm$  0.6  $\mu$ g/mL). With respect to hypoglycemic effect, the aqueous and ethanolic extracts showed IC<sub>50</sub> values of 119.7  $\pm$  2.15  $\mu$ g/mL and 102  $\pm$  3.63  $\mu$ g/mL for  $\alpha$ -amylase and 121.4  $\pm$  1.88 and 56.6  $\pm$  1.12  $\mu$ g/mL for  $\alpha$ -glucosidase, respectively, and the extracts lowered blood glucose levels in normal starch-loaded rats. Additionally, lipase inhibition was observed with aqueous (IC<sub>50</sub> = 25.3  $\pm$  1.53  $\mu$ g/mL) and ethanolic (IC<sub>50</sub> = 13.7  $\pm$  3.03  $\mu$ g/mL) extracts.

**Conclusion:** These findings show the antioxidant, hypoglycemic, and hyperlipidemic effects of *E. resinifera* extracts, which should be investigated further to validate their medicinal uses and their pharmaceutical applications.

**Keywords:** euphorbia resinifera, antioxidant activity, hypoglycemic effects, enzyme inhibitory

## INTRODUCTION

Diabetes mellitus, obesity, and oxidative stress are metabolic diseases having long-term effects [1]. Due to modern lifestyles and the increased consumption of high-fat and high-carbohydrate foods, the global prevalence of diabetes and obesity has increased dramatically. Hyperglycemia is characterized by abnormally high plasma glucose levels. In type 2 diabetes, insulin resistance can be caused by numerous signal transduction defects, including defects in insulin receptors or glucose transporters. High blood glucose levels will increase free radical

production while reducing endogenous antioxidant levels [2]. Obesity is characterized by a large amount of fat preserved in adipose tissue, resulting in weight gain [3]. One strategy to control overweight and hyperglycemia is to inhibit the absorption of dietary carbohydrates and fats. The inhibition of digestive enzymes such as  $\alpha$ -amylase and pancreatic lipase responsible for the degradation of complex carbohydrates and lipids into simple molecules is one of the most well-studied therapeutic actions to evaluate the antiobesity and antidiabetic effects of natural products [4].

Since ancient times, medicinal plants have been used for the

treatment of diabetes, especially in developing countries. Currently, scientific research is focused on the control of diabetes [5]. Remedies based on natural products have been found to be more active and cause less adverse effects compared to oral synthetic products [6]. Plants are rich in phenolic compounds that possess insulin-like activity and exert hypoglycemic effects by inhibiting digestive enzymes and lipid peroxidation at the cellular level. These inhibitory effects are associated with their antioxidant activity [7].

*Euphorbia* is among the most important genera of the family Euphorbiaceae, with approximately 2,000 species identified worldwide, of which 80 species are found in China. *Euphorbia* plants are characterized by latex that is irritating, and several species are traditionally used for the treatment of skin diseases, edema, and tuberculosis [8]. Among these species, *Euphorbia resinifera* is one of the oldest “drugs” of the Western medical tradition, frequently used by Moroccan herbalists and therapists. The Arabic name of *E. resinifera* is “Zagoume,” a species endemic to Morocco generally distributed in the center of the country, in the regions of Azilal and Beni Mellal (Middle Atlas), with some scattered populations in the High Atlas Mountains and the Anti-Atlas [9]. Studies have reported that it contains triterpenoids, diterpenoids, and phenolic acids among other constituents. Moreover, euphane triterpenes, tirucallane triterpenes, tirucallane-type spirotriterpenoids and nortriterpenes have been identified from the latex [10]. In this study, the in vitro antioxidant, hypoglycemic, and antiobesity effects of *Euphorbia resinifera* extracts were evaluated and the acute toxicity and phenolic content of these extracts investigated.

## MATERIALS AND METHODS

### 1. Plant material and extraction

The aerial part of *E. resinifera* was collected from the regions of Beni-Mellal, Morocco (geographic coordinates: 32°12'16.6"33'36.6" W). The specimens of *E. resinifera* have been deposited in herbarium HUMPO at University Mohammed First, Oujda, Morocco, under the number HUMPOM 10052. The aerial part of *E. resinifera* was dried at room temperature, powdered, and stored in a dark and humid place until use.

The aqueous extract was prepared by infusion, wherein 30 g of *E. resinifera* was mixed with 300 mL of distilled water for 1 hour. The mixture was filtered and evaporated at 50°C using

a rotary evaporator. The extract was finally freeze-dried to remove traces of water and stored for further use. To prepare the ethanolic extract, 30 g of the plant material was macerated for 48 hours under stirring and at room temperature. The extract was filtered and evaporated at 40°C using a rotary evaporator.

### 2. Determination of phenolic content

The phenolic content was determined by the method described by Spanos and Wrolstad [11], and the flavonoid content according to the method of Dewanto et al. [12].

### 3. Antioxidant activity

#### 1) DPPH, ABTS, H<sub>2</sub>O<sub>2</sub>, and xanthine oxidase assays

The antioxidant capacity by the DPPH method was determined according to the protocol described by Huang et al. [13], the ABTS test by the protocol of Tuberoso et al. [14] H<sub>2</sub>O<sub>2</sub> by Muruhan et al. [15], and xanthine oxidase by Umamaheswari et al. [16].

### 4. Enzyme inhibitory activities

#### 1) $\alpha$ -Amylase, $\alpha$ -glucosidase, and lipase inhibition tests

The method described by Chakrabarti et al. [17] was used to determine the percentage inhibition of  $\alpha$ -amylase, the protocol described by Kee et al. [18] for the determination of  $\alpha$ -glucosidase, and the protocol described by Hu et al. [19] for the inhibition of lipase.

#### 2) Oral starch tolerance in normal rats

The evaluation of oral starch tolerance in normal rats was performed according to the protocol described by Beejmohun et al. [20].

### 5. HR-MS analysis

Analysis was performed by using electrospray ionization (ESI) mass spectrometry (MS). The samples were dissolved in methanol to a final concentration of 1-2 pmol/ $\mu$ L. All compounds were then measured in negative and positive modes by total mass scanning ( $m/z$  50-1,000) using a Thermo Scientific Orbitrap Exactive mass spectrometer with a heated ESI source (HESI-II). Mass spectra were collected at a resolution of 100,000. The instrument parameters were as follows: sheath gas

10 in positive mode and 20 in negative mode (arbitrary units), sputtering voltage 3.5 kV in positive mode (3 kV in negative), capillary temperature 275°C. Data processing was performed using the associated software Xcalibur 2.2 and Exactive 1.1.

## 6. Acute toxicity

Acute oral toxicity was investigated according to the method described in OECD-423 [21]. Swiss albino mice weighing 25–35 g were used in this study. The groups received the extracts orally at the dose of 2 g/kg. After administration, the animals were observed for 14 days to assess the toxic and behavioral effects.

# RESULTS

## 1. Phytochemical analysis

The results of the phytochemical analysis are summarized in Table 1; the ethanolic extract was found to be richer in polyphenols and flavonoids than the aqueous extract. The TPC and TFC were  $99 \pm 0.56$  mg GEA/g extract and  $55.22 \pm 0.17$  mg RE/g extract in the ethanolic extract and  $89.31 \pm 0.42$  and  $35.24 \pm 0.89$  mg RE/g extract in the aqueous extract, respectively.

## 2. Antioxidant activity

### 1) DPPH, ABTS, H<sub>2</sub>O<sub>2</sub>, and xanthine oxidase activity

The results of the antioxidant activity of the aqueous and ethanolic extracts using the DPPH, ABTS, H<sub>2</sub>O<sub>2</sub>, and xanthine

oxidase methods are summarized in Table 2. Indeed, the ethanolic extract showed higher antioxidant activity in all the methods. In the DPPH assay, both the aqueous and ethanolic extracts showed inhibitory activity with  $IC_{50} = 149 \pm 1.55$  µg/mL and  $IC_{50} = 53.81 \pm 1.83$  µg/mL, respectively. Similarly, the ABTS<sup>+</sup> radical-scavenging activity of the extracts revealed that the aqueous and ethanolic extracts exhibited inhibitory values of  $86.6 \pm 1.23$  mg TE/g extract and  $111.4 \pm 2.64$  mg TE/g extract, respectively. The inhibitory effects of *E. resinifera* extracts on H<sub>2</sub>O<sub>2</sub> were evidenced by  $IC_{50} = 98.15 \pm 0.68$  µg/mL for the ethanolic extract compared with  $IC_{50} = 224.6 \pm 0.7$  µg/mL for the aqueous extract. Regarding the inhibition of xanthine oxidase, the ethanolic extract showed a significantly greater inhibitory effect ( $IC_{50} = 10.26 \pm 0.6$  µg/mL) than the aqueous extract ( $IC_{50} = 69.83 \pm 1$  µg/mL).

## 3. Enzyme inhibitory activity

### 1) α-Amylase, α-glucosidase, and lipase inhibition

The results of in vitro antihyperglycemic activity against α-amylase and α-glucosidase enzymes are summarized in Table 3. Both extracts possessed inhibitory activity against α-amylase and α-glucosidase, showing higher activity than acarbose (used as a reference compound), which showed inhibitory values of  $IC_{50} = 44.75 \pm 0.54$  µg/mL and  $IC_{50} = 18.01 \pm 2.00$  µg/mL against α-amylase and α-glucosidase, respectively. The ethanolic extract showed an  $IC_{50}$  of  $102 \pm 3.63$  µg/mL against α-amylase and an  $IC_{50}$  of  $56.6 \pm 1.12$  µg/mL against α-glucosidase. The aqueous extract had lower inhibitory activity, with an  $IC_{50}$  of  $119.7 \pm 2.15$  µg/mL against α-amylase and an  $IC_{50}$  of  $121.4 \pm$

**Table 1.** Total phenols and flavonoids content of *E. resinifera*

	Aqueous extract		Ethanol extract	
	TPC (mg GEA/g extract)	TFC (mg RE/g extract)	TPC (mg GEA/g extract)	TFC (mg RE/g extract)
<i>Euphorbia resinifera</i>	$89.31 \pm 0.42$	$35.24 \pm 0.89$	$99 \pm 0.56$	$55.22 \pm 0.17$

**Table 2.** Antioxidant activity by DPPH, ABTS, H<sub>2</sub>O<sub>2</sub> and xanthine oxidase (XO) methods of *E. resinifera* extracts


	DPPH (IC <sub>50</sub> )	ABTS (mg TE/g extract)	H <sub>2</sub> O <sub>2</sub> (IC <sub>50</sub> )	Xanthine oxidase (IC <sub>50</sub> )
Aqueous extract	$149 \pm 1.55$	$86.6 \pm 1.23$	$224.6 \pm 0.7$	$69.83 \pm 1$
Ethanol extract	$53.81 \pm 1.83$	$111.4 \pm 2.64$	$98.15 \pm 0.68$	$10.26 \pm 0.6$
BHT	$3.28 \pm 0.79$	-	-	-
Ascorbic acid	-	-	$5.98 \pm 0.47$	-
Allopurinol	-	-	-	$0.78 \pm 0.01$

mg TE/g extract: mg Trolox equivalent per gram of extract, IC<sub>50</sub>: µg/mL.

**Table 3.** IC<sub>50</sub> values of *E. resinifera* extract on  $\alpha$ -amylase,  $\alpha$ -glucosidase and lipase inhibition

	IC <sub>50</sub> ( $\mu$ g/mL)		
	$\alpha$ -amylase	$\alpha$ -glucosidase	Lipase
ERA	119.7 $\pm$ 2.15	121.4 $\pm$ 1.88	25.3 $\pm$ 1.53
ERE	102 $\pm$ 3.63	56.6 $\pm$ 1.12	13.7 $\pm$ 3.03
Acarbose	44.75 $\pm$ 0.54	18.01 $\pm$ 2.00	-
Orlistat	-	-	12.55 $\pm$ 4.17

ERA, aqueous extract of *E. resinifera*; ERE, ethanolic extract of *E. resinifera*.



**Figure 1.** Effect of *Euphorbia resinifera* on blood glucose after starch loading in normal rats (A) and with presentation in the area under curve (B). Values are means  $\pm$  SEM (n = 5). Ns, not significant to the normal controls; ERA, aqueous extract of *E. resinifera*; ERE, ethanolic extract of *E. resinifera*; AUC, area under the curve.

1.88  $\mu$ g/mL against  $\alpha$ -glucosidase. In addition, the ethanolic extract displayed an IC<sub>50</sub> = 13.7  $\pm$  3.03  $\mu$ g/mL against lipase, which was higher than that of the aqueous extract.

## 2) Oral starch tolerance in normal rats

The results of the effects of *E. resinifera* extracts in an oral starch tolerance test in normal rats are shown in Fig. 1. Acarbose and ERA decreased blood glucose after 30 minutes; however, ERE reduced blood glucose only after 60 minutes at 1.07 g/L. However, at 30 minutes, acarbose and ERA ( $p > 0.05$ ) reduced blood glucose to 0.93 and 0.81 g/L, respectively, while ERA ( $p > 0.05$ ) reduced blood glucose to 0.88 g/L only after 120 min. Similarly, ERA reduced blood glucose compared with acarbose and control.

## 3) HR-MS analysis

ESI-HRMS analysis was able to identify saponarin in the aqueous extract with a molecular formula C<sub>27</sub>H<sub>29</sub>O<sub>15</sub>, m/z 593.15, and an RDB unsaturation value of 13.5 (Fig. 2).


## 4) Acute toxicity

The oral administration of aqueous and ethanolic extracts of *E. resinifera* at a dose of 2 g/kg showed no mortality in the treated mice. Further, the treatment with the two extracts did not induce any behavioral disorder during the 14 days of observation. Therefore, we concluded that the oral LD<sub>50</sub> of *E. resinifera* is greater than 2,000 mg/kg.

## DISCUSSION

The phytochemical analysis revealed that the extracts of *E. resinifera*, particularly the ethanolic extract, are rich in polyphenols and flavonoids. However, a study by Farah et al. [22] on the same species reported lower content of polyphenols and flavonoids than that obtained in our study. The methanolic extract of the stems showed TPC of 94.83  $\pm$  1,027  $\mu$ g GAE/g extract and TFC of 39.14  $\pm$  0.48  $\mu$ g RE/g extract. These results can be influenced by several factors such as the type of extraction, the solvent used, the extraction time, the extraction temperature, the particle size of the plant materials, the solvent/solid ratio, the environment, and the harvest period [23]. Therefore, ex-

Euphorbia Resinifera extrait aqueux neg mode full scan 300-1000 amu 240821 #2-25 RT: 0.02-0.47 AV: 24 NL: 4.09E4  
T: FTMS - p ESI Full ms [300.00-1000.00]



**Figure 2.** HRMS mass spectrum obtained in negative mode for Saponarin in aqueous extract.

traction methods such as infusion and maceration are popular methods used in traditional herbal preparations due to their short times, simplicity, and high yields of bioactive compounds.

The study of the antioxidant activity of the *E. resinifera* extracts by four methods (DPPH, ABTS,  $H_2O_2$ , and xanthine oxidase) showed that the two extracts have both antioxidant and antiradical effects. The wavelength absorption of the ABTS<sup>+</sup> at 734 nm eliminated color interference by the extracts [24].  $H_2O_2$  is an oxidant, weakening the activity of enzymes by the oxidation of thiol groups (-SH); it can also react with  $Fe^{2+}$  and or  $Cu^{2+}$  to form a hydroxyl radical, causing cellular poisoning. Therefore, the inactivation of hydrogen peroxide is essential [25]. Compared to another study performed on the same species, the ethanolic extract in our study showed higher antioxidant activity against  $H_2O_2$  than that reported ( $IC_{50} = 65.01 \pm 0.32$   $\mu$ g/mL) for DPPH [26]. This may be due to many factors such as the presence of several compounds or the synergistic effect

of bioactive compounds and the position of hydroxyl groups and binding of phenolic compounds [27]. Indeed, the estimated phenolic content indicates that there may be a correlation of phenolic constituents with this antioxidant activity.

*E. resinifera* aqueous and ethanolic extracts were also tested for their inhibitory activity against the enzymes  $\alpha$ -amylase,  $\alpha$ -glucosidase, and lipase. Both extracts exhibited an inhibitory effect against  $\alpha$ -amylase,  $\alpha$ -glucosidase, and lipase, with higher activity obtained with the ethanolic extract. Additionally, extracts of *E. resinifera* lowered blood sugar in normal starch-loaded rats. These effects could be attributed to the phytochemical constituents in the plant. Ameer et al. [28] and Birari and Bhutani [29] showed that polyphenols, terpenoids, and their derivatives have great promise as antidiabetic and antiobesity agents. Indeed, it has been reported that plants acting on digestive enzymes such as  $\alpha$ -amylase and  $\alpha$ -glucosidase, that react with proteins, blocking enzymatic activity [3], or that act on

hepatic enzymes by stimulating glycogenesis or inhibiting glycogenolysis, can also inhibit glucose transporters at the level of the intestinal barrier to limit the intestinal absorption of glucose [30]. However natural products can also inhibit the pancreatic lipase responsible for the degradation of triglycerides by physically blocking the interaction between triacylglycerol (in the oily phase) and the enzyme (in the aqueous phase) at the water-oil interface. Binding of a lipase substrate and its inhibitor will prevent the substrate from entering the active site of the lipase [31]. However, our results suggest that the *E. resinifera* extracts inhibit enzymes ( $\alpha$ -amylase,  $\alpha$ -glucosidase, and lipase) and glucose transporters. Similarly, our study showed the high levels of phenolic compounds in *E. resinifera* extracts with a major proportion consisting of terpenoids [8, 32]. Terpene compounds such as carvacrol, thymol,  $\alpha$ -pinene,  $\beta$ -pinene, and  $\alpha$ -terpineol have shown antioxidant power and effects on enzyme expression [33, 34]. In addition, certain terpene derivatives have been identified and isolated from plants and their mechanisms of action elucidated. These terpene derivatives have been used as activators to facilitate permeability and transdermal absorption of drugs [35]. Indeed, Ćavar Zeljković et al. [36] reported that the phenolic and volatile terpenoid content is closely related to the phenological stage of the plant. Similarly, studies have reported the antioxidant and antidiabetic activities of natural products [37, 38]. Thus, bioactive compounds isolated from natural products offer great promise for developing new therapies.

ESI-HRMS analysis identified the saponarin compound in the aqueous extract but not the terpenoid compounds cited by other studies. Simeonova et al. [39] and Sengupta et al. [40] have shown strong antioxidant and antidiabetic effects of the saponarin molecule. Treatment with aqueous and ethanolic extracts of *E. resinifera* for 14 days showed no signs of intoxication or weight loss. Therefore, we concluded that the oral LD<sub>50</sub> of *E. resinifera* is greater than 2,000 mg/kg.

## CONCLUSION

*Euphorbia resinifera* was found to be rich in phenolic compounds, possessing antioxidant, antiobesity, and hypoglycemic activities. Indeed, the aqueous and ethanolic extracts showed antioxidant activity in DPPH and ABTS assays and against xanthine oxidase enzyme and H<sub>2</sub>O<sub>2</sub>. Similarly, the extracts inhibited digestive enzymes ( $\alpha$ -amylase,  $\alpha$ -glucosidase, and lipase) and improved postprandial glycemia in normal rats. However, further antioxidant and hypoglycemic studies must be performed

in vivo to confirm these effects.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## AUTHORS' CONTRIBUTION

Authors are expected to present author contributions statement to their manuscript such as; Kaoutar Benrahou: Investigation, Resources, and Writing – original draft. Otman El Guourami: Methodology, Supervision, and Validation. Hanae Naceiri Mrabti: Visualization, Software, Formal Analysis. Yahia Cherrah: Software, Formal Analysis. My El Abbes Faouzi: Software, Formal Analysis.

## ORCID

Kaoutar Benrahou, <https://orcid.org/0000-0002-1391-6232>

Otman El Guourami, <https://orcid.org/0000-0002-5315-6063>

Hanae Naceiri Mrabti, <https://orcid.org/0000-0002-8145-2572>

Yahia Cherrah, <https://orcid.org/0000-0001-7890-9028>

My El Abbes Faouzi, <https://orcid.org/0000-0003-3863-4677>

## REFERENCES

- Noor ZI, Ahmed D, Rehman HM, Qamar MT, Froeyen M, Ahmad S, et al. In vitro antidiabetic, anti-obesity and antioxidant analysis of *Ocimum basilicum* aerial biomass and in silico molecular docking simulations with alpha-amylase and lipase enzymes. *Biology (Basel)*. 2019;8(4):92.
- Ajiboye BO, Ojo OA, Okesola MA, Akinyemi AJ, Talabi JY, Idowu OT, et al. In vitro antioxidant activities and inhibitory effects of phenolic extract of *Senecio bialbrae* (Oliv and Hiern) against key enzymes linked with type II diabetes mellitus and Alzheimer's disease. *Food Sci Nutr*. 2018;6(7):1803-10.
- Balasubramaniam V, Mustar S, Mustafa Khalid N, Abd Rashed A, Mohd Noh MF, Wilcox MD, et al. Inhibitory activities of three Malaysian edible seaweeds on lipase and  $\alpha$ -amylase. *J Appl Phycol*. 2013;25(5):1405-12.
- Worsztynowicz P, Napierała M, Białas W, Grajek W, Olkowicz

- M. Pancreatic  $\alpha$ -amylase and lipase inhibitory activity of polyphenolic compounds present in the extract of black chokeberry (*Aronia melanocarpa* L.). *Process Biochem.* 2014;49(9):1457-63.
5. Oboh G, Akinyemi AJ, Ademiluyi AO, Adefegha SA. Inhibitory effects of aqueous extracts of two varieties of ginger on some key enzymes linked to type-2 diabetes in vitro. *J Food Nutr Res.* 2010;49(1):14-20.
  6. Poongothai K, Ponmurugan P, Ahmed KS, Kumar BS, Sheriff SA. Antihyperglycemic and antioxidant effects of *Solanum xanthocarpum* leaves (field grown & in vitro raised) extracts on alloxan induced diabetic rats. *Asian Pac J Trop Med.* 2011;4(10):778-85.
  7. Nair SS, Kavrekar V, Mishra A. In vitro studies on alpha amylase and alpha glucosidase inhibitory activities of selected plant extracts. *Eur J Exp Biol.* 2013;3(1):128-32.
  8. Wang S, Liang H, Zhao Y, Wang G, Yao H, Kasimu R, et al. New triterpenoids from the latex of *Euphorbia resinifera* Berg. *Fitoterapia.* 2016;108:33-40.
  9. Boutoub O, El-Guendouz S, Estevinho LM, Paula VB, Aazza S, El Ghadraoui L, et al. Antioxidant activity and enzyme inhibitory potential of *Euphorbia resinifera* and *E. officinarum* honeys from Morocco and plant aqueous extracts. *Environ Sci Pollut Res Int.* 2021;28(1):503-17.
  10. Wang SY, Li GY, Zhang K, Wang HY, Liang HG, Huang C, et al. New ingol-type diterpenes from the latex of *Euphorbia resinifera*. *J Asian Nat Prod Res.* 2019;21(11):1075-82.
  11. Spanos GA, Wrolstad RE. Influence of processing and storage on the phenolic composition of Thompson Seedless grape juice. *J Agric Food Chem.* 1990;38(7):1565-71.
  12. Dewanto V, Wu X, Liu RH. Processed sweet corn has higher antioxidant activity. *J Agric Food Chem.* 2002;50(17):4959-64.
  13. Huang B, Ke H, He J, Ban X, Zeng H, Wang Y. Extracts of *Halenia elliptica* exhibit antioxidant properties in vitro and in vivo. *Food Chem Toxicol.* 2011;49(1):185-90.
  14. Tuberoso CI, Boban M, Bifulco E, Budimir D, Pirisi FM. Antioxidant capacity and vasodilatory properties of Mediterranean food: the case of Cannonau wine, myrtle berries liqueur and strawberry-tree honey. *Food Chem.* 2013;140(4):686-91.
  15. Muruhan S, Selvaraj S, Viswanathan PK. In vitro antioxidant activities of *Solanum surattense* leaf extract. *Asian Pac J Trop Biomed.* 2013;3(1):28-34.
  16. Umamaheswari M, AsokKumar K, Somasundaram A, Sivashanmugam T, Subhadradevi V, Ravi TK. Xanthine oxidase inhibitory activity of some Indian medical plants. *J Ethnopharmacol.* 2007;109(3):547-51.
  17. Chakrabarti R, Singh B, Prakrith VN, Vanchhawng L, Thirumurugan K. Screening of nine herbal plants for in vitro  $\alpha$ -amylase inhibition. *Asian J Pharm Clin Res.* 2014;7(4):84-9.
  18. Kee KT, Koh M, Oong LX, Ng K. Screening culinary herbs for antioxidant and  $\alpha$ -glucosidase inhibitory activities. *Int J Food Sci Technol.* 2013;48(9):1884-91.
  19. Hu B, Cui F, Yin F, Zeng X, Sun Y, Li Y. Caffeoylquinic acids competitively inhibit pancreatic lipase through binding to the catalytic triad. *Int J Biol Macromol.* 2015;80:529-35.
  20. Beejmohun V, Peytavy-Izard M, Mignon C, Muscente-Paque D, Deplanque X, Ripoll C, et al. Acute effect of Ceylon cinnamon extract on postprandial glycemia: alpha-amylase inhibition, starch tolerance test in rats, and randomized crossover clinical trial in healthy volunteers. *BMC Complement Altern Med.* 2014;14:351.
  21. Botham PA. Acute systemic toxicity--prospects for tiered testing strategies. *Toxicol In Vitro.* 2004;18(2):227-30.
  22. Farah H, Ech-chahad A, Lamiri A. Antioxidant, antimicrobial and phytochemical investigations of polar extracts of *Euphorbia resinifera* Beg., roots, stems and flowers. *Am J Adv Drug Deliv.* 2014;2(6):776-85.
  23. Peter EL, Nagendrappa PB, Ajayi CO, Sesaaazi CD. Total polyphenols and antihyperglycemic activity of aqueous fruits extract of *Abelmoschus esculentus*: modeling and optimization of extraction conditions. *PLoS One.* 2021;16(4):e0250405.
  24. Dudonné S, Vitrac X, Coutière P, Woillez M, Mérillon JM. Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORAC assays. *J Agric Food Chem.* 2009;57(5):1768-74.
  25. Boulfia M, Lamchouri F, Senhaji S, Lachkar N, Bouabid K, Toufik H. Mineral content, chemical analysis, *In Vitro* antidiabetic and antioxidant activities, and antibacterial power of aqueous and organic extracts of Moroccan *Leopoldia comosa* (L.) Parl. Bulbs. *Evid Based Complement Alternat Med.* 2021;2021:9932291.
  26. Hanane F, Abdellah EC, Abdeslam L. In vitro antioxidant and antibacterial activity of the root extract of *Euphorbia resinifera*. *J Pharmacogn Phytochem.* 2014;2(5):161-3.
  27. Sayah K, Marmouzi I, Naceiri Mrabti H, Cherrah Y, Faouzi ME. Antioxidant activity and inhibitory potential of *Cistus salvifolius* (L.) and *Cistus monspeliensis* (L.) aerial parts extracts against key enzymes linked to hyperglycemia. *Biomed Res Int.* 2017;2017:2789482.
  28. Ameer K, Shahbaz HM, Kwon JH. Green extraction methods for polyphenols from plant matrices and their byproducts: a review. *Compr Rev Food Sci Food Saf.* 2017;16(2):295-315.
  29. Birari RB, Bhutani KK. Pancreatic lipase inhibitors from natural sources: unexplored potential. *Drug Discov Today.* 2007;12(19-20):879-89.
  30. Singh U, Singh S, Kochhar A. Therapeutic potential of antidia-

- betic nutraceuticals. *Phytopharmacology*. 2012;2(1):144-69.
31. Chater PI, Wilcox M, Cherry P, Herford A, Mustar S, Wheeler H, et al. Inhibitory activity of extracts of Hebridean brown seaweeds on lipase activity. *J Appl Phycol*. 2016;28:1303-13.
  32. Wang SY, Huang C, Sun RK, Lu LN, Liang HG, Gao L, et al. New tirucallane triterpenoids from the dried latex of *Euphorbia resinifera*. *Phytochem Lett*. 2019;29:220-4.
  33. Gutiérrez-Del-Río I, López-Ibáñez S, Magadán-Corpas P, Fernández-Calleja L, Pérez-Valero Á, Tuñón-Granda M, et al. Terpenoids and polyphenols as natural antioxidant agents in food preservation. *Antioxidants (Basel)*. 2021;10(8):1264.
  34. Chen YZ, Zhang BW, Yang J, Zou CS, Li T, Zhang GC, et al. Detoxification, antioxidant, and digestive enzyme activities and gene expression analysis of *Lymantria dispar* larvae under carvacrol. *J Asia Pac Entomol*. 2021;24(1):208-16.
  35. Yang W, Chen X, Li Y, Guo S, Wang Z, Yu X. Advances in pharmacological activities of terpenoids. *Nat Prod Commun*. 2020. doi: 10.1177/1934578X20903555. [Epub ahead of print]
  36. Čavar Zeljković S, Šišková J, Komzákova K, De Diego N, Kafková K, Tarkowski P. Phenolic compounds and biological activity of selected *Mentha* species. *Plants (Basel)*. 2021;10(3):550.
  37. Fettach S, Mrabti HN, Sayah K, Bouyahya A, Salhi N, Cherrah Y, et al. Phenolic content, acute toxicity of *Ajuga iva* extracts and assessment of their antioxidant and carbohydrate digestive enzyme inhibitory effects. *South Afr J Bot*. 2019;125:381-5.
  38. Naceiri Mrabti H, Marmouzi I, Sayah K, Chemlal L, El Ouadi Y, Elmsellem H, et al. *Arbutus unedo* L aqueous extract is associated with in vitro and in vivo antioxidant activity. *J Mater Environ Sci*. 2017;8(1):217-24.
  39. Simeonova R, Vitcheva V, Krasteva I, Zdraveva P, Konstantinov S, Ionkova I. Antidiabetic and antioxidant effects of saponarin from *Gypsophila trichotoma* on streptozotocin-induced diabetic normotensive and hypertensive rats. *Phytomedicine*. 2016; 23(5):483-90.
  40. Sengupta S, Mukherjee A, Goswami R, Basu S. Hypoglycemic activity of the antioxidant saponarin, characterized as alpha-glucosidase inhibitor present in *Tinospora cordifolia*. *J Enzyme Inhib Med Chem*. 2009;24(3):684-90.